

DIMENSION MAXIMIZING MEASURES FOR SELF-AFFINE SYSTEMS

BALÁZS BÁRÁNY AND MICHAŁ RAMS

ABSTRACT. In this paper we study the dimension theory of planar self-affine sets satisfying dominated splitting in the linear parts and strong separation condition. The main results of this paper is the existence of dimension maximizing Gibbs measures (Käenmäki measures). To prove this phenomena, we show that the Ledrappier-Young formula holds for Gibbs measures and we introduce a transversality type condition for the strong-stable directions on the projective space.

1. INTRODUCTION AND STATEMENTS

Let $\mathcal{A} := (A_1, A_2, \dots, A_N)$ be a finite set of contracting, non-singular 2×2 matrices, and let $\Phi := \{f_i : \underline{x} \mapsto A_i \underline{x} + \underline{t}_i\}_{i=1}^N$ be an *iterated function system* (IFS) on the plane with affine mappings, where $\|A_i\| < 1$ and $\underline{t}_i \in \mathbb{R}^2$ for $i = 1, \dots, N$. It is a well-known fact that there exists a unique non-empty compact subset Λ of \mathbb{R}^2 such that

$$\Lambda = \bigcup_{i=1}^N f_i(\Lambda).$$

We call the set Λ the *attractor* of Φ or *self-affine set*.

Let us denote the Hausdorff dimension of a set X by $\dim_H X$. Moreover, denote by $\underline{\dim}_B X$ and by $\overline{\dim}_B X$ the lower and upper box dimension. If the upper and lower box dimensions coincide then we call the common value the box dimension and denoted by $\dim_B X$. For the definitions and basic properties, we refer to Falconer [7].

Denote by $\alpha_i(A)$ the *i th singular value* of a 2×2 non-singular matrix A , i.e. the positive square root of the *i th eigenvalue* of AA^* , where A^* is the transpose of A . We note that in this case, $\alpha_1(A) = \|A\|$ and $\alpha_2(A) = \|A^{-1}\|^{-1}$, where $\|\cdot\|$ is the usual matrix norm induced by the Euclidean norm on \mathbb{R}^2 . Moreover, $\alpha_1(A)\alpha_2(A) = |\det A|$. For $s \geq 0$ define the *singular value function* ϕ^s as follows

$$\phi^s(A) := \begin{cases} \alpha_1(A)^s & 0 \leq s \leq 1 \\ \alpha_1(A)\alpha_2(A)^{s-1} & 1 < s \leq 2 \\ (\alpha_1(A)\alpha_2(A))^{s/2} & s > 2. \end{cases} \quad (1.1)$$

Falconer [6] introduced the *subadditive pressure*

$$P_{\mathcal{A}}(s) := \lim_{n \rightarrow \infty} \frac{1}{n} \log \sum_{i_1, \dots, i_n=1}^N \phi^s(A_{i_1} \cdots A_{i_n}). \quad (1.2)$$

The function $P_{\mathcal{A}} : [0, \infty) \mapsto \mathbb{R}$ is continuous, strictly monotone decreasing on $[0, \infty)$, moreover $P_{\mathcal{A}}(0) = \log N$ and $\lim_{s \rightarrow \infty} P_{\mathcal{A}}(s) = -\infty$. Falconer [6] showed that for the unique root $s_0 := s_0(\mathcal{A})$ of

Date: 24th August 2015.

2010 Mathematics Subject Classification. Primary 28A80 Secondary 37C45.

Key words and phrases. Self-affine measures, self-affine sets, Hausdorff dimension.

The research of Balázs Bárány was supported by the grants EP/J013560/1 and OTKA K104745. Michał Rams was supported by National Science Centre grant 2014/13/B/ST1/01033 (Poland).

the subadditive pressure function $\overline{\dim}_B \Lambda \leq \min \{2, s_0\}$ and if $\|A_i\| < 1/3$ for every $i = 1, \dots, N$ then

$$\dim_H \Lambda = \dim_B \Lambda = \min \{2, s_0\} \text{ for Lebesgue-almost every } \mathbf{t} = (t_1, \dots, t_N) \in \mathbb{R}^{2N}.$$

The condition was later weakened to $\|A_i\| < 1/2$ by Solomyak, see [15]. We call the value s_0 the *affinity dimension* of Φ . Käenmäki [11] showed that for Lebesgue-almost every $\mathbf{t} = (t_1, \dots, t_N) \in \mathbb{R}^{2N}$ there exists a measure ν^K (not necessarily self-affine) supported on Λ such that $\dim_H \mu^K = \dim_H \Lambda = \min \{2, s_0\}$.

In this paper we consider IFSs of affinities which satisfy the *strong separation condition* (SSC), i.e.

$$f_i(\Lambda) \cap f_j(\Lambda) = \emptyset \text{ for every } i \neq j.$$

Falconer [8] proved that if Φ satisfies a separation condition (milder than SSC) and the projection of Λ in every direction contains an interval separation then the box dimension of a self-affine set is equal to the affinity dimension. Hueter and Lalley [10] gave conditions, which ensure that the Hausdorff and box dimension of a self-affine set coincide and equal to the affinity dimension.

In the recent paper of Bárány [2], the result of Hueter and Lalley [10] was generalised for self-affine measures. That is, under the same conditions of Hueter and Lalley [10] the Hausdorff dimension of any self-affine measure is equal to its Lyapunov dimension. In particular, in [2] the author proved that under slightly more general conditions any self-affine measure is exact dimensional and gave a formula, which connects entropy, Lyapunov exponents and the projection of the measure (Ledrappier-Young formula).

Recently, Falconer and Kempton [9] used methods from ergodic theory along with properties of the Furstenberg measure and obtained conditions under which certain classes of plane self-affine sets have Hausdorff and box dimension equal to the affinity dimension. By adapting the conditions of Falconer and Kempton [9] and Bárány [2] we prove that for "typical" linear parts ($\{A_i\}_{i=1}^N$) if the SSC holds then the dimension of self-affine set is equal to the affinity dimension. Precisely, let

$$\mathfrak{M} := \left\{ A \in \mathbb{R}_+^{2 \times 2} \cup \mathbb{R}_-^{2 \times 2} : 0 < \frac{|\det A|}{\|A\|^2} < \frac{1}{2} \text{ and } \|A\| < 1 \right\}, \quad (1.3)$$

where

$$\|A\| = \min \{|a| + |b|, |c| + |d|\} \text{ for } A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}.$$

Let us define the following sets

$$\mathfrak{N} := \{A \in \mathfrak{M} : \|A^{-1}\| \|A\|^2 \leq 1\} \text{ and } \mathfrak{D}_N := \{\mathcal{A} \in \mathfrak{M}^N : s_0(\mathcal{A}) > 5/3\}, \quad (1.4)$$

for every $N \geq 2$.

Theorem 1.1. *Let $N \geq 2$. For \mathcal{L}_{4N} -almost every $\mathcal{A} \in \mathfrak{N}^N \cup \mathfrak{D}_N$, if $\mathbf{t} = (t_1, \dots, t_N) \in \mathbb{R}^{2N}$ is chosen such that $\Phi := \{f_i : \underline{x} \mapsto A_i \underline{x} + \underline{t}_i\}_{i=1}^N$ satisfies the SSC then there exists a measure ν^K supported on the attractor Λ of Φ such that*

$$\dim_H \nu^K = \dim_H \Lambda = \dim_B \Lambda = s_0(\mathcal{A}).$$

We call the measure ν^K the *Käenmäki measure*.

To prove Theorem 1.1, we will need a more detailed study of the dimension of invariant measures. More precisely, we extend the results of [2] for the natural projections of Gibbs measures. Theorem 1.1 is studied in higher generality.

Structure of the paper. After the Preliminaries (Section 2) we introduce the main technical result of the paper, the Ledrappier-Young formula generalised for Gibbs measures (Section 3). In Section 4 we introduce the strong-stable transversality condition (Definition 4.1) and show that under this condition there exists a dimension maximizing Gibbs measure (Käenmäki measure) almost surely. In the last section we show Theorem 1.1 as a consequence of the previous studies.

2. PRELIMINARIES

Let $\Sigma = \{1, \dots, N\}^{\mathbb{Z}}$ be the symbolic space of two side infinite sequences, $\Sigma^+ = \{1, \dots, N\}^{\mathbb{N}}$ be the set of right side and $\Sigma^- = \{1, \dots, N\}^{\mathbb{Z}^-}$ be the set of left side infinite words. Denote the left shift operator on Σ and Σ^+ by σ and denote the right shift operator on Σ and Σ^- by σ_- . Thus, σ and σ_- are invertible on Σ and $\sigma^{-1} = \sigma_-$. For any $\mathbf{i} \in \Sigma$ (or $\mathbf{j} \in \Sigma^\pm$)

$$[\mathbf{i}_m^n] := \{\mathbf{j} \in \Sigma \text{ (or } \mathbf{j} \in \Sigma^\pm) : i_k = j_k \text{ for } m \leq k \leq n\}.$$

For an $\mathbf{i} = (\dots i_{-2}i_{-1}i_0i_1\dots) \in \Sigma$, denote by $\mathbf{i}_+ = (i_0i_1\dots)$ the right-hand side and by $\mathbf{i}_- = (\dots i_{-2}i_{-1})$ the left-hand side of \mathbf{i} . To avoid confusion, we write also \mathbf{i}_+ if $\mathbf{i}_+ \in \Sigma^+$ and \mathbf{i}_- if $\mathbf{i}_- \in \Sigma^-$.

For any $\mathbf{i}_+, \mathbf{j}_+ \in \Sigma^+$ let $\mathbf{i}_+ \wedge \mathbf{j}_+ = \min\{n \geq 0 : i_n \neq j_n\}$. We define $\mathbf{i}_- \wedge \mathbf{j}_- = \min\{n-1 \geq 0 : i_{-n} \neq j_{-n}\}$ similarly.

Let us denote the set of finite length words by $\Sigma^* = \bigcup_{n=0}^{\infty} \{1, \dots, N\}^n$, and for every $\bar{i} = (i_1, \dots, i_n) \in \Sigma^*$ denote the reversed word by $\bar{i}^\rightarrow = (i_n, \dots, i_1)$. Sometimes, we may also write $(\Sigma^-)^*$ for finite length words to emphasize the negative indexes.

If $\Phi := \{f_i(\underline{x}) = A_i \underline{x} + \underline{t}_i\}_{i=1}^N$ is an iterated function system on \mathbb{R}^2 with affine mappings such that $\|A_i\| < 1$ for $i = 1, \dots, N$, we define the *natural projection* π_- from Σ^- to Λ in a natural way

$$\pi_-(\dots i_{-2}i_{-1}) = \lim_{n \rightarrow \infty} f_{i_{-1}} \circ \dots \circ f_{i_{-n}}(\underline{0}). \quad (2.1)$$

Let $\mathcal{A} := \{A_1, A_2, \dots, A_N\}$ be a finite set of non-singular 2×2 real matrices. Define a map from Σ to \mathcal{A} in a natural way, i.e. $A(\mathbf{i}) := A_{i_0}$. Let $A^{(n)}(\mathbf{i}) := A(\sigma^{n-1}\mathbf{i}) \dots A(\mathbf{i})$ for $\mathbf{i} \in \Sigma$ and $n \geq 1$.

Definition 2.1. We say that a set $\mathcal{A} = \{A_i\}_{i=1}^N$ of matrices satisfies the dominated splitting if there are constants $C, \beta > 0$ such that for every $n \geq 1$ and every $i_0, \dots, i_{n-1} \in \{1, \dots, N\}$

$$\frac{\alpha_1(A_{i_0} \dots A_{i_{n-1}})}{\alpha_2(A_{i_0} \dots A_{i_{n-1}})} \geq C e^{n\beta}.$$

Let $C_+ := \{(x, y) \in \mathbb{R}^2 \setminus \{(0, 0)\} : xy \geq 0\}$ be the *standard positive cone*. A *cone* is an image of C_+ under a linear isomorphism and a *multicone* is a disjoint union of finitely many cones. We say that a multicone M is *backward invariant* w.r.t. \mathcal{A} if $\bigcup_{A \in \mathcal{A}} A^{-1}(M) \subset M^\circ$, where M° denotes the interior of M .

Lemma 2.2 ([1], [3],[4], [17]). *The set \mathcal{A} of matrices satisfies the dominated splitting then for every $\mathbf{i} \in \Sigma$ there are two one-dimensional subspaces $e^{ss}(\mathbf{i}), e^s(\mathbf{i})$ of \mathbb{R}^2 such that*

- (1) $A_{i_0} e^j(\mathbf{i}) = e^j(\sigma \mathbf{i})$ for every $\mathbf{i} \in \Sigma$ and $j = s, ss$,
- (2) there is a constant $C > 0$ such that for every $n \geq 1$ and $\mathbf{i} \in \Sigma$

$$C^{-1} \|A^{(n)}(\mathbf{i})|e^s(\mathbf{i})\| \leq \alpha_1(A^{(n)}(\mathbf{i})) \leq C \|A^{(n)}(\mathbf{i})|e^s(\mathbf{i})\| \text{ and}$$

$$C^{-1} \|A^{(n)}(\mathbf{i})|e^{ss}(\mathbf{i})\| \leq \alpha_2(A^{(n)}(\mathbf{i})) \leq C \|A^{(n)}(\mathbf{i})|e^{ss}(\mathbf{i})\|,$$

- (3) there is a backward-invariant multicone M that

$$e^s(\mathbf{i}) = \bigcap_{n=1}^{\infty} A_{i_{-1}} \dots A_{i_{-n}}(\overline{M^c}) \text{ and } e^{ss}(\mathbf{i}) = \bigcap_{n=1}^{\infty} A_{i_0}^{-1} \dots A_{i_{n-1}}^{-1}(M),$$

where $\overline{M^c}$ denotes the closure of the complement of M .

We call the family of subspaces $e^s(\mathbf{i})$ stable directions and $e^{ss}(\mathbf{i})$ strong stable directions.

Let us observe that $e^s(\mathbf{i})$ depends only on \mathbf{i}_- and $e^{ss}(\mathbf{i}_+)$ depends only on \mathbf{i}_+ , so e^{ss} can be considered as a natural projection from Σ^+ to \mathbf{P}^1 , where \mathbf{P}^1 denotes the projective space. In the later analysis, the dimension of strong stable directions in \mathbf{P}^1 plays an important role. For $x, y \in \mathbf{P}^1$

denote by $\sphericalangle(x, y)$ the usual metric on \mathbf{P}^1 , that is the angle between the subspaces corresponding to x and y . For any $\underline{v}, \underline{w} \in \mathbb{R}^2$ denote by $\text{Area}(\underline{v}, \underline{w})$ the area of parallelogram formed by $\underline{v}, \underline{w}$.

Lemma 2.3. *For every $x, y \in \mathbf{P}^1$*

$$\frac{\text{Area}(\underline{v}, \underline{w})}{\|\underline{v}\|\|\underline{w}\|} \leq \sphericalangle(x, y) \leq \frac{2\text{Area}(\underline{v}, \underline{w})}{\|\underline{v}\|\|\underline{w}\|},$$

where $\underline{v}, \underline{w} \in \mathbb{R}^2$ are arbitrary non-zero vectors from the subspaces corresponding to x and y .

The proof of the lemma is straightforward.

Lemma 2.4. *There exists a constant $C > 0$ such that for every $\mathbf{i}, \mathbf{j} \in \Sigma$*

$$\sphericalangle(e^{ss}(\mathbf{i}), e^{ss}(\mathbf{j})) \leq Ce^{-\beta(\mathbf{i}_+ \wedge \mathbf{j}_+)} \quad \text{and} \quad \sphericalangle(e^s(\mathbf{i}), e^s(\mathbf{j})) \leq Ce^{-\beta(\mathbf{i}_- \wedge \mathbf{j}_-)}$$

where β is the domination exponent in Definition 2.1. Thus, the maps $\mathbf{i}_+ \in \Sigma^+ \mapsto e^{ss}(\mathbf{i}_+)$ and $\mathbf{i}_- \in \Sigma^- \mapsto \log \|A_{i_{-1}}|e^s(\sigma_{-}\mathbf{i}_-)\|$ are Hölder continuous.

Proof. We prove only the inequality for e^{ss} , for e^s the argument is similar. Fix $\mathbf{i}, \mathbf{j} \in \Sigma$ with $\mathbf{i}_+ \wedge \mathbf{j}_+ = n$. Let $\underline{v} \in e^{ss}(\sigma^n \mathbf{i}_+)$ and $\underline{w} \in e^{ss}(\sigma^n \mathbf{j}_+)$ be arbitrary such that $\|\underline{v}\| = \|\underline{w}\| = 1$. Then by Lemma 2.3,

$$\begin{aligned} \sphericalangle(e^{ss}(\mathbf{i}), e^{ss}(\mathbf{j})) &\leq 2 \frac{\text{Area}(A_{i_0}^{-1} \cdots A_{i_{n-1}}^{-1} \underline{v}, A_{i_0}^{-1} \cdots A_{i_{n-1}}^{-1} \underline{w})}{\|A_{i_0}^{-1} \cdots A_{i_{n-1}}^{-1} |e^{ss}(\sigma^n \mathbf{i}_+)\| \|A_{i_0}^{-1} \cdots A_{i_{n-1}}^{-1} |e^{ss}(\sigma^n \mathbf{j}_+)\|} \leq \\ &2C^2 \frac{|\det(A_{i_0}^{-1} \cdots A_{i_{n-1}}^{-1})|}{\|A_{i_0}^{-1} \cdots A_{i_{n-1}}^{-1}\|^2} \text{Area}(\underline{v}, \underline{w}) \leq 2C^2 e^{-\beta n}. \end{aligned}$$

□

Let $\varphi : \Sigma^- \mapsto \mathbb{R}$ be a Hölder continuous potential function. Then there exist a constants $C > 0, P \in \mathbb{R}$ and σ_- -invariant Borel probability measures μ_- and μ on Σ^- and Σ such that

$$C^{-1} \leq \frac{\mu_-([\mathbf{i}_-]_{-n}^{-1})}{e^{-nP + \sum_{k=0}^{n-1} \varphi(\sigma^k \mathbf{i}_-)}} \leq C, \quad \text{for every } \mathbf{i}_- \in \Sigma^-, \quad (2.2)$$

$$C^{-1} \leq \frac{\mu([\mathbf{i}]_{-n}^{-1})}{e^{-nP + \sum_{k=0}^{n-1} \varphi(\sigma^k \mathbf{i})}} \leq C, \quad \text{for every } \mathbf{i} \in \Sigma. \quad (2.3)$$

We call the measures μ_- and μ the *Gibbs measures* of the potential φ on Σ^- and Σ . Moreover, μ_- and μ are ergodic, see [5, Chapter 1]. Let $\nu = (\pi^-)_* \mu_-$, where π^- is defined in (2.1). Let us denote the projection from Σ to Σ^+ by $p_+ : \Sigma \mapsto \Sigma^+$, and similarly, the projection from Σ to Σ^- by $p_- : \Sigma \mapsto \Sigma^-$. It is easy to see that $(p_-)_* \mu = \mu_-$.

Lemma 2.5. *The measure $\mu_+ := (p_+)_* \mu$ is σ -invariant, ergodic quasi-Bernoulli measure on Σ^+ with entropy $h_{\mu_+} = h_\mu = h_{\mu_-} = P - \int \varphi(\mathbf{i}) d\mu(\mathbf{i})$.*

Proof. First, we prove invariance. Let $A \subseteq \Sigma^+$ be measurable set. Then by using that μ is σ -invariant we get

$$\mu_+(\sigma^{-1}A) = \mu_+ \left(\bigcup_{i=1}^N iA \right) = \mu \left(\Sigma^- \times \bigcup_{i=1}^N iA \right) = \mu(\Sigma^- \times A) = \mu_+(A).$$

Let $A \subseteq \Sigma^+$ be an arbitrary σ -invariant subset of Σ^+ . Then $\sigma^{-1}\Sigma^- \times A = \Sigma^- \times \left(\bigcup_{i=1}^N iA \right) = \Sigma^- \times \sigma^{-1}A = \Sigma^- \times A$. Therefore, $\mu(\Sigma^- \times A) = 0$ or 1 , which implies the ergodicity of μ_+ .

Finally, let $(i_0, \dots, i_{n+m+1}) \in (\Sigma^+)^*$ be arbitrary and let $\mathbf{j} \in \Sigma^-$ be such that $j_{-1} = i_{n+m+1}, \dots, j_{-(n+m+2)} = i_0$. Then by (2.3)

$$\begin{aligned} \mu_+([i_0, \dots, i_{n+m+1}]) &= \mu(\Sigma^- \times [i_0, \dots, i_{n+m+1}]) = \mu([\mathbf{j}]_{-(n+m+2)}^{-1}) \leq C e^{-(n+m+2)P + \sum_{k=0}^{n+m+1} \varphi(\sigma^k \mathbf{j})} = \\ &= C e^{-(n+1)P + \sum_{k=0}^n \varphi(\sigma^k \mathbf{j})} e^{-(m+1)P + \sum_{k=0}^m \varphi(\sigma^k(\sigma^{n+1} \mathbf{j}))} \leq C^3 \mu([\mathbf{j}]_{-(n+1)}^{-1}) \mu([\sigma_-^{n+1} \mathbf{j}]_{-(m+1)}^{-1}) = \\ &= C^3 \mu(\Sigma^- \times [i_0, \dots, i_n]) \mu(\Sigma^- \times [i_{n+1}, \dots, i_{n+m+1}]) = C^3 \mu_+([i_0, \dots, i_n]) \mu_+([i_{n+1}, \dots, i_{n+m+1}]). \end{aligned}$$

The inequality $\mu_+([i_0, \dots, i_{n+m+1}]) \geq C^{-3} \mu_+([i_0, \dots, i_n]) \mu_+([i_{n+1}, \dots, i_{n+m+1}])$ can be proven similarly. By using the definition of entropy, see [16, Theorem 4.10, Theorem 4.18],

$$\begin{aligned} h_{\mu_+} &= \lim_{n \rightarrow \infty} -\frac{1}{n} \sum_{\bar{i} \in \mathcal{S}^n} \mu_+([\bar{i}]) \log \mu_+([\bar{i}]) \leq P - \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{\bar{i} \in \mathcal{S}^n} \mu_+([\bar{i}]) \varphi(\bar{i} \mathbf{j}) = \\ &= P - \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{\bar{i} \in \mathcal{S}^n} \mu_-([\bar{i}]) \varphi(\bar{i} \mathbf{j}) = P - \int \varphi(\mathbf{i}) d\mu(\mathbf{i}). \end{aligned}$$

□

By Oseledec's multiplicative ergodic theorem, there are constants $0 < \chi_\mu^s \leq \chi_\mu^{ss}$ that

$$\begin{aligned} \lim_{n \rightarrow \infty} -\frac{1}{n} \log \alpha_1(A_{i_0} \cdots A_{i_{n-1}}) &= \chi_\mu^s \text{ and} \\ \lim_{n \rightarrow \infty} -\frac{1}{n} \log \alpha_2(A_{i_0} \cdots A_{i_{n-1}}) &= \chi_\mu^{ss} \text{ for } \mu\text{-a.e. } \mathbf{i} \in \Sigma \text{ (or } \mu_+\text{-a.e. } \mathbf{i}_+ \in \Sigma^+). \end{aligned}$$

We call the values χ_μ^s the *stable* and χ_μ^{ss} the *strong stable Lyapunov exponent* of μ . We define the Lyapunov exponents for μ_- similarly.

In our further analysis, a special Gibbs measure plays an important role.

Definition 2.6. Let $\mathcal{A} = \{A_1, A_2, \dots, A_N\}$ be a finite set of contracting, non-singular 2×2 matrices such that \mathcal{A} satisfies the dominated splitting. Moreover, let $s_0 = s_0(\mathcal{A})$ be the unique root of the subadditive pressure (1.2). We define $\varphi : \Sigma^- \mapsto \mathbb{R}$ be Hölder continuous potential function as follows,

$$\varphi(\mathbf{i}_-) = \begin{cases} \log \|A_{i_{-1}}|e^s(\sigma_- \mathbf{i}_-)\|^{s_0} & \text{if } 0 \leq s_0 \leq 1, \\ \log (|\det A_{i_{-1}}|^{s_0-1} \|A_{i_{-1}}|e^s(\sigma_- \mathbf{i}_-)\|^{2-s_0}) & \text{if } 1 < s_0 \leq 2, \\ \log |\det A_{i_{-1}}|^{s_0/2} & \text{if } 2 < s_0. \end{cases} \quad (2.4)$$

Then we call the Gibbs measure μ^K with potential φ the Käenmäki measure on Σ^- . In particular, there exists a constant $C > 0$ such that

$$C^{-1} \leq \frac{\mu^K([\mathbf{i}_-]_{-n}^{-1})}{\phi^{s_0}(A_{i_{-1}} \cdots A_{i_{-n}})} \leq C, \text{ for every } \mathbf{i}_- \in \Sigma^-,$$

where ϕ^s is the singular value function (1.1).

The Hölder continuity of potential φ in (2.4) follows by Lemma 2.4.

3. LEDRAPPIER-YOUNG FORMULA FOR GIBBS MEASURES

In this section, we extend the result [2, Theorem 2.7] for Gibbs-measures. For every $\theta \in \mathbf{P}^1$ we denote the orthogonal projection in the direction of θ by proj_θ . Let us define the transversal measure for every $\mathbf{i}_+ \in \Sigma^+$ by $\nu_{\mathbf{i}_+}^T = \nu \circ (\text{proj}_{e^{ss}(\mathbf{i}_+)})^{-1}$.

Theorem 3.1. *Let $\mathcal{A} = \{A_1, A_2, \dots, A_N\}$ be a finite set of contracting, non-singular 2×2 matrices, and let $\Phi = \{f_i(\underline{x}) = A_i \underline{x} + \underline{t}_i\}_{i=1}^N$ be an iterated function system on the plane with affine mappings. Let μ_- be a right-shift invariant and ergodic Gibbs measure on Σ^- defined in (2.2), and $\nu = (\pi^-)_* \mu_-$ be the push-down measure of μ_- . If*

- (1) \mathcal{A} satisfies the dominated splitting,
- (2) Φ satisfies the strong separation condition

then ν is exact dimensional and

$$\dim_H \nu = \frac{h_\mu}{\chi_\mu^s} + \left(1 - \frac{\chi_\mu^s}{\chi_\mu^{ss}}\right) \dim_H \nu_{\mathbf{i}_+}^T \text{ for } \mu_+ \text{-almost every } \mathbf{i}_+ \in \Sigma^+.$$

During the proof of Theorem 3.1, we follow the proof of [2, Theorem 2.7]. The proof of [2, Theorem 2.7] is decomposed into four propositions [2, Proposition 3.1, Proposition 3.3, Proposition 3.8 and Proposition 3.9]. However, [2, Proposition 3.1] and [2, Proposition 3.9] hold for general ergodic measures. On the other hand, [2, Proposition 3.8] follows from [2, Proposition 3.3] exactly in the same way for Gibbs measures as for Bernoulli measures. So, we extend in the rest of the section [2, Proposition 3.3] for Gibbs measures.

Let F be the dynamical system defined in [2, Section 3] acting on $\overline{\mathcal{O}} \times \Sigma^+$. Namely,

$$F(\underline{x}, \mathbf{i}) := (f_{i_0}(\underline{x}), \sigma \mathbf{i}),$$

where \mathcal{O} is the open and bounded set such that

$$\bigcup_{i=1}^N f_i(\mathcal{O}) \subseteq \mathcal{O} \text{ and } f_i(\overline{\mathcal{O}}) \cap f_j(\overline{\mathcal{O}}) = \emptyset \text{ for } i \neq j.$$

Since F is a hyperbolic map acting $\overline{\mathcal{O}} \times \Sigma^+$, its unique non-empty and compact F -invariant set is $\bigcap_{n=0}^{\infty} F^n(\overline{\mathcal{O}} \times \Sigma^+) = \Lambda \times \Sigma^+$. It is easy to see that F is conjugate to σ by the projection $\pi : \Sigma \mapsto \Lambda \times \Sigma^+$, where $\pi(\mathbf{i}) := (\pi^-(\mathbf{i}_-), \mathbf{i}_+)$. That is, $\pi \circ \sigma = F \circ \pi$. Denote the measure $\pi_* \mu$ by $\hat{\nu}$. Then $\hat{\nu}$ is F -invariant ergodic measure.

Since e^{ss} depends only on \mathbf{i}_+ , it defines a foliation on $\overline{\mathcal{O}}$ for every $\mathbf{i}_+ \in \Sigma^+$. Hence, it defines a foliation ξ^{ss} on $\Lambda \times \Sigma^+$. Namely, for a $\underline{\mathbf{y}} = (\underline{x}, \mathbf{i}_+) \in \Lambda \times \Sigma^+$ let $l_{ss}(\underline{\mathbf{y}})$ be the line through \underline{x} parallel to $e_{ss}(\mathbf{i}_+)$ on $\mathbb{R}^2 \times \{\mathbf{i}_+\}$. Let the partition element $\xi^{ss}(\underline{\mathbf{y}})$ be the intersection of the line $l_{ss}(\underline{\mathbf{y}})$ with $\Lambda \times \{\mathbf{i}_+\}$. It is easy to see that $F\xi^{ss}$ is a refinement of ξ^{ss} , that is, for every $\underline{\mathbf{y}}$, $(F\xi^{ss})(\underline{\mathbf{y}}) \subset \xi^{ss}(\underline{\mathbf{y}})$.

Applying Rokhlin's Theorem [13], there exists a canonical system of conditional measures, i.e. for $\hat{\nu}$ -a.e. $\underline{\mathbf{y}} \in \Lambda \times \Sigma^+$ there exists a measure $\hat{\nu}_{\underline{\mathbf{y}}}^{ss}$ supported on $\xi^{ss}(\underline{\mathbf{y}})$ such that the measures are uniquely defined up to a set of zero measure and for every measurable set A the function $\underline{\mathbf{y}} \mapsto \hat{\nu}_{\underline{\mathbf{y}}}^{ss}(A)$ is measurable. Moreover,

$$\hat{\nu}(A) = \int \hat{\nu}_{\underline{\mathbf{y}}}^{ss}(A) d\hat{\nu}(\underline{\mathbf{y}}). \quad (3.1)$$

Let us define the conditional entropy of $F\xi^{ss}$ with respect to ξ^{ss} in the usual way,

$$H(F\xi^{ss}|\xi^{ss}) := - \int \log \hat{\nu}_{\underline{\mathbf{y}}}^{ss}((F\xi^{ss})(\underline{\mathbf{y}})) d\hat{\nu}(\underline{\mathbf{y}}).$$

Applying again Rokhlin's Theorem [13], for μ_+ -a.e. $\mathbf{i}_+ \in \Sigma^+$ there exists a uniquely defined system of conditional measures $\mu_{\mathbf{i}_+}$ up to a set of zero measure, supported on $\Sigma^- \times \{\mathbf{i}_+\}$ and

$$\mu(A) = \int \mu_{\mathbf{i}_+}(A) d\mu_+(\mathbf{i}_+).$$

By defining $\hat{\nu}_{\mathbf{i}_+} := (\pi^-)_* \mu_{\mathbf{i}_+}$, we get

$$\hat{\nu} = \int \hat{\nu}_{\mathbf{i}_+} d\mu_+(\mathbf{i}_+).$$

For μ_+ -a.e. $\mathbf{i}_+ \in \Sigma^+$, let us define the transversal measure of $\widehat{\nu}_{\mathbf{i}_+}$ by

$$\widehat{\nu}_{\mathbf{i}_+}^T := (\text{proj}_{e_{ss}(\mathbf{i}_+)})_* \widehat{\nu}_{\mathbf{i}_+}.$$

By using the uniqueness of the conditional measures, we get

$$\widehat{\nu}_{\mathbf{i}_+} = \int \widehat{\nu}_{(\underline{x}, \mathbf{i}_+)}^{ss} d\widehat{\nu}_{\mathbf{i}_+}^T(\underline{x}) \text{ for } \mu_+\text{-a.e. } \mathbf{i}_+ \in \Sigma^+.$$

Lemma 3.2. *There exists a constant $C > 0$ such that $C^{-1}\mu_- \times \mu_+ \leq \mu \leq C\mu_- \times \mu_+$. In particular,*

$$C^{-1}\mu_- \leq \mu_{\mathbf{i}_+} \leq C\mu_- \text{ for } \mu_+\text{-a.e. } \mathbf{i}_+ \in \Sigma^+. \quad (3.2)$$

Proof. It is enough to show that there exists a $C > 0$ such that for every $\mathbf{i} \in \Sigma$ and $n, m \geq 0$

$$C^{-1}\mu_-([\mathbf{i}|_{-n}^{-1}])\mu_+([\mathbf{i}|_0^m]) \leq \mu([\mathbf{i}|_{-n}^m]) \leq C\mu_-([\mathbf{i}|_{-n}^{-1}])\mu_+([\mathbf{i}|_0^m]).$$

Indeed, every set A in the σ -algebra can be approximated by cylinder sets. By the definition of Gibbs measure μ

$$\begin{aligned} \mu([\mathbf{i}|_{-n}^m]) &= \mu([\sigma^{m+1}\mathbf{i}|_{-(n+m+1)}^{-1}]) \leq C e^{-(n+m+1)P + \sum_{k=0}^{n+m} \varphi(\sigma^k \sigma^{m+1}\mathbf{i})} = \\ &C e^{-nP + \sum_{k=0}^{n-1} \varphi \sigma^k \mathbf{i}} e^{-(m+1)P + \sum_{k=0}^m \varphi(\sigma^k \sigma^{m+1}\mathbf{i})} \leq C^2 \mu_-([\mathbf{i}|_{-n}^{-1}]) \mu([\sigma^{m+1}\mathbf{i}|_{-(m+1)}^{-1}]) = \\ &C^2 \mu_-([\mathbf{i}|_{-n}^{-1}]) \mu([\mathbf{i}|_0^m]) = C^2 \mu_-([\mathbf{i}|_{-n}^{-1}]) \mu_+([\mathbf{i}|_0^m]). \end{aligned}$$

The other inequality can be proven similarly. The relation (3.2) follows by the fact that the conditional measures are uniquely defined up to a set of zero measure. \square

By Lemma 3.2, the measures $\widehat{\nu}_{\mathbf{i}_+}$ and ν are equivalent for μ_+ -a.e. $\mathbf{i}_+ \in \Sigma^+$. Similarly, the measures $\widehat{\nu}_{\mathbf{i}_+}^T$ and $\nu_{\mathbf{i}_+}^T$ are equivalent for μ_+ -a.e. $\mathbf{i}_+ \in \Sigma^+$.

For the examination of the local dimension of the projected measure, instead of looking at balls on lines we introduce the transversal stable balls associated to the projection. Let $B_r^t(\underline{x}, \mathbf{i})$ be transversal stable ball with radius r , i.e

$$B_r^t(\underline{x}, \mathbf{i}) = \{(\underline{y}, \mathbf{j}) : \mathbf{i} = \mathbf{j} \ \& \ \text{dist}(l_{ss}(\underline{x}, \mathbf{i}), l_{ss}(\underline{y}, \mathbf{j})) \leq 2r\},$$

where $l_{ss}(\underline{x}, \mathbf{i})$ denotes the line trough \underline{x} parallel to $e_{ss}(\mathbf{i})$.

For technical reasons, we also have to introduce the modified transversal stable ball. Since the IFS Φ satisfies the SSC, for an $\underline{y} = (\underline{x}, \mathbf{i}) \in \Lambda \times \Sigma^+$ we can define the stable direction $e_s(\underline{y})$ of \underline{y} by $e_s(\underline{y}) := e_s(\underline{x}) := e_s(\mathbf{i}_-)$, where $\pi_-(\mathbf{i}_-) = \underline{x}$. Denote $\text{dist}_{e_s(\underline{y})}$ the natural Euclidean distance on the subspace $e_s(\underline{y})$.

Then for an $(\underline{x}, \mathbf{i}) \in \Lambda \times \Sigma^+$, we define the modified transversal stable ball with radius δ by

$$B_\delta^T(\underline{x}, \mathbf{i}) = \{(\underline{y}, \mathbf{j}) \in \Lambda \times \Sigma^+ : \mathbf{i} = \mathbf{j} \ \& \ \text{dist}_{e_s(\underline{x}, \mathbf{i})}(l_{ss}(\underline{x}, \mathbf{i}), l_{ss}(\underline{y}, \mathbf{j})) \leq \delta\},$$

where $\text{dist}_{e_s(\underline{x}, \mathbf{i})}(l_{ss}(\underline{x}, \mathbf{i}), l_{ss}(\underline{y}, \mathbf{j}))$ means the distance of the intersections of the lines $l_{ss}(\underline{x}, \mathbf{i}), l_{ss}(\underline{y}, \mathbf{j})$ with the subspace $e_s(\underline{x}, \mathbf{i})$ with respect to the distance $\text{dist}_{e_s(\underline{x}, \mathbf{i})}$. Since there exists a constant $\alpha > 0$ such that

$$\angle(e_s(\mathbf{i}_-), e_{ss}(\mathbf{i}_+)) \geq \alpha > 0, \text{ for every } \mathbf{i}_- \in \Sigma^- \text{ and } \mathbf{i}_+ \in \Sigma^+,$$

there exists a constant $c > 0$ that for every $\underline{y} \in \Lambda \times \Sigma^+$ and $r > 0$

$$B_{c^{-1}r}^T(\underline{x}, \mathbf{i}) \subseteq B_r^t(\underline{x}, \mathbf{i}) \subseteq B_{cr}^T(\underline{x}, \mathbf{i}). \quad (3.3)$$

We are going to prove the following proposition.

Proposition 3.3. For μ_+ -a.e. $\mathbf{i}_+ \in \Sigma^+$ the measure $\nu_{\mathbf{i}_+}^T$ is exact dimensional and

$$\dim_H \nu_{\mathbf{i}_+}^T = \frac{h_\mu - H(F\xi^{ss}|\xi^{ss})}{\chi_\mu^s}.$$

In particular,

$$\lim_{r \rightarrow 0^+} \frac{\nu(B_r^T(\underline{x}, \mathbf{i}_+))}{\log r} = \frac{h_\mu - H(F\xi^{ss}|\xi^{ss})}{\chi_\mu^s} \text{ for } \hat{\nu}\text{-a.e. } (\underline{x}, \mathbf{i}_+).$$

Let \mathcal{P} be the natural partition, i.e. $\mathcal{P} = \{f_i(\Lambda) \times \Sigma^+\}_{i=1}^N$. Denote the k th refinement of \mathcal{P} by \mathcal{P}_1^k , i.e. for every $\underline{\mathbf{y}} \in \Lambda \times \Sigma^+$, $\mathcal{P}_1^k(\underline{\mathbf{y}}) = \left(\bigvee_{i=1}^k F^i(\mathcal{P}) \right) (\underline{\mathbf{y}}) = \mathcal{P}(\underline{\mathbf{y}}) \cap F(\mathcal{P}(F^{-1}(\underline{\mathbf{y}}))) \cap \dots \cap F^k(\mathcal{P}(F^{-k}(\underline{\mathbf{y}})))$.

Let us define almost everywhere the measurable functions $g_k(\underline{\mathbf{y}}) := \hat{\nu}_{\underline{\mathbf{y}}}^{ss}(\mathcal{P}_1^k(\underline{\mathbf{y}}))$ and

$$g_{\delta,k}(\underline{\mathbf{y}}) := \frac{\hat{\nu}_{\mathbf{i}_+}(B_\delta^T(\underline{\mathbf{y}}) \cap \mathcal{P}_1^k(\underline{\mathbf{y}}))}{\hat{\nu}_{\mathbf{i}_+}(B_\delta^T(\underline{\mathbf{y}}))}.$$

By definition, $g_{\delta,k} \rightarrow g_k$ as $\delta \rightarrow 0^+$ for $\hat{\nu}$ almost everywhere and, since $g_{\delta,k}$ is uniformly bounded, (3.1) implies $g_{\delta,k} \rightarrow g_k$ in $L^1(\hat{\nu})$ as $\delta \rightarrow 0^+$.

Lemma 3.4. The function $\sup_{\delta > 0} \{-\log g_{\delta,k}\}$ is in $L^1(\hat{\nu})$ for every $k \geq 1$.

The proof of Lemma 3.4 coincides to [2, Lemma 3.6].

Lemma 3.5. For every $\underline{x} = \pi^-(i_{-1}, i_{-2}, \dots) \in \Lambda$, $\mathbf{i}_+ \in \Sigma^+$, $\delta > 0$ and $k \geq 1$

$$F^k \left(B_\delta^T(F^{-k}(\underline{\mathbf{y}})) \times [i_{-k}, \dots, i_{-1}] \right) = \left(B_{\|A_{i_{-1}} \dots A_{i_{-k}}|_{e_s(F^{-k}(\underline{\mathbf{y}}))\|}^T(\underline{\mathbf{y}})} \cap \mathcal{P}_1^k(\underline{\mathbf{y}}) \right) \times \Sigma^+,$$

where $\underline{\mathbf{y}} = (\underline{x}, \mathbf{i}_+)$.

By using the fact that $\nu = (\pi^-)_* \mu_- = (\pi^-)_*(p_-)_* \mu$, we have

$$\begin{aligned} \nu(B_\delta^T(\underline{\mathbf{y}}) \cap \mathcal{P}_1^k(\underline{\mathbf{y}})) &= \hat{\nu} \left(B_\delta^T(\underline{\mathbf{y}}) \cap \mathcal{P}_1^k \times \Sigma^+ \right) = \\ &= \hat{\nu} \left(F^{-k} \left(B_{\|A_{i_{-1}} \dots A_{i_{-k}}|_{e_s(F^{-k}(\underline{\mathbf{y}}))\|}^T(\underline{\mathbf{y}})} \cap \mathcal{P}_1^k \times \Sigma^+ \right) \right) = \hat{\nu} \left(B_{\|A_{i_{-1}} \dots A_{i_{-k}}|_{e_s(F^{-k}(\underline{\mathbf{y}}))\|}^T(F^{-k}(\underline{\mathbf{y}}))} \times [i_{-k}, \dots, i_{-1}] \right), \end{aligned}$$

where in the last equation we used Lemma 3.5. By Lemma 3.2,

$$\begin{aligned} \nu(B_\delta^T(\underline{\mathbf{y}}) \cap \mathcal{P}_1^k(\underline{\mathbf{y}})) &= \hat{\nu} \left(B_{\|A_{i_{-1}} \dots A_{i_{-k}}|_{e_s(F^{-k}(\underline{\mathbf{y}}))\|}^T(F^{-k}(\underline{\mathbf{y}}))} \times [i_{-k}, \dots, i_{-1}] \right) \leq \\ &= C \nu \left(B_{\|A_{i_{-1}} \dots A_{i_{-k}}|_{e_s(F^{-k}(\underline{\mathbf{y}}))\|}^T(F^{-k}(\underline{\mathbf{y}}))} \right) \mu_+([i_{-k}, \dots, i_{-1}]), \end{aligned} \quad (3.4)$$

and

$$\nu(B_\delta^T(\underline{\mathbf{y}}) \cap \mathcal{P}_1^k(\underline{\mathbf{y}})) \geq C^{-1} \nu \left(B_{\|A_{i_{-1}} \dots A_{i_{-k}}|_{e_s(F^{-k}(\underline{\mathbf{y}}))\|}^T(F^{-k}(\underline{\mathbf{y}}))} \right) \mu_+([i_{-k}, \dots, i_{-1}]) \quad (3.5)$$

for every $\delta > 0$, $k \geq 1$, and $\underline{\mathbf{y}} \in \Lambda \times \Sigma^+$.

Proof of Proposition 3.3. By the definition of the transversal measure, the statement of the proposition is equivalent to

$$\lim_{\delta \rightarrow 0^+} \frac{\log \nu(B_\delta^t(\underline{x}, \mathbf{i}_+))}{\log \delta} = \frac{h_\nu - H(F\xi^{ss}|\xi^{ss})}{\chi_\mu^s} \text{ for } \nu \times \mu_+\text{-a.e. } (\underline{x}, \mathbf{i}_+).$$

Hence, by (3.3) and by Lemma 2.2, it is enough to show that if $\underline{\mathbf{y}} = (\underline{x}, \mathbf{i}_+) \in \Lambda \times \Sigma^+$ with $\underline{x} = \pi_-(i_{-1}, i_{-2}, \dots)$,

$$\lim_{p \rightarrow \infty} \frac{\log \nu \left(B_{\|A_{i_{-1}} \cdots A_{i_{-pk}}\|e_s(F^{-pk}(\underline{\mathbf{y}}))}^T(\underline{\mathbf{y}}) \right)}{\log \alpha_1(A_{i_{-1}} \cdots A_{i_{-pk}})} = \frac{h_\nu - H(F\xi^{ss}|\xi^{ss})}{\chi_\mu^s} \text{ for } \nu \times \mu_+ \text{-a.e } \underline{\mathbf{y}}.$$

By Oseledec's Theorem, we have

$$\lim_{p \rightarrow \infty} \frac{1}{p} \log \alpha_1(A_{i_{-1}} \cdots A_{i_{-pk}}) = -k\chi_\mu^s \text{ for } \mu_- \text{-a.e } \mathbf{i}_-. \quad (3.6)$$

By applying (3.4), (3.5) and Lemma 3.2,

$$\begin{aligned} \nu \left(B_{\|A_{i_{-1}} \cdots A_{i_{-pk}}\|e_s(F^{-pk}(\underline{\mathbf{y}}))}^T(\underline{\mathbf{y}}) \right) &= \nu \left(B_1^T(F^{-pk}) \right) \prod_{l=1}^p \frac{\nu \left(B_{\|A_{i_{-(l-1)k-1}} \cdots A_{i_{-pk}}\|e_s(F^{-pk}(\underline{\mathbf{y}}))}^T(F^{-(l-1)k}(\underline{\mathbf{y}})) \right)}{\nu \left(B_{\|A_{i_{-(l-1)k-1}} \cdots A_{i_{-pk}}\|e_s(F^{-pk}(\underline{\mathbf{y}}))}^T(F^{-lk}(\underline{\mathbf{y}})) \right)} \leq \\ &C^p \nu \left(B_1^T(F^{-pk}) \right) \prod_{l=1}^p \frac{\nu \left(B_{\|A_{i_{-(l-1)k-1}} \cdots A_{i_{-pk}}\|e_s(F^{-pk}(\underline{\mathbf{y}}))}^T(F^{-(l-1)k}(\underline{\mathbf{y}})) \right) \mu_+([i_{-(l-1)k-1}, \dots, i_{-lk}])}{\nu \left(B_{\|A_{i_{-(l-1)k-1}} \cdots A_{i_{-pk}}\|e_s(F^{-pk}(\underline{\mathbf{y}}))}^T(F^{-(l-1)k}(\underline{\mathbf{y}})) \cap \mathcal{P}_1^k(F^{-(l-1)k}(\underline{\mathbf{y}})) \right)} \leq \\ &C^{3p} \nu \left(B_1^T(F^{-pk}) \right) \prod_{l=1}^p \frac{\hat{\nu}_{F^{-(l-1)k}(\underline{\mathbf{y}})} \left(B_{\|A_{i_{-(l-1)k-1}} \cdots A_{i_{-pk}}\|e_s(F^{-pk}(\underline{\mathbf{y}}))}^T(F^{-(l-1)k}(\underline{\mathbf{y}})) \right) \mu_+([i_{-(l-1)k-1}, \dots, i_{-lk}])}{\hat{\nu}_{F^{-(l-1)k}(\underline{\mathbf{y}})} \left(B_{\|A_{i_{-(l-1)k-1}} \cdots A_{i_{-pk}}\|e_s(F^{-pk}(\underline{\mathbf{y}}))}^T(F^{-(l-1)k}(\underline{\mathbf{y}})) \cap \mathcal{P}_1^k(F^{-(l-1)k}(\underline{\mathbf{y}})) \right)}. \end{aligned}$$

Similarly,

$$\begin{aligned} \nu \left(B_{\|A_{i_{-1}} \cdots A_{i_{-pk}}\|e_s(F^{-pk}(\underline{\mathbf{y}}))}^T(\underline{\mathbf{y}}) \right) &\geq C^{-3p} \nu \left(B_1^T(F^{-pk}) \right) \cdot \\ &\prod_{l=1}^p \frac{\hat{\nu}_{F^{-(l-1)k}(\underline{\mathbf{y}})} \left(B_{\|A_{i_{-(l-1)k-1}} \cdots A_{i_{-pk}}\|e_s(F^{-pk}(\underline{\mathbf{y}}))}^T(F^{-(l-1)k}(\underline{\mathbf{y}})) \right) \mu_+([i_{-(l-1)k-1}, \dots, i_{-lk}])}{\hat{\nu}_{F^{-(l-1)k}(\underline{\mathbf{y}})} \left(B_{\|A_{i_{-(l-1)k-1}} \cdots A_{i_{-pk}}\|e_s(F^{-pk}(\underline{\mathbf{y}}))}^T(F^{-(l-1)k}(\underline{\mathbf{y}})) \cap \mathcal{P}_1^k(F^{-(l-1)k}(\underline{\mathbf{y}})) \right)}. \end{aligned}$$

By taking logarithm and dividing by p we get

$$\begin{aligned} \frac{1}{p} \log \nu \left(B_1^T(F^{-pk}) \right) - 3 \log C - \frac{1}{p} \sum_{l=1}^p \log g_{\|A_{i_{-(l-1)k-1}} \cdots A_{i_{-pk}}\|e_s(F^{-pk}(\underline{\mathbf{y}}))\|,k}(F^{-lk}(\underline{\mathbf{y}})) + \\ \frac{1}{p} \sum_{l=1}^p \log \mu_+([i_{-(l-1)k-1}, \dots, i_{-lk}]) &\leq \frac{1}{p} \log \nu \left(B_{\|A_{i_{-1}} \cdots A_{i_{-pk}}\|e_s(F^{-pk}(\underline{\mathbf{y}}))}^T(\underline{\mathbf{y}}) \right) \leq \\ \frac{1}{p} \log \nu \left(B_1^T(F^{-pk}) \right) + 3 \log C - \frac{1}{p} \sum_{l=1}^p \log g_{\|A_{i_{-(l-1)k-1}} \cdots A_{i_{-pk}}\|e_s(F^{-pk}(\underline{\mathbf{y}}))\|,k}(F^{-lk}(\underline{\mathbf{y}})) + \\ &\frac{1}{p} \sum_{l=1}^p \log \mu_+([i_{-(l-1)k-1}, \dots, i_{-lk}]). \end{aligned}$$

By Lemma 3.4, we may apply the result of Maker [12, Theorem 1], so we get

$$\lim_{p \rightarrow \infty} -\frac{1}{p} \sum_{l=1}^p \log g_{\|A_{i-lk-1} \cdots A_{i-pk}\| e_s(F^{-pk}(\underline{y}))\|, k}(F^{-lk}(\underline{y})) = - \int \log g_k(\underline{y}) d\hat{\nu}(\underline{y}) = kH(F\xi^{ss}|\xi^{ss})$$

for $\hat{\nu}$ -a.e. \underline{y} . Applying Birkhoff's ergodic theorem and (3.6) we get

$$\frac{-3 \log C - kH(F\xi^{ss}|\xi^{ss}) - \sum_{\bar{i} \in \mathcal{S}^k} \mu_+([\bar{i}]) \log \mu_+([\bar{i}])}{k\chi_\mu^s} \leq \underline{d}_{\nu_{\mathbf{i}_+}^T}(\underline{x}) \leq \bar{d}_{\nu_{\mathbf{i}_+}^T}(\underline{x}) \leq \frac{3 \log C - kH(F\xi^{ss}|\xi^{ss}) - \sum_{\bar{i} \in \mathcal{S}^k} \mu_+([\bar{i}]) \log \mu_+([\bar{i}])}{k\chi_\mu^s} \text{ for } \hat{\nu}\text{-a.e. } \underline{y} \text{ and every } k \geq 1.$$

By taking the limit $k \rightarrow \infty$, we get that

$$\underline{d}_{\nu_{\mathbf{i}_+}^T}(\underline{x}) = \bar{d}_{\nu_{\mathbf{i}_+}^T}(\underline{x}) = \frac{h_\mu - H(F\xi^{ss}|\xi^{ss})}{\chi_\mu^s} \text{ for } \hat{\nu}\text{-a.e. } \underline{y}.$$

Since $\hat{\nu}$ is equivalent to $\nu \times \mu_+$, the statement follows. \square

Proof of Theorem 3.1. Since the proofs of [2, [Proposition 3.1, Proposition 3.8 and Proposition 3.9] do not use that the examined measure is Bernoulli, one can modify them to show that for $\hat{\nu}$ -a.e. $\underline{y} \in \Lambda \times \Sigma^+$ the measure $\hat{\nu}_{\underline{y}}^{ss}$ is exact dimensional and

$$\dim_H \hat{\nu}_{\underline{y}}^{ss} = \frac{H(F\xi^{ss}|\xi^{ss})}{\chi_\mu^{ss}}.$$

Moreover,

$$\liminf_{r \rightarrow \infty} \frac{\hat{\nu}_{\mathbf{i}_+}(B_r(\underline{x}))}{\log r} \geq \frac{H(F\xi^{ss}|\xi^{ss})}{\chi_\mu^{ss}} + \frac{h_\mu - H(F\xi^{ss}|\xi^{ss})}{\chi_\mu^s} \text{ for } \hat{\nu}\text{-a.e. } (\underline{x}, \mathbf{i}_+)$$

and by using that $\nu = (p_-)_* \hat{\nu}$

$$\limsup_{r \rightarrow \infty} \frac{\nu(B_r(\underline{x}))}{\log r} \leq \frac{H(F\xi^{ss}|\xi^{ss})}{\chi_\mu^{ss}} + \frac{h_\mu - H(F\xi^{ss}|\xi^{ss})}{\chi_\mu^s} \text{ for } \nu\text{-a.e. } \underline{x}.$$

Since the measure ν is equivalent to $\hat{\nu}_{\mathbf{i}_+}$ for μ_+ -a.e. \mathbf{i}_+ , the statement follows by Proposition 3.3. \square

As a corollary of Theorem 3.1, we are able to give two conditions which ensure that the dimension of a Gibbs measure is equal to its Lyapunov dimension. The second part of condition (iii) in the next theorem appears in [9], as well, for the Gibbs measure generated by the subadditive pressure.

Theorem 3.6. *Let $\mathcal{A} = \{A_k\}_{k=1}^N$ be a family of 2×2 real non-singular matrices and $\Phi = \{A_k \underline{x} + \underline{t}_k\}_{k=1}^N$ be an IFS of affinities on the plane. Moreover, let μ_- be a σ_- -invariant ergodic Gibbs measures on Σ^- , let μ be its unique extension to Σ and let μ_+ be the quasi-Bernoulli measure defined in Lemma 2.5. Assume that*

- (i) *the IFS Φ satisfies the strong separation condition,*
- (ii) *\mathcal{A} satisfies dominated splitting condition*
- (iii) *either $\dim_H(e^{ss})_* \mu_+ \geq \min\{1, \dim_{\text{Lyp}} \mu_-\}$ or $\dim_H(e^{ss})_* \mu_+ + \dim_H(\pi^-)_* \mu_- > 2$*

Then

$$\dim_H(\pi^-)_* \mu = \min \left\{ \frac{h_\mu}{\chi_\mu^s}, 1 + \frac{h_\mu - \chi_\mu^s}{\chi_\mu^{ss}} \right\}.$$

By Theorem 3.1, the proof is similar to the proofs of [2, Theorem 2.8 and Theorem 2.9].

4. DIMENSION OF GIBBS MEASURES AND TRANSVERSALITY CONDITION OF STRONG STABLE DIRECTIONS

In this section and the rest of the paper, we are going to study the dimension of Gibbs measures. To be able to calculate the dimension of Gibbs measure, we have to handle the dimension of strong stable directions, see (iii) of Theorem 3.6. In the case, when the matrices satisfies the backward non-overlapping condition, i.e. there exists a backward invariant multicone M such that $A_i^{-1}(M^o) \subseteq M^o$ and $A_i^{-1}(M^o) \cap A_j^{-1}(M^o) = \emptyset$ for every $i \neq j$, it is possible to calculate the dimension of strong stable directions. Namely, by [2, Lemma 4.2], for every σ -invariant ergodic measure μ on Σ^+

$$\dim_H(e^{ss})_*\mu = \frac{h_\mu}{\chi_\mu^{ss} - \chi_\mu^s},$$

where h_μ denotes the entropy of μ .

In general a set of matrices does not satisfy this phenomena. In this section we introduce a condition, which makes us able to handle the problem of overlaps. Namely, we consider a parametrized family of matrices $\mathcal{A}(\boldsymbol{\lambda})$ with the corresponding map of stable- and strong stable directions e_λ^s and e_λ^{ss} .

Definition 4.1. *Let $U \subset \mathbb{R}^d$ be open and bounded. We say that a parametrized family of matrices $\mathcal{A}(\boldsymbol{\lambda}) = \{A_i(\boldsymbol{\lambda})\}_{i=1}^N$ satisfies the strong-stable transversality on U if*

- *the parametrisation $\boldsymbol{\lambda} \mapsto A_i(\boldsymbol{\lambda})$ is continuous for every $i = 1, \dots, N$ on an open neighbourhood of \bar{U}*
- *for every $\boldsymbol{\lambda} \in \bar{U}$ the set $\mathcal{A}(\boldsymbol{\lambda})$ satisfies the dominated splitting*
- *there exists a constant $C > 0$ that for every $\mathbf{i}, \mathbf{j} \in \Sigma^+$ with $i_0 \neq j_0$*

$$\mathcal{L}_d \{ \boldsymbol{\lambda} \in U : \angle(e_\lambda^{ss}(\mathbf{i}), e_\lambda^{ss}(\mathbf{j})) < r \} \leq Cr \text{ for every } r > 0.$$

The definition of strong-stable transversality is a natural generalisation of the transversality condition for iterated function systems, see [14, (2.9)].

Theorem 4.2. *Let $U \subset \mathbb{R}^d$ be an open and bounded set and let $\mathcal{A}(\boldsymbol{\lambda}) = \{A_k(\boldsymbol{\lambda})\}_{k=1}^N$ be a parametrized family of 2×2 real matrices and $\Phi(\boldsymbol{\lambda}) = \{A_k(\boldsymbol{\lambda})\underline{x} + \underline{t}_k(\boldsymbol{\lambda})\}_{k=1}^N$ be a parametrized family affine IFSs on the real plane such that*

- (i) *for every $\boldsymbol{\lambda} \in U$ the IFS $\Phi(\boldsymbol{\lambda})$ satisfies the strong separation condition,*
- (ii) *$\mathcal{A}(\boldsymbol{\lambda})$ satisfies the strong-stable transversality on U .*

Let $\{\mu_\lambda\}_{\lambda \in U}$ be a parametrized family of σ_- -invariant ergodic Gibbs measures on Σ^- such that the family of the corresponding Hölder continuous potential functions $\{\phi_\lambda\}_{\lambda \in U}$ is continuously parametrized, moreover,

$$(iii) \text{ either } \frac{h_{\mu_\lambda}}{\chi_{\mu_\lambda}^{ss}(\boldsymbol{\lambda}) - \chi_{\mu_\lambda}^s(\boldsymbol{\lambda})} \geq \min \left\{ 1, \frac{h_{\mu_\lambda}}{\chi_{\mu_\lambda}^s(\boldsymbol{\lambda})} \right\} \text{ or } \frac{h_{\mu_\lambda}}{\chi_{\mu_\lambda}^{ss}(\boldsymbol{\lambda}) - \chi_{\mu_\lambda}^s(\boldsymbol{\lambda})} + 2 \frac{h_{\mu_\lambda}}{\chi_{\mu_\lambda}^{ss}(\boldsymbol{\lambda})} > 2$$

Then

$$\dim_H(\pi_\lambda^-)_*\mu_\lambda = \min \left\{ \frac{h_{\mu_\lambda}}{\chi_{\mu_\lambda}^s(\boldsymbol{\lambda})}, 1 + \frac{h_{\mu_\lambda} - \chi_{\mu_\lambda}^s(\boldsymbol{\lambda})}{\chi_{\mu_\lambda}^{ss}(\boldsymbol{\lambda})} \right\} \text{ for } \mathcal{L}_d\text{-a.e. } \boldsymbol{\lambda} \in U.$$

The proof of Theorem 4.2 is based on the combination of Theorem 3.6 and the following theorem.

Theorem 4.3. *Let $U \subset \mathbb{R}^d$ be an open and bounded set and let $\mathcal{A}(\boldsymbol{\lambda}) = \{A_k(\boldsymbol{\lambda})\}_{k=1}^N$ be a parametrized family of 2×2 real matrices such that $\mathcal{A}(\boldsymbol{\lambda})$ satisfies the strong-stable transversality on U . Moreover, let $\{\mu_\lambda\}_{\lambda \in U}$ be a family of σ -invariant quasi-Bernoulli ergodic measures on Σ^+ that $\boldsymbol{\lambda} \mapsto h_{\mu_\lambda}$ is*

continuous and for every $\lambda_0 \in U$ and $\varepsilon > 0$ there exists a $\delta = \delta(\varepsilon, \lambda_0) > 0$ that for every $\mathbf{i} \in \Sigma$, every $n \geq 1$ and every $\|\lambda - \lambda_0\| < \delta$

$$e^{-\varepsilon n} \leq \frac{\mu_\lambda([\mathbf{i}]_0^{n-1})}{\mu_{\lambda_0}([\mathbf{i}]_0^{n-1})} \leq e^{\varepsilon n}. \quad (4.1)$$

Then

$$\dim_H(e_\lambda^{ss})_* \mu_\lambda = \min \left\{ \frac{h_{\mu_\lambda}}{\chi_{\mu_\lambda}^{ss}(\lambda) - \chi_{\mu_\lambda}^s(\lambda)}, 1 \right\} \text{ for } \mathcal{L}_d\text{-a.e. } \lambda \in U.$$

The proof uses the standard transversality method but for completeness we present it here. First, we give an upper bound for the dimension.

Lemma 4.4. *Let $\mathcal{A} = \{A_i\}_{i=1}^N$ be a set of matrices satisfying the dominated splitting and let $e^{ss} : \Sigma^+ \mapsto \mathbb{P}^1$ be the map to strong-stable directions. Then for every μ σ -invariant ergodic measure on Σ^+ ,*

$$\dim_H(e^{ss})_* \mu \leq \min \left\{ 1, \frac{h_\mu}{\chi_\mu^{ss} - \chi_\mu^s} \right\}.$$

Proof of Lemma 4.4. For any $x \in \mathbf{P}^1$ let $B_r^\times(x) := \{y \in \mathbf{P}^1 : \times(x, y) < r\}$. It is enough to show that

$$\liminf_{r \rightarrow 0^+} \frac{\log(e^{ss})_* \mu(B_r^\times(e^{ss}(\mathbf{i})))}{\log r} \leq \frac{h_\mu}{\chi_\mu^{ss} - \chi_\mu^s} \text{ for } \mu\text{-a.e. } \mathbf{i} \in \Sigma^+.$$

By Lemma 2.3 and Lemma 2.2(2), if $\mathbf{i}, \mathbf{j} \in \Sigma^+$ that $i_k = j_k$ for $k = 0, \dots, n$

$$\times(e^{ss}(\mathbf{i}), e^{ss}(\mathbf{j})) \leq \frac{\text{Area}(A_{i_0}^{-1} \cdots A_{i_n}^{-1} \underline{v}, A_{i_0}^{-1} \cdots A_{i_n}^{-1} \underline{w})}{\|A_{i_0}^{-1} \cdots A_{i_n}^{-1}\| e^{ss}(\sigma^{n+1} \mathbf{j})} \leq C \frac{|\det(A_{i_0}^{-1} \cdots A_{i_n}^{-1})|}{\|A_{i_0}^{-1} \cdots A_{i_n}^{-1}\|^2},$$

where $\underline{v} \in e^{ss}(\sigma^{n+1} \mathbf{i})$ and $\underline{w} \in e^{ss}(\sigma^{n+1} \mathbf{j})$ such that $\|\underline{v}\| = \|\underline{w}\| = 1$. Let $n(r, \mathbf{i}) \in \mathbb{N}$ be the smallest number such that

$$\frac{|\det(A_{i_0}^{-1} \cdots A_{i_n}^{-1})|}{\|A_{i_0}^{-1} \cdots A_{i_n}^{-1}\|^2} < C^{-1} r.$$

Hence, $(e^{ss})_* \mu(B_r^\times(e^{ss}(\mathbf{i}))) \geq \mu([\mathbf{i}]_0^{n(r, \mathbf{i})})$. Therefore,

$$\frac{\log(e^{ss})_* \mu(B_r^\times(e^{ss}(\mathbf{i})))}{\log r} \leq \frac{\log \mu([\mathbf{i}]_0^{n(r, \mathbf{i})})}{\log C + \log |\det(A_{i_0}^{-1} \cdots A_{i_{n(r, \mathbf{i})-1}}^{-1})| - 2 \log \|A_{i_0}^{-1} \cdots A_{i_{n(r, \mathbf{i})-1}}^{-1}\|}$$

By ergodicity and Lemma 2.2(2),

$$\begin{aligned} \lim_{n \rightarrow \infty} -\frac{1}{n} \log \mu([\mathbf{i}]_0^n) &= h_\mu \\ \lim_{n \rightarrow \infty} -\frac{1}{n} \log |\det(A_{i_0}^{-1} \cdots A_{i_{n-1}}^{-1})| &= -\chi_\mu^{ss} - \chi_\mu^s \\ \lim_{n \rightarrow \infty} \frac{1}{n} \log \|A_{i_0}^{-1} \cdots A_{i_{n-1}}^{-1}\| &= \chi_\mu^{ss} \text{ for } \mu\text{-a.e. } \mathbf{i} \in \Sigma^+. \end{aligned}$$

□

Lemma 4.5. *Let $U \subset \mathbb{R}^d$ be open and bounded and let $\mathcal{A}(\lambda) = \{A_i(\lambda)\}_{i=1}^N$ be a parametrized family of matrices that the map $\lambda \mapsto A_i(\lambda)$ is continuous for any $i = 1, \dots, N$ in an open neighbourhood of \bar{U} , and $\mathcal{A}(\lambda)$ satisfies the dominated splitting on \bar{U} . Then the map $\lambda \mapsto e_\lambda^{ss}(\mathbf{i})$ is continuous uniformly for every $\mathbf{i} \in \Sigma^+$. That is, for every $\lambda_0 \in U$ and every $\varepsilon > 0$ there exists a $\delta = \delta(\lambda_0, \varepsilon) > 0$ that*

$$\|\lambda - \lambda_0\| < \delta \implies \times(e_\lambda^{ss}(\mathbf{i}), e_{\lambda_0}^{ss}(\mathbf{i})) < \varepsilon \text{ for every } \mathbf{i} \in \Sigma^+.$$

Proof. Let $\lambda_0 \in U$ and $\varepsilon > 0$ be arbitrary but fixed. Let M be the backward invariant multicone of $\mathcal{A}(\lambda_0)$. By definition of backward invariant multicone, there exists a $\delta' = \delta'(\lambda_0) > 0$ that for every λ with $\|\lambda - \lambda_0\| < \delta'$, M is a backward invariant multicone for $\mathcal{A}(\lambda)$. Hence, the angles between the directions of the dominated splitting are uniformly bounded from below. Thus, by Lemma 2.2(2) and Lemma 2.3, there exists a constant $C = C(\lambda_0) > 0$ that for every for every, $m \geq 0$ integer we have

$$\begin{aligned} \angle(e_{\lambda}^{ss}(\mathbf{i}), e_{\lambda_0}^{ss}(\mathbf{i})) &\leq \\ &\angle(A_{i_0}^{-1}(\lambda_0) \cdots A_{i_m}^{-1}(\lambda_0) e_{\lambda_0}^{ss}(\sigma^{m+1}\mathbf{i}), A_{i_0}^{-1}(\lambda_0) \cdots A_{i_m}^{-1}(\lambda_0) e_{\lambda}^{ss}(\sigma^{m+1}\mathbf{i})) + \\ &\angle(A_{i_0}^{-1}(\lambda_0) \cdots A_{i_m}^{-1}(\lambda_0) e_{\lambda}^{ss}(\sigma^{m+1}\mathbf{i}), A_{i_0}^{-1}(\lambda) \cdots A_{i_m}^{-1}(\lambda) e_{\lambda}^{ss}(\sigma^{m+1}\mathbf{i})) \leq \\ &C(\lambda_0)^2 2 \frac{|\det(A_{i_0}^{-1}(\lambda_0) \cdots A_{i_m}^{-1}(\lambda_0))|}{\|A_{i_0}^{-1}(\lambda_0) \cdots A_{i_m}^{-1}(\lambda_0)\|^2} \angle(e_{\lambda}^{ss}(\sigma^{m+1}\mathbf{i}), e_{\lambda_0}^{ss}(\sigma^{m+1}\mathbf{i})) + \\ &\frac{\sum_{i=1}^2 |A_{i_0}^{-1}(\lambda) \cdots A_{i_m}^{-1}(\lambda) \underline{u}_i \times A_{i_0}^{-1}(\lambda_0) \cdots A_{i_m}^{-1}(\lambda_0) \underline{u}_i| + |\sum_{i=1}^2 A_{i_0}^{-1}(\lambda) \cdots A_{i_m}^{-1}(\lambda) \underline{u}_i \times A_{i_0}^{-1}(\lambda_0) \cdots A_{i_m}^{-1}(\lambda_0) \underline{u}_{3-i}|}{\|A_{i_0}(\lambda) \cdots A_{i_m}(\lambda)\|^{-1} \|A_{i_0}(\lambda_0) \cdots A_{i_m}(\lambda_0)\|^{-1}}, \end{aligned}$$

where $\underline{u}_1, \underline{u}_2$ is the standard basis of \mathbb{R}^2 . Since $\mathcal{A}(\lambda)$ satisfies the dominated splitting on \bar{U} , there exists an integer $m = m(\lambda_0) > 0$ that

$$C(\lambda_0)^2 2 \frac{|\det(A_{i_0}^{-1}(\lambda_0) \cdots A_{i_m}^{-1}(\lambda_0))|}{\|A_{i_0}^{-1}(\lambda_0) \cdots A_{i_m}^{-1}(\lambda_0)\|^2} < \frac{1}{2},$$

for every $i_0, \dots, i_m \in \{1, \dots, N\}$. Let $f(\lambda, \lambda_0) := \sup_{\mathbf{i} \in \Sigma^+} \angle(e_{\lambda}^{ss}(\mathbf{i}), e_{\lambda_0}^{ss}(\mathbf{i}))$, then

$$f(\lambda, \lambda_0) \leq 2 \max_{i_0, \dots, i_m} \left\{ \frac{\sum_{i=1}^2 |A_{i_0}^{-1}(\lambda) \cdots A_{i_m}^{-1}(\lambda) \underline{u}_i \times A_{i_0}^{-1}(\lambda_0) \cdots A_{i_m}^{-1}(\lambda_0) \underline{u}_i|}{\|A_{i_0}(\lambda) \cdots A_{i_m}(\lambda)\|^{-1} \|A_{i_0}(\lambda_0) \cdots A_{i_m}(\lambda_0)\|^{-1}} + \frac{|\sum_{i=1}^2 A_{i_0}^{-1}(\lambda) \cdots A_{i_m}^{-1}(\lambda) \underline{u}_i \times A_{i_0}^{-1}(\lambda_0) \cdots A_{i_m}^{-1}(\lambda_0) \underline{u}_{3-i}|}{\|A_{i_0}(\lambda) \cdots A_{i_m}(\lambda)\|^{-1} \|A_{i_0}(\lambda_0) \cdots A_{i_m}(\lambda_0)\|^{-1}} \right\}.$$

Since the maps $\lambda \mapsto A_i(\lambda)$ are continuous, there exists a $\delta = \delta(\lambda_0, \varepsilon) > 0$ that the right hand side is less than $\varepsilon > 0$ for every λ with $\|\lambda - \lambda_0\| < \delta$. \square

Lemma 4.6. *Let $U \subset \mathbb{R}^d$ be open and bounded and let $\{\mu_{\lambda}\}_{\lambda \in U}$ be a family of σ -invariant quasi-Bernoulli ergodic measures on Σ^+ that (4.1) holds. Then the map $\lambda \mapsto \mu_{\lambda}$ is continuous in weak*-topology. Moreover, if $\mathcal{A}(\lambda) = \{A_i(\lambda)\}_{i=1}^N$ is a parametrized family of matrices that the map $\lambda \mapsto A_i(\lambda)$ is continuous for any $i = 1, \dots, N$ in an open neighbourhood of \bar{U} , and for every $\lambda \in \bar{U}$ the set $\mathcal{A}(\lambda)$ satisfies the dominated splitting then the maps $\lambda \mapsto \chi_{\mu_{\lambda}}^{ss}(\lambda)$ and $\lambda \mapsto \chi_{\mu_{\lambda}}^s(\lambda)$ are continuous.*

Proof. To prove the first assertion of the lemma it is enough to show that for every $O \subseteq \Sigma^+$ open set and every $\lambda_0 \in U$

$$\liminf_{\lambda \rightarrow \lambda_0} \mu_{\lambda}(O) \geq \mu_{\lambda_0}(O). \quad (4.2)$$

Since the cylinder sets form a base of open sets we get $O = \bigcup_{k=1}^{\infty} [\mathbf{i}_k|_{n_k}^{m_k}]$. Since for every cylinder $[\mathbf{i}_k|_{n_k}^{m_k}] = \bigcup_{|j|=n_k} [j\sigma^{n_k}\mathbf{i}_k|_0^{m_k}]$ without loss of generality we may write $O = \bigcup_{k=1}^{\infty} [\mathbf{i}_k|_0^{m_k}]$. On the other hand, for every pair of cylinder sets of the form $[\mathbf{i}_k|_0^{m_k}]$ either they are disjoint or one contains the other, thus, we may assume that $[\mathbf{i}_k|_0^{m_k}] \cap [\mathbf{i}_l|_0^{m_l}] = \emptyset$ if $k \neq l$. Hence,

$$\mu_{\lambda_0}(O) = \lim_{n \rightarrow \infty} \sum_{\substack{|\underline{i}|=n \\ [\underline{i}] \subseteq O}} \mu_{\lambda_0}([\underline{i}]).$$

Therefore, by (4.1) for every $n \geq 1$

$$\liminf_{\lambda \rightarrow \lambda_0} \mu_\lambda(O) \geq \liminf_{\lambda \rightarrow \lambda_0} \sum_{\substack{[i]=n \\ [i] \subseteq O}} \mu_\lambda([i]) = \sum_{\substack{[i]=n \\ [i] \subseteq O}} \mu_{\lambda_0}([i]).$$

Since $n \geq 1$ was arbitrary we get (4.2).

To prove the second assertion, by Lemma 2.2(2) and multiplicative ergodic theorem

$$\chi_{\mu_\lambda}^{ss}(\lambda) = \int \log \|A_{i_0}^{-1}(\lambda) |e_\lambda^{ss}(\sigma \mathbf{i})\| d\mu_\lambda(\mathbf{i}) \text{ and } \chi_{\mu_\lambda}^{ss}(\lambda) + \chi_{\mu_\lambda}^s(\lambda) = \int \log |\det(A_{i_0}^{-1}(\lambda))| d\mu_\lambda(\mathbf{i}).$$

By Lemma 4.5, the map $\lambda \mapsto \log \|A_{i_0}^{-1}(\lambda) |e_\lambda^{ss}(\sigma \mathbf{i})\|$ is continuous, thus by the weak*-continuity of $\lambda \mapsto \mu_\lambda$, the map $\lambda \mapsto \chi_{\mu_\lambda}^{ss}(\lambda)$ is continuous. The continuity of $\lambda \mapsto \chi_{\mu_\lambda}^s(\lambda)$ follows by the continuity of $\lambda \mapsto \mu_\lambda$, $\lambda \mapsto \chi_{\mu_\lambda}^{ss}(\lambda)$ and $\lambda \mapsto \log |\det(A_{i_0}^{-1}(\lambda))|$. \square

Proposition 4.7. *Assume that the assumptions of Theorem 4.3 hold. Then for every $\lambda_0 \in U$ and $\varepsilon > 0$ there exists a $\delta > 0$ such that*

$$\dim_H(e_\lambda^{ss})_* \mu_\lambda \geq \min \left\{ 1, \frac{h_{\mu_{\lambda_0}}}{\chi_{\mu_{\lambda_0}}^{ss}(\lambda_0) - \chi_{\mu_{\lambda_0}}^s(\lambda_0)} \right\} - \varepsilon \text{ for } \mathcal{L}_d\text{-a.e. } \lambda \in B_\delta(\lambda_0).$$

Before we prove Proposition 4.7, we prove that for every $\lambda \in U$ the map $\mathbf{i} \mapsto e_\lambda^{ss}(\mathbf{i})$ is Hölder continuous.

Lemma 4.8. *For every $\lambda_0 \in U$ there exists a $\delta = \delta(\lambda_0) > 0$ and for every $r > 0$ there exists a positive integer $N = N(\lambda_0, r)$ that for every $\lambda \in U$ with $\|\lambda - \lambda_0\| < \delta$ and for every $\mathbf{i}, \mathbf{j} \in \Sigma^+$ with $i_0 \neq j_0$*

$$\mathbb{I} \{ \angle(e_\lambda^{ss}(\mathbf{i}), e_\lambda^{ss}(\mathbf{j})) < r \} \leq \mathbb{I} \{ \angle(e_\lambda^{ss}(\mathbf{i}|_0^N \bar{\mathbf{1}}), e_\lambda^{ss}(\mathbf{j}|_0^N \bar{\mathbf{1}})) < 2r \},$$

where $\bar{\mathbf{1}} = (1, 1, \dots) \in \Sigma^+$. Precisely, $N(\lambda_0, r) = \lceil \frac{2 \log r}{-\beta(\lambda_0)} + c(\lambda_0) \rceil$, where $\beta(\lambda_0)$ is the domination exponent in Definition 2.1 and $c(\lambda_0)$ is some constant depending only on λ_0 .

Proof. Fix $\lambda_0 \in U$. Then by Lemma 2.3 for every N and every $\mathbf{i}, \mathbf{j} \in \Sigma^+$ with $i_0 \neq j_0$

$$\begin{aligned} |\angle(e_\lambda^{ss}(\mathbf{i}), e_\lambda^{ss}(\mathbf{j})) - \angle(e_\lambda^{ss}(\mathbf{i}|_0^N \bar{\mathbf{1}}), e_\lambda^{ss}(\mathbf{j}|_0^N \bar{\mathbf{1}}))| &\leq \angle(e_\lambda^{ss}(\mathbf{i}), e_\lambda^{ss}(\mathbf{i}|_0^N \bar{\mathbf{1}})) + \angle(e_\lambda^{ss}(\mathbf{j}), e_\lambda^{ss}(\mathbf{j}|_0^N \bar{\mathbf{1}})) \leq \\ &2 \frac{|\det(A_{i_0}^{-1}(\lambda) \cdots A_{i_N}^{-1}(\lambda))|}{\|A_{j_0}^{-1}(\lambda) \cdots A_{j_N}^{-1}(\lambda) |e_\lambda^{ss}(\sigma^{N+1} \mathbf{i})\| \|A_{j_0}^{-1}(\lambda) \cdots A_{j_N}^{-1}(\lambda) |e_\lambda^{ss}(\bar{\mathbf{1}})\|} \angle(e_\lambda^{ss}(\sigma^{N+1} \mathbf{i}), e_\lambda^{ss}(\bar{\mathbf{1}})) + \\ &2 \frac{|\det(A_{j_0}^{-1}(\lambda) \cdots A_{j_N}^{-1}(\lambda))|}{\|A_{j_0}^{-1}(\lambda) \cdots A_{j_N}^{-1}(\lambda) |e_\lambda^{ss}(\sigma^{N+1} \mathbf{j})\| \|A_{j_0}^{-1}(\lambda) \cdots A_{j_N}^{-1}(\lambda) |e_\lambda^{ss}(\bar{\mathbf{1}})\|} \angle(e_\lambda^{ss}(\sigma^{N+1} \mathbf{j}), e_\lambda^{ss}(\bar{\mathbf{1}})). \end{aligned}$$

Since $\lambda \mapsto A_i(\lambda)$ is continuous, by Lemma 4.5, there exists a $\delta = \delta(\lambda_0) > 0$ that

$$\begin{aligned} &\frac{|\det(A_{j_0}^{-1}(\lambda) \cdots A_{j_N}^{-1}(\lambda))|}{\|A_{j_0}^{-1}(\lambda) \cdots A_{j_N}^{-1}(\lambda) |e_\lambda^{ss}(\sigma^{N+1} \mathbf{j})\| \|A_{j_0}^{-1}(\lambda) \cdots A_{j_N}^{-1}(\lambda) |e_\lambda^{ss}(\bar{\mathbf{1}})\|} \leq \\ &e^{\frac{\delta(\lambda_0)}{2} N} \frac{|\det(A_{j_0}^{-1}(\lambda_0) \cdots A_{j_N}^{-1}(\lambda_0))|}{\|A_{j_0}^{-1}(\lambda_0) \cdots A_{j_N}^{-1}(\lambda_0) |e_{\lambda_0}^{ss}(\sigma^{N+1} \mathbf{j})\| \|A_{j_0}^{-1}(\lambda_0) \cdots A_{j_N}^{-1}(\lambda_0) |e_{\lambda_0}^{ss}(\bar{\mathbf{1}})\|} \end{aligned}$$

for every $\mathbf{j} \in \Sigma^+$. Thus, by Lemma 2.2(2)

$$|\angle(e_\lambda^{ss}(\mathbf{i}), e_\lambda^{ss}(\mathbf{j})) - \angle(e_\lambda^{ss}(\mathbf{i}|_0^N \bar{\mathbf{1}}), e_\lambda^{ss}(\mathbf{j}|_0^N \bar{\mathbf{1}}))| \leq 2\pi e^{\frac{\delta(\lambda_0)}{2} N} C(\lambda_0)^2 \max_{j_0, \dots, j_N} \left\{ \frac{|\det(A_{j_0}^{-1}(\lambda_0) \cdots A_{j_N}^{-1}(\lambda_0))|}{\|A_{j_0}^{-1}(\lambda_0) \cdots A_{j_N}^{-1}(\lambda_0)\|^2} \right\}$$

By Definition 2.1, there exists an $N = N(\boldsymbol{\lambda}_0, r)$ that the right hand side of the inequality is less than r , thus the statement follows. \square

Proof of Proposition 4.7. Let $\boldsymbol{\lambda}_0 \in U$ and $\varepsilon > 0$ be arbitrary but fixed. Let $\delta = \delta(\boldsymbol{\lambda}_0, \varepsilon) > 0$ be chosen according to Lemma 4.5, Lemma 4.8 and (4.1). By Shannon-McMillan-Breiman Theorem and (4.1), for every $\boldsymbol{\lambda} \in B_\delta(\boldsymbol{\lambda}_0)$

$$h_{\mu_{\boldsymbol{\lambda}_0}} - \varepsilon \leq \liminf_{n \rightarrow \infty} -\frac{1}{n} \log \mu_{\boldsymbol{\lambda}}([\mathbf{i}|_0^{n-1}]) \leq \limsup_{n \rightarrow \infty} -\frac{1}{n} \log \mu_{\boldsymbol{\lambda}}([\mathbf{i}|_0^{n-1}]) \leq h_{\mu_{\boldsymbol{\lambda}_0}} + \varepsilon \text{ for } \mu_{\boldsymbol{\lambda}}\text{-a.e. } \mathbf{i} \in \Sigma^+.$$

Moreover, by ergodic theorem and weak*-continuity of $\boldsymbol{\lambda} \mapsto \mu_{\boldsymbol{\lambda}}$

$$\chi_{\mu_{\boldsymbol{\lambda}_0}}^{ss}(\boldsymbol{\lambda}_0) + \chi_{\mu_{\boldsymbol{\lambda}_0}}^s(\boldsymbol{\lambda}_0) - \varepsilon \leq \lim_{n \rightarrow \infty} \frac{1}{n} \log |\det(A_{i_0}^{-1}(\boldsymbol{\lambda}) \cdots A_{i_{n-1}}^{-1}(\boldsymbol{\lambda}))| = \chi_{\mu_{\boldsymbol{\lambda}_0}}^{ss}(\boldsymbol{\lambda}_0) + \chi_{\mu_{\boldsymbol{\lambda}_0}}^s(\boldsymbol{\lambda}_0) + \varepsilon,$$

$$\chi_{\mu_{\boldsymbol{\lambda}_0}}^{ss}(\boldsymbol{\lambda}_0) - \varepsilon \leq \lim_{n \rightarrow \infty} \frac{1}{n} \log \|A_{i_0}^{-1}(\boldsymbol{\lambda}) \cdots A_{i_{n-1}}^{-1}(\boldsymbol{\lambda}) e_{\boldsymbol{\lambda}}^{ss}(\sigma^n \mathbf{i})\| \leq \chi_{\mu_{\boldsymbol{\lambda}_0}}^{ss}(\boldsymbol{\lambda}_0) + \varepsilon$$

for $\mu_{\boldsymbol{\lambda}}$ -a.e. $\mathbf{i} \in \Sigma^+$. By Egorov's theorem for every $\boldsymbol{\lambda} \in B_\delta(\boldsymbol{\lambda}_0)$ there exists a set $\Omega_{\boldsymbol{\lambda}} \subseteq \Sigma^+$ that $\mu(\Omega_{\boldsymbol{\lambda}}) > 1 - \varepsilon$ and there exist a constant $C(\boldsymbol{\lambda}) > 1$ that for every $\mathbf{i} \in \Sigma^+$ and every $n, m \geq 1$

$$C(\boldsymbol{\lambda})^{-1} \mu_{\boldsymbol{\lambda}}([\mathbf{i}|_0^{n-1}]) \mu_{\boldsymbol{\lambda}}([\sigma^n \mathbf{i}|_0^{m-1}]) \leq \mu_{\boldsymbol{\lambda}}([\mathbf{i}|_0^{n+m-1}]) \leq C(\boldsymbol{\lambda}) \mu_{\boldsymbol{\lambda}}([\mathbf{i}|_0^{n-1}]) \mu_{\boldsymbol{\lambda}}([\sigma^n \mathbf{i}|_0^{m-1}])$$

and for every $\mathbf{i} \in \Omega_{\boldsymbol{\lambda}}$ and every $n \geq 1$

$$C(\boldsymbol{\lambda})^{-1} e^{-n(h_{\mu_{\boldsymbol{\lambda}_0}} + 2\varepsilon)} \leq \mu_{\boldsymbol{\lambda}}([\mathbf{i}|_0^{n-1}]) \leq C(\boldsymbol{\lambda}) e^{-n(h_{\mu_{\boldsymbol{\lambda}_0}} - 2\varepsilon)}, \quad (4.3)$$

$$C(\boldsymbol{\lambda})^{-1} e^{-n(\chi_{\mu_{\boldsymbol{\lambda}_0}}^{ss}(\boldsymbol{\lambda}_0) - \chi_{\mu_{\boldsymbol{\lambda}_0}}^s(\boldsymbol{\lambda}_0) + 6\varepsilon)} \leq \frac{|\det(A_{i_0}^{-1}(\boldsymbol{\lambda}) \cdots A_{i_{n-1}}^{-1}(\boldsymbol{\lambda}))|}{\|A_{i_0}^{-1}(\boldsymbol{\lambda}) \cdots A_{i_{n-1}}^{-1}(\boldsymbol{\lambda})\|^2} \leq C(\boldsymbol{\lambda}) e^{-n(\chi_{\mu_{\boldsymbol{\lambda}_0}}^{ss}(\boldsymbol{\lambda}_0) - \chi_{\mu_{\boldsymbol{\lambda}_0}}^s(\boldsymbol{\lambda}_0) - 6\varepsilon)}. \quad (4.4)$$

By Lusin's theorem for every $\varepsilon' > 0$ there exists a set $J_\delta(\boldsymbol{\lambda}_0) \subseteq B_\delta(\boldsymbol{\lambda}_0)$ that $\mathcal{L}_d(B_\delta(\boldsymbol{\lambda}_0)/J_\delta(\boldsymbol{\lambda}_0)) < \varepsilon'$ and there exists a $C > 1$ that $C(\boldsymbol{\lambda}) \leq C$ for every $\boldsymbol{\lambda} \in J_\delta(\boldsymbol{\lambda}_0)$. Denote the measure $\tilde{\mu}_{\boldsymbol{\lambda}} := \mu|_{\Omega_{\boldsymbol{\lambda}}}$ and for a finite length word $\underline{k} = (k_0, \dots, k_{n-1})$ denote the set

$$\Sigma_{\underline{k}} := \{(\mathbf{i}, \mathbf{j}) \in \Sigma^+ : i_m = j_m = k_m \text{ for } m = 0, \dots, n-1 \text{ and } i_n \neq j_n\}.$$

Then for every $s > 0$ by Lemma 4.5, the continuity of $\boldsymbol{\lambda} \mapsto A_i(\boldsymbol{\lambda})$ and (4.4)

$$\begin{aligned} \mathcal{I} &:= \int_{J_\delta(\boldsymbol{\lambda}_0)} \iint \times(e_{\boldsymbol{\lambda}}^{ss}(\mathbf{i}), e_{\boldsymbol{\lambda}}^{ss}(\mathbf{j}))^{-s} d\tilde{\mu}_{\boldsymbol{\lambda}}(\mathbf{i}) d\tilde{\mu}_{\boldsymbol{\lambda}}(\mathbf{j}) d\boldsymbol{\lambda} = \\ &\quad \sum_{n=0}^{\infty} \sum_{\underline{k}=n} \int_{J_\delta(\boldsymbol{\lambda}_0)} \iint_{\Sigma_{\underline{k}}} \times(e_{\boldsymbol{\lambda}}^{ss}(\mathbf{i}), e_{\boldsymbol{\lambda}}^{ss}(\mathbf{j}))^{-s} d\tilde{\mu}_{\boldsymbol{\lambda}}(\mathbf{i}) d\tilde{\mu}_{\boldsymbol{\lambda}}(\mathbf{j}) d\boldsymbol{\lambda} \leq \\ &\quad \sum_{n=0}^{\infty} \sum_{\underline{k}=n} \int_{J_\delta(\boldsymbol{\lambda}_0)} \iint_{\Sigma_{\underline{k}}} C(\boldsymbol{\lambda})^2 \left(\frac{|\det(A_{k_0}^{-1}(\boldsymbol{\lambda}) \cdots A_{k_{n-1}}^{-1}(\boldsymbol{\lambda}))|}{2 \|A_{k_0}^{-1}(\boldsymbol{\lambda}) \cdots A_{k_{n-1}}^{-1}(\boldsymbol{\lambda})\|^2} \right)^{-s} \\ &\quad \times(e_{\boldsymbol{\lambda}}^{ss}(\sigma^n \mathbf{i}), e_{\boldsymbol{\lambda}}^{ss}(\sigma^n \mathbf{j}))^{-s} d\tilde{\mu}_{\boldsymbol{\lambda}}(\mathbf{i}) d\tilde{\mu}_{\boldsymbol{\lambda}}(\mathbf{j}) d\boldsymbol{\lambda} \leq \\ &\quad \sum_{n=0}^{\infty} C' e^{sn(\chi_{\mu_{\boldsymbol{\lambda}_0}}^{ss}(\boldsymbol{\lambda}_0) - \chi_{\mu_{\boldsymbol{\lambda}_0}}^s(\boldsymbol{\lambda}_0) + 6\varepsilon)} \sum_{\underline{k}=n} \int_{J_\delta(\boldsymbol{\lambda}_0)} \iint_{\Sigma_{\underline{k}}} \times(e_{\boldsymbol{\lambda}}^{ss}(\sigma^n \mathbf{i}), e_{\boldsymbol{\lambda}}^{ss}(\sigma^n \mathbf{j}))^{-s} d\tilde{\mu}_{\boldsymbol{\lambda}}(\mathbf{i}) d\tilde{\mu}_{\boldsymbol{\lambda}}(\mathbf{j}) d\boldsymbol{\lambda}. \end{aligned}$$

By Lemma 4.8, for any \underline{k} with $|\underline{k}| = n$

$$\begin{aligned}
\mathcal{I}_{\underline{k}} &:= \int_{J_\delta(\lambda_0)} \iint_{\Sigma_{\underline{k}}} \mathfrak{x}(e_\lambda^{ss}(\sigma^n \mathbf{i}), e_\lambda^{ss}(\sigma^n \mathbf{j}))^{-s} d\tilde{\mu}_\lambda(\mathbf{i}) d\tilde{\mu}_\lambda(\mathbf{j}) d\lambda \leq \\
&\quad \sum_{m=0}^{\infty} 2^{(m+1)s} \int_{J_\delta(\lambda_0)} \iint_{\Sigma_{\underline{k}}} \mathbb{I} \left\{ \mathfrak{x}(e_\lambda^{ss}(\sigma^n \mathbf{i}), e_\lambda^{ss}(\sigma^n \mathbf{j})) < \frac{1}{2^m} \right\} d\tilde{\mu}_\lambda(\mathbf{i}) d\tilde{\mu}_\lambda(\mathbf{j}) d\lambda \leq \\
&\quad \sum_{m=0}^{\infty} 2^{(m+1)s} \int_{J_\delta(\lambda_0)} \iint_{\Sigma_{\underline{k}}} \mathbb{I} \left\{ \mathfrak{x}(e_\lambda^{ss}(\sigma^n \mathbf{i}|_0^{N(\lambda_0, m)} \bar{1}), e_\lambda^{ss}(\sigma^n \mathbf{j}|_0^{N(\lambda_0, m)} \bar{1})) < \frac{2}{2^m} \right\} d\tilde{\mu}_\lambda(\mathbf{i}) d\tilde{\mu}_\lambda(\mathbf{j}) d\lambda = \\
&\quad \sum_{m=0}^{\infty} 2^{(m+1)s} \sum_{\substack{|\underline{l}|=N(\lambda_0, m) \\ |\underline{h}|=N(\lambda_0, m)}} \int_{J_\delta(\lambda_0)} \iint_{[\underline{k}\underline{l}] \times [\underline{k}\underline{h}]} \mathbb{I} \left\{ \mathfrak{x}(e_\lambda^{ss}(\underline{h}\bar{1}), e_\lambda^{ss}(\underline{l}\bar{1})) < \frac{2}{2^m} \right\} d\tilde{\mu}_\lambda(\mathbf{i}) d\tilde{\mu}_\lambda(\mathbf{j}) d\lambda
\end{aligned} \tag{4.5}$$

By applying (4.3), the quasi-Bernoulli property of μ_{λ_0} , (4.1) and the continuity of $\lambda \mapsto h_{\mu_\lambda}$

$$\begin{aligned}
&\int_{J_\delta(\lambda_0)} \mathbb{I} \left\{ \mathfrak{x}(e_\lambda^{ss}(\underline{h}\bar{1}), e_\lambda^{ss}(\underline{l}\bar{1})) < \frac{2}{2^m} \right\} \tilde{\mu}_\lambda([\underline{k}\underline{l}]) \tilde{\mu}_\lambda([\underline{k}\underline{h}]) d\lambda \leq \\
&\quad C^2 \int_{J_\delta(\lambda_0)} \mathbb{I} \left\{ \mathfrak{x}(e_\lambda^{ss}(\underline{h}\bar{1}), e_\lambda^{ss}(\underline{l}\bar{1})) < \frac{2}{2^m} \right\} \tilde{\mu}_\lambda([\underline{k}])^2 \tilde{\mu}_\lambda([\underline{l}]) \tilde{\mu}_\lambda([\underline{h}]) d\lambda \leq \\
&\quad c' \mu_{\lambda_0}([\underline{k}]) \mu_{\lambda_0}([\underline{l}]) \mu_{\lambda_0}([\underline{h}]) e^{2\varepsilon(n+N(\lambda_0, m))} e^{-n(h_{\mu_{\lambda_0}} - 2\varepsilon)} \mathcal{L}_d \left(\lambda \in J_\delta(\lambda_0) : \mathfrak{x}(e_\lambda^{ss}(\underline{h}\bar{1}), e_\lambda^{ss}(\underline{l}\bar{1})) < \frac{2}{2^m} \right).
\end{aligned}$$

Hence, by (4.5) and the strong-stable transversality

$$\begin{aligned}
\mathcal{I}_{\underline{k}} &\leq c' \mu_{\lambda_0}([\underline{k}]) \sum_{m=0}^{\infty} 2^{(m+1)s} \sum_{\substack{|\underline{l}|=N(\lambda_0, m) \\ |\underline{h}|=N(\lambda_0, m)}} \mu_{\lambda_0}([\underline{l}]) \mu_{\lambda_0}([\underline{h}]) e^{2\varepsilon(n+N(\lambda_0, m))} e^{-n(h_{\mu_{\lambda_0}} - 2\varepsilon)} \frac{C}{2^m} = \\
&\quad c'' \mu_{\lambda_0}([\underline{k}]) e^{-n(h_{\mu_{\lambda_0}} - 4\varepsilon)} \sum_{m=0}^{\infty} 2^{m(s-1)+2\varepsilon N(\lambda_0, m)/\log 2}
\end{aligned}$$

Since $N(\lambda_0, m)/\log 2 \leq m \frac{2}{\beta(\lambda_0)} + c(\lambda_0)$

$$\mathcal{I} \leq c''' \sum_{n=0}^{\infty} e^{n(s(\chi_\mu^{ss}(\lambda_0) - \chi_\mu^s(\lambda_0)) - h_{\mu_{\lambda_0}} + 10\varepsilon)} \sum_{m=0}^{\infty} 2^{m(s-1+\varepsilon \frac{4}{\beta(\lambda_0)})}.$$

Hence, by choosing $s < \min \left\{ 1 - \varepsilon \frac{5}{\beta(\lambda_0)}, \frac{h_{\mu_{\lambda_0}} - 11\varepsilon}{\chi_\mu^{ss}(\lambda_0) - \chi_\mu^s(\lambda_0)} \right\}$ the right hand side of the inequality is finite.

By Frostman's Lemma [7, Theorem 4.13],

$$\dim_H(e_\lambda^{ss})_* \tilde{\mu}_\lambda \geq \min \left\{ 1 - \varepsilon \frac{5}{\beta(\lambda_0)}, \frac{h_{\mu_{\lambda_0}} - 11\varepsilon}{\chi_\mu^{ss}(\lambda_0) - \chi_\mu^s(\lambda_0)} \right\} \text{ for } \mathcal{L}_d\text{-a.e. } \lambda \in J_\delta(\lambda_0).$$

But for every $\lambda \in B_\delta(\lambda_0)$, $\dim_H(e_\lambda^{ss})_* \mu_\lambda \geq \dim_H(e_\lambda^{ss})_* \tilde{\mu}_\lambda$, moreover, $\mathcal{L}_d(B_\delta(\lambda_0)/J_\delta(\lambda_0))$ can be chosen arbitrary small, thus, the statement follows. \square

Proof of Theorem 4.3. By Lemma 4.4 we have

$$\dim_H(e_{\lambda}^{ss})_*\mu_{\lambda} \leq \min \left\{ 1, \frac{h_{\mu_{\lambda}}}{\chi_{\mu_{\lambda}}^{ss}(\lambda) - \chi_{\mu_{\lambda}}^s(\lambda)} \right\} \text{ for every } \lambda \in U.$$

So it is enough to establish the lower bound. Let us argue by contradiction. Assume that there exist a set $U' \subset U$ with $\mathcal{L}_d(U') > 0$ and a $\varepsilon > 0$ that

$$\dim_H(e_{\lambda}^{ss})_*\mu_{\lambda} \leq \min \left\{ 1, \frac{h_{\mu_{\lambda}}}{\chi_{\mu_{\lambda}}^{ss}(\lambda) - \chi_{\mu_{\lambda}}^s(\lambda)} \right\} - \varepsilon \text{ for } \mathcal{L}_d\text{-a.e. } \lambda \in U'.$$

Let $\lambda_0 \in U'$ a Lebesgue density point. Thus, there exists a $\delta_0 > 0$ that for every $\delta_0 > \delta > 0$

$$\mathcal{L}_d \left(\lambda \in B_{\delta}(\lambda_0) : \dim_H(e_{\lambda}^{ss})_*\mu_{\lambda} \leq \min \left\{ 1, \frac{h_{\mu_{\lambda}}}{\chi_{\mu_{\lambda}}^{ss}(\lambda) - \chi_{\mu_{\lambda}}^s(\lambda)} \right\} - \varepsilon \right) > 0.$$

By using the continuity of entropy and Lyapunov exponents we have for sufficiently small $\delta > 0$

$$\mathcal{L}_d \left(\lambda \in B_{\delta}(\lambda_0) : \dim_H(e_{\lambda}^{ss})_*\mu_{\lambda} \leq \min \left\{ 1, \frac{h_{\mu_{\lambda_0}}}{\chi_{\mu_{\lambda_0}}^{ss}(\lambda_0) - \chi_{\mu_{\lambda_0}}^s(\lambda_0)} \right\} - \frac{\varepsilon}{2} \right) > 0,$$

but this contradicts Proposition 4.7. \square

Proof of Theorem 4.2. By Theorem 4.3, we have

$$\dim_H(e_{\lambda}^{ss})_*\mu_{\lambda} = \min \left\{ \frac{h_{\mu_{\lambda}}}{\chi_{\mu_{\lambda}}^{ss}(\lambda) - \chi_{\mu_{\lambda}}^s(\lambda)}, 1 \right\} \text{ for } \mathcal{L}_d\text{-a.e. } \lambda \in U.$$

On the other hand, by Theorem 3.6, if

$$\frac{h_{\mu_{\lambda}}}{\chi_{\mu_{\lambda}}^{ss}(\lambda) - \chi_{\mu_{\lambda}}^s(\lambda)} \geq \min \left\{ \frac{h_{\mu_{\lambda}}}{\chi_{\mu_{\lambda}}^s(\lambda)}, 1 \right\}$$

the statement holds. Thus, we may assume that

$$\frac{h_{\mu_{\lambda}}}{\chi_{\mu_{\lambda}}^{ss}(\lambda) - \chi_{\mu_{\lambda}}^s(\lambda)} < 1, \chi_{\mu_{\lambda}}^{ss}(\lambda) > 2\chi_{\mu_{\lambda}}^s(\lambda) \text{ and } \frac{h_{\mu_{\lambda}}}{\chi_{\mu_{\lambda}}^{ss}(\lambda) - \chi_{\mu_{\lambda}}^s(\lambda)} + 2\frac{h_{\mu_{\lambda}}}{\chi_{\mu_{\lambda}}^{ss}(\lambda)} > 2.$$

By [2, Lemma 4.12], we get that $\dim_H(\pi_{\lambda}^{-})_*\mu_{\lambda} \geq 2\frac{h_{\mu_{\lambda}}}{\chi_{\mu_{\lambda}}^{ss}(\lambda)}$ and the statement follows by Theorem 3.6. \square

5. PROOF OF THEOREM 1.1

Finally, in this section we prove Theorem 1.1 as an application of Theorem 4.2.

For a matrix $A \in \mathbb{R}_+^{2 \times 2} \cup \mathbb{R}_-^{2 \times 2}$ let

$$S(x, A) := \frac{|a|x + |c|(1-x)}{(|a| + |b|x + (|c| + |d|)(1-x))} \text{ where } A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}. \quad (5.1)$$

Simple calculations show that the maps $S_i \in C^2[0, 1]$, Moreover,

$$\sup_{x \in [0, 1]} |S'(x, A)| = \max \{ |S'(0, A)|, |S'(1, A)| \} = \frac{|\det A|}{\|A\|_2^2}, \text{ and} \quad (5.2)$$

$$\inf_{x \in [0, 1]} |S'(x, A)| = \min \{ |S'(0, A)|, |S'(1, A)| \} = \frac{|\det A|}{\|A\|_{\infty}^2},$$

where $\|A\|_{\infty} = \max \{ |a| + |b|, |c| + |d| \}$ the usual ∞ -norm of matrices.

Lemma 5.1. *Let $\mathcal{A} = \{A_1, \dots, A_N\}$ be a set of non-singular matrices with either strictly positive or strictly negative elements such that $\frac{|\det A_i|}{\|A_i\|^2} < 1$. Let $\phi = \{S_i(\cdot) := S(\cdot, A_i)\}_{i=1}^N$ be IFS on $[0, 1]$ and let $\Pi : \Sigma^+ \mapsto [0, 1]$ be the natural projection of ϕ . Then for every $\mathbf{i}_+ \in \Sigma^+$ the vector $(\Pi(\mathbf{i}_+) - 1, \Pi(\mathbf{i}_+))^T \in e^{ss}(\mathbf{i}_+)$.*

Proof. Let $\mathcal{A} = \{A_1, \dots, A_N\}$ and the IFS $\phi = \{S_1, \dots, S_N\}$ be as required. It is easy to see that the cone $M = \{(x, y) \in \mathbb{R}^2 / \{(0, 0)\} : xy \leq 0\}$ is backward invariant. So, by [3, Theorem B], \mathcal{A} satisfies the dominated splitting.

For an $\mathbf{i}_+ \in \Sigma^+$ let $e^{ss}(\mathbf{i}_+)$ be the invariant strong stable direction defined in (5.1). By the definition of $\Pi : \Sigma^+ \mapsto [0, 1]$

$$\begin{aligned} \begin{pmatrix} \Pi(\mathbf{i}_+) - 1 \\ \Pi(\mathbf{i}_+) \end{pmatrix} &= \frac{\begin{pmatrix} -b_{i_0}\Pi(\sigma\mathbf{i}_+) - d_{i_0}(1 - \Pi(\sigma\mathbf{i}_+)) \\ a_{i_0}\Pi(\sigma\mathbf{i}_+) + c_{i_0}(1 - \Pi(\sigma\mathbf{i}_+)) \end{pmatrix}}{(|a_{i_0}| + |b_{i_0}|)\Pi(\sigma\mathbf{i}_+) + (|c_{i_0}| + |d_{i_0}|)(1 - \Pi(\sigma\mathbf{i}_+))} = \\ &= \frac{\det A_{i_0}}{(|a_{i_0}| + |b_{i_0}|)\Pi(\sigma\mathbf{i}_+) + (|c_{i_0}| + |d_{i_0}|)(1 - \Pi(\sigma\mathbf{i}_+))} A_{i_0}^{-1} \begin{pmatrix} \Pi(\sigma\mathbf{i}_+) - 1 \\ \Pi(\sigma\mathbf{i}_+) \end{pmatrix}. \end{aligned}$$

Thus, by Lemma 2.2 and uniqueness, the 1 dimensional subspace $e^{ss}(\mathbf{i}_+)$ contains $(\Pi(\mathbf{i}_+) - 1, \Pi(\mathbf{i}_+))^T$. \square

Lemma 5.2. *Let $\mathcal{A} = \{A_1, \dots, A_N\}$ be arbitrary such that $A_i \in \mathfrak{M}$, where \mathfrak{M} is defined in (1.3). Moreover, let $\mathcal{A}(\underline{t}) = \{A_1 + t_1 B_1, \dots, A_N + t_N B_N\}$, where $\underline{t} \in \mathbb{R}^N$*

$$A_i = \begin{pmatrix} a_i & b_i \\ c_i & d_i \end{pmatrix} \text{ and } B_i = \begin{pmatrix} a_i + b_i & -(a_i + b_i) \\ c_i + d_i & -(c_i + d_i) \end{pmatrix}. \quad (5.3)$$

Then there exists a $\delta = \delta(\mathcal{A}) > 0$ such that the IFS $\phi_{\underline{t}} = \{S_i^{\underline{t}}(\cdot) := S(\cdot, A_i + t_i B_i)\}_{i=1}^N$ satisfies the transversality condition on $(-\delta, \delta)^N$.

In particular, $\mathcal{A}(\underline{t})$ satisfies the strong-stable transversality condition on $(-\delta, \delta)^N$.

Proof. Since \mathfrak{M}^N is open, there exists a $\varepsilon = \varepsilon(\mathcal{A}) > 0$ that $\mathcal{A}(\underline{t}) \in \mathfrak{M}^N$ for every $\underline{t} \in (-\varepsilon, \varepsilon)^N$. Let $\phi = \{S_1, \dots, S_N\}$ be the IFS for \mathcal{A} and $\phi_{\underline{t}} = \{S_1^{\underline{t}}, \dots, S_N^{\underline{t}}\}$ be the IFS for $\mathcal{A}(\underline{t})$. Simple calculations show that $S_i^{\underline{t}}(x) = S_i(x) + t_i$ for every $i = 1, \dots, N$. By the definition of \mathfrak{M} , by (5.2) and by [14, Corollary 7.3] there exists $\delta = \delta(\mathcal{A}) > 0$ such that $\delta < \varepsilon$ and $\phi_{\underline{t}}$ satisfies the transversality condition. By Lemma 5.2 and Definition 4.1, it follows that $\mathcal{A}(\underline{t})$ satisfies the strong-stable transversality on $(-\delta, \delta)^N$. \square

Lemma 5.3. *Let us define for every $\mathcal{A} \in \mathfrak{M}^N$*

$$P(\mathcal{A}) := \mathfrak{M}^N \cap \bigcup_{\underline{t} \in \mathbb{R}^N} \mathcal{A}(\underline{t}),$$

where $\mathcal{A}(\underline{t})$ is defined in Lemma 5.2. Then P defines a measurable partition of \mathfrak{M}^N .

Proof. By the definition of P it is enough to show that if $\mathcal{A} \neq \mathcal{A}'$ then either $P(\mathcal{A}) = P(\mathcal{A}')$ or $P(\mathcal{A}) \cap P(\mathcal{A}') = \emptyset$.

Let us fix $\mathcal{A} \neq \mathcal{A}'$ and suppose that $P(\mathcal{A}) \cap P(\mathcal{A}') \neq \emptyset$. Then there exist $t_1, \dots, t_N \in \mathbb{R}$ and $t'_1, \dots, t'_N \in \mathbb{R}$ that $A_i + t_i B_i = A'_i + t'_i B'_i$ for every $i = 1, \dots, N$, where B_i and B'_i defined in (5.3). Thus $a_i + b_i = a'_i + b'_i$ and $c_i + d_i = c'_i + d'_i$. Hence, $P(\mathcal{A}) = P(\mathcal{A}')$. The measurability is straightforward. \square

Proof of Theorem 1.1. First we show that if $\mathcal{A} \in \mathfrak{N}^N \cup \mathfrak{D}_N$, where \mathfrak{N}^N and \mathfrak{D}_N are defined in (1.4), then condition (iii) of Theorem 4.2 holds for the Käenmäki measure μ^K of \mathcal{A} , defined in Definition 2.6.

Indeed, if $\mathcal{A} \in \mathfrak{N}^N$ then $\frac{h_{\mu^K}}{\chi_{\mu^K}^{ss} - \chi_{\mu^K}^s} \geq \frac{h_{\mu^K}}{\chi_{\mu^K}^s}$ and on the other hand, if $\mathcal{A} \in \mathfrak{D}_N$ then

$$\begin{aligned} \frac{h_{\mu^K}}{\chi_{\mu^K}^{ss} - \chi_{\mu^K}^s} + 2 \frac{h_{\mu^K}}{\chi_{\mu^K}^{ss}} &= \frac{\chi_{\mu^K}^s + (s_0 - 1)\chi_{\mu^K}^{ss}}{\chi_{\mu^K}^{ss} - \chi_{\mu^K}^s} + 2 \frac{\chi_{\mu^K}^s + (s_0 - 1)\chi_{\mu^K}^{ss}}{\chi_{\mu^K}^{ss}} = \\ &= -3 + \left(2 + \frac{1}{1 - \frac{\chi_{\mu^K}^s}{\chi_{\mu^K}^{ss}}} \right) s_0 + 2 \frac{\chi_{\mu^K}^s}{\chi_{\mu^K}^{ss}} > \frac{1}{3} + \frac{5}{3 \left(1 - \frac{\chi_{\mu^K}^s}{\chi_{\mu^K}^{ss}} \right)} + 2 \frac{\chi_{\mu^K}^s}{\chi_{\mu^K}^{ss}} > 2. \end{aligned}$$

Now, let $V \subset \mathfrak{N}^N \cup \mathfrak{D}_N \subset \mathfrak{M}^N$ be a compact set such that $\overline{V^o} = V$. Let us define for a $\mathcal{A} \in V$

$$Q(\mathcal{A}) := V \cap P(\mathcal{A}),$$

Thus, $\bigcup_{\mathcal{B} \in P(\mathcal{A})} \left\{ \bigcup_{\underline{t} \in (-\delta(\mathcal{B}), \delta(\mathcal{B}))^N} \mathcal{B}(\underline{t}) \right\}$ defines an open cover of $Q(\mathcal{A})$. Since $Q(\mathcal{A})$ is compact there is a finite set $\{\mathcal{B}_1, \dots, \mathcal{B}_n\}$ that $\bigcup_{i=1}^n \left\{ \bigcup_{\underline{t} \in (-\delta(\mathcal{B}_i), \delta(\mathcal{B}_i))^N} \mathcal{B}_i(\underline{t}) \right\}$ is a cover for $Q(\mathcal{A})$. But by Lemma 5.2, for every $i = 1, \dots, n$ the parametrized family of matrices $\mathcal{B}_i(\underline{t})$ satisfies the strong-stable transversality condition on $(-\delta(\mathcal{B}_i), \delta(\mathcal{B}_i))^N$. Thus, by Theorem 4.2 for every $i = 1, \dots, n$

$$\dim_H \mu_{\underline{t}}^K = \dim_H \Lambda_{\underline{t}} = \dim_B \Lambda_{\underline{t}} = s_0(\underline{t}) \text{ for } \mathcal{L}_N\text{-a.e } \underline{t} \in (-\delta(\mathcal{B}_i), \delta(\mathcal{B}_i))^N,$$

where $\mu_{\underline{t}}^K$ is the Käenmäki measure of the system $\mathcal{B}_i(\underline{t})$ and $s_0(\underline{t})$ is the affinity dimension. In particular, for every $\mathcal{A} \in V$

$$\dim_H \mu^K = \dim_H \Lambda = \dim_B \Lambda = s_0(\mathcal{B}) \text{ for } \mathcal{L}_N\text{-a.e } \mathcal{B} \in Q(\mathcal{A}).$$

By Lemma 5.3, Q is a measurable foliation of V , thus, by Rokhlin's Theorem

$$\dim_H \mu^K = \dim_H \Lambda = \dim_B \Lambda = s_0(\mathcal{A}) \text{ for } \mathcal{L}_{4N}\text{-a.e. } \mathcal{A} \in V.$$

Since V was arbitrary, the statement follows. \square

REFERENCES

- [1] A. Avila, J. Bochi and J.-C. Yoccoz: Uniformly hyperbolic finite-valued $SL(2, \mathbb{R})$ cocycles, *Comment. Math. Helv.* **85** (2010), 813-884.
- [2] B. Bárány: On the Ledrappier-Young formula for self-affine measures, preprint, 2015, available at arXiv:1503.00892.
- [3] J. Bochi and N. Gourmelon: Some characterizations of domination, *Math. Z.* **263** (2009), no. 1, 221-231.
- [4] J. Bochi and M. Rams: The entropy of Lyapunov-optimizing measures of some matrix cocycles, preprint, 2014, available at arXiv:1312.6718.
- [5] R. Bowen: *Equilibrium states and the ergodic theory of Anosov diffeomorphisms*, Second revised edition. With a preface by David Ruelle. Edited by Jean-Ren Chazottes. Lecture Notes in Mathematics, 470. Springer-Verlag, Berlin, 2008.
- [6] K. Falconer: The Hausdorff dimension of self-affine fractals, *Math. Proc. Camb. Phil. Soc.* **103** (1988), 339-350.
- [7] K. Falconer: *Fractal Geometry: Mathematical Foundations and Applications*, John Wiley and Sons, 1990.
- [8] K. Falconer: The dimension of self-affine fractals II, *Math. Proc. Camb. Phil. Soc.* **111** (1992), 169-179.
- [9] K. Falconer and T. Kempton: Planar self-affine sets with equal Hausdorff, box and affinity dimensions, preprint, 2015, available at arXiv:1503.01270.
- [10] I. Hueter and S. P. Lalley: Falconer's formula for the Hausdorff dimension of a self-affine set in \mathbb{R}^2 , *Erg. Th. & Dyn. Sys.* **15** (1995), no. 1, 77-97.
- [11] A. Käenmäki: Measures of full dimension on self-affine sets, *Acta Univ. Carolin. Math. Phys.* **45** (2004), no. 2, 45-53.
- [12] P. T. Maker: Ergodic Theorem for a sequence of functions, *Duke Math. J.* **6** (1940), 27-30.
- [13] V. A. Rokhlin: On the fundamental ideas of measure theory, *AMS Trans.* **10** (1962), 1-52.
- [14] K. Simon, B. Solomyak, and M. Urbański: Invariant measures for parabolic IFS with overlaps and random continued fractions, *Trans. Amer. Math. Soc.* **353** (2001), 5145-5164.

- [15] B. Solomyak: Measure and dimension for some fractal families, *Math. Proc. Camb. Phil. Soc.* **124**, (1998), no. 3, 531-546.
- [16] P. Walters: *An introduction to ergodic theory*, volume 79 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1982.
- [17] J.-C. Yoccoz: Some questions and remarks about $SL(2, \mathbb{R})$ cocycles, *Modern dynamical systems and applications*, Cambridge University Press, Cambridge 2004, 447-458.

(Balázs Bárány) BUDAPEST UNIVERSITY OF TECHNOLOGY AND ECONOMICS, DEPARTMENT OF STOCHASTICS, P.O.BOX 91, 1521 BUDAPEST, HUNGARY & MATHEMATICS INSTITUTE, UNIVERSITY OF WARWICK, COVENTRY CV4 7AL, UK
E-mail address: balubsheep@gmail.com

(Michał Rams) INSTITUTE OF MATHEMATICS, POLISH ACADEMY OF SCIENCES, UL. ŚNIADECKICH 8, 00-656 WARSZAWA, POLAND
E-mail address: rams@impan.pl