

Amenability, unique ergodicity and random orderings

Alexander S. Kechris


WARSAW, JULY 10, 2012

I will discuss some aspects of the ergodic theory of automorphism groups of countable structures and its connections with finite Ramsey theory and probability theory. This is joint work with Omer Angel and Russell Lyons.

Throughout I will consider countable first-order languages and countable (finite or infinite) structures for such languages. Recall first some standard concepts of Fraïssé theory.

Definition

A class \mathcal{K} of finite structures of the same language is called a **Fraïssé class** if it satisfies the following properties:


- (HP) Hereditary property.
- (JEP) Joint embedding property.
- (AP) Amalgamation property. 
- It is countable (up to \cong).
- It is unbounded.



Throughout I will consider countable first-order languages and countable (finite or infinite) structures for such languages. Recall first some standard concepts of Fraïssé theory.

Definition


A class \mathcal{K} of finite structures of the same language is called a **Fraïssé class** if it satisfies the following properties:

- (HP) Hereditary property.
- (JEP) Joint embedding property.
- (AP) Amalgamation property. 
- It is countable (up to \cong).
- It is unbounded.




Fraïssé theory

Joint embedding property (JEP)



Amalgamation property (AP)



Definition

A countable structure \mathbf{K} is a **Fraïssé structure** if it satisfies the following properties:

- It is infinite.
- It is locally finite.
- It is **ultrahomogeneous** (i.e., an isomorphism between finite substructures can be extended to an automorphism of the whole structure).

Definition

For a structure \mathbf{A} , its **age**, denoted by $\text{Age}(\mathbf{A})$, is the class of finite structures that can be embedded in \mathbf{A} .

Definition

A countable structure \mathbf{K} is a **Fraïssé structure** if it satisfies the following properties:

- It is infinite.
- It is locally finite.
- It is **ultrahomogeneous** (i.e., an isomorphism between finite substructures can be extended to an automorphism of the whole structure).

Definition

For a structure \mathbf{A} , its **age**, denoted by $\text{Age}(\mathbf{A})$, is the class of finite structures that can be embedded in \mathbf{A} .

The age of a Fraïssé structure is a Fraïssé class and Fraïssé showed that one can associate to each Fraïssé class \mathcal{K} a canonical Fraïssé structure $\mathbf{K} = \text{Frlim}(\mathcal{K})$, called its **Fraïssé limit**, which is the unique Fraïssé structure whose age is equal to \mathcal{K} . Therefore one has a canonical one-to-one correspondence:

$$\mathcal{K} \mapsto \text{Frlim}(\mathcal{K})$$

between Fraïssé classes and Fraïssé structures whose inverse is:

$$\mathbf{K} \mapsto \text{Age}(\mathbf{K}).$$

Examples

- finite graphs \rightleftarrows random graph
- finite linear orderings \rightleftarrows $\langle \mathbb{Q}, < \rangle$
- f.d. vector spaces \rightleftarrows (countable) infinite-dimensional vector space (over a finite field)
- finite Boolean algebras \rightleftarrows countable atomless Boolean algebra
- finite rational metric spaces \rightleftarrows rational Urysohn space

Examples

- finite graphs \rightleftarrows random graph
- finite linear orderings \rightleftarrows $\langle \mathbb{Q}, < \rangle$
- f.d. vector spaces \rightleftarrows (countable) infinite-dimensional vector space (over a finite field)
- finite Boolean algebras \rightleftarrows countable atomless Boolean algebra
- finite rational metric spaces \rightleftarrows rational Urysohn space

Examples

- finite graphs \rightleftarrows random graph
- finite linear orderings \rightleftarrows $\langle \mathbb{Q}, < \rangle$
- f.d. vector spaces \rightleftarrows (countable) infinite-dimensional vector space (over a finite field)
- finite Boolean algebras \rightleftarrows countable atomless Boolean algebra
- finite rational metric spaces \rightleftarrows rational Urysohn space

Examples

- finite graphs \rightleftarrows random graph
- finite linear orderings \rightleftarrows $\langle \mathbb{Q}, < \rangle$
- f.d. vector spaces \rightleftarrows (countable) infinite-dimensional vector space (over a finite field)
- finite Boolean algebras \rightleftarrows countable atomless Boolean algebra
- finite rational metric spaces \rightleftarrows rational Urysohn space

Examples

- finite graphs \rightleftarrows random graph
- finite linear orderings \rightleftarrows $\langle \mathbb{Q}, < \rangle$
- f.d. vector spaces \rightleftarrows (countable) infinite-dimensional vector space (over a finite field)
- finite Boolean algebras \rightleftarrows countable atomless Boolean algebra
- finite rational metric spaces \rightleftarrows rational Urysohn space

Examples

- finite graphs \rightleftarrows random graph
- finite linear orderings \rightleftarrows $\langle \mathbb{Q}, < \rangle$
- f.d. vector spaces \rightleftarrows (countable) infinite-dimensional vector space (over a finite field)
- finite Boolean algebras \rightleftarrows countable atomless Boolean algebra
- finite rational metric spaces \rightleftarrows rational Urysohn space

$\text{Aut}(\mathbf{A})$ as a topological group

For a countable structure \mathbf{A} , we view $\text{Aut}(\mathbf{A})$ as a topological group with the pointwise convergence topology. It is not hard to check then that it becomes a Polish group. In fact one can characterize these groups as follows:

Theorem

For any Polish group G , the following are equivalent:

- *G is isomorphic to a closed subgroup of S_∞ , the permutation group of \mathbb{N} with the pointwise convergence topology.*
- *G is non-Archimedean, i.e., admits a basis at the identity consisting of open subgroups.*
- *$G \cong \text{Aut}(\mathbf{A})$, for a countable structure \mathbf{A} .*
- *$G \cong \text{Aut}(\mathbf{K})$, for a Fraïssé structure \mathbf{K} .*

$\text{Aut}(\mathbf{A})$ as a topological group

For a countable structure \mathbf{A} , we view $\text{Aut}(\mathbf{A})$ as a topological group with the pointwise convergence topology. It is not hard to check then that it becomes a Polish group. In fact one can characterize these groups as follows:

Theorem

For any Polish group G , the following are equivalent:

- *G is isomorphic to a closed subgroup of S_∞ , the permutation group of \mathbb{N} with the pointwise convergence topology.*
- *G is non-Archimedean, i.e., admits a basis at the identity consisting of open subgroups.*
- *$G \cong \text{Aut}(\mathbf{A})$, for a countable structure \mathbf{A} .*
- *$G \cong \text{Aut}(\mathbf{K})$, for a Fraïssé structure \mathbf{K} .*

Amenability of $\text{Aut}(\mathbf{A})$

We will now consider some aspects of the dynamics of automorphism groups, especially the concept of amenability.

Definition

Let G be a topological group. A G -flow is a continuous action of G on a compact Hausdorff space. A group G is called **amenable** if every G -flow admits an invariant (Borel probability) measure. It is called **extremely amenable** if every G -flow admits an invariant point.

Amenability of $\text{Aut}(\mathbf{A})$

We will now consider some aspects of the dynamics of automorphism groups, especially the concept of amenability.

Definition

Let G be a topological group. A G -flow is a continuous action of G on a compact Hausdorff space. A group G is called **amenable** if every G -flow admits an invariant (Borel probability) measure. It is called **extremely amenable** if every G -flow admits an invariant point.

In a paper of K-Pestov-Todorčević (2005) a duality theory was developed that relates the Ramsey theory of Fraïssé classes (sometimes called **structural Ramsey theory**) to the topological dynamics of the automorphism groups of their Fraïssé limits.

Structural Ramsey theory is a vast generalization of the classical Ramsey theorem to classes of finite structures. It was developed primarily in the 1970's by: Graham, Leeb, Rothchild, Nešetřil-Rödl, Prömel, Voigt, Abramson-Harrington, ...

Extreme amenability and Ramsey theory

In a paper of K-Pestov-Todorčević (2005) a duality theory was developed that relates the Ramsey theory of Fraïssé classes (sometimes called **structural Ramsey theory**) to the topological dynamics of the automorphism groups of their Fraïssé limits.

Structural Ramsey theory is a vast generalization of the classical Ramsey theorem to classes of finite structures. It was developed primarily in the 1970's by: Graham, Leeb, Rothchild, Nešetřil-Rödl, Prömel, Voigt, Abramson-Harrington, ...

Definition

A class \mathcal{K} of finite structures (in the same language) has the **Ramsey property (RP)** if for any $A \leq B$ in \mathcal{K} , and any $n \geq 1$, there is $C \geq B$ in \mathcal{K} , such that

$$C \rightarrow (B)_n^A.$$

Examples of classes with Ramsey property:

- finite linear orderings (Ramsey)
- finite Boolean algebras (Graham-Rothschild)
- finite-dimensional vector spaces over a given finite field (Graham-Leeb-Rothschild)
- finite ordered graphs (Nešetřil-Rödl)
- finite rational metric spaces (Nešetřil)

However, the class of finite graphs does not have the Ramsey property!

Definition

A class \mathcal{K} of finite structures (in the same language) has the **Ramsey property (RP)** if for any $A \leq B$ in \mathcal{K} , and any $n \geq 1$, there is $C \geq B$ in \mathcal{K} , such that

$$C \rightarrow (B)_n^A.$$

Examples of classes with Ramsey property:

- finite linear orderings (Ramsey)
- finite Boolean algebras (Graham-Rothschild)
- finite-dimensional vector spaces over a given finite field (Graham-Leeb-Rothschild)
- finite ordered graphs (Nešetřil-Rödl)
- finite rational metric spaces (Nešetřil)

However, the class of finite graphs does not have the Ramsey property!

Definition

A class \mathcal{K} of finite structures (in the same language) has the **Ramsey property (RP)** if for any $A \leq B$ in \mathcal{K} , and any $n \geq 1$, there is $C \geq B$ in \mathcal{K} , such that

$$C \rightarrow (B)_n^A.$$

Examples of classes with Ramsey property:

- finite linear orderings (Ramsey)
- finite Boolean algebras (Graham-Rothschild)
- finite-dimensional vector spaces over a given finite field (Graham-Leeb-Rothschild)
- finite ordered graphs (Nešetřil-Rödl)
- finite rational metric spaces (Nešetřil)

However, the class of finite graphs does not have the Ramsey property!

Definition

A class \mathcal{K} of finite structures (in the same language) has the **Ramsey property (RP)** if for any $A \leq B$ in \mathcal{K} , and any $n \geq 1$, there is $C \geq B$ in \mathcal{K} , such that

$$C \rightarrow (B)_n^A.$$

Examples of classes with Ramsey property:

- finite linear orderings (Ramsey)
- finite Boolean algebras (Graham-Rothschild)
- finite-dimensional vector spaces over a given finite field (Graham-Leeb-Rothschild)
- finite ordered graphs (Nešetřil-Rödl)
- finite rational metric spaces (Nešetřil)

However, the class of finite graphs does not have the Ramsey property!

Definition

A class \mathcal{K} of finite structures (in the same language) has the **Ramsey property (RP)** if for any $A \leq B$ in \mathcal{K} , and any $n \geq 1$, there is $C \geq B$ in \mathcal{K} , such that

$$C \rightarrow (B)_n^A.$$

Examples of classes with Ramsey property:

- finite linear orderings (Ramsey)
- finite Boolean algebras (Graham-Rothschild)
- finite-dimensional vector spaces over a given finite field (Graham-Leeb-Rothschild)
- finite ordered graphs (Nešetřil-Rödl)
- finite rational metric spaces (Nešetřil)

However, the class of finite graphs does not have the Ramsey property!

Definition

A class \mathcal{K} of finite structures (in the same language) has the **Ramsey property (RP)** if for any $A \leq B$ in \mathcal{K} , and any $n \geq 1$, there is $C \geq B$ in \mathcal{K} , such that

$$C \rightarrow (B)_n^A.$$

Examples of classes with Ramsey property:

- finite linear orderings (Ramsey)
- finite Boolean algebras (Graham-Rothschild)
- finite-dimensional vector spaces over a given finite field (Graham-Leeb-Rothschild)
- finite ordered graphs (Nešetřil-Rödl)
- finite rational metric spaces (Nešetřil)

However, the class of finite graphs does not have the Ramsey property!

Definition

A class \mathcal{K} of finite structures (in the same language) has the **Ramsey property (RP)** if for any $A \leq B$ in \mathcal{K} , and any $n \geq 1$, there is $C \geq B$ in \mathcal{K} , such that

$$C \rightarrow (B)_n^A.$$

Examples of classes with Ramsey property:

- finite linear orderings (Ramsey)
- finite Boolean algebras (Graham-Rothschild)
- finite-dimensional vector spaces over a given finite field (Graham-Leeb-Rothschild)
- finite ordered graphs (Nešetřil-Rödl)
- finite rational metric spaces (Nešetřil)

However, the class of finite graphs does not have the Ramsey property!

Definition

A class \mathcal{K} of finite structures (in the same language) has the **Ramsey property (RP)** if for any $A \leq B$ in \mathcal{K} , and any $n \geq 1$, there is $C \geq B$ in \mathcal{K} , such that

$$C \rightarrow (B)_n^A.$$

Examples of classes with Ramsey property:

- finite linear orderings (Ramsey)
- finite Boolean algebras (Graham-Rothschild)
- finite-dimensional vector spaces over a given finite field (Graham-Leeb-Rothschild)
- finite ordered graphs (Nešetřil-Rödl)
- finite rational metric spaces (Nešetřil)

However, the class of finite graphs does not have the Ramsey property!

Extreme amenability and Ramsey theory

One of the consequences of the duality theory is the following characterization of extreme amenability of automorphism groups.

Theorem (KPT)

The extremely amenable automorphism groups are exactly the automorphism groups of ordered Fraïssé structures whose age satisfies the Ramsey Property.

Examples

The automorphism groups of the following structures are extremely amenable:

- random ordered graph
- (Pestov) rational order
- lex. ordered infinite-dimensional vector space (over a finite field)
- lex. ordered countable atomless Boolean algebra
- rational ordered Urysohn space

Extreme amenability and Ramsey theory

One of the consequences of the duality theory is the following characterization of extreme amenability of automorphism groups.

Theorem (KPT)

The extremely amenable automorphism groups are exactly the automorphism groups of ordered Fraïssé structures whose age satisfies the Ramsey Property.

Examples

The automorphism groups of the following structures are extremely amenable:

- random ordered graph
- (Pestov) rational order
- lex. ordered infinite-dimensional vector space (over a finite field)
- lex. ordered countable atomless Boolean algebra
- rational ordered Urysohn space

Extreme amenability and Ramsey theory

One of the consequences of the duality theory is the following characterization of extreme amenability of automorphism groups.

Theorem (KPT)

The extremely amenable automorphism groups are exactly the automorphism groups of ordered Fraïssé structures whose age satisfies the Ramsey Property.

Examples

The automorphism groups of the following structures are extremely amenable:

- random ordered graph
- (Pestov) rational order
- lex. ordered infinite-dimensional vector space (over a finite field)
- lex. ordered countable atomless Boolean algebra
- rational ordered Urysohn space

Extreme amenability and Ramsey theory

One of the consequences of the duality theory is the following characterization of extreme amenability of automorphism groups.

Theorem (KPT)

The extremely amenable automorphism groups are exactly the automorphism groups of ordered Fraïssé structures whose age satisfies the Ramsey Property.

Examples

The automorphism groups of the following structures are extremely amenable:

- random ordered graph
- (Pestov) rational order
- lex. ordered infinite-dimensional vector space (over a finite field)
- lex. ordered countable atomless Boolean algebra
- rational ordered Urysohn space

Extreme amenability and Ramsey theory

One of the consequences of the duality theory is the following characterization of extreme amenability of automorphism groups.

Theorem (KPT)

The extremely amenable automorphism groups are exactly the automorphism groups of ordered Fraïssé structures whose age satisfies the Ramsey Property.

Examples

The automorphism groups of the following structures are extremely amenable:

- random ordered graph
- (Pestov) rational order
- lex. ordered infinite-dimensional vector space (over a finite field)
- lex. ordered countable atomless Boolean algebra
- rational ordered Urysohn space

Extreme amenability and Ramsey theory

One of the consequences of the duality theory is the following characterization of extreme amenability of automorphism groups.

Theorem (KPT)

The extremely amenable automorphism groups are exactly the automorphism groups of ordered Fraïssé structures whose age satisfies the Ramsey Property.

Examples

The automorphism groups of the following structures are extremely amenable:

- random ordered graph
- (Pestov) rational order
- lex. ordered infinite-dimensional vector space (over a finite field)
- lex. ordered countable atomless Boolean algebra
- rational ordered Urysohn space

Extreme amenability and Ramsey theory

One of the consequences of the duality theory is the following characterization of extreme amenability of automorphism groups.

Theorem (KPT)

The extremely amenable automorphism groups are exactly the automorphism groups of ordered Fraïssé structures whose age satisfies the Ramsey Property.

Examples

The automorphism groups of the following structures are extremely amenable:

- random ordered graph
- (Pestov) rational order
- lex. ordered infinite-dimensional vector space (over a finite field)
- lex. ordered countable atomless Boolean algebra
- rational ordered Urysohn space

Extreme amenability and Ramsey theory

One of the consequences of the duality theory is the following characterization of extreme amenability of automorphism groups.

Theorem (KPT)

The extremely amenable automorphism groups are exactly the automorphism groups of ordered Fraïssé structures whose age satisfies the Ramsey Property.

Examples

The automorphism groups of the following structures are extremely amenable:

- random ordered graph
- (Pestov) rational order
- lex. ordered infinite-dimensional vector space (over a finite field)
- lex. ordered countable atomless Boolean algebra
- rational ordered Urysohn space

Extreme amenability and Ramsey theory

One of the consequences of the duality theory is the following characterization of extreme amenability of automorphism groups.

Theorem (KPT)

The extremely amenable automorphism groups are exactly the automorphism groups of ordered Fraïssé structures whose age satisfies the Ramsey Property.

Examples

The automorphism groups of the following structures are extremely amenable:

- random ordered graph
- (Pestov) rational order
- lex. ordered infinite-dimensional vector space (over a finite field)
- lex. ordered countable atomless Boolean algebra
- rational ordered Urysohn space

Hrushovski structures

Clearly every extremely amenable group is amenable. There are however many amenable automorphism groups that are not extremely amenable. Many such examples arise in the context of the Hrushovski Property.

Definition

Let \mathcal{K} be a Fraïssé class of finite structures. We say that \mathcal{K} is a **Hrushovski class** if for any A in \mathcal{K} there is B in \mathcal{K} containing A such that any partial automorphism of A extends to an automorphism of B .

Some basic examples of such classes are the pure sets, graphs (Hrushovski), hypergraphs and K_n -free graphs (Herwig), rational valued metric spaces (Solecki), finite dimensional vector spaces over finite fields, etc.

Definition

Let \mathcal{K} be a Fraïssé class of finite structures and K its Fraïssé limit. If \mathcal{K} is a Hrushovski class, then we say that K is a **Hrushovski structure**.

Hrushovski structures

Clearly every extremely amenable group is amenable. There are however many amenable automorphism groups that are not extremely amenable. Many such examples arise in the context of the Hrushovski Property.

Definition

Let \mathcal{K} be a Fraïssé class of finite structures. We say that \mathcal{K} is a **Hrushovski class** if for any A in \mathcal{K} there is B in \mathcal{K} containing A such that any partial automorphism of A extends to an automorphism of B .

Some basic examples of such classes are the pure sets, graphs (Hrushovski), hypergraphs and K_n -free graphs (Herwig), rational valued metric spaces (Solecki), finite dimensional vector spaces over finite fields, etc.

Definition

Let \mathcal{K} be a Fraïssé class of finite structures and K its Fraïssé limit. If \mathcal{K} is a Hrushovski class, then we say that K is a **Hrushovski structure**.

Hrushovski structures

Clearly every extremely amenable group is amenable. There are however many amenable automorphism groups that are not extremely amenable. Many such examples arise in the context of the Hrushovski Property.

Definition

Let \mathcal{K} be a Fraïssé class of finite structures. We say that \mathcal{K} is a **Hrushovski class** if for any A in \mathcal{K} there is B in \mathcal{K} containing A such that any partial automorphism of A extends to an automorphism of B .

Some basic examples of such classes are the pure sets, graphs (Hrushovski), hypergraphs and K_n -free graphs (Herwig), rational valued metric spaces (Solecki), finite dimensional vector spaces over finite fields, etc.

Definition

Let \mathcal{K} be a Fraïssé class of finite structures and K its Fraïssé limit. If \mathcal{K} is a Hrushovski class, then we say that K is a **Hrushovski structure**.

Hrushovski structures

Clearly every extremely amenable group is amenable. There are however many amenable automorphism groups that are not extremely amenable. Many such examples arise in the context of the Hrushovski Property.

Definition

Let \mathcal{K} be a Fraïssé class of finite structures. We say that \mathcal{K} is a **Hrushovski class** if for any A in \mathcal{K} there is B in \mathcal{K} containing A such that any partial automorphism of A extends to an automorphism of B .

Some basic examples of such classes are the pure sets, graphs (Hrushovski), hypergraphs and K_n -free graphs (Herwig), rational valued metric spaces (Solecki), finite dimensional vector spaces over finite fields, etc.

Definition

Let \mathcal{K} be a Fraïssé class of finite structures and \mathbf{K} its Fraïssé limit. If \mathcal{K} is a Hrushovski class, then we say that \mathbf{K} is a **Hrushovski structure**.

Hrushovski structures

This turns out to be a property of automorphism groups:

Proposition (K-Rosendal)

Let \mathcal{K} be a Fraïssé class of finite structures and \mathbf{K} its Fraïssé limit. Then the following are equivalent

- *\mathbf{K} is a Hrushovski structure.*
- *$\text{Aut}(\mathbf{K})$ is compactly approximable, i.e., there is a increasing sequence K_n of compact subgroups whose union is dense in the automorphism group.*

In particular the automorphism group of a Hrushovski structure is amenable. Thus S_∞ and the automorphism groups of the random graph, random n -uniform hypergraph, random K_n -free graph, rational Urysohn space, (countably) infinite-dimensional vector space over a finite field, etc., are amenable (but not extremely amenable).

This turns out to be a property of automorphism groups:

Proposition (K-Rosendal)

Let \mathcal{K} be a Fraïssé class of finite structures and \mathbf{K} its Fraïssé limit. Then the following are equivalent

- *\mathbf{K} is a Hrushovski structure.*
- *$\text{Aut}(\mathbf{K})$ is compactly approximable, i.e., there is an increasing sequence K_n of compact subgroups whose union is dense in the automorphism group.*

In particular the automorphism group of a Hrushovski structure is amenable. Thus S_∞ and the automorphism groups of the random graph, random n -uniform hypergraph, random K_n -free graph, rational Urysohn space, (countably) infinite-dimensional vector space over a finite field, etc., are amenable (but not extremely amenable).

This turns out to be a property of automorphism groups:

Proposition (K-Rosendal)

Let \mathcal{K} be a Fraïssé class of finite structures and \mathbf{K} its Fraïssé limit. Then the following are equivalent

- *\mathbf{K} is a Hrushovski structure.*
- *$\text{Aut}(\mathbf{K})$ is compactly approximable, i.e., there is an increasing sequence K_n of compact subgroups whose union is dense in the automorphism group.*

In particular the automorphism group of a Hrushovski structure is amenable. Thus S_∞ and the automorphism groups of the random graph, random n -uniform hypergraph, random K_n -free graph, rational Urysohn space, (countably) infinite-dimensional vector space over a finite field, etc., are amenable (but not extremely amenable).

At the other end of the spectrum there are also automorphism groups that are not amenable. These include the following:

Theorem (K-Sokić)

The automorphism groups of the random poset, random distributive lattice and the dense local order are not amenable.

Unique ergodicity

I am interested here in the ergodic theory of flows of automorphism groups and especially in the phenomenon of unique ergodicity. Let G be a topological group and X a G -flow. Consider G -invariant (Borel probability) measures in such a flow.

Definition

A G -flow is **uniquely ergodic** if it admits a unique invariant measure (which must then be ergodic).

I am interested here in the ergodic theory of flows of automorphism groups and especially in the phenomenon of unique ergodicity. Let G be a topological group and X a G -flow. Consider G -invariant (Borel probability) measures in such a flow.

Definition

A G -flow is **uniquely ergodic** if it admits a unique invariant measure (which must then be ergodic).

Unique ergodicity

Recall that a flow is called **minimal** if every orbit is dense or equivalently if it has no proper subflows. Every flow contains a minimal subflow.

Definition

Let G be a topological group. We call G **uniquely ergodic** if every minimal flow admits a unique invariant measure (which must then be ergodic).

Remark: The assumption of minimality is necessary because in general a flow has many minimal subflows which are of course pairwise disjoint. Note also that every uniquely ergodic group is amenable.

Clearly every extremely amenable Polish group is uniquely ergodic and so is every compact Polish group. On the other hand Weiss has shown that no infinite countable (discrete) group can be uniquely ergodic and he believes that this extends to Polish locally compact, non-compact groups although this has not been verified in detail.

Unique ergodicity

Recall that a flow is called **minimal** if every orbit is dense or equivalently if it has no proper subflows. Every flow contains a minimal subflow.

Definition

Let G be a topological group. We call G **uniquely ergodic** if every minimal flow admits a unique invariant measure (which must then be ergodic).

Remark: The assumption of minimality is necessary because in general a flow has many minimal subflows which are of course pairwise disjoint. Note also that every uniquely ergodic group is amenable.

Clearly every extremely amenable Polish group is uniquely ergodic and so is every compact Polish group. On the other hand Weiss has shown that no infinite countable (discrete) group can be uniquely ergodic and he believes that this extends to Polish locally compact, non-compact groups although this has not been verified in detail.

Unique ergodicity

Recall that a flow is called **minimal** if every orbit is dense or equivalently if it has no proper subflows. Every flow contains a minimal subflow.

Definition

Let G be a topological group. We call G **uniquely ergodic** if every minimal flow admits a unique invariant measure (which must then be ergodic).

Remark: The assumption of minimality is necessary because in general a flow has many minimal subflows which are of course pairwise disjoint. Note also that every uniquely ergodic group is amenable.

Clearly every extremely amenable Polish group is uniquely ergodic and so is every compact Polish group. On the other hand Weiss has shown that no infinite countable (discrete) group can be uniquely ergodic and he believes that this extends to Polish locally compact, non-compact groups although this has not been verified in detail.

Unique ergodicity

Recall that a flow is called **minimal** if every orbit is dense or equivalently if it has no proper subflows. Every flow contains a minimal subflow.

Definition

Let G be a topological group. We call G **uniquely ergodic** if every minimal flow admits a unique invariant measure (which must then be ergodic).

Remark: The assumption of minimality is necessary because in general a flow has many minimal subflows which are of course pairwise disjoint. Note also that every uniquely ergodic group is amenable.

Clearly every extremely amenable Polish group is uniquely ergodic and so is every compact Polish group. On the other hand Weiss has shown that no infinite countable (discrete) group can be uniquely ergodic and he believes that this extends to Polish locally compact, non-compact groups although this has not been verified in detail.

Universal minimal flows

In order to understand better the concept of unique ergodicity we need to discuss first the idea of a universal minimal flow.

A **homomorphism** between two G -flows X, Y is a continuous G -map $\pi : X \rightarrow Y$. If Y is minimal, then π must be onto. An **isomorphism** is a bijective homomorphism.

Theorem

*For any G , there is a minimal G -flow, $M(G)$, called its **universal minimal flow** with the following property: For any minimal G -flow X , there is a homomorphism $\pi : M(G) \rightarrow X$. Moreover $M(G)$ is uniquely determined up to isomorphism by this property.*

Universal minimal flows

In order to understand better the concept of unique ergodicity we need to discuss first the idea of a universal minimal flow.

A **homomorphism** between two G -flows X, Y is a continuous G -map $\pi : X \rightarrow Y$. If Y is minimal, then π must be onto. An **isomorphism** is a bijective homomorphism.

Theorem

*For any G , there is a minimal G -flow, $M(G)$, called its **universal minimal flow** with the following property: For any minimal G -flow X , there is a homomorphism $\pi : M(G) \rightarrow X$. Moreover $M(G)$ is uniquely determined up to isomorphism by this property.*

Universal minimal flows

In order to understand better the concept of unique ergodicity we need to discuss first the idea of a universal minimal flow.

A **homomorphism** between two G -flows X, Y is a continuous G -map $\pi : X \rightarrow Y$. If Y is minimal, then π must be onto. An **isomorphism** is a bijective homomorphism.

Theorem

*For any G , there is a minimal G -flow, $M(G)$, called its **universal minimal flow** with the following property: For any minimal G -flow X , there is a homomorphism $\pi : M(G) \rightarrow X$. Moreover $M(G)$ is uniquely determined up to isomorphism by this property.*

The following is a consequence of the Hahn-Banach Theorem.

Proposition

Let G be an amenable group. Then G is uniquely ergodic iff $M(G)$ is uniquely ergodic.

So it is enough to concentrate on the universal minimal flow.

The following is a consequence of the Hahn-Banach Theorem.

Proposition

Let G be an amenable group. Then G is uniquely ergodic iff $M(G)$ is uniquely ergodic.

So it is enough to concentrate on the universal minimal flow.

The following is a consequence of the Hahn-Banach Theorem.

Proposition

Let G be an amenable group. Then G is uniquely ergodic iff $M(G)$ is uniquely ergodic.

So it is enough to concentrate on the universal minimal flow.

Universal minimal flows

If G is compact, then $M(G) = G$. If G is non-compact but locally compact, then $M(G)$ is extremely complicated, e.g., it is non-metrizable. However, by definition G is extremely amenable iff $M(G)$ trivializes!

This leads to a general problem in topological dynamics:

For a given G , can one explicitly determine $M(G)$ and show that it is metrizable?

The first example of calculation of a metrizable but non-trivial universal minimal flow is due to Pestov (1998): The universal minimal flow of $H_+(\mathbb{T})$ is \mathbb{T} . Two more examples were found later by Glasner-Weiss (2002,2003): The universal minimal flow of S_∞ is the space LO of linear orderings of \mathbb{N} . The universal minimal flow of $H(2^{\mathbb{N}})$ is the Uspenskii space of maximal chains of closed subsets of the Cantor space. These all used Ramsey techniques.

Universal minimal flows

If G is compact, then $M(G) = G$. If G is non-compact but locally compact, then $M(G)$ is extremely complicated, e.g., it is non-metrizable. However, by definition G is extremely amenable iff $M(G)$ trivializes!

This leads to a general problem in topological dynamics:

For a given G , can one explicitly determine $M(G)$ and show that it is metrizable?

The first example of calculation of a metrizable but non-trivial universal minimal flow is due to Pestov (1998): The universal minimal flow of $H_+(\mathbb{T})$ is \mathbb{T} . Two more examples were found later by Glasner-Weiss (2002,2003): The universal minimal flow of S_∞ is the space LO of linear orderings of \mathbb{N} . The universal minimal flow of $H(2^{\mathbb{N}})$ is the Uspenskii space of maximal chains of closed subsets of the Cantor space. These all used Ramsey techniques.

Universal minimal flows

If G is compact, then $M(G) = G$. If G is non-compact but locally compact, then $M(G)$ is extremely complicated, e.g., it is non-metrizable. However, by definition G is extremely amenable iff $M(G)$ trivializes!

This leads to a general problem in topological dynamics:

For a given G , can one explicitly determine $M(G)$ and show that it is metrizable?

The first example of calculation of a metrizable but non-trivial universal minimal flow is due to Pestov (1998): The universal minimal flow of $H_+(\mathbb{T})$ is \mathbb{T} . Two more examples were found later by Glasner-Weiss (2002,2003): The universal minimal flow of S_∞ is the space LO of linear orderings of \mathbb{N} . The universal minimal flow of $H(2^{\mathbb{N}})$ is the Uspenskii space of maximal chains of closed subsets of the Cantor space. These all used Ramsey techniques.

Universal minimal flows

If G is compact, then $M(G) = G$. If G is non-compact but locally compact, then $M(G)$ is extremely complicated, e.g., it is non-metrizable. However, by definition G is extremely amenable iff $M(G)$ trivializes!

This leads to a general problem in topological dynamics:

For a given G , can one explicitly determine $M(G)$ and show that it is metrizable?

The first example of calculation of a metrizable but non-trivial universal minimal flow is due to Pestov (1998): The universal minimal flow of $H_+(\mathbb{T})$ is \mathbb{T} . Two more examples were found later by Glasner-Weiss (2002,2003): The universal minimal flow of S_∞ is the space LO of linear orderings of \mathbb{N} . The universal minimal flow of $H(2^{\mathbb{N}})$ is the Uspenskii space of maximal chains of closed subsets of the Cantor space. These all used Ramsey techniques.

Universal minimal flows of automorphism groups

The duality theory of K-Pestov-Todorčević provides tools for computing the universal minimal flows of automorphism groups of Fraïssé structures. We will discuss this next.

Order expansions of Fraïssé classes

Consider a Fraïssé class \mathcal{K} in a language L . Let $L^* = L \cup \{<\}$ be the language obtained by adding a binary relation symbol $<$ to L . A structure \mathbf{A}^* for L^* has the form $\mathbf{A}^* = \langle \mathbf{A}, < \rangle$, where \mathbf{A} is a structure for L and $<$ is a binary relation on A ($=$ the universe of \mathbf{A}). A class \mathcal{K}^* of finite structures on L^* is called an *order class* if ($\langle \mathbf{A}, < \rangle \in \mathcal{K}^* \Rightarrow <$ is a linear ordering on A). For such $\mathbf{A}^* = \langle \mathbf{A}, < \rangle$, let $\mathbf{A}^*|L = \mathbf{A}$.

We say that a Fraïssé order class \mathcal{K}^* on L^* is an *order expansion* of \mathcal{K} if $\mathcal{K} = \mathcal{K}^*|L = \{\mathbf{A}^*|L : \mathbf{A}^* \in \mathcal{K}^*\}$. In this case, if $\mathbf{A} \in \mathcal{K}$ and $\mathbf{A}^* = \langle \mathbf{A}, < \rangle \in \mathcal{K}^*$, we say that $<$ is a \mathcal{K}^* -**admissible ordering** of \mathbf{A} . The order expansion \mathcal{K}^* of \mathcal{K} is *reasonable* if for every $\mathbf{A}, \mathbf{B} \in \mathcal{K}$, with $\mathbf{A} \subseteq \mathbf{B}$ and any \mathcal{K}^* -admissible ordering $<$ on \mathbf{A} , there is a \mathcal{K}^* -admissible ordering $<'$ on \mathbf{B} such that $< \subseteq <'$.

Order expansions of Fraïssé classes

If \mathcal{K} is a Fraïssé class with $\mathbf{K} = \text{Flim}(\mathcal{K})$ and \mathcal{K}^* is a reasonable, order expansion of \mathcal{K} , we denote by $X_{\mathcal{K}^*}$ the space of linear orderings $<$ on K such that for any finite substructure A of \mathbf{K} , $<|_A$ is \mathcal{K}^* -admissible on A . We call these the \mathcal{K}^* -admissible orderings on \mathbf{K} . They form a compact, metrizable, non-empty subspace of 2^{K^2} (with the product topology) on which the group $G = \text{Aut}(\mathbf{K})$ acts continuously, thus $X_{\mathcal{K}^*}$ is a G -flow.

Examples

- \mathcal{K} = finite graphs, $\mathbf{K} = \mathbf{R}$; \mathcal{K}^* = finite ordered graphs. Then $X_{\mathcal{K}^*}$ is the space of all linear orderings of the random graph.
- \mathcal{K} = finite sets, $\mathbf{K} = \langle \mathbb{N} \rangle$; \mathcal{K}^* = finite orderings. Then $X_{\mathcal{K}^*}$ is the space of all linear orderings on \mathbb{N} .
- \mathcal{K} = f.d. vector spaces over a fixed finite field, $\mathbf{K} = \mathbf{V}_\infty$; \mathcal{K}^* = lex. ordered f.d. vector spaces. Then $X_{\mathcal{K}^*}$ is the space of all "lex. orderings" on \mathbf{V}_∞ .
- \mathcal{K} = finite posets, $\mathbf{K} = \mathbf{P}$; \mathcal{K}^* = finite posets with linear extensions. Then $X_{\mathcal{K}^*}$ is the space of all linear extensions of the random poset.

Examples

- \mathcal{K} = finite graphs, $\mathbf{K} = \mathbf{R}$; \mathcal{K}^* = finite ordered graphs. Then $X_{\mathcal{K}^*}$ is the space of all linear orderings of the random graph.
- \mathcal{K} = finite sets, $\mathbf{K} = \langle \mathbb{N} \rangle$; \mathcal{K}^* = finite orderings. Then $X_{\mathcal{K}^*}$ is the space of all linear orderings on \mathbb{N} .
- \mathcal{K} = f.d. vector spaces over a fixed finite field, $\mathbf{K} = V_\infty$; \mathcal{K}^* = lex. ordered f.d. vector spaces. Then $X_{\mathcal{K}^*}$ is the space of all "lex. orderings" on V_∞ .
- \mathcal{K} = finite posets, $\mathbf{K} = P$; \mathcal{K}^* = finite posets with linear extensions. Then $X_{\mathcal{K}^*}$ is the space of all linear extensions of the random poset.

Order expansions of Fraïssé classes

Examples

- \mathcal{K} = finite graphs, $\mathbf{K} = \mathbf{R}$; \mathcal{K}^* = finite ordered graphs. Then $X_{\mathcal{K}^*}$ is the space of all linear orderings of the random graph.
- \mathcal{K} = finite sets, $\mathbf{K} = \langle \mathbb{N} \rangle$; \mathcal{K}^* = finite orderings. Then $X_{\mathcal{K}^*}$ is the space of all linear orderings on \mathbb{N} .
- \mathcal{K} = f.d. vector spaces over a fixed finite field, $\mathbf{K} = V_\infty$; \mathcal{K}^* = lex. ordered f.d. vector spaces. Then $X_{\mathcal{K}^*}$ is the space of all "lex. orderings" on V_∞ .
- \mathcal{K} = finite posets, $\mathbf{K} = P$; \mathcal{K}^* = finite posets with linear extensions. Then $X_{\mathcal{K}^*}$ is the space of all linear extensions of the random poset.

Order expansions of Fraïssé classes

Examples

- \mathcal{K} = finite graphs, $\mathbf{K} = \mathbf{R}$; \mathcal{K}^* = finite ordered graphs. Then $X_{\mathcal{K}^*}$ is the space of all linear orderings of the random graph.
- \mathcal{K} = finite sets, $\mathbf{K} = \langle \mathbb{N} \rangle$; \mathcal{K}^* = finite orderings. Then $X_{\mathcal{K}^*}$ is the space of all linear orderings on \mathbb{N} .
- \mathcal{K} = f.d. vector spaces over a fixed finite field, $\mathbf{K} = \mathbf{V}_\infty$; \mathcal{K}^* = lex. ordered f.d. vector spaces. Then $X_{\mathcal{K}^*}$ is the space of all “lex. orderings” on \mathbf{V}_∞ .
- \mathcal{K} = finite posets, $\mathbf{K} = \mathbf{P}$; \mathcal{K}^* = finite posets with linear extensions. Then $X_{\mathcal{K}^*}$ is the space of all linear extensions of the random poset.

Examples

- \mathcal{K} = finite graphs, $\mathbf{K} = \mathbf{R}$; \mathcal{K}^* = finite ordered graphs. Then $X_{\mathcal{K}^*}$ is the space of all linear orderings of the random graph.
- \mathcal{K} = finite sets, $\mathbf{K} = \langle \mathbb{N} \rangle$; \mathcal{K}^* = finite orderings. Then $X_{\mathcal{K}^*}$ is the space of all linear orderings on \mathbb{N} .
- \mathcal{K} = f.d. vector spaces over a fixed finite field, $\mathbf{K} = \mathbf{V}_\infty$; \mathcal{K}^* = lex. ordered f.d. vector spaces. Then $X_{\mathcal{K}^*}$ is the space of all “lex. orderings” on \mathbf{V}_∞ .
- \mathcal{K} = finite posets, $\mathbf{K} = \mathbf{P}$; \mathcal{K}^* = finite posets with linear extensions. Then $X_{\mathcal{K}^*}$ is the space of all linear extensions of the random poset.

Beyond the Ramsey Property, there is an additional property of classes of finite structures that was introduced by Nešetřil and Rödl in the 1970's and played an important role in the structural Ramsey theory.

Definition

If \mathcal{K}^* is an order expansion of \mathcal{K} , we say that \mathcal{K}^* satisfies the **ordering property (OP)** if for every $A \in \mathcal{K}$, there is $B \in \mathcal{K}$ such that for every \mathcal{K}^* -admissible orderings $<$ on A and $<'$ on B , $\langle A, < \rangle$ can be embedded in $\langle B, <' \rangle$.

In all the examples of the previous page we have the ordering property.

Order expansions of Fraïssé classes

Beyond the Ramsey Property, there is an additional property of classes of finite structures that was introduced by Nešetřil and Rödl in the 1970's and played an important role in the structural Ramsey theory.

Definition

If \mathcal{K}^* is an order expansion of \mathcal{K} , we say that \mathcal{K}^* satisfies the **ordering property (OP)** if for every $A \in \mathcal{K}$, there is $B \in \mathcal{K}$ such that for every \mathcal{K}^* -admissible orderings $<$ on A and $<'$ on B , $\langle A, < \rangle$ can be embedded in $\langle B, <' \rangle$.

In all the examples of the previous page we have the ordering property.

Beyond the Ramsey Property, there is an additional property of classes of finite structures that was introduced by Nešetřil and Rödl in the 1970's and played an important role in the structural Ramsey theory.

Definition

If \mathcal{K}^* is an order expansion of \mathcal{K} , we say that \mathcal{K}^* satisfies the **ordering property (OP)** if for every $A \in \mathcal{K}$, there is $B \in \mathcal{K}$ such that for every \mathcal{K}^* -admissible orderings $<$ on A and $<'$ on B , $\langle A, < \rangle$ can be embedded in $\langle B, <' \rangle$.

In all the examples of the previous page we have the ordering property.

Theorem (KPT)

Let \mathcal{K} be a Fraïssé class and \mathcal{K}^ a reasonable order expansion of \mathcal{K} . Then if G is the automorphism group of the Fraïssé limit of \mathcal{K} the following are equivalent:*

- *$X_{\mathcal{K}^*}$ is the universal minimal flow of the automorphism group of G .*
- *\mathcal{K}^* has the Ramsey Property and the Ordering Property.*

Examples

- \mathcal{K} = finite graphs, $\mathbf{K} = \mathbf{R}$; \mathcal{K}^* = finite ordered graphs. Then the space of all linear orderings of the random graph is the UMF of its automorphism group.
- \mathcal{K} = finite sets, $\mathbf{K} = \langle \mathbb{N} \rangle$; \mathcal{K}^* = finite orderings. Then the space of all linear orderings on \mathbb{N} is the UMF of S_∞ (Glasner-Weiss).
- \mathcal{K} = f.d. vector spaces over a fixed finite field, $\mathbf{K} = \mathbf{V}_\infty$; \mathcal{K}^* = lex. ordered f.d. vector spaces. Then the space of all “lex. orderings” on \mathbf{V}_∞ is the UMF of its general linear group.
- \mathcal{K} = finite posets, $\mathbf{K} = \mathbf{P}$; \mathcal{K}^* = finite posets with linear extensions. Then the space of all linear extensions of the random poset is the UMF of its automorphism group.

Examples

- \mathcal{K} = finite graphs, $\mathbf{K} = \mathbf{R}$; \mathcal{K}^* = finite ordered graphs. Then the space of all linear orderings of the random graph is the UMF of its automorphism group.
- \mathcal{K} = finite sets, $\mathbf{K} = \langle \mathbb{N} \rangle$; \mathcal{K}^* = finite orderings. Then the space of all linear orderings on \mathbb{N} is the UMF of S_∞ (Glasner-Weiss).
- \mathcal{K} = f.d. vector spaces over a fixed finite field, $\mathbf{K} = V_\infty$; \mathcal{K}^* = lex. ordered f.d. vector spaces. Then the space of all “lex. orderings” on V_∞ is the UMF of its general linear group.
- \mathcal{K} = finite posets, $\mathbf{K} = \mathbf{P}$; \mathcal{K}^* = finite posets with linear extensions. Then the space of all linear extensions of the random poset is the UMF of its automorphism group.

Calculation of universal minimal flows

Examples

- \mathcal{K} = finite graphs, $\mathbf{K} = \mathbf{R}$; \mathcal{K}^* = finite ordered graphs. Then the space of all linear orderings of the random graph is the UMF of its automorphism group.
- \mathcal{K} = finite sets, $\mathbf{K} = \langle \mathbb{N} \rangle$; \mathcal{K}^* = finite orderings. Then the space of all linear orderings on \mathbb{N} is the UMF of S_∞ (Glasner-Weiss).
- \mathcal{K} = f.d. vector spaces over a fixed finite field, $\mathbf{K} = \mathbf{V}_\infty$; \mathcal{K}^* = lex. ordered f.d. vector spaces. Then the space of all “lex. orderings” on \mathbf{V}_∞ is the UMF of its general linear group.
- \mathcal{K} = finite posets, $\mathbf{K} = \mathbf{P}$; \mathcal{K}^* = finite posets with linear extensions. Then the space of all linear extensions of the random poset is the UMF of its automorphism group.

Calculation of universal minimal flows

Examples

- \mathcal{K} = finite graphs, $\mathbf{K} = \mathbf{R}$; \mathcal{K}^* = finite ordered graphs. Then the space of all linear orderings of the random graph is the UMF of its automorphism group.
- \mathcal{K} = finite sets, $\mathbf{K} = \langle \mathbb{N} \rangle$; \mathcal{K}^* = finite orderings. Then the space of all linear orderings on \mathbb{N} is the UMF of S_∞ (Glasner-Weiss).
- \mathcal{K} = f.d. vector spaces over a fixed finite field, $\mathbf{K} = \mathbf{V}_\infty$; \mathcal{K}^* = lex. ordered f.d. vector spaces. Then the space of all “lex. orderings” on \mathbf{V}_∞ is the UMF of its general linear group.
- \mathcal{K} = finite posets, $\mathbf{K} = \mathbf{P}$; \mathcal{K}^* = finite posets with linear extensions. Then the space of all linear extensions of the random poset is the UMF of its automorphism group.

Unique ergodicity revisited

Let \mathcal{K} be a Fraïssé class and \mathcal{K}^* a reasonable order expansion of \mathcal{K} that has the Ramsey Property and the Ordering Property. We will say then that \mathcal{K}^* is a **companion** of \mathcal{K} . It was shown in the paper of KPT that such a companion, when it exists, is essentially unique.

Thus we have seen that when \mathcal{K} has a companion class \mathcal{K}^* , and this happens for many important examples, then the UMF of the automorphism group G of its Fraïssé limit is the compact, metrizable space $X_{\mathcal{K}^*}$. Thus the unique ergodicity of G is equivalent to the unique ergodicity of $X_{\mathcal{K}^*}$. This can then be seen to be equivalent to the following probabilistic notion.

Unique ergodicity revisited

Let \mathcal{K} be a Fraïssé class and \mathcal{K}^* a reasonable order expansion of \mathcal{K} that has the Ramsey Property and the Ordering Property. We will say then that \mathcal{K}^* is a **companion** of \mathcal{K} . It was shown in the paper of KPT that such a companion, when it exists, is essentially unique.

Thus we have seen that when \mathcal{K} has a companion class \mathcal{K}^* , and this happens for many important examples, then the UMF of the automorphism group G of its Fraïssé limit is the compact, metrizable space $X_{\mathcal{K}^*}$. Thus the unique ergodicity of G is equivalent to the unique ergodicity of $X_{\mathcal{K}^*}$. This can then be seen to be equivalent to the following probabilistic notion.

Unique ergodicity revisited

Definition

Let \mathcal{K}^* be a companion of \mathcal{K} . A **random, consistent** \mathcal{K}^* -admissible ordering is a map that assigns to each structure $\mathbf{A} \in \mathcal{K}$ a probability measure $\mu_{\mathbf{A}}$ on the (finite) space of \mathcal{K}^* -admissible orderings on \mathbf{A} , which is isomorphism invariant and has the property that if $\mathbf{A} \subseteq \mathbf{B}$, then $\mu_{\mathbf{B}}$ projects by the restriction map to $\mu_{\mathbf{A}}$.

Example: graphs

We now have:

Proposition (AKL)

Let \mathcal{K}^ be a companion of \mathcal{K} . Then amenability of the automorphism group G of the Fraïssé limit of \mathcal{K} is equivalent to the existence of a random, consistent \mathcal{K}^* -admissible ordering and unique ergodicity of G is equivalent to the uniqueness of a random, consistent \mathcal{K}^* -admissible ordering.*

Unique ergodicity revisited

Definition

Let \mathcal{K}^* be a companion of \mathcal{K} . A **random, consistent** \mathcal{K}^* -admissible ordering is a map that assigns to each structure $\mathbf{A} \in \mathcal{K}$ a probability measure $\mu_{\mathbf{A}}$ on the (finite) space of \mathcal{K}^* -admissible orderings on \mathbf{A} , which is isomorphism invariant and has the property that if $\mathbf{A} \subseteq \mathbf{B}$, then $\mu_{\mathbf{B}}$ projects by the restriction map to $\mu_{\mathbf{A}}$.

Example: graphs

We now have:

Proposition (AKL)

Let \mathcal{K}^ be a companion of \mathcal{K} . Then amenability of the automorphism group G of the Fraïssé limit of \mathcal{K} is equivalent to the existence of a random, consistent \mathcal{K}^* -admissible ordering and unique ergodicity of G is equivalent to the uniqueness of a random, consistent \mathcal{K}^* -admissible ordering.*

Unique ergodicity revisited

Definition

Let \mathcal{K}^* be a companion of \mathcal{K} . A **random, consistent** \mathcal{K}^* -admissible ordering is a map that assigns to each structure $\mathbf{A} \in \mathcal{K}$ a probability measure $\mu_{\mathbf{A}}$ on the (finite) space of \mathcal{K}^* -admissible orderings on \mathbf{A} , which is isomorphism invariant and has the property that if $\mathbf{A} \subseteq \mathbf{B}$, then $\mu_{\mathbf{B}}$ projects by the restriction map to $\mu_{\mathbf{A}}$.

Example: graphs

We now have:

Proposition (AKL)

Let \mathcal{K}^ be a companion of \mathcal{K} . Then amenability of the automorphism group G of the Fraïssé limit of \mathcal{K} is equivalent to the existence of a random, consistent \mathcal{K}^* -admissible ordering and unique ergodicity of G is equivalent to the uniqueness of a random, consistent \mathcal{K}^* -admissible ordering.*

Unique ergodicity as a quantitative version of the Ordering Property

Interestingly it turns out that unique ergodicity fits well in the framework of the duality theory of KPT (which originally was developed in the context of topological dynamics). In many cases it can simply be viewed as a quantitative version of the Ordering Property.

Definition (AKL)

Let \mathcal{K}^* be a companion of \mathcal{K} . We say that \mathcal{K}^* satisfies the **Quantitative Ordering Property (QOP)** if the following holds:

There is an isomorphism invariant map that assigns to each structure $A^* = \langle A, \langle \rangle \rangle \in \mathcal{K}^*$ a real number $\rho(A^*)$ in $(0, 1]$ such that for every $A \in \mathcal{K}$ and each $\epsilon > 0$, there is a $B \in \mathcal{K}$ and a nonempty set of embeddings $E(A, B)$ of A into B with the property that for each \mathcal{K}^* -admissible ordering \langle of A and each \mathcal{K}^* -admissible ordering \langle' of B the proportion of embeddings in $E(A, B)$ that preserve \langle, \langle' is equal to $\rho(\langle A, \langle \rangle)$, within ϵ .

Unique ergodicity as a quantitative version of the Ordering Property

Interestingly it turns out that unique ergodicity fits well in the framework of the duality theory of KPT (which originally was developed in the context of topological dynamics). In many cases it can simply be viewed as a quantitative version of the Ordering Property.

Definition (AKL)

Let \mathcal{K}^* be a companion of \mathcal{K} . We say that \mathcal{K}^* satisfies the **Quantitative Ordering Property (QOP)** if the following holds:

There is an isomorphism invariant map that assigns to each structure $\mathbf{A}^* = \langle \mathbf{A}, \langle \rangle \rangle \in \mathcal{K}^*$ a real number $\rho(\mathbf{A}^*)$ in $(0, 1]$ such that for every $\mathbf{A} \in \mathcal{K}$ and each $\epsilon > 0$, there is a $\mathbf{B} \in \mathcal{K}$ and a nonempty set of embeddings $E(\mathbf{A}, \mathbf{B})$ of \mathbf{A} into \mathbf{B} with the property that for each \mathcal{K}^* -admissible ordering $\langle \rangle$ of \mathbf{A} and each \mathcal{K}^* -admissible ordering $\langle' \rangle$ of \mathbf{B} the proportion of embeddings in $E(\mathbf{A}, \mathbf{B})$ that preserve $\langle \rangle, \langle' \rangle$ is equal to $\rho(\langle \mathbf{A}, \langle \rangle \rangle)$, within ϵ .

Unique ergodicity as a quantitative version of the Ordering Property

For example, if \mathcal{K} is the class of finite graphs, one can establish QOP by showing that for any finite graph \mathbf{A} with n vertices and $\epsilon > 0$, there is a graph \mathbf{B} , containing a copy of \mathbf{A} , such that given any orderings $<$ on \mathbf{A} and $<'$ on \mathbf{B} , the proportion of all embeddings of \mathbf{A} into \mathbf{B} that preserve the orderings $<, <'$ is, up to ϵ , equal to $1/n!$.

Unique ergodicity as a quantitative version of the Ordering Property

Theorem (AKL)

Let \mathcal{K}^ be a companion of \mathcal{K} , let G be the automorphism group of the Fraïssé limit of \mathcal{K} and assume that G is amenable. Then QOP implies the unique ergodicity of G . Moreover, if \mathcal{K} is a Hrushovski class, QOP is equivalent to the unique ergodicity of G .*

Proving unique ergodicity

By more direct means (but still using the calculation of the UMF), one can show that the following automorphism groups are uniquely ergodic:

- S_∞ (Glasner-Weiss)
- The isometry group of the Baire space and various ultrametric Urysohn spaces (AKL)
- The general linear group of the (countably) infinite-dimensional vector space over a finite field (AKL)

By applying now the preceding QOP criterion and probabilistic arguments (deviation inequalities) one can now also show the following:

Theorem (AKL)

The automorphism groups of the following structures are uniquely ergodic

- *The random graph*
- *The random K_n -free graph*
- *The random n -uniform hypergraph*
- *The rational Urysohn space*

Proving unique ergodicity

By more direct means (but still using the calculation of the UMF), one can show that the following automorphism groups are uniquely ergodic:

- S_∞ (Glasner-Weiss)
- The isometry group of the Baire space and various ultrametric Urysohn spaces (AKL)
- The general linear group of the (countably) infinite-dimensional vector space over a finite field (AKL)

By applying now the preceding QOP criterion and probabilistic arguments (deviation inequalities) one can now also show the following:

Theorem (AKL)

The automorphism groups of the following structures are uniquely ergodic

- *The random graph*
- *The random K_n -free graph*
- *The random n -uniform hypergraph*
- *The rational Urysohn space*

Proving unique ergodicity

By more direct means (but still using the calculation of the UMF), one can show that the following automorphism groups are uniquely ergodic:

- S_∞ (Glasner-Weiss)
- The isometry group of the Baire space and various ultrametric Urysohn spaces (AKL)
- The general linear group of the (countably) infinite-dimensional vector space over a finite field (AKL)

By applying now the preceding QOP criterion and probabilistic arguments (deviation inequalities) one can now also show the following:

Theorem (AKL)

The automorphism groups of the following structures are uniquely ergodic

- *The random graph*
- *The random K_n -free graph*
- *The random n -uniform hypergraph*
- *The rational Urysohn space*

Proving unique ergodicity

By more direct means (but still using the calculation of the UMF), one can show that the following automorphism groups are uniquely ergodic:

- S_∞ (Glasner-Weiss)
- The isometry group of the Baire space and various ultrametric Urysohn spaces (AKL)
- The general linear group of the (countably) infinite-dimensional vector space over a finite field (AKL)

By applying now the preceding QOP criterion and probabilistic arguments (deviation inequalities) one can now also show the following:

Theorem (AKL)

The automorphism groups of the following structures are uniquely ergodic

- *The random graph*
- *The random K_n -free graph*
- *The random n -uniform hypergraph*
- *The rational Urysohn space*

Proving unique ergodicity

By more direct means (but still using the calculation of the UMF), one can show that the following automorphism groups are uniquely ergodic:

- S_∞ (Glasner-Weiss)
- The isometry group of the Baire space and various ultrametric Urysohn spaces (AKL)
- The general linear group of the (countably) infinite-dimensional vector space over a finite field (AKL)

By applying now the preceding QOP criterion and probabilistic arguments (deviation inequalities) one can now also show the following:

Theorem (AKL)

The automorphism groups of the following structures are uniquely ergodic

- *The random graph*
- *The random K_n -free graph*
- *The random n -uniform hypergraph*
- *The rational Urysohn space*

Proving unique ergodicity

By more direct means (but still using the calculation of the UMF), one can show that the following automorphism groups are uniquely ergodic:

- S_∞ (Glasner-Weiss)
- The isometry group of the Baire space and various ultrametric Urysohn spaces (AKL)
- The general linear group of the (countably) infinite-dimensional vector space over a finite field (AKL)

By applying now the preceding QOP criterion and probabilistic arguments (deviation inequalities) one can now also show the following:

Theorem (AKL)

The automorphism groups of the following structures are uniquely ergodic

- *The random graph*
- *The random K_n -free graph*
- *The random n -uniform hypergraph*
- *The rational Urysohn space*

Unique Ergodicity Problem

In fact I do not know any counterexample to the following problem:

Problem (Unique Ergodicity Problem)

Let G be an automorphism group of a countable structure with a metrizable universal minimal flow. If G is amenable, then is it uniquely ergodic?

In fact one can even consider an even stronger form of this problem:

Problem (Unique Ergodicity Problem-Strong Form)

Let G be an automorphism group of a countable structure with a metrizable universal minimal flow. If G is amenable, then is it uniquely ergodic and in every minimal G -flow the unique invariant measure is supported by a single comeager orbit?

A positive answer to this stronger form has been obtained in many cases, e.g., S_∞ (Glasner-Weiss); the automorphism group of the random graph, random n -uniform hypergraph, random K_n -free graph, rational Urysohn space, etc. (AKL)

Unique Ergodicity Problem

In fact I do not know any counterexample to the following problem:

Problem (Unique Ergodicity Problem)

Let G be an automorphism group of a countable structure with a metrizable universal minimal flow. If G is amenable, then is it uniquely ergodic?

In fact one can even consider an even stronger form of this problem:

Problem (Unique Ergodicity Problem-Strong Form)

Let G be an automorphism group of a countable structure with a metrizable universal minimal flow. If G is amenable, then is it uniquely ergodic and in every minimal G -flow the unique invariant measure is supported by a single comeager orbit?

A positive answer to this stronger form has been obtained in many cases, e.g., S_∞ (Glasner-Weiss); the automorphism group of the random graph, random n -uniform hypergraph, random K_n -free graph, rational Urysohn space, etc. (AKL)

Unique Ergodicity Problem

In fact I do not know any counterexample to the following problem:

Problem (Unique Ergodicity Problem)

Let G be an automorphism group of a countable structure with a metrizable universal minimal flow. If G is amenable, then is it uniquely ergodic?

In fact one can even consider an even stronger form of this problem:

Problem (Unique Ergodicity Problem-Strong Form)

Let G be an automorphism group of a countable structure with a metrizable universal minimal flow. If G is amenable, then is it uniquely ergodic and in every minimal G -flow the unique invariant measure is supported by a single comeager orbit?

A positive answer to this stronger form has been obtained in many cases, e.g., S_∞ (Glasner-Weiss); the automorphism group of the random graph, random n -uniform hypergraph, random K_n -free graph, rational Urysohn space, etc. (AKL)

Unique Ergodicity Problem

In fact I do not know any counterexample to the following problem:

Problem (Unique Ergodicity Problem)

Let G be an automorphism group of a countable structure with a metrizable universal minimal flow. If G is amenable, then is it uniquely ergodic?

In fact one can even consider an even stronger form of this problem:

Problem (Unique Ergodicity Problem-Strong Form)

Let G be an automorphism group of a countable structure with a metrizable universal minimal flow. If G is amenable, then is it uniquely ergodic and in every minimal G -flow the unique invariant measure is supported by a single comeager orbit?

A positive answer to this stronger form has been obtained in many cases, e.g., S_∞ (Glasner-Weiss); the automorphism group of the random graph, random n -uniform hypergraph, random K_n -free graph, rational Urysohn space, etc. (AKL)

Unique Ergodicity Problem

In fact I do not know any counterexample to the following problem:

Problem (Unique Ergodicity Problem)

Let G be an automorphism group of a countable structure with a metrizable universal minimal flow. If G is amenable, then is it uniquely ergodic?

In fact one can even consider an even stronger form of this problem:

Problem (Unique Ergodicity Problem-Strong Form)

Let G be an automorphism group of a countable structure with a metrizable universal minimal flow. If G is amenable, then is it uniquely ergodic and in every minimal G -flow the unique invariant measure is supported by a single comeager orbit?

A positive answer to this stronger form has been obtained in many cases, e.g., S_∞ (Glasner-Weiss); the automorphism group of the random graph, random n -uniform hypergraph, random K_n -free graph, rational Urysohn space, etc. (AKL)