

Polish G -spaces similar to logic G -spaces of continuous structures

Aleksander Ivanov and Barbara Majcher-Iwanow

Institute of Mathematics
University of Wrocław

July 23, 2012

Logic S_∞ -space

Let $L = (R_i^{n_i})_{i \in I}$ be a countable relational language and

$$X_L = \prod_{i \in I} 2^{\omega^{n_i}}$$

be the corresponding topol. space under the product topology τ .

We view X_L as the space of all L -structures on ω identifying every $\mathbf{x} = (\dots x_i \dots) \in X_L$ with the structure $(\omega, R_i)_{i \in I}$ where R_i is the n_i -ary relatin defined by the charactrstc functn $x_i : \omega^{n_i} \rightarrow 2$.

The **logic action** of the group S_∞ of all permutations of ω is defined on X_L by the rule:

$$g \circ \mathbf{x} = \mathbf{y} \Leftrightarrow \forall i \forall \bar{s} (y_i(\bar{s}) = x_i(g^{-1}(\bar{s})).$$

Logic S_∞ -space

Let $L = (R_i^{n_i})_{i \in I}$ be a countable relational language and

$$X_L = \prod_{i \in I} 2^{\omega^{n_i}}$$

be the corresponding topol. space under the product topology τ .

We view X_L as the space of all L -structures on ω identifying every $\mathbf{x} = (\dots x_i \dots) \in X_L$ with the structure $(\omega, R_i)_{i \in I}$ where R_i is the n_i -ary relatin defined by the charactrstc functn $x_i : \omega^{n_i} \rightarrow 2$.

The **logic action** of the group S_∞ of all permutations of ω is defined on X_L by the rule:

$$g \circ \mathbf{x} = \mathbf{y} \Leftrightarrow \forall i \forall \bar{s} (y_i(\bar{s}) = x_i(g^{-1}(\bar{s})).$$

Logic S_∞ -space

Let $L = (R_i^{n_i})_{i \in I}$ be a countable relational language and

$$X_L = \prod_{i \in I} 2^{\omega^{n_i}}$$

be the corresponding topol. space under the product topology τ .

We view X_L as the space of all L -structures on ω identifying every $\mathbf{x} = (\dots x_i \dots) \in X_L$ with the structure $(\omega, R_i)_{i \in I}$ where R_i is the n_i -ary relatin defined by the charactrstc functn $x_i : \omega^{n_i} \rightarrow 2$.

The **logic action** of the group S_∞ of all permutations of ω is defined on X_L by the rule:

$$g \circ \mathbf{x} = \mathbf{y} \Leftrightarrow \forall i \forall \bar{s} (y_i(\bar{s}) = x_i(g^{-1}(\bar{s})).$$

Canonical partition

Let $(\langle X, \tau \rangle, G)$ be a Polish G -space with a countable basis $\{C_j\}$.

H.Becker: there exists a unique partition of X ,

$$X = \bigcup \{Y_t : t \in T\}$$

into invariant G_δ -sets Y_t s. t. every orbit from Y_t is dense in Y_t .

To construct it take $\{C_j\}$ and for any $t \in 2^{\mathbb{N}}$ define

$$Y_t = \left(\bigcap \{GC_j : t(j) = 1\} \right) \cap \left(\bigcap \{X \setminus GC_j : t(j) = 0\} \right)$$

and take $T = \{t \in 2^{\mathbb{N}} : Y_t \neq \emptyset\}$.

In the case of the logic action of S_∞ on the space X_L each piece consists of structures which satisfy the same \forall -sentences and \exists -sentences.

Canonical partition

Let $(\langle X, \tau \rangle, G)$ be a Polish G -space with a countable basis $\{C_j\}$.

H.Becker: there exists a unique partition of X ,

$$X = \bigcup \{Y_t : t \in T\}$$

into invariant G_δ -sets Y_t s. t. every orbit from Y_t is dense in Y_t .

To construct it take $\{C_j\}$ and for any $t \in 2^{\mathbb{N}}$ define

$$Y_t = \left(\bigcap \{GC_j : t(j) = 1\} \right) \cap \left(\bigcap \{X \setminus GC_j : t(j) = 0\} \right)$$

and take $T = \{t \in 2^{\mathbb{N}} : Y_t \neq \emptyset\}$.

In the case of the logic action of S_∞ on the space X_L each piece consists of structures which satisfy the same \forall -sentences and \exists -sentences.

Canonical partition

Let $(\langle X, \tau \rangle, G)$ be a Polish G -space with a countable basis $\{C_j\}$.

H.Becker: there exists a unique partition of X ,

$$X = \bigcup \{Y_t : t \in T\}$$

into invariant G_δ -sets Y_t s. t. every orbit from Y_t is dense in Y_t .

To construct it take $\{C_j\}$ and for any $t \in 2^{\mathbb{N}}$ define

$$Y_t = \left(\bigcap \{GC_j : t(j) = 1\} \right) \cap \left(\bigcap \{X \setminus GC_j : t(j) = 0\} \right)$$

and take $T = \{t \in 2^{\mathbb{N}} : Y_t \neq \emptyset\}$.

In the case of the logic action of S_∞ on the space X_L each piece consists of structures which satisfy the same \forall -sentences and \exists -sentences.

Canonical partition

Let $(\langle X, \tau \rangle, G)$ be a Polish G -space with a countable basis $\{C_j\}$.

H.Becker: there exists a unique partition of X ,

$$X = \bigcup \{Y_t : t \in T\}$$

into invariant G_δ -sets Y_t s. t. every orbit from Y_t is dense in Y_t .

To construct it take $\{C_j\}$ and for any $t \in 2^{\mathbb{N}}$ define

$$Y_t = \left(\bigcap \{GC_j : t(j) = 1\} \right) \cap \left(\bigcap \{X \setminus GC_j : t(j) = 0\} \right)$$

and take $T = \{t \in 2^{\mathbb{N}} : Y_t \neq \emptyset\}$.

In the case of the logic action of S_∞ on the space X_L each piece consists of structures which satisfy the same \forall -sentences and \exists -sentences.

Vaught transforms

Let X be a Polish G -space, $B \subset X$ and $u \subset G$ is open.

Vaught transforms:

$$B^{*u} = \{x \in X : \{g \in u : gx \in B\} \text{ is comeagre in } u\}$$

$$B^{\Delta u} = \{x \in X : \{g \in u : gx \in B\} \text{ is not meagre in } u\}.$$

In the case of the **logic action** of S_∞ on the space X_L if

$$B = \{M \in X_L : M \models \phi(s)\} \text{ with } s \in \omega$$

then

$$B^{*S_\infty} = \{M \in X_L : M \models \forall x \phi(x)\}.$$

Vaught transforms

Let X be a Polish G -space, $B \subset X$ and $u \subset G$ is open.

Vaught transforms:

$$B^{*u} = \{x \in X : \{g \in u : gx \in B\} \text{ is comeagre in } u\}$$

$$B^{\Delta u} = \{x \in X : \{g \in u : gx \in B\} \text{ is not meagre in } u\}.$$

In the case of the **logic action** of S_∞ on the space X_L if

$$B = \{M \in X_L : M \models \phi(s)\} \text{ with } s \in \omega$$

then

$$B^{*S_\infty} = \{M \in X_L : M \models \forall x \phi(x)\}.$$

Vaught transforms

Let X be a Polish G -space, $B \subset X$ and $u \subset G$ is open.

Vaught transforms:

$$B^{*u} = \{x \in X : \{g \in u : gx \in B\} \text{ is comeagre in } u\}$$

$$B^{\Delta u} = \{x \in X : \{g \in u : gx \in B\} \text{ is not meagre in } u\}.$$

In the case of the **logic action** of S_∞ on the space X_L if

$$B = \{M \in X_L : M \models \phi(s)\} \text{ with } s \in \omega$$

then

$$B^{*S_\infty} = \{M \in X_L : M \models \forall x \phi(x)\}.$$

Vaught transforms 2

- If $B \in \Sigma_\alpha$, then $B^{\Delta H} \in \Sigma_\alpha$ and
if $B \in \Pi_\alpha$, then $B^{*H} \in \Pi_\alpha$.
- For any open $B \subseteq X$ and any open $K < G$ we have
 $B^{\Delta K} = KB$, where $KB = \{gx : g \in K, x \in B\}$.

Vaught transforms 2

- If $B \in \Sigma_\alpha$, then $B^{\Delta H} \in \Sigma_\alpha$ and
if $B \in \Pi_\alpha$, then $B^{*H} \in \Pi_\alpha$.
- For any open $B \subseteq X$ and any open $K < G$ we have
 $B^{\Delta K} = KB$, where $KB = \{gx : g \in K, x \in B\}$.

Actions of closed subgroups of S_∞

Let G be a closed subgroup of S_∞ .

Let \mathcal{N}^G be the standard basis of the topology of G consisting of cosets of pointwise stabilisers of finite subsets of ω .

Let $(\langle X, \tau \rangle, G)$ be a Polish G -space with a countable basis \mathcal{A} .
Along with τ we shall consider another topology on X .

Nice topology:

Actions of closed subgroups of S_∞

Let G be a closed subgroup of S_∞ .

Let \mathcal{N}^G be the standard basis of the topology of G consisting of cosets of pointwise stabilisers of finite subsets of ω .

Let $(\langle X, \tau \rangle, G)$ be a Polish G -space with a countable basis \mathcal{A} .
Along with τ we shall consider another topology on X .

Nice topology:

Actions of closed subgroups of S_∞

Let G be a closed subgroup of S_∞ .

Let \mathcal{N}^G be the standard basis of the topology of G consisting of cosets of pointwise stabilisers of finite subsets of ω .

Let $(\langle X, \tau \rangle, G)$ be a Polish G -space with a countable basis \mathcal{A} .
Along with τ we shall consider another topology on X .

Nice topology:

Nice topology

Definition (H.Becker) A topology \mathbf{t} on X is **nice** for the G -space $(\langle X, \tau \rangle, G)$ if the following conditions are satisfied.

(A) \mathbf{t} is a Polish topology, \mathbf{t} is finer than τ and the G -action remains continuous with respect to \mathbf{t} .

(B) There exists a basis \mathcal{B} for \mathbf{t} (called **nice**) such that:

- 1 \mathcal{B} is countable;
- 2 for all $B_1, B_2 \in \mathcal{B}$, $B_1 \cap B_2 \in \mathcal{B}$;
- 3 for all $B \in \mathcal{B}$, $X \setminus B \in \mathcal{B}$;
- 4 for all $B \in \mathcal{B}$ and $u \in \mathcal{N}^G$, $B^{\Delta u}, B^{*u} \in \mathcal{B}$;
- 5 for any $B \in \mathcal{B}$ there exists an open subgroup $H < G$ such that B is invariant under the corresponding H -action.

Example

Logic action

For any countable fragment F of $L_{\omega_1\omega}$, which is closed under quantifiers, all sets

$$\text{Mod}(\phi, \bar{s}) = \{M \in X_L : M \models \phi(\bar{s})\} \text{ with } \bar{s} \subset \omega$$

form a nice basis defining a nice topology (denoted by \mathbf{t}_F) of the S_∞ -space X_L .

Each piece of the canonical partition corresponding to \mathbf{t}_F consists of structures which satisfy the same F -sentences (without parameters).

Example

Logic action

For any countable fragment F of $L_{\omega_1\omega}$, which is closed under quantifiers, all sets

$$\text{Mod}(\phi, \bar{s}) = \{M \in X_L : M \models \phi(\bar{s})\} \text{ with } \bar{s} \subset \omega$$

form a nice basis defining a nice topology (denoted by \mathbf{t}_F) of the S_∞ -space X_L .

Each piece of the canonical partition corresponding to \mathbf{t}_F consists of structures which satisfy the same F -sentences (without parameters).

Example and illustration

Let G be a closed subgroup of S_∞ and (X, τ) be a Polish G -space. Let \mathbf{t} be a nice topology for $(\langle X, \tau \rangle, G)$.

A generalized version of Lindström's model completeness theorem:

Theorem (B.M-I)

For any $x_1 \in X$ if $X_1 = Gx_1$ is a G_δ -subset of X , then both topologies τ and \mathbf{t} coincide on X_1 .

H.Becker: **J.Amer.Math.Soc**, 11(1998), 397 - 449 and **APAL**, 111(2001), 145 - 184

Example and illustration

Let G be a closed subgroup of S_∞ and (X, τ) be a Polish G -space. Let \mathbf{t} be a nice topology for $(\langle X, \tau \rangle, G)$.

A generalized version of Lindström's model completeness theorem:

Theorem (B.M-I)

For any $x_1 \in X$ if $X_1 = Gx_1$ is a G_δ -subset of X , then both topologies τ and \mathbf{t} coincide on X_1 .

H.Becker: **J.Amer.Math.Soc**, 11(1998), 397 - 449 and **APAL**, 111(2001), 145 - 184

Example and illustration

Let G be a closed subgroup of S_∞ and (X, τ) be a Polish G -space. Let \mathbf{t} be a nice topology for $(\langle X, \tau \rangle, G)$.

A generalized version of Lindström's model completeness theorem:

Theorem (B.M-I)

For any $x_1 \in X$ if $X_1 = Gx_1$ is a G_δ -subset of X , then both topologies τ and \mathbf{t} coincide on X_1 .

H.Becker: **J.Amer.Math.Soc**, 11(1998), 397 - 449 and **APAL**, 111(2001), 145 - 184

Existence

Theorem

(H.Becker) Let G be a closed subgroup of S_∞ and (X, τ) be a Polish G -space.

Let \mathbf{t}' be a topology on X finer than τ , such that the action remains continuous with respect to \mathbf{t}' .

Then there is a nice topology \mathbf{t} for $(\langle X, \tau \rangle, G)$ such that \mathbf{t} is finer than \mathbf{t}' .

Remark: All elements of \mathbf{t} are τ -Borel.

Question

Question:

Is it possible to extend the generalised model theory of H.Becker to actions of Polish groups (without the assumption $G \leq S_\infty$) ?

Continuous structures

A countable continuous signature:

$$L = \{d, R_1, \dots, R_k, \dots, F_1, \dots, F_l, \dots\}.$$

Definition

A **metric L -structure** is a complete metric space (M, d) with d bounded by 1, along with a family of uniformly continuous operations F_k on M and a family of predicates R_l , i.e. uniformly continuous maps from appropriate M^{k_l} to $[0, 1]$.

It is usually assumed that to a predicate symbol R_i a continuity modulus γ_i is assigned so that when $d(x_j, x'_j) < \gamma_i(\varepsilon)$ with $1 \leq j \leq k_i$ the corresponding predicate of M satisfies

$$|R_i(x_1, \dots, x_j, \dots, x_{k_i}) - R_i(x_1, \dots, x'_j, \dots, x_{k_i})| < \varepsilon.$$

Continuous structures

A countable continuous signature:

$$L = \{d, R_1, \dots, R_k, \dots, F_1, \dots, F_l, \dots\}.$$

Definition

A **metric L -structure** is a complete metric space (M, d) with d bounded by 1, along with a family of uniformly continuous operations F_k on M and a family of predicates R_l , i.e. uniformly continuous maps from appropriate M^{k_l} to $[0, 1]$.

It is usually assumed that to a predicate symbol R_i a continuity modulus γ_i is assigned so that when $d(x_j, x'_j) < \gamma_i(\varepsilon)$ with $1 \leq j \leq k_i$ the corresponding predicate of M satisfies

$$|R_i(x_1, \dots, x_j, \dots, x_{k_i}) - R_i(x_1, \dots, x'_j, \dots, x_{k_i})| < \varepsilon.$$

Canonical structure

Let (G, d) be a Polish group with a left invariant metric ≤ 1 .

If (\mathbf{X}, d) is its completion, then $G \leq Iso(\mathbf{X})$.

Let S be a countable dense subset of \mathbf{X} .

Enumerate all orbits of G of finite tuples of S .

For the closure of such an n -orbit C define a predicate $R_{\bar{C}}$ on (\mathbf{X}, d) by

$$R_{\bar{C}}(y_1, \dots, y_n) = d((y_1, \dots, y_n), \bar{C}) \text{ (i.e. } \inf \{d(\bar{y}, \bar{c}) : \bar{c} \in \bar{C}\} \text{)}.$$

It is observed by J.Melleray that G is the automorphism group of the continuous structure M of all these predicates on \mathbf{X} , with continuous moduli = id .

Canonical structure

Let (G, d) be a Polish group with a left invariant metric ≤ 1 .
If (\mathbf{X}, d) is its completion, then $G \leq Iso(\mathbf{X})$.

Let S be a countable dense subset of \mathbf{X} .

Enumerate all orbits of G of finite tuples of S .

For the closure of such an n -orbit C define a predicate $R_{\bar{C}}$ on (\mathbf{X}, d) by

$$R_{\bar{C}}(y_1, \dots, y_n) = d((y_1, \dots, y_n), \bar{C}) \text{ (i.e. } \inf \{d(\bar{y}, \bar{c}) : \bar{c} \in \bar{C}\} \text{)}.$$

It is observed by J.Melleray that G is the automorphism group of the continuous structure M of all these predicates on \mathbf{X} , with continuous moduli = id .

The space of continuous structures

Fix a relational continuous signature L and a Polish space (\mathbf{Y}, d) .
Let S be a dense countable subset of \mathbf{Y} .

Define the space \mathbf{Y}_L of continuous L -structures on (\mathbf{Y}, d) as follows.

Metric on the set of L -structures: Enumerate all tuples of the form (j, \bar{s}) , where \bar{s} is a tuple $\in S$ of the length of the arity of R_j .

For L -structures M and N on \mathbf{Y} let

$$\delta(M, N) = \sum_{i=1}^{\infty} \{2^{-i} |R_j^M(\bar{s}) - R_j^N(\bar{s})| : i \text{ is the number of } (j, \bar{s})\}.$$

Logic action

- the space \mathbf{Y}_L is Polish;
- the Polish group $\text{Iso}(\mathbf{Y})$ acts on \mathbf{Y}_L continuously

The space of continuous structures

Fix a relational continuous signature L and a Polish space (\mathbf{Y}, d) .
Let S be a dense countable subset of \mathbf{Y} .

Define the space \mathbf{Y}_L of continuous L -structures on (\mathbf{Y}, d) as follows.

Metric on the set of L -structures: Enumerate all tuples of the form (j, \bar{s}) , where \bar{s} is a tuple $\in S$ of the length of the arity of R_j .

For L -structures M and N on \mathbf{Y} let

$$\delta(M, N) = \sum_{i=1}^{\infty} \{2^{-i} |R_j^M(\bar{s}) - R_j^N(\bar{s})| : i \text{ is the number of } (j, \bar{s})\}.$$

Logic action

- the space \mathbf{Y}_L is Polish;
- the Polish group $\text{Iso}(\mathbf{Y})$ acts on \mathbf{Y}_L continuously

The space of continuous structures

Fix a relational continuous signature L and a Polish space (\mathbf{Y}, d) .
Let S be a dense countable subset of \mathbf{Y} .

Define the space \mathbf{Y}_L of continuous L -structures on (\mathbf{Y}, d) as follows.

Metric on the set of L -structures: Enumerate all tuples of the form (j, \bar{s}) , where \bar{s} is a tuple $\in S$ of the length of the arity of R_j .

For L -structures M and N on \mathbf{Y} let

$$\delta(M, N) = \sum_{i=1}^{\infty} \{2^{-i} |R_j^M(\bar{s}) - R_j^N(\bar{s})| : i \text{ is the number of } (j, \bar{s})\}.$$

Logic action

- the space \mathbf{Y}_L is Polish;
- the Polish group $\text{Iso}(\mathbf{Y})$ acts on \mathbf{Y}_L continuously

Universality

Theorem

For any Polish group G there is a Polish space (\mathbf{Y}, d) and a continuous relational signature L such that

- $G < Iso(\mathbf{Y})$
- for any Polish G -space \mathbf{X} there is a Borel 1-1-map $\mathcal{M} : \mathbf{X} \rightarrow \mathbf{Y}_L$ such that for any $x, x' \in \mathbf{X}$ structures $\mathcal{M}(x)$ and $\mathcal{M}(x')$ are isomorphic if and only if x and x' are in the same G -orbit,

The map \mathcal{M} is a Borel G -invariant 1-1-reduction of the G -orbit equivalence relation on \mathbf{X} to the $Iso(\mathbf{Y})$ -orbit equivalence relation on the space \mathbf{Y}_L of all L -structures.

Grades subsets and subgroups

A **graded subset** of \mathbf{X} , denoted $\phi \sqsubseteq \mathbf{X}$, is a function $\mathbf{X} \rightarrow [0, 1]$.

It is **open (closed)**, $\phi \in \Sigma_1$ (resp. $\phi \in \Pi_1$), if it is upper (lower) semi-continuous, i.e. the set $\phi_{<r}$ (resp. $\phi_{>r}$) is open for all $r \in [0, 1]$ (here $\phi_{<r} = \{z \in \mathbf{X} : \phi(z) < r\}$).

When G is a Polish group, then a graded subset $H \sqsubseteq G$ is called a **graded subgroup** if $H(1) = 0$, $\forall g \in G (H(g) = H(g^{-1}))$ and $\forall g, g' \in G (H(gg') \leq H(g) + H(g'))$.

We also define Borel classes $\Sigma_\alpha, \Pi_\alpha$ so that ϕ is Σ_α if $\phi = \inf \Phi$ for some countable $\Phi \subset \bigcup \{\Pi_\gamma : \gamma < \alpha\}$ and $\Pi_\alpha = \{1 - \phi : \phi \in \Sigma_\alpha\}$.

Grades subsets and subgroups

A **graded subset** of \mathbf{X} , denoted $\phi \sqsubseteq \mathbf{X}$, is a function $\mathbf{X} \rightarrow [0, 1]$.

It is **open (closed)**, $\phi \in \Sigma_1$ (resp. $\phi \in \Pi_1$), if it is upper (lower) semi-continuous, i.e. the set $\phi_{<r}$ (resp. $\phi_{>r}$) is open for all $r \in [0, 1]$ (here $\phi_{<r} = \{z \in \mathbf{X} : \phi(z) < r\}$).

When G is a Polish group, then a graded subset $H \sqsubseteq G$ is called a **graded subgroup** if $H(1) = 0$, $\forall g \in G (H(g) = H(g^{-1}))$ and $\forall g, g' \in G (H(gg') \leq H(g) + H(g'))$.

We also define Borel classes $\Sigma_\alpha, \Pi_\alpha$ so that ϕ is Σ_α if $\phi = \inf \Phi$ for some countable $\Phi \subset \bigcup \{\Pi_\gamma : \gamma < \alpha\}$ and $\Pi_\alpha = \{1 - \phi : \phi \in \Sigma_\alpha\}$.

Grades subsets and subgroups

A **graded subset** of \mathbf{X} , denoted $\phi \sqsubseteq \mathbf{X}$, is a function $\mathbf{X} \rightarrow [0, 1]$.

It is **open (closed)**, $\phi \in \Sigma_1$ (resp. $\phi \in \Pi_1$), if it is upper (lower) semi-continuous, i.e. the set $\phi_{<r}$ (resp. $\phi_{>r}$) is open for all $r \in [0, 1]$ (here $\phi_{<r} = \{z \in \mathbf{X} : \phi(z) < r\}$).

When G is a Polish group, then a graded subset $H \sqsubseteq G$ is called a **graded subgroup** if $H(1) = 0$, $\forall g \in G (H(g) = H(g^{-1}))$ and $\forall g, g' \in G (H(gg') \leq H(g) + H(g'))$.

We also define Borel classes $\Sigma_\alpha, \Pi_\alpha$ so that ϕ is Σ_α if $\phi = \inf \Phi$ for some countable $\Phi \subset \bigcup \{\Pi_\gamma : \gamma < \alpha\}$ and $\Pi_\alpha = \{1 - \phi : \phi \in \Sigma_\alpha\}$.

Grades subsets and subgroups

A **graded subset** of \mathbf{X} , denoted $\phi \sqsubseteq \mathbf{X}$, is a function $\mathbf{X} \rightarrow [0, 1]$.

It is **open (closed)**, $\phi \in \Sigma_1$ (resp. $\phi \in \Pi_1$), if it is upper (lower) semi-continuous, i.e. the set $\phi_{<r}$ (resp. $\phi_{>r}$) is open for all $r \in [0, 1]$ (here $\phi_{<r} = \{z \in \mathbf{X} : \phi(z) < r\}$).

When G is a Polish group, then a graded subset $H \sqsubseteq G$ is called a **graded subgroup** if $H(1) = 0$, $\forall g \in G (H(g) = H(g^{-1}))$ and $\forall g, g' \in G (H(gg') \leq H(g) + H(g'))$.

We also define Borel classes $\Sigma_\alpha, \Pi_\alpha$ so that ϕ is Σ_α if $\phi = \inf \Phi$ for some countable $\Phi \subset \bigcup \{\Pi_\gamma : \gamma < \alpha\}$ and $\Pi_\alpha = \{1 - \phi : \phi \in \Sigma_\alpha\}$.

Graded subsets of \mathbf{Y}_L

For \bar{c} from (\mathbf{Y}, d) and a linear δ with $\delta(0) = 0$

graded subgroup $H_{\delta, \bar{c}} \sqsubseteq Iso(\mathbf{Y})$:

$$H_{\delta, \bar{c}}(g) = \delta(\max(d(c_1, g(c_1)), \dots, d(c_n, g(c_n))))), \text{ where } g \in Iso(\mathbf{Y}).$$

A **continuous formula** is an expression built from 0,1 and atomic formulas by applications of the following functions:

$$x/2, x \dot{-} y = \max(x - y, 0), \min(x, y), \max(x, y), |x - y|,$$

$$\neg(x) = 1 - x, x \dot{+} y = \min(x + y, 1), \sup_x \text{ and } \inf_x.$$

Any continuous sentence $\phi(\bar{c})$ defines a graded subset of \mathbf{Y}_L which belongs to Σ_n for some n :

$$\phi(\bar{c}) \text{ takes } M \text{ to the value } \phi^M(\bar{c}).$$

Graded subsets of \mathbf{Y}_L

For \bar{c} from (\mathbf{Y}, d) and a linear δ with $\delta(0) = 0$

graded subgroup $H_{\delta, \bar{c}} \sqsubseteq Iso(\mathbf{Y})$:

$H_{\delta, \bar{c}}(g) = \delta(\max(d(c_1, g(c_1)), \dots, d(c_n, g(c_n))))$, where $g \in Iso(\mathbf{Y})$.

A **continuous formula** is an expression built from 0,1 and atomic formulas by applications of the following functions:

$$x/2, x \dot{-} y = \max(x - y, 0), \min(x, y), \max(x, y), |x - y|,$$

$$\neg(x) = 1 - x, x \dot{+} y = \min(x + y, 1), \sup_x \text{ and } \inf_x.$$

Any continuous sentence $\phi(\bar{c})$ defines a graded subset of \mathbf{Y}_L which belongs to Σ_n for some n :

$\phi(\bar{c})$ takes M to the value $\phi^M(\bar{c})$.

Graded subsets of \mathbf{Y}_L

For \bar{c} from (\mathbf{Y}, d) and a linear δ with $\delta(0) = 0$

graded subgroup $H_{\delta, \bar{c}} \sqsubseteq Iso(\mathbf{Y})$:

$$H_{\delta, \bar{c}}(g) = \delta(\max(d(c_1, g(c_1)), \dots, d(c_n, g(c_n))))), \text{ where } g \in Iso(\mathbf{Y}).$$

A **continuous formula** is an expression built from 0,1 and atomic formulas by applications of the following functions:

$$x/2, x \dot{-} y = \max(x - y, 0), \min(x, y), \max(x, y), |x - y|,$$

$$\neg(x) = 1 - x, x \dot{+} y = \min(x + y, 1), \sup_x \text{ and } \inf_x.$$

Any continuous sentence $\phi(\bar{c})$ defines a graded subset of \mathbf{Y}_L which belongs to Σ_n for some n :

$$\phi(\bar{c}) \text{ takes } M \text{ to the value } \phi^M(\bar{c}).$$

Invariant graded subsets

Assuming that continuity moduli of L -symbols are **id** for any $\phi(\bar{x})$ as above we find a linear function δ such that the graded subgroup

$$H_{\delta, \bar{c}}(g) = \delta(\max(d(c_1, g(c_1)), \dots, d(c_n, g(c_n))))), \text{ where } g \in \text{Iso}(\mathbf{Y}).$$

and the graded subset $\phi(\bar{c}) \sqsubseteq \mathbf{Y}_L$ satisfy

$$\phi^{g(M)}(\bar{c}) \leq \phi^M(\bar{c}) \dot{+} H_{\delta, \bar{c}}(g).$$

Definition

Let \mathbf{X} be a continuous G -space. A graded subset $\phi \sqsubseteq \mathbf{X}$ is called invariant with respect to a graded subgroup $H \sqsubseteq G$ if for any $g \in G$ we have $\phi(g(x)) \leq \phi(x) \dot{+} H(g)$.

Invariant graded subsets

Assuming that continuity moduli of L -symbols are **id** for any $\phi(\bar{x})$ as above we find a linear function δ such that the graded subgroup

$$H_{\delta, \bar{c}}(g) = \delta(\max(d(c_1, g(c_1)), \dots, d(c_n, g(c_n))))), \text{ where } g \in \text{Iso}(\mathbf{Y}).$$

and the graded subset $\phi(\bar{c}) \sqsubseteq \mathbf{Y}_L$ satisfy

$$\phi^{g(M)}(\bar{c}) \leq \phi^M(\bar{c}) \dot{+} H_{\delta, \bar{c}}(g).$$

Definition

Let \mathbf{X} be a continuous G -space. A graded subset $\phi \sqsubseteq \mathbf{X}$ is called invariant with respect to a graded subgroup $H \sqsubseteq G$ if for any $g \in G$ we have $\phi(g(x)) \leq \phi(x) \dot{+} H(g)$.

Vaught transforms 3

For any non-empty open $J \sqsubseteq G$ let

$\phi^{\Delta J}(x) = \inf\{r \dot{+} s : \{h : \phi(h(x)) < r\} \text{ is not meagre in } J_{<s}\}$.

$\phi^{*J}(x) = \sup\{r \dot{-} s : \{h : \phi(h(x)) \leq r\} \text{ is not comeagre in } J_{<s}\}$,

Theorem

- $\phi^{*J}(x) = 1 - (1 - \phi)^{\Delta J}(x)$ for all $x \in \mathbf{X}$.
- $\phi^{\Delta J}(x) \leq \phi^{*J}(x)$ for all $x \in \mathbf{X}$.
- If ϕ is a graded Σ_α -subset, then $\phi^{\Delta J}$ is also Σ_α .
 If ϕ is a graded Π_α -subset, then $\phi^{*J}(x)$ is also Π_α .
- Vaught transforms of Borel graded subsets are Borel.

Vaught transforms 3

For any non-empty open $J \sqsubseteq G$ let

$\phi^{\Delta J}(x) = \inf\{r \dot{+} s : \{h : \phi(h(x)) < r\} \text{ is not meagre in } J_{<s}\}$.

$\phi^{*J}(x) = \sup\{r \dot{-} s : \{h : \phi(h(x)) \leq r\} \text{ is not comeagre in } J_{<s}\}$,

Theorem

- $\phi^{*J}(x) = 1 - (1 - \phi)^{\Delta J}(x)$ for all $x \in \mathbf{X}$.
- $\phi^{\Delta J}(x) \leq \phi^{*J}(x)$ for all $x \in \mathbf{X}$.
- If ϕ is a graded Σ_α -subset, then $\phi^{\Delta J}$ is also Σ_α .
 If ϕ is a graded Π_α -subset, then $\phi^{*J}(x)$ is also Π_α .
- Vaught transforms of Borel graded subsets are Borel.

Vaught transforms 4

Theorem

If H is a graded subgroup of G , then both $\phi^{*H}(x)$ and $\phi^{\Delta H}(x)$ are H -invariant:

$$\phi^{*H}(x) \dot{-} H(h) \leq \phi^{*H}(h(x)) \leq \phi^{*H}(x) \dot{+} H(h) \text{ and}$$

$$\phi^{\Delta H}(x) \dot{-} H(h) \leq \phi^{\Delta H}(h(x)) \leq \phi^{\Delta H}(x) \dot{+} H(h).$$

Moreover if $\phi(x) \leq \phi(h(x)) \dot{+} H(h)$ for all x and h , then

$$\phi^{*H}(x) = \phi(x) = \phi^{\Delta H}(x).$$

Graded bases

We consider G together with a distinguished countable family of open graded subsets \mathcal{R} so that all $\rho < r$ for $\rho \in \mathcal{R}$ and $r \in \mathbb{Q}$, form a basis of the topology of G .

We usually assume that \mathcal{R} consists of **graded cosets**, i.e. for such $\rho \in \mathcal{R}$ there is a graded subgroup $H \in \mathcal{R}$ and an element $g_0 \in G$ so that for any $g \in G$, $\rho(g) = H(gg_0^{-1})$.

(For every Polish group G there is a countable family of open graded subsets \mathcal{R} as above.)

Considering a (G, \mathcal{R}) -space \mathbf{X} we distinguish a similar family too: a cntble family \mathcal{U} of open graded sbsts of \mathbf{X} generating the topol.

Graded bases

We consider G together with a distinguished countable family of open graded subsets \mathcal{R} so that all $\rho < r$ for $\rho \in \mathcal{R}$ and $r \in \mathbb{Q}$, form a basis of the topology of G .

We usually assume that \mathcal{R} consists of **graded cosets**, i.e. for such $\rho \in \mathcal{R}$ there is a graded subgroup $H \in \mathcal{R}$ and an element $g_0 \in G$ so that for any $g \in G$, $\rho(g) = H(gg_0^{-1})$.

(For every Polish group G there is a countable family of open graded subsets \mathcal{R} as above.)

Considering a (G, \mathcal{R}) -space \mathbf{X} we distinguish a similar family too: a cntble family \mathcal{U} of open graded sbsts of \mathbf{X} generating the topol.

Graded bases

We consider G together with a distinguished countable family of open graded subsets \mathcal{R} so that all $\rho < r$ for $\rho \in \mathcal{R}$ and $r \in \mathbb{Q}$, form a basis of the topology of G .

We usually assume that \mathcal{R} consists of **graded cosets**, i.e. for such $\rho \in \mathcal{R}$ there is a graded subgroup $H \in \mathcal{R}$ and an element $g_0 \in G$ so that for any $g \in G$, $\rho(g) = H(gg_0^{-1})$.

(For every Polish group G there is a countable family of open graded subsets \mathcal{R} as above.)

Considering a (G, \mathcal{R}) -space \mathbf{X} we distinguish a similar family too: a cntble family \mathcal{U} of open graded sbsts of \mathbf{X} generating the topol.

Graded bases

We consider G together with a distinguished countable family of open graded subsets \mathcal{R} so that all $\rho < r$ for $\rho \in \mathcal{R}$ and $r \in \mathbb{Q}$, form a basis of the topology of G .

We usually assume that \mathcal{R} consists of **graded cosets**, i.e. for such $\rho \in \mathcal{R}$ there is a graded subgroup $H \in \mathcal{R}$ and an element $g_0 \in G$ so that for any $g \in G$, $\rho(g) = H(gg_0^{-1})$.

(For every Polish group G there is a countable family of open graded subsets \mathcal{R} as above.)

Considering a (G, \mathcal{R}) -space \mathbf{X} we distinguish a similar family too: a cntble family \mathcal{U} of open graded sbsts of \mathbf{X} generating the topol.

Nice basis

Definition. A family \mathcal{B} of Borel graded subsets of the G -space \mathbf{X} is a **nice basis** w.r.to \mathcal{R} if:

- \mathcal{B} is countable and generates the topol. finer than τ ;
- for all $\phi_1, \phi_2 \in \mathcal{B}$, the functions $\min(\phi_1, \phi_2)$, $\max(\phi_1, \phi_2)$, $|\phi_1 - \phi_2|$, $\phi_1 \dot{-} \phi_2$, $\phi_1 \dot{+} \phi_2$ belong to \mathcal{B} ;
- for all $\phi \in \mathcal{B}$ and rational $r \in [0, 1]$, $r\phi$ and $1 - \phi \in \mathcal{B}$;
- for all $\phi \in \mathcal{B}$ and $\rho \in \mathcal{R}$, $\phi^{*\rho}, \phi^{\Delta\rho} \in \mathcal{B}$;
- for any $\phi \in \mathcal{B}$ there exists an open graded subgroup $H \in \mathcal{R}$ such that ϕ is invariant under the corresponding H -action.

A topology \mathbf{t} on \mathbf{X} is \mathcal{R} -nice for the G -space $\langle \mathbf{X}, \tau \rangle$ if:

- (a) \mathbf{t} is a Polish topology, \mathbf{t} is finer than τ and the G -action remains continuous with respect to \mathbf{t} ;
- (b) there exists a nice basis \mathcal{B} so that \mathbf{t} is generated by all $\phi_{<q}$ with $\phi \in \mathcal{B}$ and $q \in \mathbb{Q} \cap (0, 1]$.

Nice basis

Definition. A family \mathcal{B} of Borel graded subsets of the G -space \mathbf{X} is a **nice basis** w.r.to \mathcal{R} if:

- \mathcal{B} is countable and generates the topol. finer than τ ;
- for all $\phi_1, \phi_2 \in \mathcal{B}$, the functions $\min(\phi_1, \phi_2)$, $\max(\phi_1, \phi_2)$, $|\phi_1 - \phi_2|$, $\phi_1 \dot{-} \phi_2$, $\phi_1 \dot{+} \phi_2$ belong to \mathcal{B} ;
- for all $\phi \in \mathcal{B}$ and rational $r \in [0, 1]$, $r\phi$ and $1 - \phi \in \mathcal{B}$;
- for all $\phi \in \mathcal{B}$ and $\rho \in \mathcal{R}$, $\phi^{*\rho}, \phi^{\Delta\rho} \in \mathcal{B}$;
- for any $\phi \in \mathcal{B}$ there exists an open graded subgroup $H \in \mathcal{R}$ such that ϕ is invariant under the corresponding H -action.

A topology \mathbf{t} on \mathbf{X} is **\mathcal{R} -nice** for the G -space $\langle \mathbf{X}, \tau \rangle$ if:

- \mathbf{t} is a Polish topology, \mathbf{t} is finer than τ and the G -action remains continuous with respect to \mathbf{t} ;
- there exists a nice basis \mathcal{B} so that \mathbf{t} is generated by all $\phi_{<q}$ with $\phi \in \mathcal{B}$ and $q \in \mathbb{Q} \cap (0, 1]$.

The case of \mathbf{U}_L

Let \mathbf{U} be **Urysohn** spce of diameter 1: This is the unique Polish mtrc space which is universal and ultrahomogeneous, i.e. every isometry between fnte substs of \mathbf{U} extends to an isometry of \mathbf{U} .

There is a ctble family \mathcal{R} consisting of cosets of clopen graded subgroups of $Iso(\mathbf{U})$ of the form

$$H_s : g \rightarrow d(g(\mathbf{s}), \mathbf{s}), \text{ where } \mathbf{s} \subset S \text{ (ctble, dense) ,}$$

which generates the topology of $Iso(\mathbf{U})$.

Let L be a continuous signature of continuity moduli id .

Then the family of all continuous L -sentences

$$\phi(\mathbf{s}) : M \rightarrow \phi^M(\mathbf{s}), \text{ where } \bar{\mathbf{s}} \in S,$$

forms an \mathcal{R} -nice basis \mathcal{B} of the G -space \mathbf{U}_L .

The case of \mathbf{U}_L

Let \mathbf{U} be **Urysohn** spce of diameter 1: This is the unique Polish mtrc space which is universal and ultrahomogeneous, i.e. every isometry between fnte substs of \mathbf{U} extends to an isometry of \mathbf{U} .

There is a ctble family \mathcal{R} consisting of cosets of clopen graded subgroups of $Iso(\mathbf{U})$ of the form

$$H_{\mathbf{s}} : g \rightarrow d(g(\mathbf{s}), \mathbf{s}), \text{ where } \mathbf{s} \subset S \text{ (ctble, dense) ,}$$

which generates the topology of $Iso(\mathbf{U})$.

Let L be a continuous signature of continuity moduli id .
 Then the family of all continuous L -sentences

$$\phi(\mathbf{s}) : M \rightarrow \phi^M(\mathbf{s}), \text{ where } \bar{\mathbf{s}} \in S,$$

forms an \mathcal{R} -nice basis \mathcal{B} of the G -space \mathbf{U}_L .

The case of \mathbf{U}_L

Let \mathbf{U} be **Urysohn** spce of diameter 1: This is the unique Polish mtrc space which is universal and ultrahomogeneous, i.e. every isometry between fnte substs of \mathbf{U} extends to an isometry of \mathbf{U} .

There is a ctble family \mathcal{R} consisting of cosets of clopen graded subgroups of $Iso(\mathbf{U})$ of the form

$$H_{\mathbf{s}} : g \rightarrow d(g(\mathbf{s}), \mathbf{s}), \text{ where } \mathbf{s} \subset S \text{ (ctble, dense) ,}$$

which generates the topology of $Iso(\mathbf{U})$.

Let L be a continuous signature of continuity moduli id .

Then the family of all continuous L -sentences

$$\phi(\mathbf{s}) : M \rightarrow \phi^M(\mathbf{s}), \text{ where } \bar{\mathbf{s}} \in S,$$

forms an \mathcal{R} -nice basis \mathcal{B} of the G -space \mathbf{U}_L .

The case of \mathbf{U}_L

Let \mathbf{U} be **Urysohn** spce of diameter 1: This is the unique Polish mtrc space which is universal and ultrahomogeneous, i.e. every isometry between fnte substs of \mathbf{U} extends to an isometry of \mathbf{U} .

There is a ctble family \mathcal{R} consisting of cosets of clopen graded subgroups of $Iso(\mathbf{U})$ of the form

$$H_{\mathbf{s}} : g \rightarrow d(g(\mathbf{s}), \mathbf{s}), \text{ where } \mathbf{s} \subset S \text{ (ctble, dense) ,}$$

which generates the topology of $Iso(\mathbf{U})$.

Let L be a continuous signature of continuity moduli id .

Then the family of all continuous L -sentences

$$\phi(\mathbf{s}) : M \rightarrow \phi^M(\mathbf{s}), \text{ where } \bar{\mathbf{s}} \in S,$$

forms an \mathcal{R} -nice basis \mathcal{B} of the G -space \mathbf{U}_L .

Existence 2

Theorem. Let (G, \mathcal{R}) be a Polish group with \mathcal{R} satisfying
 (i) for every graded subgroup $H \in \mathcal{R}$ if $gH \in \mathcal{R}$, then $H^g \in \mathcal{R}$;
 (ii) \mathcal{R} is closed under **max** and multiplying by rationals.

Let (\mathbf{X}, τ) be a G -space and \mathcal{U} be a countable family of Borel graded subsets of \mathbf{X} generating a topology finer than τ , so that each $\phi \in \mathcal{U}$ is invariant with respect to some graded subgroup $H \in \mathcal{R}$.

Then there is an \mathcal{R} -nice topology for $(\langle \mathbf{X}, \tau \rangle, G)$ so that \mathcal{U} consists of open graded subsets.

Existence 2

Theorem. Let (G, \mathcal{R}) be a Polish group with \mathcal{R} satisfying
 (i) for every graded subgroup $H \in \mathcal{R}$ if $gH \in \mathcal{R}$, then $H^g \in \mathcal{R}$;
 (ii) \mathcal{R} is closed under **max** and multiplying by rationals.

Let $\langle \mathbf{X}, \tau \rangle$ be a G -space and \mathcal{U} be a countable family of Borel graded subsets of \mathbf{X} generating a topology finer than τ , so that each $\phi \in \mathcal{U}$ is invariant with respect to some graded subgroup $H \in \mathcal{R}$.

Then there is an \mathcal{R} -nice topology for $(\langle \mathbf{X}, \tau \rangle, G)$ so that \mathcal{U} consists of open graded subsets.

Existence 2

Theorem. Let (G, \mathcal{R}) be a Polish group with \mathcal{R} satisfying
 (i) for every graded subgroup $H \in \mathcal{R}$ if $gH \in \mathcal{R}$, then $H^g \in \mathcal{R}$;
 (ii) \mathcal{R} is closed under **max** and multiplying by rationals.

Let $\langle \mathbf{X}, \tau \rangle$ be a G -space and \mathcal{U} be a countable family of Borel graded subsets of \mathbf{X} generating a topology finer than τ , so that each $\phi \in \mathcal{U}$ is invariant with respect to some graded subgroup $H \in \mathcal{R}$.

Then there is an \mathcal{R} -nice topology for $(\langle \mathbf{X}, \tau \rangle, G)$ so that \mathcal{U} consists of open graded subsets.

G is a Polish group with a graded basis \mathcal{R} consisting of graded cosets,
 $\langle \mathbf{X}, \tau \rangle$ is a Polish G -space, ect.

Theorem

Let \mathfrak{t} be \mathcal{R} -nice.

Let $X = Gx_0$ for some (any) $x_0 \in X$ and X be a G_δ -subset of \mathbf{X} .

Then both topologies τ and \mathfrak{t} are equal on X .

G is a Polish group with a graded basis \mathcal{R} consisting of graded cosets,
 $\langle \mathbf{X}, \tau \rangle$ is a Polish G -space, ect.

Theorem

Let \mathbf{t} be \mathcal{R} -nice.

Let $X = Gx_0$ for some (any) $x_0 \in X$ and X be a G_δ -subset of \mathbf{X} .

Then both topologies τ and \mathbf{t} are equal on X .

Categoricity

Let \mathcal{B} be a nice basis defining \mathcal{R} -nice \mathbf{t} ,
 H be an open graded subgroup from \mathcal{R} ,
 X be an invariant G_δ -subset of \mathbf{X} with respect to \mathbf{t} .

(1) A family \mathcal{F} of subsets of the form $\phi_{<r}$ with H -invariant $\phi \in \mathcal{B}$ is called an **H -type** in X , if it is maximal w.r. to the condition $X \cap \bigcap \mathcal{F} \neq \emptyset$.

(2) An H -type \mathcal{F} is called **principal** if there is an H -invariant graded basic set $\phi \in \mathcal{B}$ and there is r such that $\phi_{<r} \in \mathcal{F}$ and $\bigcap \{\bar{B} : B \in \mathcal{F}\} \cap X$ coincides with the closure of $\phi_{<r} \cap X$.

Then we say that $\phi_{<r}$ **defines** \mathcal{F} .

Categoricity

Let \mathcal{B} be a nice basis defining \mathcal{R} -nice \mathbf{t} ,
 H be an open graded subgroup from \mathcal{R} ,
 X be an invariant G_δ -subset of \mathbf{X} with respect to \mathbf{t} .

(1) A family \mathcal{F} of subsets of the form $\phi_{<r}$ with H -invariant $\phi \in \mathcal{B}$ is called an **H -type** in X , if it is maximal w.r. to the condition $X \cap \bigcap \mathcal{F} \neq \emptyset$.

(2) An H -type \mathcal{F} is called **principal** if there is an H -invariant graded basic set $\phi \in \mathcal{B}$ and there is r such that $\phi_{<r} \in \mathcal{F}$ and $\bigcap \{\bar{B} : B \in \mathcal{F}\} \cap X$ coincides with the closure of $\phi_{<r} \cap X$.

Then we say that $\phi_{<r}$ **defines** \mathcal{F} .

Ryll-Nardzewski

Let \mathcal{R} consist of clopen graded cosets.

Let \mathcal{B} be an \mathcal{R} -nice basis of a G -space $\langle \mathbf{X}, \tau \rangle$ and \mathbf{t} be the corresponding nice topology,

Theorem

*Assume that the action satisfies the **approximation property** for graded subgroups.*

A piece X of the canonical partition with respect to the topology \mathbf{t} is a G -orbit if and only if for any basic open graded subgroup $H \sqsubset G$ any H -type of X is principal.

Ryll-Nardzewski

Let \mathcal{R} consist of clopen graded cosets.

Let \mathcal{B} be an \mathcal{R} -nice basis of a G -space $\langle \mathbf{X}, \tau \rangle$ and \mathfrak{t} be the corresponding nice topology,

Theorem

*Assume that the action satisfies the **approximation property** for graded subgroups.*

A piece X of the canonical partition with respect to the topology \mathfrak{t} is a G -orbit if and only if for any basic open graded subgroup $H \sqsubset G$ any H -type of X is principal.

Ryll-Nardzewski

Let \mathcal{R} consist of clopen graded cosets.

Let \mathcal{B} be an \mathcal{R} -nice basis of a G -space $\langle \mathbf{X}, \tau \rangle$ and \mathbf{t} be the corresponding nice topology,

Theorem

*Assume that the action satisfies the **approximation property** for graded subgroups.*

A piece X of the canonical partition with respect to the topology \mathbf{t} is a G -orbit if and only if for any basic open graded subgroup $H \sqsubset G$ any H -type of X is principal.

Approximation property for graded subgroups

Definition

The (G, \mathcal{R}) -space $(\mathbf{X}, \mathcal{U})$ has the **approximation property for graded subgroups** if for any $\varepsilon > 0$

for any graded subgrp $H \in \mathcal{R}$, any c and $c' \in \mathbf{X}$ of the same G -orbit

if c, c' belong to the same subsets of the form $\phi_{\leq t}$ for H -invariant $\phi \in \mathcal{U}$, then

c' can be approximated by the values $g(c)$ with $H(g) < \varepsilon$.

When $G = \text{Aut}(M)$, where M is an approximately ultrahomogeneous separably categorical structure on \mathbf{Y} , then this holds in the space of all L -expansions of M .

Approximation property for graded subgroups

Definition

The (G, \mathcal{R}) -space $(\mathbf{X}, \mathcal{U})$ has the **approximation property for graded subgroups** if for any $\varepsilon > 0$
 for any graded subgrp $H \in \mathcal{R}$, any c and $c' \in \mathbf{X}$ of the same G -orbit
 if c, c' belong to the same subsets of the form $\phi_{\leq t}$ for H -invariant $\phi \in \mathcal{U}$, then
 c' can be approximated by the values $g(c)$ with $H(g) < \varepsilon$.

When $G = \text{Aut}(M)$, where M is an approximately ultrahomogeneous separably categorical structure on \mathbf{Y} , then this holds in the space of all L -expansions of M .

Approximation property for graded subgroups

Definition

The (G, \mathcal{R}) -space $(\mathbf{X}, \mathcal{U})$ has the **approximation property for graded subgroups** if for any $\varepsilon > 0$
 for any graded subgrp $H \in \mathcal{R}$, any c and $c' \in \mathbf{X}$ of the same G -orbit
 if c, c' belong to the same subsets of the form $\phi_{\leq t}$ for H -invariant $\phi \in \mathcal{U}$, then
 c' can be approximated by the values $g(c)$ with $H(g) < \varepsilon$.

When $G = \text{Aut}(M)$, where M is an approximately ultrahomogeneous separably categorical structure on \mathbf{Y} , then this holds in the space of all L -expansions of M .

Approximation property for graded subgroups

Definition

The (G, \mathcal{R}) -space $(\mathbf{X}, \mathcal{U})$ has the **approximation property for graded subgroups** if for any $\varepsilon > 0$
 for any graded subgrp $H \in \mathcal{R}$, any c and $c' \in \mathbf{X}$ of the same G -orbit
 if c, c' belong to the same subsets of the form $\phi_{\leq t}$ for H -invariant $\phi \in \mathcal{U}$, then
 c' can be approximated by the values $g(c)$ with $H(g) < \varepsilon$.

When $G = \text{Aut}(M)$, where M is an approximately ultrahomogeneous separably categorical structure on \mathbf{Y} , then this holds in the space of all L -expansions of M .

Ultrahomogeneity

A relational continuous structure M is **approximately ultrahomogeneous** if for any n -tuples (a_1, \dots, a_n) and (b_1, \dots, b_n) with the same quantifier-free type (i.e. with the same values of predicates for corresponding subtuples) and any $\varepsilon > 0$ there exists $g \in \text{Aut}(M)$ such that

$$\max\{d(g(a_j), b_j) : 1 \leq j \leq n\} \leq \varepsilon.$$

Any Polish group can be chosen as the automorphism group of a continuous metric structure which is approximately ultrahomogeneous.

Ultrahomogeneity

A relational continuous structure M is **approximately ultrahomogeneous** if for any n -tuples (a_1, \dots, a_n) and (b_1, \dots, b_n) with the same quantifier-free type (i.e. with the same values of predicates for corresponding subtuples) and any $\varepsilon > 0$ there exists $g \in \text{Aut}(M)$ such that

$$\max\{d(g(a_j), b_j) : 1 \leq j \leq n\} \leq \varepsilon.$$

Any Polish group can be chosen as the automorphism group of a continuous metric structure which is approximately ultrahomogeneous.

Complexity

Let (\mathbf{Y}, d) be a Polish space.

Theorem

- *There is a Borel subset $SC \subset \mathbf{Y}_L$ of separably categorical continuous L -structures on (\mathbf{Y}, d) so that any separably categorical continuous structure from \mathbf{Y}_L is isomorphic to a structure from SC .*
- *There is a Borel subset $SCU \subset \mathbf{Y}_L$ of separably categorical approximately ultrahomogeneous continuous structures on \mathbf{Y} so that any sep.cat., appr. ultrhom. structure from \mathbf{Y}_L is isomorphic to a structure from SCU .*

Complexity

Let (\mathbf{Y}, d) be a Polish space.

Theorem

- *There is a Borel subset $SC \subset \mathbf{Y}_L$ of separably categorical continuous L -structures on (\mathbf{Y}, d) so that any separably categorical continuous structure from \mathbf{Y}_L is isomorphic to a structure from SC .*
- *There is a Borel subset $SCU \subset \mathbf{Y}_L$ of separably categorical approximately ultrahomogeneous continuous structures on \mathbf{Y} so that any sep.cat., appr. ultrhom. structure from \mathbf{Y}_L is isomorphic to a structure from SCU .*

Complexity

Let (\mathbf{Y}, d) be a Polish space.

Theorem

- *There is a Borel subset $SC \subset \mathbf{Y}_L$ of separably categorical continuous L -structures on (\mathbf{Y}, d) so that any separably categorical continuous structure from \mathbf{Y}_L is isomorphic to a structure from SC .*
- *There is a Borel subset $SCU \subset \mathbf{Y}_L$ of separably categorical approximately ultrahomogeneous continuous structures on \mathbf{Y} so that any sep.cat., appr. ultrhom. structure from \mathbf{Y}_L is isomorphic to a structure from SCU .*

CLI metrics

Observation.

Let M be a Polish approximatly ultrahomogeneous continuous.
Then $Aut(M)$ admits a compatible complete left-invariant metric if
and only if there is no proper embedding of M into itself.

The subset of \mathbf{Y}_L consisting of structures M so that $Aut(M)$
admits compatible complete left-invariant metric, is coanalytic in
any Borel subset of \mathbf{Y}_L .
It does not have any member in \mathcal{SC} .

CLI metrics

Observation.

Let M be a Polish approximatly ultrahomogeneous continuous.
Then $Aut(M)$ admits a compatible complete left-invariant metric if
and only if there is no proper embedding of M into itself.

The subset of \mathbf{Y}_L consisting of structures M so that $Aut(M)$
admits compatible complete left-invariant metric, is coanalytic in
any Borel subset of \mathbf{Y}_L .

It does not have any member in \mathcal{SC} .

CLI metrics

Observation.

Let M be a Polish approximatly ultrahomogeneous continuous.
Then $Aut(M)$ admits a compatible complete left-invariant metric if and only if there is no proper embedding of M into itself.

The subset of \mathbf{Y}_L consisting of structures M so that $Aut(M)$ admits compatible complete left-invariant metric, is coanalytic in any Borel subset of \mathbf{Y}_L .

It does not have any member in \mathcal{SC} .