Local homogeneity of measures: Results

Ville Suomala

University of Jyväskylä, Finland

Warsaw, October 2010

Joint work with Antti Käenmäki (Jyväskylä) and Tapio Rajala (Pisa)

On the definition of the dimension

On the definition of the dimension

Let X be a metric space and $A \subset X$, $x \in X$, $0 < \delta < 1$ and $r > 0$.

On the definition of the dimension

Let X be a metric space and $A \subset X$, $x \in X$, $0 < \delta < 1$ and $r > 0$.

Question

How many balls of radius δr are needed to cover $B(x, r) \cap A$?





















On the definition of the dimension

Let X be a metric space and $A \subset X$, $x \in X$, $0 < \delta < 1$ and $r > 0$.

Question

How many balls of radius δr are needed to cover $B(x, r) \cap A$?

On the definition of the dimension

Let X be a metric space and $A \subset X$, $x \in X$, $0 < \delta < 1$ and $r > 0$.

Question

How many balls of radius δr are needed to cover $B(x, r) \cap A$?

Heuristic answer

If $\dim A < s$ we typically need at most $C\delta^{-s}$ such balls.

On the definition of the dimension

Let X be a metric space and $A \subset X$, $x \in X$, $0 < \delta < 1$ and $r > 0$.

Question

How many balls of radius δr are needed to cover $B(x, r) \cap A$?

Heuristic answer

If $\dim A < s$ we typically need at most $C\delta^{-s}$ such balls.
If $\dim A > s$ we typically need at least $c\delta^{-s}$ such balls.

On the definition of the dimension

Let X be a metric space and $A \subset X$, $x \in X$, $0 < \delta < 1$ and $r > 0$.

Question

How many balls of radius δr are needed to cover $B(x, r) \cap A$?

Heuristic answer

If $\dim A < s$ we typically need at most $C\delta^{-s}$ such balls.
If $\dim A > s$ we typically need at least $c\delta^{-s}$ such balls.

In general, the above answer is of course false

On the definition of the dimension

Let X be a metric space and $A \subset X$, $x \in X$, $0 < \delta < 1$ and $r > 0$.

Question

How many balls of radius δr are needed to cover $B(x, r) \cap A$?

Heuristic answer

If $\dim A < s$ we typically need at most $C\delta^{-s}$ such balls.
If $\dim A > s$ we typically need at least $c\delta^{-s}$ such balls.

In general, the above answer is of course false(and it depends on the definition of the dimension how false...)

On the definition of the dimension

On the definition of the dimension

Let μ be a measure on X, $x \in X$, $r > 0$, and $0 < \varepsilon, \delta < 1$.

On the definition of the dimension

Let μ be a measure on X, $x \in X$, $r > 0$, and $0 < \varepsilon, \delta < 1$.

Question

How many disjoint balls $B_i \subset B(x, r)$ of radius δr can we find so that $\mu(B_i) > \varepsilon \mu(B(x, r))$?















On the definition of the dimension

Let μ be a measure on X, $x \in X$, $r > 0$, and $0 < \varepsilon, \delta < 1$.

Question

How many disjoint balls $B_i \subset B(x, r)$ of radius δr can we find so that $\mu(B_i) \geq \varepsilon \mu(B(x, r))$?

On the definition of the dimension

Let μ be a measure on X, $x \in X$, $r > 0$, and $0 < \varepsilon, \delta < 1$.

Question

How many disjoint balls $B_i \subset B(x, r)$ of radius δr can we find so that $\mu(B_i) \geq \varepsilon \mu(B(x, r))$?

Heuristic guess

If ε is small and $\dim \mu = s$ the answer should be roughly δ^{-s}.

Notation

Notation

A metric space X is *doubling* if there is $N < \infty$ such that for all $x \in X$ and $r > 0$, the ball $B(x, 2r)$ may be covered by N balls of radius r.

Notation

A metric space X is *doubling* if there is $N < \infty$ such that for all $x \in X$ and $r > 0$, the ball $B(x, 2r)$ may be covered by N balls of radius r.

If μ is a measure on X and $x \in X$, we define the *upper and lower local dimensions of μ at x* by

$$\overline{\dim}_{\mathrm{loc}}(\mu, x) = \limsup_{r \downarrow 0} \frac{\log \mu\big(B(x, r)\big)}{\log r},$$

$$\underline{\dim}_{\mathrm{loc}}(\mu, x) = \liminf_{r \downarrow 0} \frac{\log \mu\big(B(x, r)\big)}{\log r}.$$

Notation

A metric space X is *doubling* if there is $N < \infty$ such that for all $x \in X$ and $r > 0$, the ball $B(x, 2r)$ may be covered by N balls of radius r.

If μ is a measure on X and $x \in X$, we define the *upper and lower local dimensions of μ at x* by

$$\overline{\dim}_{\mathrm{loc}}(\mu, x) = \limsup_{r \downarrow 0} \frac{\log \mu\big(B(x,r)\big)}{\log r},$$

$$\underline{\dim}_{\mathrm{loc}}(\mu, x) = \liminf_{r \downarrow 0} \frac{\log \mu\big(B(x,r)\big)}{\log r}.$$

A measure μ on X has the *density point property* if

$$\lim_{r \downarrow 0} \frac{\mu\big(A \cap B(x,r)\big)}{\mu\big(B(x,r)\big)} = 1$$

for μ-almost all $x \in A$ whenever $A \subset X$ is μ-measurable.

Local homogeneity and the homogeneity dimension

Let μ be a measure on X, $x \in X$, and $\delta, \varepsilon, r > 0$.

Local homogeneity and the homogeneity dimension

Let μ be a measure on X, $x \in X$, and $\delta, \varepsilon, r > 0$. Define

$$\hom_{\delta,\varepsilon,r}(\mu, x) = \sup\{\#\mathcal{B} : \mathcal{B} \text{ is a } (\delta r)\text{-packing of } B(x, r) \text{ so that } \mu(B) > \varepsilon\mu\big(B(x, 5r)\big) \text{ for all } B \in \mathcal{B}\}$$

Local homogeneity and the homogeneity dimension

Let μ be a measure on X, $x \in X$, and $\delta, \varepsilon, r > 0$. Define

$$\hom_{\delta,\varepsilon,r}(\mu, x) = \sup\{\#\mathcal{B} : \mathcal{B} \text{ is a } (\delta r)\text{-packing of } B(x,r) \text{ so that } \mu(B) > \varepsilon\mu\big(B(x,5r)\big) \text{ for all } B \in \mathcal{B}\}$$

and from this let the *local δ-homogeneity of a measure μ at x* be

$$\hom_{\delta}(\mu, x) = \lim_{\varepsilon \downarrow 0} \limsup_{r \downarrow 0} \hom_{\delta,\varepsilon,r}(\mu, x).$$

Local homogeneity and the homogeneity dimension

Let μ be a measure on X, $x \in X$, and $\delta, \varepsilon, r > 0$. Define

$$\hom_{\delta,\varepsilon,r}(\mu, x) = \sup\{\#\mathcal{B} : \mathcal{B} \text{ is a } (\delta r)\text{-packing of } B(x,r) \text{ so that } \mu(B) > \varepsilon\mu\big(B(x,5r)\big) \text{ for all } B \in \mathcal{B}\}$$

and from this let the *local δ-homogeneity of a measure μ at x* be

$$\hom_\delta(\mu, x) = \lim_{\varepsilon \downarrow 0} \limsup_{r \downarrow 0} \hom_{\delta,\varepsilon,r}(\mu, x).$$

The *local homogeneity dimension of a measure μ at x* is then defined as

$$\dim_{\hom}(\mu, x) = \liminf_{\delta \downarrow 0} \frac{\log^+ \hom_\delta(\mu, x)}{-\log \delta},$$

where $\log^+(t) = \max\{0, \log t\}$.

Remarks on the definition

Remarks on the definition

- $\dim_{\mathrm{hom}}(\mu, x)$ is roughly the least possible exponent s so that "large parts" of $B(x, r)$ in terms of μ can always be covered by δ^{-s} balls of radius δr for all small $r, \delta > 0$.

Remarks on the definition

- $\dim_{\mathrm{hom}}(\mu, x)$ is roughly the least possible exponent s so that "large parts" of $B(x, r)$ in terms of μ can always be covered by δ^{-s} balls of radius δr for all small $r, \delta > 0$.
- May be considered as a kind of local Assouad dimension for the measure μ around x.

Remarks on the definition

- $\dim_{\mathrm{hom}}(\mu, x)$ is roughly the least possible exponent s so that "large parts" of $B(x, r)$ in terms of μ can always be covered by δ^{-s} balls of radius δr for all small $r, \delta > 0$.
- May be considered as a kind of local Assouad dimension for the measure μ around x.
- It is not essential that the constant in the definition of $\mathrm{hom}_{\delta,\varepsilon,r}$ is 5, but it is important that it is strictly greater than 1!

Main results

Main results

Theorem A

If μ is a Radon measure on a doubling metric space X, then

$$\overline{\dim}_{\mathrm{loc}}(\mu, x) \leq \dim_{\mathrm{hom}}(\mu, x)$$

for μ-almost all $x \in X$.

A quantitative version

A quantitative version

Theorem B

Suppose X is a doubling metric space with a doubling constant N. If $0 < m < s$, then there exists a constant $\delta_0 = \delta_0(m, s, N) > 0$ such that for every $0 < \delta < \delta_0$ there is $\varepsilon_0 = \varepsilon_0(m, s, N, \delta) > 0$ so that for every Radon measure μ on X we have

$$\limsup_{r \downarrow 0} \hom_{\delta,\varepsilon,r}(\mu, x) \geq \delta^{-m}$$

for all $0 < \varepsilon \leq \varepsilon_0$ and for μ-almost all $x \in X$ that satisfy $\overline{\dim}_{\mathrm{loc}}(\mu, x) > s$.

A quantitative version

Theorem B

Suppose X is a doubling metric space with a doubling constant N. If $0 < m < s$, then there exists a constant $\delta_0 = \delta_0(m, s, N) > 0$ such that for every $0 < \delta < \delta_0$ there is $\varepsilon_0 = \varepsilon_0(m, s, N, \delta) > 0$ so that for every Radon measure μ on X we have

$$\limsup_{r \downarrow 0} \hom_{\delta,\varepsilon,r}(\mu, x) \geq \delta^{-m}$$

for all $0 < \varepsilon \leq \varepsilon_0$ and for μ-almost all $x \in X$ that satisfy $\overline{\dim}_{\mathrm{loc}}(\mu, x) > s$.

Remark: Theorem B easily implies Theorem A.

About the proof

About the proof

Key Lemma

Suppose μ is a Radon measure on a doubling metric space X with bounded support and $A \subset X$ is a Borel set with $\mu(A) > 0$. If there are $0 < \delta < 1$, $0 < t < s$, and $k_0 \in \mathbb{N}$ such that for every integer $k \geq k_0$ and for every δ^k-packing $\mathcal{B}$ of A there is an δ^{k-1}-packing $\mathcal{B}'$ of A so that

$$\delta^t \sum_{B \in \mathcal{B}} \mu(B)^{1-t/s} \leq \tfrac{1}{2} \sum_{B \in \mathcal{B}'} \mu(B)^{1-t/s},$$

then $\mu(\{x \in A : \overline{\dim}_{\mathrm{loc}}(\mu, x) \leq s\}) > 0$.

About the proof

Key Lemma

Suppose μ is a Radon measure on a doubling metric space X with bounded support and $A \subset X$ is a Borel set with $\mu(A) > 0$. If there are $0 < \delta < 1$, $0 < t < s$, and $k_0 \in \mathbb{N}$ such that for every integer $k \geq k_0$ and for every δ^k-packing $\mathcal{B}$ of A there is an δ^{k-1}-packing $\mathcal{B}'$ of A so that

$$\delta^t \sum_{B \in \mathcal{B}} \mu(B)^{1-t/s} \leq \tfrac{1}{2} \sum_{B \in \mathcal{B}'} \mu(B)^{1-t/s},$$

then $\mu(\{x \in A : \overline{\dim}_{\mathrm{loc}}(\mu, x) \leq s\}) > 0$.

It is somewhat easier to prove such an estimate for dyadic mesh cubes of $\mathbb{R}^n$, or more generally for certain generalised nested cubes of a metric space X.

About the proof

Key Lemma

Suppose μ is a Radon measure on a doubling metric space X with bounded support and $A \subset X$ is a Borel set with $\mu(A) > 0$. If there are $0 < \delta < 1$, $0 < t < s$, and $k_0 \in \mathbb{N}$ such that for every integer $k \geq k_0$ and for every δ^k-packing $\mathcal{B}$ of A there is an δ^{k-1}-packing $\mathcal{B}'$ of A so that

$$\delta^t \sum_{B \in \mathcal{B}} \mu(B)^{1-t/s} \leq \tfrac{1}{2} \sum_{B \in \mathcal{B}'} \mu(B)^{1-t/s},$$

then $\mu(\{x \in A : \overline{\dim}_{\mathrm{loc}}(\mu, x) \leq s\}) > 0$.

It is somewhat easier to prove such an estimate for dyadic mesh cubes of $\mathbb{R}^n$, or more generally for certain generalised nested cubes of a metric space X. One could also define the local homogeneity and prove results analogous to Theorems A and B with respect to such a grid structure.

About the proof

Key Lemma

Suppose μ is a Radon measure on a doubling metric space X with bounded support and $A \subset X$ is a Borel set with $\mu(A) > 0$. If there are $0 < \delta < 1$, $0 < t < s$, and $k_0 \in \mathbb{N}$ such that for every integer $k \geq k_0$ and for every δ^k-packing $\mathcal{B}$ of A there is an δ^{k-1}-packing $\mathcal{B}'$ of A so that

$$\delta^t \sum_{B \in \mathcal{B}} \mu(B)^{1-t/s} \leq \tfrac{1}{2} \sum_{B \in \mathcal{B}'} \mu(B)^{1-t/s},$$

then $\mu(\{x \in A : \overline{\dim}_{\mathrm{loc}}(\mu, x) \leq s\}) > 0$.

It is somewhat easier to prove such an estimate for dyadic mesh cubes of $\mathbb{R}^n$, or more generally for certain generalised nested cubes of a metric space X. One could also define the local homogeneity and prove results analogous to Theorems A and B with respect to such a grid structure. **However, it is essential for our applications that our concepts of local homogeneity and homogeneity dimension enjoy a "spherically symmetric" definition.**

Local L^q-spectrum and dimension

A (2^{-n})-partition of X is a partition of X into Borel sets Q such that for each such Q there is $x \in Q$ for which $B(x, 2^{-n}) \subset Q \subset B(x, C2^{-n})$.

Local L^q-spectrum and dimension

A (2^{-n})-partition of X is a partition of X into Borel sets Q such that for each such Q there is $x \in Q$ for which $B(x, 2^{-n}) \subset Q \subset B(x, C2^{-n})$.

For each $n \in \mathbb{N}$, let $\mathcal{Q}_n$ be a (2^{-n})-partition of X and let $\mathcal{Q}_n(x, r) = \{Q \cap B(x, r) \,:\, Q \in \mathcal{Q}_n\}$. For a measure μ on X, $x \in X$ and $q \in \mathbb{R}$ $(q \geq 0)$, we set

$$\tau_q(\mu, x) = \lim_{r \downarrow 0} \liminf_{n \to \infty} \frac{\log \sum_{Q \in \mathcal{Q}_n(x,r)} \mu(Q)^q}{\log 2^{-n}}.$$

Local L^q-spectrum and dimension

A (2^{-n})-partition of X is a partition of X into Borel sets Q such that for each such Q there is $x \in Q$ for which $B(x, 2^{-n}) \subset Q \subset B(x, C2^{-n})$.

For each $n \in \mathbb{N}$, let $\mathcal{Q}_n$ be a (2^{-n})-partition of X and let $\mathcal{Q}_n(x, r) = \{Q \cap B(x, r) \,:\, Q \in \mathcal{Q}_n\}$. For a measure μ on X, $x \in X$ and $q \in \mathbb{R}$ $(q \geq 0)$, we set

$$\tau_q(\mu, x) = \lim_{r \downarrow 0} \liminf_{n \to \infty} \frac{\log \sum_{Q \in \mathcal{Q}_n(x,r)} \mu(Q)^q}{\log 2^{-n}}.$$

If $q \neq 1$, we define

$$\dim_q(\mu, x) = \tau_q(\mu, x)/(q - 1).$$

Local L^q-spectrum and dimension

A (2^{-n})-partition of X is a partition of X into Borel sets Q such that for each such Q there is $x \in Q$ for which $B(x, 2^{-n}) \subset Q \subset B(x, C2^{-n})$.

For each $n \in \mathbb{N}$, let $\mathcal{Q}_n$ be a (2^{-n})-partition of X and let $\mathcal{Q}_n(x,r) = \{Q \cap B(x,r) \,:\, Q \in \mathcal{Q}_n\}$. For a measure μ on X, $x \in X$ and $q \in \mathbb{R}$ $(q \geq 0)$, we set

$$\tau_q(\mu, x) = \lim_{r \downarrow 0} \liminf_{n \to \infty} \frac{\log \sum_{Q \in \mathcal{Q}_n(x,r)} \mu(Q)^q}{\log 2^{-n}}.$$

If $q \neq 1$, we define

$$\dim_q(\mu, x) = \tau_q(\mu, x)/(q-1).$$

Remark

If we forget (x, r), then we arrive at the (classical) global L^q-spectrum and dimension, $\tau_q(\mu)$ and $\dim_q(\mu)$ (provided that $0 < \mu(X) < \infty$).

The case $q = 1$

The case $q = 1$

When $q = 1$, we set

$$\overline{\dim}_1(\mu, x) = \limsup_{r \downarrow 0} \limsup_{n \to \infty} \frac{\sum_{Q \in \mathcal{Q}_n(x,r)} \mu(Q) \log \mu(Q)}{\mu\big(B(x,r)\big) \log 2^{-n}},$$

$$\underline{\dim}_1(\mu, x) = \liminf_{r \downarrow 0} \liminf_{n \to \infty} \frac{\sum_{Q \in \mathcal{Q}_n(x,r)} \mu(Q) \log \mu(Q)}{\mu\big(B(x,r)\big) \log 2^{-n}}.$$

Basic results

Basic results

Proposition

The above definitions are all independent of the choice of the partitions. (Observe that the partitions do not have to be nested!)

Basic results

Proposition

The above definitions are all independent of the choice of the partitions. (Observe that the partitions do not have to be nested!)

Theorem C

If μ is a Radon measure on a doubling metric space X, then

$$\lim_{q\downarrow 1} \dim_q(\mu, x) \leq \underline{\dim}_{\mathrm{loc}}(\mu, x) \leq \overline{\dim}_{\mathrm{loc}}(\mu, x) \leq \lim_{q\uparrow 1} \dim_q(\mu, x)$$

for μ-almost all $x \in X$.

Basic results

Proposition

The above definitions are all independent of the choice of the partitions. (Observe that the partitions do not have to be nested!)

Theorem C

If μ is a Radon measure on a doubling metric space X, then

$$\lim_{q\downarrow 1} \dim_q(\mu,x) \le \underline{\dim}_{\mathrm{loc}}(\mu,x) \le \overline{\dim}_{\mathrm{loc}}(\mu,x) \le \lim_{q\uparrow 1} \dim_q(\mu,x)$$

for μ-almost all $x \in X$. In addition, if μ has the density point property, then

$$\underline{\dim}_{\mathrm{loc}}(\mu,x) \le \underline{\dim}_1(\mu,x) \le \overline{\dim}_1(\mu,x) \le \overline{\dim}_{\mathrm{loc}}(\mu,x)$$

for μ-almost all $x \in X$.

Basic results

Proposition

The above definitions are all independent of the choice of the partitions. (Observe that the partitions do not have to be nested!)

Theorem C

If μ is a Radon measure on a doubling metric space X, then

$$\lim_{q\downarrow 1} \dim_q(\mu,x) \le \underline{\dim}_{\mathrm{loc}}(\mu,x) \le \overline{\dim}_{\mathrm{loc}}(\mu,x) \le \lim_{q\uparrow 1} \dim_q(\mu,x)$$

for μ-almost all $x \in X$. In addition, if μ has the density point property, then

$$\underline{\dim}_{\mathrm{loc}}(\mu,x) \le \underline{\dim}_1(\mu,x) \le \overline{\dim}_1(\mu,x) \le \overline{\dim}_{\mathrm{loc}}(\mu,x)$$

for μ-almost all $x \in X$.

Note

There is an analogous result for the global L^q-dimensions.

Relation between local and global concepts

Relation between local and global concepts

Proposition

Let μ be a Radon measure on a doubling metric space X with a compact support. Then

$$\tau_q(\mu) = \min_{x \in \operatorname{spt}(\mu)} \tau_q(\mu, x)$$

for every $q \in \mathbb{R}$.

Relation between local and global concepts

Proposition

Let μ be a Radon measure on a doubling metric space X with a compact support. Then

$$\tau_q(\mu) = \min_{x \in \operatorname{spt}(\mu)} \tau_q(\mu, x)$$

for every $q \in \mathbb{R}$. In particular,

$$\dim_q(\mu) = \begin{cases} \max_{x \in \operatorname{spt}(\mu)} \dim_q(\mu, x), & \text{if } q < 1, \\ \min_{x \in \operatorname{spt}(\mu)} \dim_q(\mu, x), & \text{if } q > 1. \end{cases}$$