

# SPINE DECOMPOSITION AND $L \log L$ CRITERION FOR SUPERPROCESSES WITH NON-LOCAL BRANCHING MECHANISMS

RENMING SONG

In this talk, I will present a pathwise spine decomposition for superprocesses with both local and non-local branching mechanisms under a martingale change of measure. This result complements the related results obtained in [2, 3, 4] for superprocesses with purely local branching mechanisms and in [1, 5] for multitype superprocesses. As an application of this decomposition, we obtain necessary/sufficient conditions for the limit of the fundamental martingale to be non-degenerate. In particular, we obtain extinction properties of superprocesses with non-local branching mechanisms as well as a Kesten-Stigum  $L \log L$  theorem for the fundamental martingale.

This talk is based a joint paper with Y.-X. Ren and T. Yang.

## REFERENCES

- [1] Chen, Z.-Q., Ren, Y.-X., and Song, R.:  $L \log L$  criterion for a class of multitype superdiffusions with non-local branching mechanism. Preprint, 2016.
- [2] Evans, S. N.: Two representations of a conditioned superprocess. *Proc. R. Soc. Edinburgh A*, **123** (1993), 959–971.
- [3] Kyprianou, A. E., Liu, R.-L., Murillo-Salas, A. and Ren, Y.-X.: Supercritical super-Brownian motion with a general branching mechanism and travelling waves. *Ann. Inst. Henri Poincaré Probab. Stat.*, **48**(2012), 661-687.
- [4] Liu, R.-L., Ren, Y.-X. and Song, R.:  $L \log L$  criteria for a class of superdiffusions. *J. Appl. Probab.*, **46** (2009), 479–496.
- [5] Kyprianou, A.E. and Palau, S.: Extinction properties of multi-type continuous-state branching processes. Preprint, 2016. <https://arxiv.org/abs/1604.04129v2>.