

# Spike dynamics near spacelike singularities

Woei Chet Lim

Department of Mathematics and Statistics, University of Waikato

April 2018, KTH

# Abstract

It is widely known that, near spacelike singularities, most worldlines would undergo the so-called BKL singularity dynamics, which is an infinite sequence of Kasner saddle states connected by Taub transitions.


Some worldlines however would undergo highly inhomogeneous spike dynamics, which is an infinite sequence of Kasner saddle states connected by spike transitions.

My contributions in this area are the discovery of the exact solutions which describe the spike transitions, and the development of a numerical zooming technique specially designed for spike simulations.

I will illustrate the spike transitions with snapshots of the Hubble-normalised state space trajectories, and point out interesting features.

# The BKL Conjecture

How does a general cosmological solution of Einstein's equations behave near a spacelike initial singularity?



Conjecture by Vladimir Belinski, Isaak Khalatnikov and Evgeny Lifshitz:  
[LK 1963, BKL 1970,1982]

A generic singularity is

1. **Vacuum-dominated** ("matter doesn't matter")
2. **Local** (spatial derivatives are negligible)
3. **Oscillatory** (an infinite sequence of Kasner states)

Background: BKL dynamics in spatially homogeneous spacetimes

# Building blocks of the BKL dynamics are made of

1. the Kasner saddle states (Bianchi type I) [Kasner 1925]
2. the Taub transitions (Bianchi type II) [Taub 1951]

They are the two simplest vacuum, anisotropic, spatially homogeneous solutions of the EFE.

Q: How do these solutions behave?

The EFE for spatially homogeneous spacetimes can be written as a **system of first order ODEs**.

We will explain the evolution of the solution in the language of **dynamical systems** (state space, equilibrium points, orbits, attractor).

The evolution of a solution is represented by an **orbit** in the state space. The majority of the orbits may approach a special subset of the state space asymptotically. We call that subset the **attractor**.

**Self-similar** solutions can be represented by **equilibrium points** if you use the right state space variables.

# Orthonormal frame formalism of EFE

[Ehlers 1961, Ellis 1971, MacCallum 1973, van Elst & Uggla 1997]

An appropriate choice for the state space variables is the Hubble-normalized scale-invariant variables of the orthonormal frame formalism.

Timelike congruence  $\mathbf{u}$ . Decompose  $u_{a;b}$  into irreducible parts.

$$u_{a;b} = \sigma_{ab} + \omega_{ab} + \frac{1}{3}\Theta(g_{ab} + u_a u_b) - \dot{u}_a u_b$$

$\Theta$  rate of expansion scalar.

$\sigma_{ab}$  rate of shear tensor. Symmetric traceless.

$\omega_{ab}$  vorticity tensor. Antisymmetric.

$\dot{u}_a$  acceleration vector.

In cosmological context, Hubble scalar  $H = \frac{1}{3}\Theta$  is used.

We usually use  $\mathbf{u}$  with zero vorticity.

Construct an orthonormal frame  $\{\mathbf{e}_0, \mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3\}$  with  $\mathbf{e}_0 = \mathbf{u}$  and 3 spatial frame vector  $\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3$ .

Orthonormal frame vector fields (assuming  $\mathbf{e}_0$  has zero vorticity)

$$\text{timelike} \quad \mathbf{e}_0 = \frac{1}{N} \frac{\partial}{\partial t}$$

$$\text{spacelike} \quad \mathbf{e}_\alpha = e_\alpha^i \frac{\partial}{\partial x^i}, \quad \alpha = 1, 2, 3, \quad i = 1, 2, 3.$$

$N$  is the lapse function, and  $e_\alpha^i$  the frame variables.

Orthonormality:  $e_\alpha^i e_\beta^j g_{ij} = \delta_{\alpha\beta}$

$e_\alpha^i$  has inverse  $e^\alpha_i$ .

Related to the metric as follows

$$ds^2 = -N^2 dt^2 + \delta_{\alpha\beta} e^\alpha_i e^\beta_j dx^i dx^j.$$

Further reduce the degrees of freedom by aligning the orthonormal frame vectors to the coordinate vectors as much as possible. Later, for Gowdy spacetimes, we align  $\mathbf{e}_2$  with  $\partial_y$  (no  $x$  and  $z$  components), and align  $\mathbf{e}_3$  with a linear combination of  $\partial_y$  and  $\partial_z$  (no  $x$  component).

The orthonormal frame vector fields are linear combinations of partial differential operators with non-constant coefficients, and they do not commute.

$$[\mathbf{e}_0, \mathbf{e}_\alpha] \neq 0, \quad [\mathbf{e}_\alpha, \mathbf{e}_\beta] \neq 0.$$

# Frame commutator coefficients

The orthonormal frame vectors do not commute as operators. The commutator coefficients are

$$\begin{aligned}[\mathbf{e}_0, \mathbf{e}_\alpha] &= \dot{u}_\alpha \mathbf{e}_0 - (H\delta_\alpha^\beta + \sigma_\alpha^\beta - \epsilon_{\alpha\gamma}{}^\beta \Omega^\gamma) \mathbf{e}_\beta \\ [\mathbf{e}_\alpha, \mathbf{e}_\beta] &= -(2a_{[\alpha} \delta_{\beta]}{}^\gamma + \epsilon_{\alpha\beta\delta} n^{\delta\gamma}) \mathbf{e}_\gamma\end{aligned}$$

$\Omega^\alpha$  is the angular velocity of the spatial frame  $\mathbf{e}_\alpha$ , and is determined by the alignment of the frame vectors with the coordinate vectors.

$a_\alpha$  and  $n_{\alpha\beta}$  (symmetric) determine the curvature of the spacelike hypersurface  $t = \text{const}$ .

The commutator coefficients are essentially partial derivatives of the lapse  $N$  and the frame variables  $e_\alpha^i$ .

## Choosing the right state space variables

Divide these variables by  $H$  to give the Hubble-normalized variables.

$$\begin{aligned}\frac{1}{\mathcal{N}^H} &= \frac{1}{H} \frac{1}{N}, & (E_\alpha^i)^H &= \frac{1}{H} e_\alpha^i, \\ \Sigma_{\alpha\beta}^H &= \frac{1}{H} \sigma_{\alpha\beta}, & \dot{U}_\alpha^H &= \frac{1}{H} \dot{u}_\alpha, & R_\alpha^H &= \frac{1}{H} \Omega_\alpha, \\ A_\alpha^H &= \frac{1}{H} a_\alpha, & N_{\alpha\beta}^H &= \frac{1}{H} n_{\alpha\beta}\end{aligned}$$

The Hubble-normalized variables have the nice property of being **constant for self-similar solutions**. (A self-similar solution is a scaled version of itself at anytime.)

Gauge choice is to choose timelike congruence orthogonal to the spatially homogeneous slices, and  $\mathcal{N}^H = 1$ . This leads to  $\dot{u}_\alpha = 0$ . Orthonormal frame alignment earlier gives  $R_1^H = -\Sigma_{23}^H$ ,  $R_2^H = -\Sigma_{31}^H$ ,  $R_3^H = \Sigma_{12}^H$ .

$(E_\alpha^i)^H$  decouple from the ODEs.  $H$  also decouples.

Reduced state space variables are  $\Sigma_{\alpha\beta}^H$ ,  $A_\alpha^H$ ,  $N_{\alpha\beta}^H$ .

This means that the self-similar solutions are represented by **equilibrium points** in the reduced state space of Hubble-normalized variables.

# Spatially homogeneous solutions

These are models with (at least) 3 spacelike Killing vector fields acting on 3D manifolds. As a result, the state space variables  $\Sigma_{\alpha\beta}^H, A_{\alpha}^H, N_{\alpha\beta}^H$  are functions of time only.

Spatially homogeneous solutions play an important role in describing asymptotically local dynamics.

# Spatially homogeneous solutions are classified into different Bianchi types

In 1898, Luigi Bianchi classified three-dimensional Lie groups of isometries of a Riemannian manifold.

Bianchi class	Bianchi type	Eigenvalues of $n_{\alpha\beta}$		
A ( $a_\alpha$ is zero)	I	0	0	0
	II	0	0	+
	VI <sub>0</sub>	0	+	-
	VII <sub>0</sub>	0	+	+
	VIII	-	+	+
	IX	+	+	+
B ( $a_\alpha$ is nonzero)	V	0	0	0
	IV	0	0	+
	VI <sub>h</sub>	0	+	-
	VII <sub>h</sub>	0	+	+

Bianchi VI<sub>h</sub> has  $h < 0$  while Bianchi VII<sub>h</sub> has  $h > 0$ . Bianchi III is Bianchi VI<sub>-1</sub>. Bianchi VI<sub>-1/9</sub> has an exceptional class denoted Bianchi VI<sub>-1/9</sub>\*

# The Kasner solution is a self-similar solution

For the Kasner solution,

$$\Sigma_{\alpha\beta}^H = \text{diag}(-2\Sigma_+^H, \Sigma_+^H + \sqrt{3}\Sigma_-^H, \Sigma_+^H - \sqrt{3}\Sigma_-^H).$$

where  $(\Sigma_+^H, \Sigma_-^H)$  are constant and satisfy  $\Sigma_+^{H^2} + \Sigma_-^{H^2} = 1$ . All other Hubble-normalized variables are zero.

In the Hubble-normalized state space, each Kasner solution is represented by an **equilibrium point** on the unit circle  $\Sigma_+^{H^2} + \Sigma_-^{H^2} = 1$ .

# The Taub solution is asymptotic to Kasner solutions

For the Taub solution,

$$\begin{aligned}\Sigma_{\alpha\beta}^H &= \text{diag}(-2\Sigma_+^H, \Sigma_+^H + \sqrt{3}\Sigma_-^H, \Sigma_+^H - \sqrt{3}\Sigma_-^H), \\ N_{\alpha\beta}^H &= \text{diag}(N_{11}^H, 0, 0).\end{aligned}$$


where  $(\Sigma_+^H, \Sigma_-^H, N_{11}^H)$  satisfy  $\Sigma_+^{H^2} + \Sigma_-^{H^2} + \frac{1}{12}(N_{11}^H)^2 = 1$ . All other Hubble-normalized variables are zero. The explicit expression for  $(\Sigma_+^H, \Sigma_-^H, N_{11}^H)$  is not needed here.

In the Hubble-normalized state space, each Taub solution is represented by an **orbit** connecting two Kasner equilibrium points (one source and one sink).

There are two other orientations  $\text{diag}(0, N_{22}^H, 0)$  and  $\text{diag}(0, 0, N_{33}^H)$ .

# Kasner points and Taub orbits

3 orientations of Taub orbits projected onto the  $(\Sigma_+^H, \Sigma_-^H)$  plane.  
Arrows indicate evolution towards singularity.




The Kasner points represent Kasner **epochs** and the Taub orbits represent the transitions between two Kasner epochs.

# Vacuum Bianchi VIII and IX – the original BKL archetype

For Bianchi VIII and IX spatially homogeneous solutions,  $n_{\alpha\beta}$  has 3 non-zero eigenvalues and their eigenvectors do not rotate.

All Kasner equilibrium points are saddle points.



A Bianchi VIII or IX BKL orbit is built from consecutive Taub orbits connecting the Kasner points.

The BKL regime consists of quick Taub transitions between long Kasner epochs.

A sequence of Kasner epochs with alternatingly active pair of  $n_{\alpha\beta}$  eigenvalues defines a Kasner era.

# The BKL index $u$ tracks the Kasner epochs within each Kasner era

BKL introduced an index  $u$  that characterizes a Kasner solution that is independent of spatial frame orientation. It can be defined implicitly through

$$\det \Sigma_{\alpha\beta}^H = 2 - \frac{27(1+u)(1+\frac{1}{u})}{(1+u+\frac{1}{u})^3}.$$

$u$  satisfies  $u \geq 1$ .

The Kasner map for each Taub transition is  $u \rightarrow u - 1$  if  $u \geq 2$ .

A Kasner era ends when  $1 \leq u < 2$ .

Then the next Kasner era begins with the map  $u \rightarrow \frac{1}{u-1}$ .

## Vacuum Bianchi VI and VII – a single Kasner era


Vacuum Bianchi VI and VII can only sustain the BKL oscillation for a single Kasner era, terminating at a final Kasner epoch.

$n_{\alpha\beta}$  has 2 non-zero eigenvalues and their eigenvectors rotates on one axis every Kasner epoch.


The spatial frame has to rotate with the eigenvectors, so that the circle of Kasner points remain on the  $(\Sigma_+^H, \Sigma_-^H)$  plane.

This introduces a **frame transition**, whose orbit is a vertical line when projected onto the  $(\Sigma_+^H, \Sigma_-^H)$  plane.



# Bianchi VI<sub>0</sub>'s Taub transition orbits




# Bianchi VI<sub>0</sub>'s frame transition orbits



# Bianchi VI<sub>0</sub>'s Taub and frame transition orbits projected



# Bianchi VI<sub>0</sub> orbit during a Kasner era



# Vacuum Bianchi VI $^*_{-1/9}$ – the other BKL archetype

[Hewitt, Horwood & Wainwright 2003]

Vacuum Bianchi VI $^*_{-1/9}$  can sustain the BKL oscillation indefinitely.


$n_{\alpha\beta}$  has 2 non-zero eigenvalues and their eigenvectors rotate on 2 axes – one rotation occurs once every Kasner epoch, and the other rotation occurs only once between two Kasner eras.

The spatial frame has to rotate with the eigenvectors to maintain the circle of Kasner points on the  $(\Sigma^H_+, \Sigma^H_-)$  plane.


This introduces two frame transitions.

Both BKL archetypes differ in details but the Kasner map is the same:  
 $u \rightarrow u - 1$ .


# Bianchi VI $^*_{-1/9}$ 's Taub transition orbits




# Bianchi VI $^*_{-1/9}$ 's frame transition orbits




# Bianchi VI<sup>\*</sup><sub>-1/9</sub>'s second frame transition orbits



# Bianchi VI $^*_{-1/9}$ 's Taub and frame transition orbits projected



# Bianchi $VI_{-1/9}^*$ 's BKL orbit



Having understood BKL dynamics in spatially homogeneous models, we now turn to BKL dynamics in spatially **inhomogeneous** models.

# OT $G_2$ class of inhomogeneous spacetimes

## OT $G_2$ spacetimes

- ▶ are vacuum models
- ▶ admit 2 Abelian Killing vector fields (spacelike,  $G_2$  group action is orthogonally transitive)
- ▶ contain Bianchi VI<sub>0</sub> as spatially homogeneous case.

This class can sustain the BKL dynamics for only a single Kasner era.

Surprise inhomogeneous feature: **spikes**.

# Discovery of spikes in OT $G_2$ spacetimes

Berger & Moncrief 1993 carried out numerical study of Gowdy spacetimes and found that for certain initial conditions, **small-scale spatial structures** develop on the final approach to the singularity. These turn out to be **permanent spikes**.

There are isolated worldlines along which the BKL index  $u$  tends to  $u > 2$  for their final Kasner epoch, while along other typical worldlines  $u$  tends to  $u < 2$ .

i.e.  $u$  tends to its limit pointwise but not uniformly.

# Transient spikes in OT $G_2$ spacetimes

Furthermore, before the final approach, at these same isolated worldlines, **transient spikes** form, and they smooth out after two Kasner epochs, and then they form again at the next Kasner epoch.

Q: What is so special about these isolated worldlines?

Two possible explanations:

1. The active eigenvalue of  $n_{\alpha\beta}$  changes sign here.
2. The trace of  $n_{\alpha\beta}$  changes sign here.

## Transient spikes in general (non-OT) $G_2$ spacetimes

In general (non-OT)  $G_2$  spacetimes, an additional shear component is non-zero. This provides sufficient degree of freedom to the rotation of the  $n_{\alpha\beta}$  eigenvector to sustain an infinite sequence of BKL dynamics.

The BKL orbits are similar to those in Bianchi VI $^*_{-1/9}$ .

All spikes are transient here. The permanent OT  $G_2$  spike is a result of unfinished spike transition.

# Exact spike solutions


The spike transitions are described by the exact spike solutions, which were discovered in two stages.

The OT  $G_2$  spike solution [Lim 2008] was found by applying the Rendall-Weaver transformation on a Kasner seed solution.


The non-OT  $G_2$  spike solution [Lim 2015] was found by applying the Geroch transformation on a Kasner seed solution.

# OT $G_2$ spike orbits


$w=0.5$




$w=1$



$w=1.5$




$w=3.5$




# OT $G_2$ spike orbits

$\tau = -2.2$




# OT $G_2$ spike orbits

$\tau = -1.1$




# OT $G_2$ spike orbits

$\tau = -0.1$




# OT $G_2$ spike orbits

$\tau=0.1$




# OT $G_2$ spike orbits

$\tau=0.6$




# OT $G_2$ spike orbits

$\tau=1.4$




# OT $G_2$ spike orbits

$\tau=3.4$




# OT $G_2$ spike orbits

$\tau=5$




# non-OT $G_2$ spike orbits

$$\tau = -3.7$$




# non-OT $G_2$ spike orbits

$$\tau = -3.5$$




# non-OT $G_2$ spike orbits

$$\tau = -1.5$$




# non-OT $G_2$ spike orbits

$$\tau = -0.4$$




# non-OT $G_2$ spike orbits

$$\tau = 0.1$$




# non-OT $G_2$ spike orbits

$$\tau = 5$$



# non-OT $G_2$ spike orbits

$$\tau = 10$$



# Quantum fates?

Towards spacelike singularities, as the spacetime reaches the quantum regime,

most common: Kasner saddle states

common: Taub transitions

rare: transient OT  $G_2$  (same-Kasner-era) spike transitions

rarer: non-OT  $G_2$  (inter-Kasner-era) spike transitions

Try to quantize these solutions to find out their quantum fates.