

Beams and Scaffolds - The art of building modular Garside groups

Carsten Dietzel



Alexander von Humboldt
Stiftung/Foundation

LOOPS 23

Bedlewo - June 30th, 2023

- 1 Introduction
- 2 Modular quasi-Garside groups

Introduction

Notation

Let (L, \leq) be a poset and $A \subseteq L$. For $z \in L$, write

- $z = \bigwedge A$, if for all $y \in L$, we have the equivalence

$$(\forall x \in A : y \leq x) \Leftrightarrow y \leq z.$$

We call z the **meet** of A .

- $z = \bigvee A$, if for all $y \in L$, we have the equivalence

$$(\forall x \in A : y \geq x) \Leftrightarrow y \geq z.$$

We call z the **join** of A .

Definition: (Bounded) Lattice

Let L be a poset.

- L is a **lattice** if for all $x, y \in L$, the elements $x \vee y = \bigvee \{x, y\}$ and $x \wedge y = \bigwedge \{x, y\}$ exist.
- L **bounded from above** (resp. **below**) if $1_L = \bigwedge \emptyset$ (resp. $0_L = \bigvee \emptyset$) exist.
- L is **bounded**, if L is bounded from above and below.

Definition: Right ℓ -group

A **right-ordered group** is a group G with a partial order \leq such that for all $x, y, z \in G$,

$$x \leq y \Rightarrow xz \leq yz.$$

If (G, \leq) is a lattice, G is a **right ℓ -group**.

The **negative cone** of G is $G^- = \{x \in G : x \leq e\}$.

Definition: Strong order unit

Let G be a right ℓ -group. An element $s \in G$ is a **strong order unit**, if

- $x \leq y \Leftrightarrow sx \leq sy$ holds for all $x, y \in G$,
- for each $g \in G$ there is a $k \in \mathbb{Z}$ such that $s^k \geq g$.

The interval $[s^{-1}, e] = \{x \in G : s \leq x \leq e\}$ is the respective **strong order interval**.

Definition: Noetherian

A right ℓ -group G is **noetherian** if every sequence $x_1 \leq x_2 \leq \dots$, all $x_i \leq e$, becomes stationary and every sequence $y_1 \geq y_2 \geq \dots$, all $y_i \geq e$, becomes stationary.

Definition: (Quasi-)Garside group

- A **quasi-Garside group** is a noetherian right ℓ -group with a strong order unit s .
- A **Garside group** is a quasi-Garside group with a *finite* strong order interval $[s^{-1}, e]$.

Examples of Garside groups: Spherical Artin-Tits groups, \mathbb{Z}^n, \dots

Some bad problems

- **Classify all Garside groups!** 😞
- **Find all Garside structures on a given torsion-free group!** 😞

Some better problems

- **Find all Garside groups with a given lattice structure!** 😊
- **Classify all Garside groups whose lattices fulfill certain identities!** 😊

A Garside-theorist's favorite lattice identities!

- **Distributivity:** $x \wedge (y \vee z) = (x \wedge y) \vee (x \wedge z)$
- **Modularity:** $x \leq z \Rightarrow x \vee (y \wedge z) = (x \vee y) \wedge z$

- Distributive lattices: $(\mathcal{P}(X), \subseteq), (\mathbb{Z}^n, \leq), \dots$
- Modular lattices: $L(R, n) = \{R\text{-submodules of } R^n\}, R \text{ unital, under } \subseteq, \dots$
- Distributivity \Rightarrow Modularity.

Definition: Cycle set

A **nondegenerate cycle set** (X, \cdot) is a (finite) set with a binary operation $(x, y) \mapsto x \cdot y$, such that

$$(x \cdot y) \cdot (x \cdot z) = (y \cdot x) \cdot (y \cdot z) \quad \forall x, y, z \in X$$

the maps $\sigma_x(y) = x \cdot y$ are bijective for all $x \in X$

the map $x \mapsto x \cdot x$ is bijective.

The **structure group** of a cycle set X is

$$G(X) = \langle X \mid (x \cdot y)x = (y \cdot x)y \rangle.$$

- Non-degenerate cycle sets are equivalent to set-theoretic solutions to the Yang-Baxter equation,
- $G(X)$ is equivalent to the structure group of a set-theoretic solution.

Theorem [Chouraqui (2011), Rump (2015)]

- Let X be a nondegenerate cycle set. Then the submonoid $G^- = \langle X^{-1} \rangle \subseteq G(X)$ is the negative cone of a Garside structure on $G(X)$, that has $s = \bigvee X$ as its strong order unit.
- Every distributive Garside group G is of this form.
- There is a lattice isomorphism $G \cong \mathbb{Z}^n$.

Modular quasi-Garside groups

Example: Paraunitary groups

Let $A \in K^{n \times n}$ be such that $v \neq 0 \Rightarrow v^\top A v \neq 0$ for $v \in K^n$ and $A^\top = A$. Then the **pure paraunitary group** associated with A is

$$\text{PPU}(A) = \left\{ M(t) \in K[t, t^{-1}]^{n \times n} : M(t^{-1})^\top A M(t) = A \wedge M(1) = E_n \right\}.$$

Theorem [D. (2019)]

The submonoid $\text{PPU}(A)^- = \text{PPU}(A) \cap K[t^{-1}]^{n \times n}$ is the negative cone of a quasi-Garside structure on $\text{PPU}(A)$.

The element $s = t \cdot E_n$ is a strong order unit and

$$[s^{-1}, e] \cong L(K, n).$$

- For a cyclic field extensions L/K of degree n , one can construct a quasi-Garside group with a strong order unit s such that $[s^{-1}, e] \cong L(K, n)$. This construction uses skew polynomial rings.
- Rings seem to play a role in the construction of modular quasi-Garside groups...

The distributive scaffold

For $x > y$, write $x \succ y$ if $x \geq z \geq y$ implies $z = x$ or $z = y$.

- Let G be a modular quasi-Garside group, where $s = \bigvee \{x \in G : x \succ e\}$ exists.
- There is a finite decomposition $[s^{-1}, e] \cong \prod_{i=1}^k L_i$ into directly irreducible bounded lattices.
- For $1 \leq i \leq k$, let $z_i \in [s^{-1}, e]$ correspond to $\varepsilon^{(i)} = (\varepsilon_j^{(i)})_{1 \leq j \leq k}$ where

$$\varepsilon_j^{(i)} = \begin{cases} 0_j & j = i \\ 1_j & j \neq i \end{cases}$$

and set $\mathcal{Z} = \{z_i : 1 \leq i \leq k\}$.

Theorem [D. (2023)]

\mathcal{Z} has the structure a nondegenerate cycle set such that the (well-defined) group homomorphism $G(\mathcal{Z}) \rightarrow G$ is an embedding of $\mathcal{D}(G) = G(\mathcal{Z})$ as a distributive sublattice.

Definition: Distributive scaffold

The subgroup $\langle \mathcal{Z} \rangle$ is the **distributive scaffold** of G .

Theorem [D. (2023)]

Let G be a modular quasi-Garside group with strong order unit $s = \vee \{x \in G : x \succ e\}$. Let

$$[s^{-1}, e] \cong \prod_{i=1}^k L_i$$

be a decomposition of $[s^{-1}, e]$ into directly irreducible lattices. Then there exist directly irreducible sublattices $\sqsupseteq_i \subseteq G$ ($1 \leq i \leq k$) - the **beams** - such that there is a *lattice-theoretic* decomposition

$$G \cong \prod_{i=1}^k \sqsupseteq_i$$

that induces the decompositions

- $[s^{-1}, e] \cong \prod_{i=1}^k L_i$
- $\mathcal{D}(G) \cong \prod_{i=1}^k \mathbb{Z}$.

⇒ The decomposition of modular quasi-Garside groups is controlled by $\mathcal{D}(G)$, the structure group of a cycle set!

Some facts

Let G be a modular quasi-Garside group with strong order unit $s = \vee \{x \in X : x \succ e\}$. Let $[s^{-1}, e] \cong \prod_{i=1}^k L_i$ be a decomposition into irreducible lattices.

- By a theorem of Rump, the L_i are bounded modular geometric lattices - i.e. isomorphic to $L(K, n)$ for some skew field K or the subspace lattice of a degenerate geometry or a non-desarguesian plane.
- The associated beams \beth_i can be shown to be primary lattices.

Theorem [D. (2023)]

Let $L_i \cong L(K, n)$ for some $n \geq 4$ and a skew field K . For the associated beam \beth_i , there is a noncommutative discrete valuation field Q with valuation ring R , such that there is a lattice-isomorphism

$$\begin{aligned}\beth_i &\cong \text{Lat}(Q, n) \\ &= \{A \subseteq Q^n : A \text{ is a finitely generated, essential } R\text{-submodule}\}.\end{aligned}$$

Isotypical components

Let $G \cong \prod_{i=1}^k \mathfrak{Q}_i$ be the decomposition into beams. Let $i \sim j \Leftrightarrow \mathfrak{Q}_i \cong \mathfrak{Q}_j$.
For $1 \leq i \leq k$, call $\mathcal{C}_i = \prod_{j \sim i} \mathfrak{Q}_j$ the **isotypical component** of \mathfrak{Q}_i .

Theorem [D. (2023)]

- \mathcal{C}_i corresponds to a convex subgroup $G_i \leq G$.
- If $\mathfrak{Q}_i \cong \text{Lat}(Q, n)$, the isotypical subgroup G_i embeds as a subgroup of $\mathcal{S}_m \wr \text{P}\Gamma L(RQ^n)$, m the number of $\mathfrak{Q}_j \cong \mathfrak{Q}_i$, where

$$\text{P}\Gamma L(RQ^n) = (\text{GL}(n, Q) \rtimes \text{Aut}(R)) / R^\times,$$

a generalized projective semilinear group. More precisely,

$$\text{P}\Gamma L(RQ^n) = G \cdot \text{P}\Gamma L(RR^n) \quad ; \quad G \cap \text{P}\Gamma L(RR^n) = 1.$$

- G is a matched product over all isotypical subgroups!

Two of my favorite questions

- Question 1: Can every lattice $L(K, n)$ appear as the strong order interval of a modular quasi-Garside group?
 - Question 2: What about the subspace lattice of a non-desarguesian plane?
-
- Only known answers to Question 1: $L(K, n)$ is realizable if there is an anisotropic hermitian form on K^n or if there is a cyclic field extension L/K of degree n .
In particular, all finite desarguesian geometries $L(\mathbb{F}_q, n)$ are strong order intervals!
 - No answers to Question 2 are known: even for the Hughes plane (91 points and lines), a computational approach fails (even with an insane amount of RAM!).

😊 Thanks for your attention! 😊