


On the attractor of piecewise linear iterated function systems


Dániel Prokaj
Joint with Peter Raith and Károly Simon

Alfréd Rényi Institute of Mathematics

Bedlewo, 16 May 2023



Self-similar IFS
 $\mathcal{S} = \{S_1, S_2, S_3\}$



CPLIFS
 $\mathcal{F} = \{f_1, f_2, f_3\}$

The Main Result

Theorem (Raith, Simon, P.)

For packing dimension typical CPLIFS \mathcal{F}

$$(1) \quad \dim_{\text{H}} \Lambda = \dim_{\text{B}} \Lambda = \min \{1, s^{\mathcal{F}}\}.$$

The meaning of "packing dimension typical": the packing dimension of the parameters of the exceptional CPLIFS is less than the dimension of the parameter space.

Introduction

Markov diagrams

Proof of Theorem 1.4

Limit-irreducibility

The **attractor Λ** of the IFS $\mathcal{F} = \{f_k\}_{k=1}^m$ is the unique non-empty compact set satisfying


$$(2) \quad \Lambda = \bigcup_{k=1}^m f_k(\Lambda).$$


Let I be the smallest non-empty compact interval such that **$f_i(I) \subset I$** for all $i \in [m] := \{1, \dots, m\}$. Then

$$(3) \quad \Lambda = \bigcap_{n=1}^{\infty} \bigcup_{(i_1, \dots, i_n) \in [m]^n} I_{i_1 \dots i_n},$$

where $I_{i_1 \dots i_n} = f_{i_1} \circ \dots \circ f_{i_n}(I)$.

The natural dimension





$$\mathcal{F} = \{f_k\}_{k=1}^m, \tau_k := f_k(0),$$

f_k has $l(k)$ breaking points
 $\{b_{k,1}, \dots, b_{k,l(k)}\}$.

The type of \mathcal{F} is

$$\ell = (l(1), \dots, l(m))$$

$$L := l(1) + \dots + l(m).$$

Packing dimension typicality



Fix a type $\ell = (l(1), \dots, l(m))$ and a vector of contractions $\rho \in ((-1, 1) \setminus \{0\})^{L+m}$. Let \mathfrak{P} be a property that makes sense for every CPLIFS, and consider the exceptional set

$$(4) \quad E_{\ell}^{\rho} =: \left\{ (\mathbf{b}, \boldsymbol{\tau}) \in \mathbb{R}^{L+m} : \mathcal{F}^{(\mathbf{b}, \boldsymbol{\tau}, \rho)} \text{ does not have property } \mathfrak{P} \right\}.$$

We say that **property \mathfrak{P} holds $\dim_{\mathbb{P}}$ -typically** if for all type ℓ and for all contraction vector ρ we have

$$(5) \quad \dim_{\mathbb{P}} E_{\ell}^{\rho} < L + m.$$

The generated self-similar IFS



We fix the vector of slopes ρ .

Lemma 1.1

There is a *non-singular linear transformation* F which depends only on ρ such that

$$F_{\rho}(\mathbf{b}, \tau) = \mathbf{t}.$$

Theorem 1.2 (Hochman¹)

$$(6) \quad \dim_{\mathbb{P}} \{ \mathbf{t} \in \mathbb{R}^M : \mathcal{S}^{\mathbf{t}} \text{ does not satisfy the ESC} \} = M - 1.$$

¹Michael Hochman. On self-similar sets with overlaps and inverse theorems for entropy in \mathbb{R}^d , 2015

Corollary 1.3

For a $\dim_{\mathbb{P}}$ -typical CPLIFS \mathcal{F} , the generated self-similar IFS $\mathcal{S}_{\mathcal{F}}$ satisfies the ESC.

Theorem 1.4 (Raith, Simon, P.)

Let \mathcal{F} be a CPLIFS with generated self-similar system $\mathcal{S}_{\mathcal{F}}$ and attractor Λ . If $\mathcal{S}_{\mathcal{F}}$ satisfies the ESC, then

$$(7) \quad \dim_{\mathbb{H}} \Lambda = \dim_{\mathbb{B}} \Lambda = \min \{1, s^{\mathcal{F}}\}.$$

Theorem 1.5 (Raith, Simon, P.)

Fix a type ℓ and a **slope vector ρ with positive entries**. For \mathcal{L}_{m+L} -almost every $(\mathbf{b}, \boldsymbol{\tau}) \in \mathfrak{B}^\ell \times \mathbb{R}^m$ we have

$$(8) \quad s^{\mathcal{F}} > 1 \implies \mathcal{L}_1(\Lambda^{(\mathbf{b}, \boldsymbol{\tau})}) > 0,$$

where $\Lambda^{(\mathbf{b}, \boldsymbol{\tau})}$ denotes the attractor of $\mathcal{F}^{(\rho, \mathbf{b}, \boldsymbol{\tau})}$.

Introduction

Markov diagrams

Proof of Theorem 1.4

Limit-irreducibility

Let $I_k := f_k(I)$ and $\mathcal{I} = \cup_{k=1}^m I_k$. We define the expanding multi-valued mapping associated to \mathcal{F} as

$$(9) \quad T : \mathcal{I} \mapsto \mathcal{P}(\mathcal{P}(I))$$

$$(10) \quad T(y) := \left\{ \left\{ x \in I : f_k(x) = y \right\} \right\}_{k=1}^m.$$

For $k \in [m]$, $j \in [l(k) + 1]$, we define $f_{k,j} : J_{k,j} \mapsto I_k$ as the unique linear function that satisfies $f_k(x) = f_{k,j}(x)$, $\forall x \in J_{k,j}$.

We refer to the linear functions

$$\forall k \in [m], \forall j \in [l(k) + 1] : f_{k,j}^{-1}$$



as the branches of T .

We define the set of critical points as

$$\mathcal{K} := \bigcup_{k=1}^m \{f_k(0), f_k(1)\} \cup \bigcup_{k=1}^m \bigcup_{j=1}^{l(k)} f_k(b_{k,j}) \cup$$


$$\{x \in \mathcal{I} \mid \exists k_1, k_2 \in [m], \exists j_1 \in [l(k_1)], \exists j_2 \in [l(k_2)] : f_{k_1, j_1}^{-1}(x) = f_{k_2, j_2}^{-1}(x)\}$$

The associated multi-valued mapping



We call the partition of \mathcal{I} into closed intervals defined by the set of critical points \mathcal{K} the **monotonicity partition \mathcal{Z}_0** of \mathcal{F} . We call its elements **monotonicity intervals**.

That is, above monotonicity intervals T is always linear, and branches can only take the same value at the boundary.



Let $Z \in \mathcal{Z}_0$. We write $Z \rightarrow D$ for the **successors** of Z .

$$\exists Z_0 \in \mathcal{Z}_0, Z' \in T(Z) :$$

$$D = Z_0 \cap Z'$$

Further, we write $Z \rightarrow_{k,j} D$ if

$$\exists Z_0 \in \mathcal{Z}_0 : D = Z_0 \cap f_{k,j}^{-1}(Z).$$

The set of successors of Z is

$$w(Z) := \{D \mid Z \rightarrow D\}.$$

Following Hofbauer and Raith, we say that $(\mathcal{D}, \rightarrow)$ is the Markov Diagram of \mathcal{F} with respect to \mathcal{Z}_0 if \mathcal{D} is the smallest set containing \mathcal{Z}_0 such that $\mathcal{D} = w(\mathcal{D})$.

We can similarly define the Markov diagram of \mathcal{F} with respect to any finite partition \mathcal{Z}'_0 of \mathcal{I} .

One can imagine the Markov diagram as a (potentially infinitely big) directed graph, with vertex set \mathcal{D} .

Between $C, D \in \mathcal{D}$, we have a directed edge $C \rightarrow D$ if and only if $D \in w(C)$. We call the Markov diagram **irreducible** if there exists a directed path between any two intervals $C, D \in \mathcal{D}$.

Since the functions of a CPLIFS are always continuous on \mathbb{R} , we can always assume that $(\mathcal{D}, \rightarrow)$ is irreducible.

Associated matrix

We define the matrix $\mathbf{F}(s) := \mathbf{F}_{\mathcal{D}}(s)$ indexed by the elements of \mathcal{D} as

$$(11) \quad [\mathbf{F}(s)]_{C,D} := \begin{cases} \sum_{(k,j):C \rightarrow (k,j)D} |f'_{k,j}|^s, & \text{if } C \rightarrow D \\ 0, & \text{otherwise.} \end{cases}$$

This matrix is often associated to self-similar graph directed iterated function systems. When the diagram is finite, our system is actually a self-similar GDIFS.

Let $\mathcal{C} \subset \mathcal{D}$. We write $\mathcal{E}_{\mathcal{C}}(n)$ for the set of n -length directed paths in the subgraph $(\mathcal{C}, \rightarrow)$.

Assume that $(\mathcal{C}, \rightarrow)$ is irreducible. Each path in $(\mathcal{C}, \rightarrow)$ of infinite length represents a point in the invariant set $\Lambda_{\mathcal{C}} \subset \Lambda$. We define the natural pressure of these sets as

$$(12) \quad \Phi_{\mathcal{C}}(s) := \limsup_{n \rightarrow \infty} \frac{1}{n} \log \sum_{\mathbf{k}} |I_{\mathbf{k}}|^s,$$

where the sum is taken over all $\mathbf{k} = (k_1, \dots, k_n)$ for which $\exists j_1, \dots, j_n : ((k_1, j_1), \dots, (k_n, j_n)) \in \mathcal{E}_{\mathcal{C}}(n)$.

As an operator, $(\mathbf{F}_{\mathcal{D}}(s))^n$ is always bounded in the l^∞ -norm. Thus we can define

$$\varrho(\mathbf{F}_{\mathcal{C}}(s)) := \lim_{n \rightarrow \infty} \|(\mathbf{F}_{\mathcal{C}}(s))^n\|_\infty^{1/n}.$$

Lemma 2.1

Let $\mathcal{C} \subset \mathcal{D}$. If $(\mathcal{C}, \rightarrow)$ is **irreducible**, then

$$(13) \quad \Phi_{\mathcal{C}}(s) \leq \log \varrho(\mathbf{F}_{\mathcal{C}}(s)).$$

If $(\mathcal{C}, \rightarrow)$ is **irreducible and finite**, then

$$(14) \quad \Phi_{\mathcal{C}}(s) = \log \varrho(\mathbf{F}_{\mathcal{C}}(s)).$$

Introduction

Markov diagrams

Proof of Theorem 1.4

Limit-irreducibility

Proof of Theorem 1.4

We need to approximate the Markov diagram of the CPLIFS with finite subdiagrams.

Since $\mathbf{F}(s)$ is always irreducible, according to Seneta's results², it can be done if our CPLIFS has the following property.

²Eugene Seneta. *Non-negative matrices and Markov chains*. Springer Science & Business Media, 2006

We say that the CPLIFS \mathcal{F} is **limit-irreducible** if there exists a \mathcal{Y} finite refinement of \mathcal{Z}_0 such that for all $s \in (0, \dim_{\mathbb{H}} \Lambda]$ the matrix **$\mathbf{F}(\mathcal{Y}, s)$ has right and left eigenvectors with nonnegative entries for the eigenvalue $\varrho(\mathbf{F}(\mathcal{Y}, s))$.**

We call this finite partition \mathcal{Y} a **limit-irreducible partition** and $(\mathcal{D}(\mathcal{Y}), \rightarrow)$ a **limit-irreducible Markov diagram** of \mathcal{F} . $\mathbf{F}(\mathcal{Y}, s)$ is the matrix associated to this diagram.

Proof of Theorem 1.4 cont.

Proposition 3.1

Let \mathcal{F} be a limit-irreducible CPLIFS, and let $(\mathcal{D}, \rightarrow)$ be its limit-irreducible Markov diagram. For any $\varepsilon > 0$ there exists a $\mathcal{C} \subset \mathcal{D}$ finite subset such that

$$(15) \quad \varrho(\mathbf{F}(s)) - \varepsilon \leq \varrho(\mathbf{F}_{\mathcal{C}}(s)) \leq \varrho(\mathbf{F}(s)),$$

where $\mathbf{F}(s)$ is the matrix associated to $(\mathcal{D}, \rightarrow)$.

Proof of Theorem 1.4 cont.

As $\dim_{\mathbb{H}} \Lambda \leq s^{\mathcal{F}}$ always holds, we only need to prove the other direction.

Choose an arbitrary $t \in (0, s^{\mathcal{F}})$. By Lemma 2.1

$$0 < \Phi(t) < \log \varrho(\mathbf{F}(t)).$$

According to Proposition 3.1

$$\exists \mathcal{C} \subset \mathcal{D} \text{ finite} : 0 < \log \varrho(\mathbf{F}_{\mathcal{C}}(t)) = \Phi_{\mathcal{C}}(t).$$

Proof of Theorem 1.4 cont.

Theorem 3.2 (Simon, P.³)

Let \mathcal{F} be a self-similar graph directed IFS with attractor Λ and generated self-similar IFS \mathcal{S} . If \mathcal{S} satisfies the ESC, then

$$\dim_{\text{H}} \Lambda = \min\{1, s^{\mathcal{F}}\}.$$

It follows, that $\dim_{\text{H}} \Lambda_{\mathcal{C}} = \min\{s_{\mathcal{C}}, 1\}$, where $s_{\mathcal{C}}$ is the unique root of $\Phi_{\mathcal{C}}(s)$.

³R Dániel Prokaj and Károly Simon. [Piecewise linear iterated function systems on the line of overlapping construction.](#)

Nonlinearity, 35(1):245, 2021

Proof of Theorem 1.4 cont.

$s^{\mathcal{F}} > 1$ implies $\dim_{\mathbb{H}} \Lambda_{\mathcal{C}} = 1$, for a suitable finite and irreducible subdiagram $(\mathcal{C}, \rightarrow)$.

$s^{\mathcal{F}} \leq 1$ implies $s_{\mathcal{C}} \leq 1$ for all $\mathcal{C} \subset \mathcal{D}$.

$$(16) \quad 0 < \Phi_{\mathcal{C}}(t) \implies t < s_{\mathcal{C}} = \dim_{\mathbb{H}} \Lambda_{\mathcal{C}} \leq \dim_{\mathbb{H}} \Lambda,$$

and it holds for any $t \in (0, s^{\mathcal{F}})$. Thus $s^{\mathcal{F}} \leq \dim_{\mathbb{H}} \Lambda$.

Introduction

Markov diagrams

Proof of Theorem 1.4

Limit-irreducibility

Although limit-irreducibility is required in the proof, we do not need to assume that our CPLIFS have this property, as it is already granted by the ESC.

Lemma 4.1 (F. Hofbauer⁴)

Let $\mathcal{F} = \{f_k\}_{k=1}^m$ be a CPLIFS with Markov diagram $(\mathcal{D}, \rightarrow)$ and associated matrix $\mathbf{F}(s)$. If $\mathbf{F}(s)$ can be written in the form

$$\mathbf{F}(s) = \begin{bmatrix} P & Q \\ R & S \end{bmatrix}$$


such that $\varrho(\mathbf{F}(s)) > \varrho(S)$, then \mathcal{F} is limit-irreducible.

⁴Franz Hofbauer. Piecewise invertible dynamical systems. *Probability theory and related fields*, 72(3):359–386, 1986


Lemma 4.1 always applies for systems without overlaps, where all the entries of $\mathbf{F}(s)$ are smaller than 1.

We have to investigate what happens in the overlapping cases, as multiple edges in $(\mathcal{D}, \rightarrow)$ might yield bigger than 1 entries in the associated matrix.

Two types of overlaps




Light overlap



Cross overlap

Light overlaps





By choosing a finite refinement of \mathcal{Z}_0 that has sufficiently small entries, we can easily avoid having multiple edges in the diagram.

Cross overlaps

The case of cross overlaps is more complicated, as they induce nested sequences of intervals for any finite refinement of \mathcal{Z}_0 .

The ESC implies that no crossing point can have a periodic orbit. Thus, $\varrho(S)$ won't grow too big if we use the branch with the largest expansion ratio among the crossing branches instead of the others.

Cross overlaps



Thank you for your attention!