

Quotients of $L_p(0, 1)$ for $0 < p < 1$

by

N. J. KALTON (Swansea) and N. T. PECK (Urbana, Ill.)

Abstract. One of the main results of this paper is a lifting theorem for operators from L_p , $0 < p < 1$, into a quotient space L_p/N . (The theorem is developed separately for L_0 and for L_p , $0 < p < 1$; the hypotheses on N are different in the two cases.) A corollary is that if N is a non-trivial finite dimensional subspace of L_p , $0 < p < 1$, then L_p/N is not isomorphic to L_p . Several similar results are obtained; at the end of the paper, the idea of a K -space (K_p -space) is introduced and studied in connection with the lifting theorems.

1. Introduction. Let $L_0 = L_0[0, 1]$ be the space of all real (or complex) measurable functions on $[0, 1]$ with the topology of convergence in measure. A. Pełczyński has asked whether the quotient of L_0 by a non-trivial finite-dimensional subspace is isomorphic to L_0 . In this paper we prove a lifting theorem for operators on L_0 ; using this theorem, we can show that if B is a non-trivial closed subspace of L_0 which is either locally bounded or which admits a continuous linear functional, then $L_0/B \not\cong L_0$. Parallel results are developed for the spaces L_p ($0 < p < 1$), where again we have that the quotient of L_p by a non-trivial finite-dimensional subspace cannot be isomorphic to L_p (contrasting of course with the case $p = 1$).

In Section 2, we show from certain general considerations that for $0 < p < 1$, $L_p/V \cong L_p/W$ whenever $\dim V = \dim W < \infty$. This enables us to define (L_p/n) to be the (unique) space obtained by forming the quotient of L_p by a subspace of dimension n . In Section 3 we prove our main lifting theorems and in Section 4 we apply them to show that $(L_p/n) \cong (L_p/m)$ if and only if $m = n$. We conclude Section 4 by giving an example of two isomorphic locally bounded subspaces of L_0 , B_1 and B_2 , such that $L_0/B_1 \not\cong L_0/B_2$.

In Section 5 we develop the idea of a K -space; this is an F -space X such that every short exact sequence of F -spaces $0 \rightarrow R \rightarrow Y \rightarrow X \rightarrow 0$ splits. Using this notion we show that $L_0/N \cong L_0$ implies that N has no non-zero continuous linear functionals. Similar ideas for p -Banach spaces are also developed.

Throughout this paper an F -space will mean a complete metric topological vector space. An F -norm $x \mapsto \|x\|$ on a space X is a mapping from X to \mathbb{R}_+ such that

- (a) $\|x+y\| \leq \|x\| + \|y\|$ if $x, y \in X$,
- (b) $\|\lambda x\| \leq \|x\|$ if $|\lambda| \leq 1$ and if $x \in X$,
- (c) $\|\lambda x\| \rightarrow 0$, as $\lambda \rightarrow 0$ for each $x \in X$,
- (d) $\|x\| = 0$ if and only if $x = 0$.

For $0 < p \leq 1$, a p -Banach space is an F -space with an F -norm $\|\cdot\|$ such that

- (e) $\|\lambda x\| = |\lambda|^p \|x\|$ for all λ and $x \in X$.

If X and Y are p -Banach spaces and if $S: X \rightarrow Y$ is a continuous linear operator, we define

$$\|S\| = \sup \{\|Sx\| : \|x\| \leq 1\}.$$

We denote by $\mathcal{L}(X)$ the space of all linear operators on X . If X is a p -Banach space, then so is $\mathcal{L}(X)$; if X is an F -space, then $\mathcal{L}(X)$ has, in general, no convenient F -norm topology. Unless otherwise stated, "linear map" and "linear operator" always refer to *continuous* maps.

We would like to thank Leonard Dor for several valuable conversations.

2. Transitive F -spaces. In this section we show that if V and W are two subspaces of L_p ($0 \leq p < 1$) of the same finite dimension, then $L_p/V \cong L_p/W$. We approach this result through some general results on transitive F -spaces. An F -space X is said to be *transitive* if given $x, y \in X$ with $x \neq 0$, there exists $T \in \mathcal{L}(X)$ with $Tx = y$. We shall say that X is *strictly transitive*, if for any $k \in \mathbb{N}$, $x_1, \dots, x_k \in X$ and $y_1, \dots, y_k \in X$ such that $\{x_1 \dots x_k\}$ is linearly independent, there exists $T \in \mathcal{L}(X)$ with $Tx_i = y_i$.

We do not know whether a transitive F -space is strictly transitive; however, it is possible to generalize standard arguments in Banach algebra theory (cf. Rickart [6], pp. 60–62) to yield the following:

PROPOSITION 2.1. *Let X be a transitive F -space; suppose that*

- (a) X is separable,
- (b) *The centre of $\mathcal{L}(X)$ consists only of scalar multiples of the identity operator.*

Then X is strictly transitive.

If X is a p -Banach space, condition (a) may be omitted; if X is a complex p -Banach space, then conditions (a) and (b) may be omitted.

Proof. By [6], Lemma 2.4.3, it is enough to show that given two linearly independent elements $v, w \in X$, there exists $T \in \mathcal{L}(X)$ such that

$Tv = 0$ and $Tw \neq 0$. If not, we may define a (not necessarily continuous) operator D on X by $Dx = Tw$ when $Tv = x$ (cf. [6], Theorem 2.4.6). Then D commutes with each $T \in \mathcal{L}(X)$.

It is necessary to show that $D \in \mathcal{L}(X)$; at this point we require condition (a) in general. Consider $\mathcal{L}(X)$ with the topology of pointwise convergence. Then $\mathcal{L}(X)$ is a Souslin space and by the Open Mapping Theorem, the map $\varepsilon_v: \mathcal{L}(X) \rightarrow X$ defined by $\varepsilon_v(T) = Tv$ is open. Since $D \circ \varepsilon_v = \varepsilon_w$, it follows that $D \in \mathcal{L}(X)$. If X is a p -Banach space, then so is $\mathcal{L}(X)$ with its usual topology and the Open Mapping Theorem may be applied to this topology.

Now by condition (b), D is a multiple of I and we have a contradiction. If X is a complex p -Banach space, then it may be shown that the centre of $\mathcal{L}(X)$ is a field and by Żelazko's extension of the Gelfand–Mazur theorem [11], condition (b) must hold.

It is easy to check that each of the spaces L_p ($p \geq 0$) satisfies conditions (a) and (b) of the proposition and is transitive (use an argument similar to [7], pp. 253–254; see also [5]), and hence is strictly transitive.

PROPOSITION 2.2. *Suppose X is a strictly transitive F -space and $X \cong X \oplus X$; then if $\{x_1 \dots x_n\}$ and $\{y_1 \dots y_n\}$ are two linearly independent sets in X , there exists an invertible $T \in \mathcal{L}(X)$ such that $Tx_i = y_i$, $1 \leq i \leq n$.*

Proof. First we prove that there exists a projection $P \in \mathcal{L}(X)$ such that $P(X) \cong X$, $(I - P)(X) \cong X$ and $\{Px_1 \dots Px_n\}$ is linearly independent. Let F be the linear span of $\{x_1 \dots x_n\}$; then we may choose a projection P so that $P(X) \cong X$, $(I - P)(X) \cong X$ and $\dim P(F)$ is maximal. Since $(I - P)(X) \cong X \cong X \oplus X$, there exists a projection $Q \in \mathcal{L}(X)$ such that $PQ = QP = 0$ and $Q(X) \cong (I - P - Q)(X) \cong X$. Then since $(P + Q)(X) \cong (I - P - Q)(X) \cong X$, we have $\dim (P + Q)(F) = \dim P(F)$. Hence P is one-one on $(P + Q)(F)$ and so if we have $\sum a_i Px_i = 0$, then also $\sum a_i (P + Q)x_i = 0$. Similarly we have $\sum a_i (I - Q)x_i = 0$; combining, $\sum a_i x_i = 0$ and $a_1 = a_2 = \dots = a_n = 0$, i.e. $\{Px_1, \dots, Px_n\}$ is linearly independent.

Now pick projections P_1 and $P_2 \in \mathcal{L}(X)$ so that $P_1(X) \cong (I - P_1)(X) \cong P_2(X) \cong (I - P_2)(X) \cong X$ and $\{P_1 x_1 \dots P_1 x_n\}, \{P_2 y_1 \dots P_2 y_n\}$ are linearly independent. Then there exists an invertible $T \in \mathcal{L}(X)$ such that $TP_1 = (I - P_2)T$. Since X is strictly transitive, there exists $S: (I - P_2)(X) \rightarrow P_2(X)$ such that $S(I - P_2)Tx_i = P_2(y_i - Tx_i)$, for $1 \leq i \leq n$. Similarly, there exists $R: P_2(X) \rightarrow (I - P_2)(X)$ so that $RP_2 y_i = (I - P_2)(y_i - Tx_i)$. Then $(I + R \circ P_2)$ and $(I + S \circ (I - P_2))$ are invertible, since $(R \circ P_2)^2 = (S \circ (I - P_2))^2 = 0$ and $(I + R \circ P_2)(I + S \circ (I - P_2))Tx_i = y_i$ ($1 \leq i \leq n$).

We now have:

THEOREM 2.3. *If $0 \leq p < 1$ and V and W are two subspaces of L_p with $\dim V = \dim W < \infty$, then $L_p/V \cong L_p/W$.*

Proof. This is immediate, since there is an invertible operator T on L_p such that $T(V) = W$.

Let us denote now by (L_p/n) the quotient of L_p by an n -dimensional subspace; of course $(L_p/0) = L_p$. Theorem 2.3 guarantees that (L_p/n) is well defined.

THEOREM 2.4. For $0 \leq p < 1$, $(L_p/(m+n)) \cong (L_p/m) \oplus (L_p/n)$ when $m, n > 0$.

Proof. Let V be a subspace of $L_p[0, \frac{1}{2}]$ of dimension m (embedded in L_p in the obvious way) and W a subspace of $L_p[\frac{1}{2}, 1]$ of dimension n . Then

$$\begin{aligned} L_p/(m+n) &\cong L_p[0, 1]/(V+W) \\ &\cong (L_p[0, \tfrac{1}{2}]/V) \oplus (L_p[\tfrac{1}{2}, 1]/W) \cong (L_p/m) \oplus (L_p/n). \end{aligned}$$

3. Lifting theorems. Let X be a p -Banach space ($0 < p \leq 1$) and N a closed subspace of X . It is easy to see that any linear operator $S: L_p \rightarrow X/N$ may be lifted to a linear operator $\tilde{S}: L_p \rightarrow X$, so that $\pi\tilde{S} = S$ where $\pi: X \rightarrow X/N$ is the quotient map. In the case $p = 1$, a similar lifting property holds if L_p is replaced by any \mathcal{L}_1 -space and N is isomorphic to a complemented subspace of a dual space (this is effectively proved by Lindenstrauss [4]). Not surprisingly there is a corresponding result for the case $p < 1$. We say that a p -Banach space Y is an \mathcal{L}_p -space ($0 < p < 1$), if there is an increasing net $\{Y_\alpha: \alpha \in A\}$ of finite-dimensional subspaces of Y such that $\bigcup (Y_\alpha: \alpha \in A)$ is dense in Y and there exist linear maps $S_\alpha: Y_\alpha \rightarrow l_p^{(n_\alpha)}$ and $T_\alpha: l_p^{(n_\alpha)} \rightarrow Y_\alpha$ (where $n_\alpha = \dim Y_\alpha$), such that $\sup_\alpha \|S_\alpha\| \|T_\alpha\| < \infty$ and $T_\alpha S_\alpha = I$ on Y_α . Clearly L_p is an \mathcal{L}_p -space.

We shall also call a p -Banach space Z *pseudo-dual* if there is a Hausdorff vector topology ϱ on Z such that the unit ball is relatively compact for ϱ . The space L_p is not pseudo-dual (see [1]), but the spaces l_p and H_p ($0 < p < 1$) are pseudo-dual.

THEOREM 3.1. Let Y be an \mathcal{L}_p -space ($0 < p \leq 1$) and X a p -Banach space. Let N be a closed subspace of X and suppose N is isomorphic to a complemented subspace of a pseudo-dual p -Banach space Z . Then any operator $S: Y \rightarrow X/N$ may be lifted to an operator $\tilde{S}: Y \rightarrow X$.

If $Y = L_p$, then the lifting is unique.

Proof. We observe that the unit ball of Z may be supposed to be ϱ -compact (by [1], Lemma 1). Then the argument is a straightforward imitation of the Lemma of [4]. We omit the details.

In the case $Y = L_p$, suppose T is any other lifting. Then $T - \tilde{S}$ maps L_p into N and there is a non-zero operator from L_p into Z . The induced map into (Z, ϱ) is compact, contradicting the results of [2].

We now give another result of a similar type for the space L_p .

THEOREM 3.2. Let X be a p -Banach space ($p < 1$) and let N be a closed subspace of X which is isomorphic to a q -Banach space, where $p < q \leq 1$. Then any linear operator $S: L_p \rightarrow X/N$ may be lifted uniquely to a linear operator $\tilde{S}: L_p \rightarrow X$.

Proof. For $n \in \mathbb{N}$, let Y_n be the linear span of the functions χ_k^n ($1 \leq k \leq 2^n$), where χ_k^n is the characteristic function of $((k-1)2^{-n}, k2^{-n})$. Each Y_n is isometric to $l_p^{(2^n)}$. Let $S_n: Y_n \rightarrow X$ be a lift of $S: Y_n \rightarrow X/N$ with $\|S_n\| \leq 2\|S\|$. We shall show that for $y \in \bigcup Y_n$, $\lim_{n \rightarrow \infty} S_n y = \tilde{S}y$ exists. Then clearly $\|\tilde{S}\| \leq 2\|S\|$ and \tilde{S} may be extended to L_p by continuity and is then a lift of S .

Since N is isomorphic to a q -Banach space, there exists a constant C such that the q -convex hull of the unit ball of N is bounded in norm by C . Hence, if x_1, \dots, x_n are in N , then

$$\left\| \sum_{i=1}^n x_i \right\| \leq C \left(\sum_{i=1}^n \|x_i\|^{q/p} \right)^{p/q}.$$

Now suppose $j \leq m \leq n$, and $1 \leq k \leq 2^j$; then

$$\begin{aligned} \|S_m \chi_k^j - S_n \chi_k^j\| &= \left\| (S_m - S_n) \left(\sum_{i=1}^{2^{m-j}} \chi_{2^{m-j}k+i}^m \right) \right\| \\ &\leq C \left(\sum_{i=1}^{2^{m-j}} \|(S_m - S_n) \chi_{2^{m-j}k+i}^m\|^{q/p} \right)^{p/q} \\ &\leq 4C \|S\| \left(\sum_{i=1}^{2^{m-j}} \|\chi_{2^{m-j}k+i}^m\|^{q/p} \right)^{p/q} \\ &= 4C \|S\| 2^{-(jp/q + m(1-p/q))} \end{aligned}$$

and so $\{S_m \chi_k^j; m \geq j\}$ is a Cauchy sequence. Hence we may find the lift \tilde{S} . As in Theorem 3.1, \tilde{S} must be unique, since there are no non-zero operators from L_p into N (see Théorème 3.4.5 of [10] or Proposition 2 of [9], or use the argument above).

We now examine the case $p = 0$, which is rather different. Suppose X is an F -space with F -norm $\|\cdot\|$. For $x \in X$ we define $\sigma: X \rightarrow \mathbb{R} \cup \{\infty\}$, by

$$\sigma(x) = \sup_{t \in \mathbb{R}} \|tx\|.$$

In the case of L_0 with the F -norm

$$\|x\| = \int_0^1 \frac{|x(t)|}{1 + |x(t)|} d\mu(t),$$

we have that $\sigma(x) = \mu(\text{supp } x)$.

In general, note that $\sigma(ax) = \sigma(x)$ if $a \neq 0$ and that $\sigma(x+y) \leq \sigma(x) + \sigma(y)$. If L is a linear subspace of X , we define $\sigma(L) = \sup\{\sigma(x) : x \in L\}$.

We shall say that X admits L_0 -structure, if for any $\varepsilon > 0$ there exist $n = n(\varepsilon)$ and subspaces X_1, \dots, X_n of X , such that $X = X_1 \oplus \dots \oplus X_n$ and $\sigma(X_i) \leq \varepsilon$, $i = 1, 2, \dots, n$. In addition to the obvious example of L_0 itself, any space of the type $L_0(Z)$ (all measurable functions from $[0, 1]$ into an F -space Z) admits L_0 -structure.

The following proposition is trivial.

PROPOSITION 3.3. *Suppose X admits L_0 -structure and B is a locally bounded space. If $T: X \rightarrow B$ is continuous, then $T = 0$.*

We next prove two lemmas before giving the main lifting theorem.

LEMMA 3.4. *Suppose X is an F -space and B is a closed locally bounded subspace of X ; let $\pi: X \rightarrow X/B$ be the quotient map. Let δ be chosen so that the set $\{b \in B: \|b\| \leq \delta\}$ is bounded.*

Then if $\xi \in X/B$ and $\sigma(\xi) \leq \delta/3$, there is a unique $x \in X$ such that $\pi x = \xi$ and $\sigma(x) \leq \delta/3$. For this x , $\sigma(x) = \sigma(\xi)$.

Proof. We can find a sequence $\{x_n\} \in X$ such that $\pi x_n = \xi$ and

$$\|n x_n\| \leq \left(1 + \frac{1}{n}\right) \|n \xi\|, \quad n \in \mathbb{N}.$$

Let $u_n = x_n - x_1$ ($n \in \mathbb{N}$); then $u_n \in B$ and if $m \geq n \geq 2$,

$$\begin{aligned} \|n(u_n - u_m)\| &= \|n(x_n - x_m)\| \\ &\leq \|n x_n\| + \|n x_m\| \\ &\leq \left(1 + \frac{1}{n}\right) \|n \xi\| + \left(1 + \frac{1}{m}\right) \|m \xi\| \\ &\leq \left(2 + \frac{1}{n} + \frac{1}{m}\right) \delta/3 \leq \delta. \end{aligned}$$

By choice of δ , this implies that $\{u_n\}$ is a Cauchy sequence and hence so is $\{x_n\}$. If $x = \lim x_n$, then $\sigma(x) = \sigma(\xi) \leq \delta/3$. If y is any other point satisfying $\pi y = \xi$ and $\sigma(y) \leq \delta/3$, then $x - y \in B$ and $\sigma(x - y) \leq \frac{2}{3} \delta$; this implies $x - y = 0$.

LEMMA 3.5. *Under the assumptions of Lemma 3.4, let Y be a linear subspace of X/B with $\sigma(Y) \leq \frac{1}{3} \delta$. Then there is a linear operator $h: Y \rightarrow X$ such that $\pi \circ h(\xi) = \xi$ for $\xi \in Y$.*

Proof. For $\xi \in Y$, define $h(\xi)$ to be the unique $x \in X$ such that $\pi x = \xi$ and $\sigma(x) = \sigma(\xi)$. If $\alpha, \beta \in \mathbb{R}$ and $\xi, \eta \in Y$, then

$$\sigma(\alpha h(\xi) + \beta h(\eta)) \leq \sigma(\xi) + \sigma(\eta) \leq \delta/3.$$

Thus

$$h(\alpha \xi + \beta \eta) = \alpha h(\xi) + \beta h(\eta),$$

and h is linear.

Now suppose $\xi_n \rightarrow 0$ in Y ; choose $x_n \in X$ such that $\pi x_n = \xi_n$ and $\|x_n\| \leq 2 \|\xi_n\|$. Then $x_n - h(\xi_n) \in B$. If $x_n - h(\xi_n) \rightarrow 0$, we may assume, by passing to a subsequence, that for some $a > 0$ we have

$$\|a(x_n - h(\xi_n))\| \geq \delta$$

(since the set $\{b \in B: \|b\| \leq \delta\}$ is bounded).

Then

$$\|a x_n\| \geq \delta - \|h(a \xi_n)\| \geq \delta - \frac{1}{3} \delta = \frac{2}{3} \delta.$$

This is a contradiction since $\|x_n\| \rightarrow 0$. Hence we have $x_n - h(\xi_n) \rightarrow 0$ and so $h(\xi_n) \rightarrow 0$.

THEOREM 3.6. *Suppose X admits L_0 -structure, Y is an F -space, and B is a closed locally bounded subspace of Y . Then if $S: X \rightarrow Y/B$ is a linear operator, there is a unique linear operator $\tilde{S}: X \rightarrow Y$ such that $\pi \circ \tilde{S} = S$ where $\pi: Y \rightarrow Y/B$ is the quotient map.*

Proof. Choose $\delta > 0$ so that $\{b \in B: \|b\| \leq \delta\}$ is bounded, and then $\varepsilon > 0$, so that if $\|x\| \leq \varepsilon$ ($x \in X$), then $\|Sx\| \leq \delta/6$. Let X_1, \dots, X_n be closed subspaces of X such that $X = X_1 \oplus \dots \oplus X_n$ and $\sigma(X_i) \leq \varepsilon$. Then $\sigma(SX_i) \leq \delta/6$, and so there exist linear operators $h_i: SX_i \rightarrow Y$, such that $\pi h_i(\xi) = \xi$, $\xi \in SX_i$. If we define $\tilde{S}: X \rightarrow Y$ by

$$\tilde{S}(x_1 + \dots + x_n) = \sum_{i=1}^n h_i S x_i, \quad x_i \in X_i,$$

then \tilde{S} is the required lifting of S .

If T is any other lifting, then $\tilde{S} - T$ maps X into B and hence $\tilde{S} = T$ by Proposition 3.3.

4. Quotient spaces of L_p ($0 \leq p < 1$). In this section we treat the case $p = 0$ first and in more detail than the case $p > 0$; the arguments are analogous.

THEOREM 4.1. *Suppose X_1 and X_2 are two F -spaces with L_0 -structure. Suppose B_1 and B_2 are closed locally bounded subspaces of X_1 and X_2 , respectively. Then $X_1/B_1 \cong X_2/B_2$ if and only if there is an isomorphism $V: X_1 \rightarrow X_2$ mapping X_1 onto X_2 and such that $V(B_1) = B_2$.*

Proof. The "if" part is clear. For the "only if" part, let $S: X_1/B_1 \rightarrow X_2/B_2$ be an isomorphism, and let π_1, π_2 be the quotient maps. Then

by Theorem 3.6, there exist lifts V, U of $S\pi_1: X_1 \rightarrow X_2/B_2$ and $S^{-1}\pi_2: X_2 \rightarrow X_1/B_1$.

$$\begin{array}{ccc} X_1 & \xrightleftharpoons[U]{V} & X_2 \\ \pi_1 \downarrow & & \downarrow \pi_2 \\ X_1/B_1 & \xrightarrow{S} & X_2/B_2 \end{array}$$

Then $UV: X_1 \rightarrow X_1$ is a lift of $\pi_1: X_1 \rightarrow X_1/B_1$. By the uniqueness, $UV = I_{X_1}$; similarly $VU = I_{X_2}$, so V is an isomorphism of X_1 onto X_2 . Clearly $V(B_1) \subset B_2$ and $U(B_2) \subset B_1$; hence $V(B_1) = B_2$.

COROLLARY 4.2. *If X admits L_0 -structure and $B \subset X$ is a locally bounded subspace, then X/B admits L_0 -structure if and only if $B = \{0\}$.*

Theorem 2.3 and Theorem 4.1 give

COROLLARY 4.3. *If B_1 and B_2 are locally bounded subspaces of L_0 , then $L_0/B_1 \cong L_0/B_2$ if and only if there is an invertible operator $T: L_0 \rightarrow L_0$ such that $T(B_1) = B_2$.*

In particular, $(L_0/m) \cong (L_0/n)$ if and only if $m = n$, and $(L_0/1) \cong L_0$.

This solves the problem of Pełczyński (see Introduction). Also in this section we shall illustrate this corollary by showing that $B_1 \cong B_2$ does not imply $L_0/B_1 \cong L_0/B_2$. First however, we state the corresponding theorems for $p > 0$; the proofs are similar.

THEOREM 4.4. *Suppose B_1 and B_2 are two closed subspaces of L_p , each of which is either isomorphic to a complemented subspace of a pseudo-dual p -Banach space or to a q -Banach space where $p < q \leq 1$.*

Then $L_p/B_1 \cong L_p/B_2$ if and only if there is invertible operator $T: L_p \rightarrow L_p$ such that $T(B_1) = B_2$.

In particular, $(L_p/m) \cong (L_p/n)$ if and only if $m = n$, and $(L_p/1) \cong L_p$.

THEOREM 4.5. *If $B \subset L_p$ is isomorphic to a complemented subspace of a pseudo-dual p -Banach space and $B \neq \{0\}$, then L_p/B is not an \mathcal{L}_p -space.*

Proof. If L_p/B is an \mathcal{L}_p -space, then the identity map $I: L_p/B \rightarrow L_p/B$ may be lifted to a map $J: L_p/B \rightarrow L_p$. Then on L_p , $I - J\pi$ maps L_p into B . Hence, by applying the results of [2] as in Theorem 3.1, $I = J\pi$ and so $B = \{0\}$.

EXAMPLE. Let $B_1 \subset L_0$ be the closed linear span of the Rademacher functions r_k on $[0, 1]$

$$(r_k(t) = \text{sgn}(\sin 2^k \pi t)),$$

and let B_2 be the closed linear span of a sequence of independent random variables normally distributed with mean zero and variance one. Then $B_1 \cong B_2 \cong L_2$; we shall show, however, that $L_0/B_1 \not\cong L_0/B_2$.

For suppose $L_0/B_1 \cong L_0/B_2$; then there is an invertible linear operator $T: L_0 \rightarrow L_0$ such that $T(B_1) = B_2$. By Kwapiński's Representation Theorem [3], T takes the form

$$Tw(t) = \sum_{i=1}^{\infty} \varphi_i(t) x(\Phi_i t) \text{ a.e.}$$

where

(i) $\varphi_n \in L_0, n \geq 1$,

(ii) $m\{t: \varphi_n(t) \neq 0 \text{ for infinitely many } n\} = 0$,

(iii) Φ_n maps $[0, 1]$ into $[0, 1]$; if A is measurable, then $\Phi_n^{-1}(A)$ is measurable; if $m(A) = 0$, then $m(\Phi_n^{-1}(A) \cap \text{Supp } \varphi_n) = 0$.

Thus for almost every $t \in [0, 1]$, the sequence $\{Tr_k(t)\}$ assumes only finitely many values. Hence for some j, k , with $j \neq k$, we must have $m\{t: Tr_j(t) = Tr_k(t)\} > 0$. However $Tr_j - Tr_k$ is normally distributed and hence $Tr_j = Tr_k$. Thus T is not injective, and we have a contradiction.

Remarks. For $p > 0$, let $x \in L_p$ be non-zero and let V be the linear span of x . Let $\mathcal{L}(L_p)$ and $\mathcal{L}(L_p/V)$ be the p -Banach algebras of all bounded linear operators on L_p and L_p/V , respectively. If $S \in \mathcal{L}(L_p/V)$, let $\hat{S}: L_p \rightarrow L_p$ be the unique lift of $S \circ \pi$. Then the map $S \rightarrow \hat{S}$ is an algebra homomorphism, and in fact an embedding of $\mathcal{L}(L_p/V)$ into $\mathcal{L}(L_p)$. Thus $\mathcal{L}(L_p/V)$ is isomorphic to the closed subalgebra of $\mathcal{L}(L_p)$ consisting of all $T \in \mathcal{L}(L_p)$ such that $Tx \in V$. We may define a multiplicative linear functional φ on $\mathcal{L}(L_p/V)$ by

$$\varphi(S)x = \hat{S}x.$$

5. K -spaces. In this section, we abstract a particular property of the spaces L_p and consider it in more generality. We restrict to the real case, but the complex case is identical.

If X is an F -space, we shall say that X is a K -space if every short exact sequence $0 \rightarrow \mathbf{R} \rightarrow Y \rightarrow X \rightarrow 0$, with Y an F -space, splits. Alternatively, if $S: Y \rightarrow X$ is onto and $\dim S^{-1}(0) = 1$, then there exists an operator $T: X \rightarrow Y$ such that $ST = I_X$.

If X is a p -Banach space ($0 < p \leq 1$), we shall say that X is a K_p -space if every short exact sequence $0 \rightarrow \mathbf{R} \rightarrow Y \rightarrow X \rightarrow 0$, with Y a p -Banach space, splits.

THEOREM 5.1. *An F -space [p -Banach space] X is a K -space [K_p -space] if and only if whenever Y and Z are F -spaces [p -Banach spaces] and $S: Y \rightarrow Z$ is a surjective operator with $\dim S^{-1}(0) = 1$, then each linear operator $T: X \rightarrow Z$ may be lifted to an operator $\hat{T}: X \rightarrow Y$ such that $\hat{S}\hat{T} = T$.*

Proof. We prove the statement for K -spaces. Suppose X is a K -space. Let $V \subset X \oplus Y$ be the subspace of all (x, y) such that $Tx = Sy$,

and define $P: V \rightarrow X$ by $P(x, y) = x$. Then $P: V \rightarrow X$ is surjective, and $\dim P^{-1}(0) = 1$. Hence there exists a linear operator $R: X \rightarrow V$ such that $PR = I_X$. Then $Rx = (x, \tilde{T}x)$; clearly $S\tilde{T} = T$.

For the converse take $Z = X$ and T to be the identity.

We remark now that if X admits L_0 -structure, then X is a K -space (Theorem 3.6), and that an \mathcal{L}_p -space is a K_p -space.

THEOREM 5.2. *If X is an F -space [p -Banach space] and N is a closed subspace of X such that X/N is a K -space [K_p -space], then N has the Hahn-Banach Extension Property in X .*

Proof. Again we restrict to the K -space case. Suppose $\varphi \in N'$ is non-zero; let $M = \varphi^{-1}(0) \subset N$. Consider the natural quotient map $\pi: X/M \rightarrow X/N$; then there is a map $S: X/N \rightarrow X/M$ such that $\pi S = I$ on X/N . Then $S(X/N)$ is a closed subspace of co-dimension one in X/M and so there exists $\psi \in (X/M)'$ such that $\psi \neq 0$ and $\psi \circ S = 0$. If $q: X \rightarrow X/M$ is the natural quotient map, then $\psi q \in X'$. If $x \in N$, then $\psi q(x) = 0$ if and only if $q(x) \in S(X/N)$; then $q(x) = S\pi q(x) = 0$. Thus $(\psi q)^{-1}(0) \cap N = M$ and so a suitable multiple of ψq extends φ .

There is also a converse to Theorem 5.2.

THEOREM 5.3. *If X is a K -space [K_p -space] and $N \subset X$ is a closed subspace with HBEP, then X/N is a K -space [K_p -space].*

Proof. Suppose we have a short exact sequence

$$0 \rightarrow R \rightarrow Z \xrightarrow{\varrho} X/N \rightarrow 0,$$

and let $\pi: X \rightarrow X/N$ be the quotient map. Then there is a lifting of $\pi, S: X \rightarrow Z$, so that $\varrho S = \pi$ (by Theorem 5.1). Suppose first S is not surjective; then $S(X)$ has co-dimension one in Z and $\varrho|S(X)$ is one-one. Define $R: X/N \rightarrow Z$ by $R\xi = z$ where $z \in S(X)$ and $\varrho z = \xi$. If $\xi_n \rightarrow 0$ in X/N , then there exists a sequence $\{x_n\}$ in X such that $x_n \rightarrow 0$ and $\pi x_n = \xi_n$. Since $Sx_n \rightarrow 0$ and $Sx_n = R\xi_n$, $R\xi_n \rightarrow 0$, i.e. R is continuous.

Now suppose S is surjective; then $S^{-1}(0)$ has co-dimension one in N . Let $\varphi \in N'$ be a non-zero linear functional with kernel $S^{-1}(0)$. Then φ may be extended to $\psi \in X'$. Now define $\tilde{S}: X \rightarrow Z$ by $\tilde{S}x = S(x - \psi(x)u)$ where $u \in N$ is chosen so that $\psi(u) = \varphi(u) = 1$. Then $\varrho \tilde{S}x = \pi(x - \psi(x)u) = \pi x$; and $\varrho \tilde{S}x = 0$ implies $x \in N$. Hence $\tilde{S}x = S(x - \varphi(x)u) = 0$. Thus ϱ is one-one on $\tilde{S}(X)$ and $\tilde{S}(X)$ has co-dimension one in Z ; we can apply the previous part of the proof.

COROLLARY 5.4. *X is a K_p -space if and only if $X \cong l_p(I)/N$ where I is some index set and $N \subset l_p(I)$ has the HBEP.*

We remark that if $l_p/N \cong L_p$, then N has HBEP and the extension is unique, since $L'_p = \{0\}$.

COROLLARY 5.5. (i) *If N is a closed subspace of L_0 , then L_0/N is a K -space if and only if $N' = \{0\}$. In particular, if L_0/N has L_0 -structure, then $N' = \{0\}$.*

(ii) *If N is a closed subspace of L_p , then L_p/N is a K_p -space if and only if $N' = \{0\}$. In particular, if L_p/N is an \mathcal{L}_p -space, then $N' = \{0\}$.*

Note here that if we take for N the closed linear span of a sequence of functions with disjoint supports in L_0 , then $N \cong \omega$, and hence $L_0/N \not\cong L_0$. However $L_0/N \cong \omega(L_0/1)$ (the countable product of copies of $L_0/1$); hence $\omega(L_0/1) \not\cong L_0$.

PROBLEM. Is L_p or l_p a K_r -space for any $r < p$, or even a K -space? In particular, is l_1 (or any Banach space) a K_p -space for any $p < 1$? This latter question is essentially the same as a problem of Stiles [8]: if l_p/N is locally convex, must N have the HBEP?

References

- [1] N. J. Kalton, *A note on the spaces L_p ($0 < p < 1$)*, Proc. Amer. Math. Soc., to appear.
- [2] — *Compact and strictly singular operators on Orlicz spaces*, Israel J. Math. 26 (1977), pp. 126–136.
- [3] S. Kwapien, *On the form of a linear operator in the space of all measurable functions*, Bull. Acad. Polon. Sci. 21 (1973), pp. 951–954.
- [4] J. Lindenstrauss, *On a certain subspace of l_p* , Bull. Acad. Polon. Sci. 12 (1964), pp. 539–542.
- [5] D. Pallaschke, *The compact endomorphisms on the metric linear space L_φ* , Studia Math. 47 (1973), pp. 123–133.
- [6] C. E. Rickart, *Banach algebras*, Van Nostrand, Princeton 1960.
- [7] S. Rolewicz, *Metric linear spaces*, PWN, Warszawa 1972.
- [8] W. J. Stiles, *Some properties of l_p , $0 < p < 1$* , Studia Math. 42 (1972), pp. 109–119.
- [9] P. Turpin, *Opérateurs linéaires entre espaces d'Orlicz non localement convexes*, Studia Math. 46 (1973), pp. 153–165.
- [10] — *Convexités dans les espaces vectoriels topologiques généraux*, Diss. Math. 131 (1976).
- [11] W. Zelazko, *On the locally bounded and m -convex topological algebras*, Studia Math. 19 (1960), pp. 333–356.

UNIVERSITY COLLEGE OF SWANSEA
SINGLETON PARK, SWANSEA
UNIVERSITY OF ILLINOIS
URBANA, ILLINOIS

Received October 10, 1976

(1212)