

An L^1 estimate for half-space discrepancy

by

WILLIAM W. L. CHEN (Sydney) and GIANCARLO TRAVAGLINI (Milano)

1. Introduction. The half-space discrepancy is a typical problem in the study of irregularities of point distribution, and represents a multi-dimensional variant of an open problem first posed by Roth; see Schmidt [8, pp. 124–125]. In its general form, it asks whether it is possible to choose N points in a given bounded convex body in such a way that after cutting it into two parts by hyperplanes in different ways, the numbers of points in the two parts essentially depend only on the relative volumes. More precisely, let \mathcal{P} denote a distribution of N points in a bounded convex body $B \subset \mathbb{R}^d$. For every unit vector $\sigma \in \Sigma_{d-1}$ and every $r \geq 0$, consider the half-space $H_{\sigma,r} = \{t \in \mathbb{R}^d : t \cdot \sigma \leq r\}$, where \cdot denotes the usual inner product in \mathbb{R}^d , and let $S_{\sigma,r} = B \cap H_{\sigma,r}$. The problem is whether ⁽¹⁾

$$(1.1) \quad \inf_{\text{card}(\mathcal{P})=N} \sup_{\substack{r \geq 0 \\ \sigma \in \Sigma_{d-1}}} |\text{card}(\mathcal{P} \cap S_{\sigma,r}) - N|B|^{-1}|S_{\sigma,r}||$$

is unbounded with N .

This question was first answered in the affirmative by Beck [2] in the case when $d = 2$ and B is the unit disc, using Fourier transform techniques; see also [3]. Subsequently, his almost sharp lower bound was improved by Alexander [1] who used integral-geometric techniques to establish the L^2 result that for every distribution \mathcal{P} of N points in the unit disc, we have ⁽²⁾

$$(1.2) \quad \int_{\Sigma_1} \int_0^{\pi^{-1/2}} |\text{card}(\mathcal{P} \cap S_{\sigma,r}) - N|S_{\sigma,r}||^2 dr d\sigma \geq cN^{1/2}.$$

2010 *Mathematics Subject Classification*: Primary 11K38; Secondary 42B10.

Key words and phrases: irregularities of distribution, decay of Fourier transforms.

⁽¹⁾ We write $|S|$ to denote the Lebesgue measure of a Lebesgue measurable set S .

⁽²⁾ Throughout this paper, the letter c denotes positive absolute constants which may vary in value from one appearance to the next. Furthermore, the symbol c with subscripts denotes positive constants whose values may depend on the subscripts displayed, and again may vary in value from one appearance to the next.

The unboundedness of (1.1) in this special case follows immediately.

In fact, this last bound (1.2) is sharp, in view of the amazing result of Matoušek [6] that there exist distributions \mathcal{P} of N points in the unit disc such that

$$\sup_{\substack{r \geq 0 \\ \sigma \in \Sigma_1}} |\text{card}(\mathcal{P} \cap S_{\sigma,r}) - N|S_{\sigma,r}|| \leq cN^{1/4},$$

from which the upper bound

$$\int_{\Sigma_1} \int_0^{\pi^{-1/2}} |\text{card}(\mathcal{P} \cap S_{\sigma,r}) - N|S_{\sigma,r}||^2 dr d\sigma \leq cN^{1/2}$$

follows immediately.

However, if one replaces the L^2 norm by the corresponding L^1 norm, one gets a rather different picture. No lower bound corresponding to (1.2) is currently known, while Beck and the first author [4] have shown that for every bounded convex body $B \subset \mathbb{R}^2$ with centre of gravity at the origin and every natural number N , there exists a distribution \mathcal{P} of N points in B such that

$$(1.3) \quad \int_{\Sigma_1} \int_0^{R(\sigma)} |\text{card}(\mathcal{P} \cap S_{\sigma,r}) - N|B|^{-1}|S_{\sigma,r}|| dr d\sigma \leq c_B(\log N)^2,$$

where $R(\sigma) = \sup\{t \cdot \sigma : t \in B\}$.

A careful description of the above and related problems can be found in Matoušek [7, Sections 3.2 and 6.6].

2. Main results. The purpose of this paper is to establish an estimate in the spirit of (1.3), in several variables and when the convex body is a cube. More precisely, let $Q = [-1, 1]^d$. For every unit vector $\sigma \in \Sigma_{d-1}$ and every $r \geq 0$,

$$P_{\sigma,r} = Q \cap \{t \in \mathbb{R}^d : t \cdot \sigma \leq r\}$$

denotes the intersection of the cube Q with one of the two half-spaces in \mathbb{R}^d created by cutting \mathbb{R}^d by the hyperplane

$$(2.1) \quad s_{\sigma,r} = \{t \in \mathbb{R}^d : t \cdot \sigma = r\}.$$

Our main result is the following.

THEOREM 1. *Let $M > 1$ be an integer, and let $N = (2M + 1)^d$. Then there exists a distribution \mathcal{P} of N points in the cube $Q = [-1, 1]^d$ such that*

$$\sup_{r \geq 0} \int_{\Sigma_{d-1}} |\text{card}(\mathcal{P} \cap P_{\sigma,r}) - N2^{-d}|P_{\sigma,r}|| d\sigma \leq c_d(\log N)^d.$$

For every integer $M > 1$, let

$$D_{\sigma,r}(M) = \text{card}((M + 1/2)P_{\sigma,r} \cap \mathbb{Z}^d) - (M + 1/2)^d |P_{\sigma,r}|.$$

Theorem 1 follows immediately from the following result on lattice points by a simple scaling argument.

THEOREM 2. *For every integer $M > 1$, we have*

$$\sup_{r \geq 0} \int_{\Sigma_{d-1}} |D_{\sigma,r}(M)| \, d\sigma \leq c_d (\log M)^d.$$

REMARK. If we replace the fraction $1/2$ in the dilation $(M + 1/2)P_{\sigma,r}$ of the set $P_{\sigma,r}$ by a different number in the interval $[0, 1)$, then we obtain the trivial conclusion that $D_{\sigma,r}(M)$ is of order M^{d-1} for every σ and r .

The remainder of the paper is organized as follows. In Section 3, we begin our proof of Theorem 2, and split our argument into two cases. We then discuss these two cases separately in Sections 4 and 5.

3. Fourier transform and divergence theorem. For every $x \in \mathbb{R}$, write

$$\varphi(x) = (2 - 4|x|)_+ = \max\{2 - 4|x|, 0\}.$$

The function φ is supported in $[-1/2, 1/2]$, satisfies

$$\int_{\mathbb{R}} \varphi(x) \, dx = 1,$$

and has Fourier transform $\widehat{\varphi}$ given by

$$(3.1) \quad \widehat{\varphi}(y) = \int_{-1/2}^{1/2} \varphi(x) e^{-2\pi iyx} \, dx = \left(\frac{2 \sin(\pi y/2)}{\pi y} \right)^2.$$

For every $t = (t_1, \dots, t_d) \in \mathbb{R}^d$, write

$$\Phi(t) = \varphi(t_1) \dots \varphi(t_d).$$

For every positive integer M , write

$$\varphi_M(x) = M^{d-1} \varphi(M^{d-1}x) \quad \text{and} \quad \Phi_M(t) = M^{d^2-d} \Phi(M^{d-1}t).$$

Then

$$\widehat{\Phi}_M(\xi) = \int_{\mathbb{R}^d} \Phi_M(t) e^{-2\pi i \xi \cdot t} \, dt = \widehat{\Phi}(M^{-d+1}\xi) = \widehat{\varphi}(M^{-d+1}\xi_1) \dots \widehat{\varphi}(M^{-d+1}\xi_d)$$

for every $\xi = (\xi_1, \dots, \xi_d) \in \mathbb{R}^d$. In particular, $\widehat{\Phi}_M(0) = 1$.

Following a classical argument, we smooth the characteristic function $\chi_{(M+1/2)P_{\sigma,r}}$ of the set $(M + 1/2)P_{\sigma,r}$ by convolving it with Φ_M . We subsequently apply the Poisson summation formula to the convolution

$$\lambda_{M,(M+1/2)P_{\sigma,r}} = \Phi_M * \chi_{(M+1/2)P_{\sigma,r}}$$

and deduce that

$$\begin{aligned}
 (3.2) \quad & \sum_{m \in \mathbb{Z}^d} \lambda_{M, (M+1/2)P_{\sigma,r}}(m) \\
 &= \sum_{m \in \mathbb{Z}^d} \widehat{\lambda}_{M, (M+1/2)P_{\sigma,r}}(m) = \sum_{m \in \mathbb{Z}^d} \widehat{\Phi}_M(m) \widehat{\chi}_{(M+1/2)P_{\sigma,r}}(m) \\
 &= (M + 1/2)^d \sum_{m \in \mathbb{Z}^d} \widehat{\Phi}_M(m) \widehat{\chi}_{P_{\sigma,r}}((M + 1/2)m).
 \end{aligned}$$

Observe next that the assumptions on Φ_M imply

$$\begin{aligned}
 \sum_{m \in \mathbb{Z}^d} \lambda_{M, (M+1/2-M^{-d+1})P_{\sigma,r}}(m) &\leq \text{card}((M + 1/2)P_{\sigma,r} \cap \mathbb{Z}^d) \\
 &\leq \sum_{m \in \mathbb{Z}^d} \lambda_{M, (M+1/2+M^{-d+1})P_{\sigma,r}}(m).
 \end{aligned}$$

It follows from (3.2) that

$$\begin{aligned}
 (3.3) \quad & D_{\sigma,r}(M) \\
 &\leq \sum_{m \in \mathbb{Z}^d} \lambda_{M, (M+1/2+M^{-d+1})P_{\sigma,r}}(m) - (M + 1/2)^d |P_{\sigma,r}| \\
 &= (M + 1/2 + M^{-d+1})^d \sum_{m \in \mathbb{Z}^d} \widehat{\Phi}_M(m) \widehat{\chi}_{P_{\sigma,r}}((M + 1/2 + M^{-d+1})m) \\
 &\quad - (M + 1/2)^d |P_{\sigma,r}| \\
 &\leq c_d M^d \sum_{0 \neq m \in \mathbb{Z}^d} \widehat{\Phi}_M(m) \widehat{\chi}_{P_{\sigma,r}}((M + 1/2 + M^{-d+1})m) + O(1),
 \end{aligned}$$

and a corresponding estimate holds from below.

We have to evaluate $\widehat{\chi}_{P_{\sigma,r}}(\xi)$ when $|\xi| \geq 1$. By the divergence theorem, we have

$$\widehat{\chi}_{P_{\sigma,r}}(\xi) = \int_{P_{\sigma,r}} e^{-2\pi i \xi \cdot t} dt = \frac{i}{2\pi |\xi|^2} \int_{\partial P_{\sigma,r}} e^{-2\pi i \xi \cdot t} \xi \cdot \nu(t) dS_t,$$

where $\nu(t)$ is the outward unit vector and dS_t is the restriction of the Lebesgue measure to the boundary $\partial P_{\sigma,r}$, consisting of a bounded number of $(d - 1)$ -dimensional faces of $P_{\sigma,r}$. Let the polyhedron $G_{\sigma,r,d-1}$ denote one of these faces, and note that $\nu(t)$ is constant on $G_{\sigma,r,d-1}$. The study of $\widehat{\chi}_{P_{\sigma,r}}(\xi)$ therefore reduces to that of a finite number of terms of the form

$$(3.4) \quad \frac{\xi \cdot \nu}{|\xi|^2} \widehat{\mu}_{G_{\sigma,r,d-1}}(\xi),$$

where $\mu_{G_{\sigma,r,d-1}}$ is the restriction of the Lebesgue measure to $G_{\sigma,r,d-1}$. We have two cases.

CASE A₁. The face $G_{\sigma,r,d-1}$ is entirely contained in the hyperplane $s_{\sigma,r}$; see (2.1).

CASE B₁. The face $G_{\sigma,r,d-1}$ is entirely contained in one of the $(d - 1)$ -dimensional faces of the cube Q .

4. Case A₁. In this section, we consider the case when the face $G_{\sigma,r,d-1}$ is entirely contained in the hyperplane $s_{\sigma,r}$.

In this case, for every r , the face $G_{\sigma,r,d-1}$ rotates with $\sigma \in \Sigma_{d-1}$, changing its shape as well as the number of its lower dimensional faces. However, the number of these lower dimensional faces and the lengths of their edges are bounded by positive constants that depend only on the dimension d . Then a mild variation of the proof of [5, Theorem 2.1(ii)] gives

$$\int_{\Sigma_{d-1}} \left| \frac{t \cdot \nu}{|t|^2} \widehat{\mu}_{G_{\sigma,r,d-1}}(t) \right| d\sigma \leq c_d \frac{(\log |t|)^{d-1}}{|t|^d}, \quad |t| \geq 2.$$

REMARK. According to [5, Theorem 2.1(ii)], a d -dimensional polyhedron P satisfies

$$(4.1) \quad \int_{\Sigma_{d-1}} |\widehat{\chi}_P(\rho\sigma)| d\sigma \leq c_P \frac{(\log \rho)^{d-1}}{\rho^d}, \quad \rho \geq 2.$$

If the diameter and the number of faces of P are bounded, then the constant c_P can be replaced by a constant c_d . The proof of (4.1) starts with the divergence theorem, and then proceeds by induction on the dimensions of the faces of P in the following way. Write $\sigma = (\cos \varphi, \eta \sin \varphi) \in \Sigma_{d-1}$, with $\eta \in \Sigma_{d-2}$ and $0 \leq \varphi \leq \pi$. By the induction assumption, we have

$$\int_{\Sigma_{d-1}} |\widehat{\chi}_P(\rho\sigma)| d\sigma \leq c_P \frac{1}{\rho} \int_0^\pi \frac{(\log \rho \sin \varphi)^{d-2}}{(\rho \sin \varphi)^{d-1}} (\sin \varphi)^{d-2} d\varphi \leq c_P \frac{(\log \rho)^{d-1}}{\rho^d},$$

with the induction starting from the simple inequality

$$\int_0^\pi \frac{|\sin(\rho \sin \varphi)|}{\rho \sin \varphi} d\varphi \leq c \frac{\log \rho}{\rho}.$$

In the present case, the edges of $P_{\sigma,r}$ change in number and lengths under rotation and translation, but this does not affect the induction argument. For the first step, let $\gamma(\varphi, r)$ denote the length of a given edge on the boundary of $P_{\sigma,r}$. Then

$$\int_0^\pi \frac{|\sin(\rho\gamma(\varphi, r) \sin \varphi)|}{\sin \varphi} d\varphi \leq 2 + 2 \int_{1/\rho}^{\pi/2} \frac{1}{\sin \varphi} d\varphi \leq c \log \rho,$$

and the contribution of Case A_1 to the estimate of

$$\int_{\Sigma_{d-1}} |D_{\sigma,r}(M)| d\sigma$$

is bounded above (see (3.3)) by

$$\begin{aligned} (4.2) \quad c_d M^d \sum_{0 \neq m \in \mathbb{Z}^d} |\widehat{\Phi}_M(m)| \frac{(\log M|m|)^{d-1}}{(M|m|)^d} \\ \leq c_d (\log M)^{d-1} \sum_{0 \leq |m| \leq M^{d-1}} |\widehat{\Phi}_M(m)| \frac{1}{|m|^d} \\ + c_d \sum_{|m| > M^{d-1}} |\widehat{\Phi}_M(m)| \frac{(\log |m|)^{d-1}}{|m|^d} \\ = \Theta_1 + \Theta_2, \end{aligned}$$

say. Recall that the constants c_d may change in value from one occurrence to the next.

We observe that $0 \leq \widehat{\Phi}_M(m) \leq \widehat{\Phi}_M(0) = 1$, and shall bound Θ_1 by showing that

$$(4.3) \quad \sum_{\substack{0 < |m| \leq M^{d-1} \\ m_1 \geq 0, \dots, m_d \geq 0}} \frac{1}{|m|^d} \leq c_d \log M,$$

where $m = (m_1, \dots, m_d)$. We shall achieve this by using induction to show that

$$(4.4) \quad \sum_{\substack{0 < |m| \leq M^{d-1} \\ m_1 \geq 0, \dots, m_k \geq 0 \\ m_{k+1} = 0, \dots, m_d = 0}} \frac{1}{|m|^d} \leq c_d \log M$$

for every $k = 1, \dots, d$.

Indeed, it is trivial to show that (4.4) holds for $k = 1$, noting that $d \geq 2$. Suppose now that this inequality holds for every $k = 1, \dots, d - 1$. Let

$$(4.5) \quad \mathcal{T} := \{m = (m_1, \dots, m_d) \in \mathbb{Z}^d : m_1 \geq 1, \dots, m_d \geq 1, \max_j m_j \geq 2\}.$$

Then

$$(4.6) \quad \sum_{\substack{0 < |m| \leq M^{d-1} \\ m_1 \geq 0, \dots, m_d \geq 0}} \frac{1}{|m|^d} = \sum_{\substack{0 < |m| \leq M^{d-1} \\ \min_j m_j = 0}} \frac{1}{|m|^d} + \frac{1}{d^{d/2}} + \sum_{\substack{0 < |m| \leq M^{d-1} \\ m \in \mathcal{T}}} \frac{1}{|m|^d}.$$

Observe that the first sum on the right hand side of (4.6) is a sum of a

bounded number of terms of the form

$$(4.7) \quad \sum_{\substack{0 < |m| \leq M^{d-1} \\ m_{j_1} \geq 0, \dots, m_{j_k} \geq 0 \\ m_j = 0 \text{ if } j \notin \{j_1, \dots, j_k\}}} \frac{1}{|m|^d}$$

with $k < d$. Their overall contribution does not exceed $c_d \log M$ by the induction hypothesis—note that the quantity (4.7) is invariant under permutation of the variables m_1, \dots, m_d , and is therefore equal to the left hand side of (4.4). To study the last term on the right hand side of (4.6), we consider the bijection (see (4.5))

$$\mathcal{T} \ni m = (m_1, \dots, m_d) \leftrightarrow (m_1 - 1, m_1] \times \dots \times (m_d - 1, m_d] =: Q_m,$$

and note that the union of the cubes Q_m satisfies

$$\bigcup_{m \in \mathcal{T}} Q_m = (0, +\infty)^d \setminus (0, 1]^d.$$

Then

$$\begin{aligned} \sum_{\substack{0 < |m| \leq M^{d-1} \\ m \in \mathcal{T}}} \frac{1}{|m|^d} &\leq \sum_{\substack{0 < |m| \leq M^{d-1} \\ m \in \mathcal{T}}} \int_{Q_m} \frac{1}{|x|^d} dx \leq \int_{1 \leq |x| \leq M^d} \frac{1}{|x|^d} dx \\ &= c_d \int_1^{M^d} \frac{1}{s} ds = c_d \log M. \end{aligned}$$

This completes the proof of the inequality (4.3).

We now conclude from (4.2) and (4.3) that

$$(4.8) \quad \Theta_1 \leq c_d (\log M)^d.$$

To study the term Θ_2 , note first of all that for every $m = (m_1, \dots, m_d)$, there exists at least one index j^* such that $|m_{j^*}| \geq |m|/\sqrt{d}$. In view of (3.1), we have

$$\begin{aligned} \widehat{\Phi}_M(m) &= \prod_{j=1}^d \widehat{\varphi}(M^{-d+1}m_j) = \prod_{j=1}^d \left(\frac{2 \sin(\pi M^{-d+1}m_j/2)}{\pi M^{-d+1}m_j} \right)^2 \\ &\leq \left(\frac{2 \sin(\pi M^{-d+1}m_{j^*}/2)}{\pi M^{-d+1}m_{j^*}} \right)^2 \leq c_d M^{2d-2} \frac{1}{|m|^2}. \end{aligned}$$

It follows that

$$(4.9) \quad \Theta_2 \leq c_d M^{2d-2} \sum_{|m| > M^{d-1}} \frac{(\log |m|)^{d-1}}{|m|^{d+2}}.$$

For $s > M^{d-1}$, the function $s \mapsto (\log s)^{d-1}/s^{d+2}$ decreases with s . We can then apply the earlier argument and control the right hand side of (4.9)

with an integral, which can then be handled using integration by parts $d - 1$ times. More precisely, we have

$$\begin{aligned}
 (4.10) \quad \Theta_2 &\leq c_d M^{2d-2} \int_{M^{d-1}}^{+\infty} \frac{(\log s)^{d-1}}{s^3} ds \\
 &\leq c_d M^{2d-2} \left(M^{2-2d} (\log M)^{d-1} + \int_{M^{d-1}}^{+\infty} \frac{(\log s)^{d-2}}{s^3} ds \right) \\
 &\leq c_d M^{2d-2} \\
 &\quad \times \left(M^{2-2d} (\log M)^{d-1} + M^{2-2d} (\log M)^{d-2} + \int_{M^{d-1}}^{+\infty} \frac{(\log s)^{d-3}}{s^3} ds \right) \\
 &\leq \dots \leq c_d (\log M)^{d-1}.
 \end{aligned}$$

Combining (4.2), (4.8) and (4.10), we conclude that the contribution of Case A_1 to the estimate of $\int_{\Sigma_{d-1}} |D_{\sigma,r}(M)| d\sigma$ is bounded above by $c_d (\log M)^d$.

5. Case B_1 . In this section, we consider the case when the face $G_{\sigma,r,d-1}$ is entirely contained in one of the $(d - 1)$ -dimensional faces of the cube Q . Our proof is inductive in nature.

Without loss of generality, we may assume that $\nu = (0, \dots, 0, 1)$, so that the face $G_{\sigma,r,d-1}$ is contained in the hyperplane $t_d = 1$. Then (3.4) becomes

$$\begin{aligned}
 (5.1) \quad \frac{\xi \cdot \nu}{|\xi|^2} \widehat{\mu}_{G_{\sigma,r,d-1}}(\xi) &= \frac{\xi_d}{|\xi|^2} e^{-2\pi i \xi_d} \int_{F_{\sigma,r,d-1}} e^{-2\pi i (\xi_1, \dots, \xi_{d-1}) \cdot (t_1, \dots, t_{d-1})} dt_1 \dots dt_{d-1} \\
 &= \frac{\xi_d}{|\xi|^2} e^{-2\pi i \xi_d} \widehat{\chi}_{F_{\sigma,r,d-1}}(\xi_1, \dots, \xi_{d-1}),
 \end{aligned}$$

where $F_{\sigma,r,d-1} = G_{\sigma,r,d-1} - \nu$ can be interpreted as a polyhedron in \mathbb{R}^{d-1} , with characteristic function $\chi_{F_{\sigma,r,d-1}}$. To study (5.1), we consider two cases.

CASE C_1 . We have $|(\xi_1, \dots, \xi_{d-1})| < 1$.

CASE D_1 . We have $|(\xi_1, \dots, \xi_{d-1})| \geq 1$.

We begin with Case C_1 . Recall that $m \in \mathbb{Z}^d$, so $|(m_1, \dots, m_{d-1})| < 1$ clearly implies $m_1 = \dots = m_{d-1} = 0$. The contribution of this case to an upper estimate for (3.3) therefore does not exceed

$$\begin{aligned}
 (5.2) \quad M^d \left| \sum_{0 \neq m_d \in \mathbb{Z}} \widehat{\varphi} \left(\frac{m_d}{M^{d-1}} \right) \frac{1}{M m_d} e^{-2\pi i (M+1/2+M^{-d+1})m_d} |F_{\sigma,r,d-1}| \right| \\
 = 2 |F_{\sigma,r,d-1}| \left| \sum_{m_d=1}^{+\infty} (-1)^{m_d} \widehat{\varphi} \left(\frac{m_d}{M^{d-1}} \right) \frac{\sin 2\pi M^{-d+1} m_d}{M^{-d+1} m_d} \right| \leq H + K,
 \end{aligned}$$

where

$$H = 2^d \left| \sum_{m_d=1}^{M^{d-1}} (-1)^{m_d} \widehat{\varphi} \left(\frac{m_d}{M^{d-1}} \right) \frac{\sin 2\pi M^{-d+1} m_d}{M^{-d+1} m_d} \right|,$$

$$K = 2^d \left| \sum_{m_d=M^{d-1}+1}^{+\infty} (-1)^{m_d} \widehat{\varphi} \left(\frac{m_d}{M^{d-1}} \right) \frac{\sin 2\pi M^{-d+1} m_d}{M^{-d+1} m_d} \right|.$$

REMARK. Note that the equality in (5.2) depends on the fraction $1/2$ in the dilation $(M + 1/2)P_{\sigma,r}$ of the set $P_{\sigma,r}$.

For the sum H , note that $0 \leq M^{-d+1} m_d \leq 1$, and that we can split the interval $0 \leq x \leq 1$ into a bounded number of subintervals where the function

$$x \mapsto \widehat{\varphi}(x) \frac{\sin 2\pi x}{x}$$

is monotone and does not change sign. It follows that H is not greater than the sum of a bounded number of Leibniz sums, and this implies $H \leq c_d$.

For the sum in K , note that

$$\begin{aligned} & \sum_{m_d=M^{d-1}+1}^{+\infty} (-1)^{m_d} \widehat{\varphi} \left(\frac{m_d}{M^{d-1}} \right) \frac{\sin 2\pi M^{-d+1} m_d}{M^{-d+1} m_d} \\ &= \sum_{j=1}^{+\infty} \sum_{m_d=jM^{d-1}+1}^{(j+1)M^{d-1}} (-1)^{m_d} \widehat{\varphi} \left(\frac{m_d}{M^{d-1}} \right) \frac{\sin 2\pi M^{-d+1} m_d}{M^{-d+1} m_d} \\ &= \sum_{j=1}^{+\infty} (-1)^{jM^{d-1}} \sum_{m_d=1}^{M^{d-1}} (-1)^{m_d} \widehat{\varphi} \left(j + \frac{m_d}{M^{d-1}} \right) \frac{\sin 2\pi M^{-d+1} m_d}{j + M^{-d+1} m_d}. \end{aligned}$$

It is an exercise in calculus to show the existence of a positive constant c such that for every index j , there are at most c subintervals of the interval $0 \leq x \leq 1$ where the function

$$x \mapsto \widehat{\varphi}(j + x) \frac{\sin 2\pi x}{j + x}$$

is monotone and does not change sign. Then

$$K \leq c_d \sum_{j=1}^{+\infty} j^{-3} \leq c_d.$$

Next, we turn our attention to Case D₁. Applying the divergence theorem to the polyhedron $F_{\sigma,r,d-1}$, we meet cases similar to Case A₁ and Case B₁. At the ℓ th step, where $1 \leq \ell \leq d - 1$, the divergence theorem leads to one of the following two cases.

CASE A_ℓ . We have a $(d - \ell)$ -dimensional face entirely contained in some hyperplane in $\mathbb{R}^{d-\ell+1}$ analogous to $s_{\sigma,r}$.

CASE B_ℓ . We have a $(d - \ell)$ -dimensional face entirely contained in one of the $(d - \ell)$ -dimensional faces of the cube $[-1, 1]^{d-\ell+1}$.

In Case A_ℓ , we proceed as in Case A_1 .

In Case B_ℓ , we need to study terms of the form

$$(5.3) \quad M^d \sum_{0 \neq m \in \mathbb{Z}^d} \left(\prod_{j=0}^{\ell-1} \Xi_j(m) \right) \widehat{\chi}_{F_{\sigma,r,d-\ell}}((M+1/2+M^{-d+1})(m_1, \dots, m_{d-\ell})),$$

where, for $0 \leq j \leq \ell - 1$,

$$\begin{aligned} \Xi_j(m) &= \widehat{\varphi} \left(\frac{m_{d-j}}{M^{d-1}} \right) \\ &\quad \times \frac{(M + 1/2 + M^{-d+1})m_{d-j}}{|(M + 1/2 + M^{-d+1})(m_1, \dots, m_{d-j})|^2} e^{-2\pi i(M+1/2+M^{-d+1})m_{d-j}}. \end{aligned}$$

We split the sum (5.3) into the following two cases.

CASE C_ℓ . We have $|(m_1, \dots, m_{d-\ell})| < 1$.

CASE D_ℓ . We have $|(m_1, \dots, m_{d-\ell})| \geq 1$.

Since $m_1 = \dots = m_{d-\ell} = 0$ in Case C_ℓ , the contribution of this case to the sum (5.3) is equal to

$$A = M^d |F_{\sigma,r,d-\ell}| \sum_{0 \neq m_{d-\ell+1} \in \mathbb{Z}} \dots \sum_{0 \neq m_d \in \mathbb{Z}} \left(\prod_{j=0}^{\ell-1} \Xi_j(m) \right).$$

To study this sum, write

$$\begin{aligned} B(m_{d-\ell+1}, \dots, m_{d-1}) &= \sum_{0 \neq m_d \in \mathbb{Z}} \Xi_0(m) \\ &= \frac{2M^{-d+1}}{M + 1/2 + M^{-d+1}} \\ &\quad \times \sum_{m_d=1}^{+\infty} \widehat{\varphi} \left(\frac{m_d}{M^{d-1}} \right) (-1)^{m_d} \frac{M^{-d+1} m_d \sin 2\pi M^{-d+1} m_d}{|M^{-d+1}(0, \dots, 0, m_{d-\ell+1}, \dots, m_d)|^2}, \end{aligned}$$

and observe that the function

$$x \mapsto \widehat{\varphi}(x) \frac{x \sin 2\pi x}{|(0, \dots, 0, M^{-d+1}m_{d-\ell+1}, \dots, M^{-d+1}m_{d-1}, x)|^2}$$

is bounded in x , uniformly for $m_{d-\ell+1}, \dots, m_{d-1}$. Applying the earlier argument for the sum H to each of these functions, we conclude that

$$|M^d B(m_{d-\ell+1}, \dots, m_{d-1})| \leq c_d.$$

Then

$$A = M^d |F_{\sigma,r,d-\ell}| \sum_{0 \neq m_{d-\ell+1} \in \mathbb{Z}} \cdots \sum_{0 \neq m_{d-1} \in \mathbb{Z}} \left(\prod_{j=1}^{\ell-1} \Xi_j(m) \right) B(m_{d-\ell+1}, \dots, m_{d-1})$$

satisfies

$$\begin{aligned} (5.4) \quad |A| &\leq c_d \sum_{0 \neq m_{d-\ell+1} \in \mathbb{Z}} \cdots \sum_{0 \neq m_{d-1} \in \mathbb{Z}} \left(\prod_{j=1}^{\ell-1} |\Xi_j(m)| \right) \\ &\leq c_d \sum_{0 \neq m_{d-\ell+1} \in \mathbb{Z}} \cdots \sum_{0 \neq m_{d-1} \in \mathbb{Z}} \left(\prod_{j=1}^{\ell-1} \left| \widehat{\varphi} \left(\frac{m_{d-j}}{M^{d-1}} \right) \right| \frac{1}{M^{m_{d-j}}} \right) \\ &\leq c_d M^{-\ell+1} \left(\sum_{k=1}^{+\infty} \left| \widehat{\varphi} \left(\frac{k}{M^{d-1}} \right) \right| \frac{1}{k} \right)^{\ell-1} \\ &\leq c_d M^{-\ell+1} \left(\sum_{k=1}^{M^{d-1}} \frac{1}{k} + \sum_{k=M^{d-1}+1}^{+\infty} \frac{M^{2d-2}}{k^3} \right)^{\ell-1} \\ &\leq c_d M^{-\ell+1} (\log M)^\ell, \end{aligned}$$

using (3.1).

REMARKS. (1) The upper bound (5.4) is more than we need. However, the problem of bounding the sum A is not entirely trivial, since simply putting absolute values inside the sums $B(m_{d-\ell+1}, \dots, m_{d-1})$ does not lead to a useful estimate. Again we have used the cancellations given by the term $1/2$ in the dilation $(M + 1/2)P_{\sigma,r}$ of the set $P_{\sigma,r}$.

(2) It appears that we are studying the contribution of the boundary of Q to the discrepancy, but we know that this contribution is actually zero, as a consequence of the term $1/2$ in the dilation, at least as far as whole faces of Q are concerned. The delicate point here is that we are not estimating the actual discrepancy arising from the boundary, but have arrived at the boundary through the Poisson summation formula and the divergence theorem. Thus this approach does not seem to allow us to state mathematically that the contribution of the boundary must be negligible.

In case D_ℓ , we again apply the divergence theorem, and meet cases similar to Case A_ℓ and Case B_ℓ . At the last step, we have part of an edge of Q , say $\{t, 1, \dots, 1\}_{b(\sigma,r) \leq t \leq 1}$. Then we need to bound the sum

$$(5.5) \quad M^d \sum_{0 \neq m \in \mathbb{Z}^d} \widehat{\Phi}_M(m) \left(\prod_{j=2}^d \Upsilon_j(m) \right) \int_{b(\sigma,r)}^1 e^{-2\pi i(M+1/2+M^{-d+1})m_1 s} ds,$$

where, for $2 \leq j \leq d$,

$$\mathcal{Y}_j(m) = \frac{(M + 1/2 + M^{-d+1})m_j}{|(M + 1/2 + M^{-d+1})(m_1, \dots, m_j)|^2} e^{-2\pi i(M+1/2+M^{-d+1})m_j}.$$

The part of the sum (5.5) where $m_1 = 0$ is Case C_{d-1} . For $m_1 \neq 0$, we compute the integral and bound the sum (5.5), uniformly in σ and t , by

$$\begin{aligned} M^d \prod_{j=1}^d \left(\sum_{m_j=1}^{+\infty} \left| \widehat{\varphi} \left(\frac{m_j}{M^{d-1}} \right) \right| \frac{1}{M m_j} \right) &= \left(\sum_{k=1}^{+\infty} \left| \widehat{\varphi} \left(\frac{k}{M^{d-1}} \right) \right| \frac{1}{k} \right)^\ell \\ &\leq c_d M^{-d} (\log M)^d, \end{aligned}$$

as in (5.4).

This completes the proof of Theorem 2.

Acknowledgements. The authors wish to express their gratitude to the referee for his careful reading of the manuscript and constructive comments.

References

- [1] R. Alexander, *Geometric methods in the study of irregularities of distribution*, *Combinatorica* 10 (1990), 115–136.
- [2] J. Beck, *On a problem of K. F. Roth concerning irregularities of point distribution*, *Invent. Math.* 74 (1983), 477–487.
- [3] J. Beck and W. W. L. Chen, *Irregularities of Distribution*, Cambridge Tracts in Math. 89, Cambridge Univ. Press, 1987.
- [4] —, —, *Irregularities of point distribution relative to half-planes I*, *Mathematika* 40 (1993), 102–126.
- [5] L. Brandolini, L. Colzani and G. Travaglini, *Average decay of Fourier transforms and integer points in polyhedra*, *Ark. Mat.* 35 (1997), 253–275.
- [6] J. Matoušek, *Tight upper bounds for the discrepancy of half-spaces*, *Discrete Comput. Geom.* 13 (1995), 593–601.
- [7] —, *Geometric Discrepancy*, Algorithms Combin. 18, Springer, 2010.
- [8] W. M. Schmidt, *Lectures on Irregularities of Distribution*, Lectures on Math. and Phys., Tata Inst. Fundam. Res., 1977.

William W. L. Chen
 Department of Mathematics
 Macquarie University
 Sydney, NSW 2109, Australia
 E-mail: william.chen@mq.edu.au

Giancarlo Travaglini
 Dipartimento di Statistica
 Università di Milano-Bicocca
 Edificio U7
 Via Bicocca degli Arcimboldi 8
 20126 Milano, Italy
 E-mail: giancarlo.travaglini@unimib.it

Received on 16.10.2008
 and in revised form on 12.6.2009

(5830)