

Division of Distributions by Locally Definable Quasianalytic Functions

by

Krzysztof Jan NOWAK

Presented by Wiesław PLEŚNIAK

Summary. We demonstrate that the Łojasiewicz theorem on the division of distributions by analytic functions carries over to the case of division by quasianalytic functions locally definable in an arbitrary polynomially bounded, o-minimal structure which admits smooth cell decomposition. Hence, in particular, the principal ideal generated by a locally definable quasianalytic function is closed in the Fréchet space of smooth functions.

In his famous paper [9], S. Łojasiewicz solved the problem of the division of distributions by analytic functions, posed by L. Schwartz [18]. The first part of his paper was devoted to distributions $\mathcal{D}(U)$ and smooth functions $\mathcal{E}(U)$, i.e. of class C^∞ on U . In the theorem from Section 10, he achieved the division of distributions by a smooth function Φ , provided that its zero locus $Z := \{\Phi = 0\}$ admits a certain finite smooth stratification which enjoys some properties of growth and regular separation. In part two, he established an inequality for analytic functions, a special case of one for subanalytic functions, now called the Łojasiewicz inequality. It was a crucial point in the proof that every analytic function Φ satisfies the assumptions of the above theorem.

Let us emphasize that the regular separation of any two closed subanalytic subsets is a direct consequence of the Łojasiewicz inequality applied to the distance functions from those two sets. Sometimes the condition of regular separation itself is called Łojasiewicz's inequality too.

We wish to recall Łojasiewicz's theorem, and next to prove that its assumptions are satisfied—after a suitable, generic, linear change of coordi-

2010 *Mathematics Subject Classification*: 14P15, 46F10, 26E10.

Key words and phrases: distributions, closed ideals, quasianalytic functions, polynomially bounded structures, smooth stratifications.

nates—by every quasianalytic function definable in a polynomially bounded, o-minimal structure \mathcal{R} which admits smooth cell decomposition. Note that examples of such structures are the expansions of the real field by restricted quasianalytic functions (including some classical Denjoy–Carleman classes) which satisfy certain natural conditions (cf. [17, 13, 14]).

For a bounded open subset $U \subset \mathbb{R}^n$, let us introduce, after Łojasiewicz, the following notation:

$\mathcal{D}'(U)$, $\mathcal{E}'(U)$ and $\mathcal{P}'(U)$ stand for the spaces of distributions, distributions with compact supports and distributions prolongable onto \mathbb{R}^n , respectively.

It is easy to check, by means of a partition of unity, that the division problem is local. Therefore, if the answer to the division problem is affirmative for one of those spaces of distributions, it is so for the remaining two.

For a bounded open subset $\Omega \subset \mathbb{R}^k$, let $\mathcal{H}(\Omega)$ denote the set of smooth functions $f : \Omega \rightarrow \mathbb{R}$ that satisfy the following growth condition:

$$\forall \alpha \in \mathbb{N}^k \exists M_\alpha, s_\alpha > 0 \quad \left| \frac{\partial^{|\alpha|} f}{\partial x^\alpha}(u) \right| \leq M_\alpha \operatorname{dist}(u, \partial\Omega)^{-s_\alpha} \quad \text{for all } u \in \Omega.$$

Further, $\mathcal{G}(\Omega)$ denotes the set of smooth functions $f : \Omega \rightarrow \mathbb{R}$ that satisfy the following growth condition:

$$\exists \varepsilon, s > 0 \quad |f(u)| > \varepsilon \operatorname{dist}(u, \partial\Omega)^s \quad \text{for all } u \in \Omega.$$

Consider a bounded smooth leaf $\Gamma \subset \mathbb{R}^n$ of the form

$$(*) \quad \Gamma = \{(u, v) \in \Omega \times \mathbb{R}_v^{n-k} : v = \eta(u)\},$$

where Ω is a bounded open subset of \mathbb{R}_u^k , $u = (x_1, \dots, x_k)$, $v = (x_{k+1}, \dots, x_n)$, $\eta(u) = (\eta_{k+1}(u), \dots, \eta_n(u))$ and $\eta_i(u) \in \mathcal{H}(\Omega)$ for $i = k+1, \dots, n$. We say that Γ satisfies *condition* (R) with respect to a subset $E \subset \mathbb{R}^n$ if the closures $\overline{\Gamma}$ and $\overline{E \setminus \Gamma}$ of Γ and $E \setminus \Gamma$, respectively, are regularly separated.

For the reader's convenience, we recall the definition of regular separation, due to Łojasiewicz (cf. [10, Sect. 18] or [9, Sect. 3]). We say that two closed subsets $E, F \subset \mathbb{R}^n$ are *regularly separated* at a point $a \in E \cap F$ if there are a neighbourhood W of a and $\varepsilon, s > 0$ such that one of the following two equivalent inequalities holds:

$$\operatorname{dist}(x, E) \geq \varepsilon \operatorname{dist}(x, E \cap F)^s \quad \text{for all } x \in F \cap W$$

or

$$\operatorname{dist}(x, E) + \operatorname{dist}(x, F) \geq \varepsilon \operatorname{dist}(x, E \cap F)^s \quad \text{for all } x \in W.$$

The closed subsets E and F are called *regularly separated* if they are so at all points $a \in E \cap F$.

Let us mention that condition (R) presented herein differs slightly from the one defined by Łojasiewicz in [9, Sect. 4]. However, it is also suitable

for his proof of the theorem in question (cf. [9, Sect. 10]), which can be formulated as follows.

THEOREM OF ŁOJASIEWICZ. *Let U be a bounded open subset of \mathbb{R}^n , Φ a smooth function in a neighbourhood of the closure \bar{U} and $Z := \{x \in U : \Phi(x) = 0\}$ its zero locus. Suppose $\Phi \in \mathcal{G}(U \setminus Z)$ and Z admits a finite smooth stratification*

$$Z = \bigcup_{k=0}^{n-1} \bigcup_i \Gamma_i^k$$

such that each stratum Γ_i^k is a k -dimensional smooth leaf of the form $(*)$ with appropriate functions

$$\eta_{k+1}^{k,i}, \dots, \eta_n^{k,i} \in \mathcal{H}(\Omega_i^k), \quad \eta^{k,i} = (\eta_{k+1}^{k,i}, \dots, \eta_n^{k,i}),$$

which satisfies condition (R) with respect to the set

$$\bigcup_{j=0}^k \bigcup_i \Gamma_i^j \cup \partial U \cup (\partial \Omega_i^k \times \mathbb{R}^{n-k}).$$

Further, assume that, for each stratum Γ_i^k , there is an integer $l = l_i^k$ such that

$$\Phi = \frac{\partial \Phi}{\partial x_n} = \frac{\partial^2 \Phi}{\partial x_n^2} = \dots = \frac{\partial^{l-1} \Phi}{\partial x_n^{l-1}} = 0 \quad \text{on } \Gamma_i^k,$$

and

$$\frac{\partial^l \Phi}{\partial x_n^l}(u, \eta^{k,i}(u)) \in \mathcal{G}(\Omega_i^k) \quad (\text{a fortiori, } \frac{\partial^l \Phi}{\partial x_n^l} \neq 0 \text{ on } \Gamma_i^k).$$

Then the mapping

$$\mathcal{P}'(U) \ni S \mapsto \Phi S \in \mathcal{P}'(U)$$

is surjective.

We now fix a polynomially bounded, o-minimal structure \mathcal{R} which admits smooth cell decomposition.

PROPOSITION. *If $f : U \rightarrow \mathbb{R}$ is a smooth definable function on a bounded open subset $U \subset \mathbb{R}^n$, then*

$$f \in \mathcal{H}(U) \quad \text{and} \quad f \in \mathcal{G}(U \setminus Z(f)),$$

where $Z(f)$ is the zero locus of f .

For the first property, it is sufficient to show that every definable function $g : U \rightarrow \mathbb{R}$ satisfies the following growth condition:

$$\exists M, s > 0 \quad g(x) \leq M \text{dist}(x, \partial U)^{-s} \quad \text{for all } x \in U.$$

For any $t > 0$, put

$$\theta(t) := \begin{cases} 0 & \text{if } U \cap \{x : \text{dist}(x, \partial U) = t\} = \emptyset, \\ \max \{|g(x)| : \text{dist}(x, \partial U) = t\} & \text{otherwise.} \end{cases}$$

Since the structure \mathcal{R} is polynomially bounded, there are $M, s > 0$ such that $\theta(t) \leq Mt^{-s}$ for all $t > 0$, as desired.

In order to prove the second property, for any $t > 0$, put

$$\zeta(t) := \min\{|f(x)| : x \in U, \text{dist}(x, \partial U \cup Z(f)) = t\} > 0.$$

Again, by polynomial boundedness, there are $\varepsilon, s > 0$ such that $\zeta(t) > \varepsilon t^s$ for all $t > 0$, which completes the proof.

The Łojasiewicz inequality (cf. [10, Sect. 18], [5, Sect. 4.14] or [3, Theorem 6.2]) yields immediately the regular separation of any two closed definable subsets. Consider a smooth function Φ definable in a neighbourhood of the closure \bar{U} of a bounded open definable subset $U \subset \mathbb{R}^n$. Then, in view of the foregoing proposition, the assumptions of the theorem in question will be satisfied for Φ , once we find a finite smooth definable stratification

$$Z = \bigcup_{k=0}^{n-1} \bigcup_i \Gamma_i^k$$

such that each stratum Γ_i^k is a k -dimensional smooth leaf of the form (*) with appropriate functions

$$\eta_{k+1}^{k,i}, \dots, \eta_n^{k,i} \in \mathcal{H}(\Omega_i^k), \quad \eta^{k,i} = (\eta_{k+1}^{k,i}, \dots, \eta_n^{k,i}),$$

for which there is an integer $l = l_i^k$ such that

$$\Phi = \frac{\partial \Phi}{\partial x_n} = \frac{\partial^2 \Phi}{\partial x_n^2} = \dots = \frac{\partial^{l-1} \Phi}{\partial x_n^{l-1}} = 0 \quad \text{on } \Gamma_i^k,$$

and

$$\frac{\partial^l \Phi}{\partial x_n^l}(u, \eta^{k,i}(u)) \in \mathcal{G}(\Omega_i^k) \quad (\text{a fortiori, } \frac{\partial^l \Phi}{\partial x_n^l} \neq 0 \text{ on } \Gamma_i^k).$$

Our procedure will be to construct a stratification just described. We need the lemma below, a generalization of the classical lemma of Koopman–Brown (cf. [8], [10, Sect. 22] or [5, Sect. 4.9]).

GOOD DIRECTIONS LEMMA. *Let E be a definable subset of \mathbb{R}^n of dimension $\leq n - 1$. Then for a generic line $\lambda \in \mathbb{P}_{n-1}$, i.e. for every line λ outside a nowhere dense, definable subset of the projective space \mathbb{P}_{n-1} , we have*

$$\sharp(E \cap (a + \lambda)) < \infty \quad \text{for all } a \in \mathbb{R}^n.$$

As a direct consequence, we obtain

COROLLARY. *Let E be a definable subset of \mathbb{R}^n of dimension $\leq n - k$. Then for a generic k -dimensional vector subspace $V \in \mathbb{G}_{n,k}$, i.e. for every subspace V outside a nowhere dense, definable subset of the Grassmannian $\mathbb{G}_{n,k}$, we have*

$$\sharp(E \cap (a + V)) < \infty \quad \text{for all } a \in \mathbb{R}^n.$$

Now, we shall demonstrate how to construct the required stratification. First, consider the decreasing sequence of quasianalytic subsets

$$\begin{aligned} Z := \{x \in U : \Phi(x) = 0\} \supset Z_1 := \left\{x \in U : \Phi(x) = \frac{\partial \Phi}{\partial x_n} = 0\right\} \supset \dots \\ \dots \supset Z_j := \left\{x \in U : \Phi(x) = \frac{\partial \Phi}{\partial x_n} = \dots = \frac{\partial^j \Phi}{\partial x_n^j} = 0\right\} \supset \dots \end{aligned}$$

By topological noetherianity (cf. [5, Sect. 4.17], [3, Theorem 6.1] or [12, Appendix]), this sequence stabilizes:

$$Z_l = Z_{l+1} = Z_{l+2} = \dots \quad \text{for an } l \in \mathbb{N}.$$

Next, choose a line $\lambda_1 \in \mathbb{P}_{n-1}$ according to the above lemma applied to the zero locus $Z := \{\Phi = 0\}$; we may assume that $\lambda_1 = \mathbb{R} \cdot e_n$. Take a smooth cell decomposition \mathcal{C} of \mathbb{R}^n compatible with the sets Z, Z_1, \dots, Z_l . The cells of dimension $n - 1$ are, of course, smooth definable leaves of the form (*). Project the cells of dimension $< n - 1$ onto \mathbb{R}^{n-1} parallel to $\mathbb{R} \cdot e_n$, again choose a line $\lambda_2 \in \mathbb{P}_{n-2}$ according to the above lemma applied to those projections, and take $\lambda_2 = \mathbb{R} \cdot e_{n-1}$. Consequently, the plane $\mathbb{R} \cdot e_n + \mathbb{R} \cdot e_{n-1}$ satisfies the conclusion of the foregoing corollary applied to the initial cells of dimension $n - 2$ in \mathbb{R}^n .

Further, refine the cell decomposition of \mathbb{R}^{n-2} induced by \mathcal{C} so that all cells of dimension $n - 1$ and $n - 2$ are smooth definable leaves of the form (*). We continue in this fashion, and eventually attain, after performing a linear change of coordinates, a new, finer, smooth definable cell decomposition such that every cell C is a smooth definable leaf of the form (*).

Finally, we must take a smooth definable stratification compatible with that new cell decomposition. We are thus led to the following quasianalytic generalization of the Łojasiewicz division theorem.

DIVISION THEOREM. *Consider a polynomially bounded, o -minimal structure \mathcal{R} which admits smooth cell decomposition. Let U be a connected open subset of \mathbb{R}^n and $\Phi : U \rightarrow \mathbb{R}$ a smooth, non-vanishing function locally definable on U with respect to the structure \mathcal{R} . Then the mapping*

$$\mathcal{D}'(U) \ni S \mapsto \Phi \cdot S \in \mathcal{D}'(U)$$

is surjective.

Hence and by routine arguments from functional analysis, we obtain

COROLLARY. *Let $\Phi : U \rightarrow \mathbb{R}$ be a function as in the above theorem. Then $\Phi \cdot \mathcal{E}(U)$ is a closed ideal of the Fréchet algebra $\mathcal{E}(U)$ of smooth functions on U , and the linear mapping*

$$\varphi : \mathcal{E}(U) \ni f \mapsto \Phi f \in \Phi \cdot \mathcal{E}(U)$$

is a homeomorphism.

Our proof starts with the observation that, by the Banach open mapping theorem, the ideal $\Phi \cdot \mathcal{E}(U)$ is closed iff the mapping φ is a homeomorphism. Suppose, on the contrary, that φ is not a homeomorphism. Then there is a sequence $(f_\nu) \subset \mathcal{E}(U)$ such that $\Phi f_\nu \rightarrow 0$ and $f_\nu \not\rightarrow 0$. We may, of course, assume that all the functions f_ν lie outside a neighbourhood W of $0 \in \mathcal{E}(U)$.

For any distribution $T \in \mathcal{E}'(U)$, take a distribution $S \in \mathcal{E}'(U)$ for which $T = \Phi \cdot S$. Then

$$T(f_\nu) = (\Phi \cdot S)(f_\nu) = S(\Phi f_\nu) \rightarrow 0,$$

and thus the numerical sequence $(T(f_\nu))$ is bounded. The sequence (f_ν) is therefore bounded in $\mathcal{E}(U)$. Indeed, any subset F of a locally convex topological vector space is bounded iff it is weakly bounded (see e.g. [7, Chap. 6, §8, Theorem 4']). This fact relies on two fundamental results from functional analysis, namely, the Banach–Steinhaus and Hahn–Banach theorems.

Since $\mathcal{E}(U)$ is a Montel space (i.e. it has the Heine–Borel property), the set $\{f_\nu\}$ is relatively compact. Consequently, we can take a subsequence (f_{ν_k}) convergent to an element $f \in \mathcal{E}(U)$. But we must have $f \neq 0$, whence

$$\Phi f_{\nu_k} \rightarrow \Phi f \neq 0.$$

This contradicts our assumption, concluding the proof.

REMARKS. 1) It follows immediately from the Hahn–Banach theorem that the converse is also valid:

If the linear mapping

$$\varphi : \mathcal{E}(U) \ni f \mapsto \Phi f \in \Phi \cdot \mathcal{E}(U)$$

is a homeomorphism, then the mapping

$$\mathcal{D}'(U) \ni S \mapsto \Phi \cdot S \in \mathcal{D}'(U)$$

is surjective.

2) B. Malgrange generalized the foregoing theorem on closed ideals to the case of ideals generated locally by finitely many analytic functions (cf. [11, Chap. 6, Theorem 1.1]). This strengthening seems not to carry over easily to the quasianalytic setting for lack of theorems about good algebraic properties of quasianalytic local rings (such as, for instance, noetherianity, flatness

properties or coherence). Actually, the problem whether quasianalytic local rings are noetherian remains open as yet.

3) The above corollary is tantamount to the quasianalytic division theorem to the effect that a smooth function f formally divisible by a quasianalytic function Φ is divisible by Φ . The latter was established by Bierstone–Milman [3, Section 6], who followed Atiyah’s proof of the analytic division theorem [1], based on transformation to normal crossings by blowing up.

4) The above results related to division by a quasianalytic function will be applied in our further research (e.g. [15, 16]) on carrying over the issues linked with Glaeser’s composite function theorem (cf. [6, 4, 2]) to the quasianalytic setting.

Acknowledgements. This research was partially supported by Research Project No. N N201 372336 from the Polish Ministry of Science and Higher Education.

References

- [1] M. F. Atiyah, *Resolution of singularities and division of distributions*, Comm. Pure Appl. Math. 23 (1970), 145–150.
- [2] E. Bierstone and P. D. Milman, *Geometric and differential properties of subanalytic sets*, Ann. of Math. 147 (1998), 731–785.
- [3] —, —, *Resolution of singularities in Denjoy–Caleman classes*, Selecta Math. (NS) 10 (2004), 1–28.
- [4] E. Bierstone, P. D. Milman and W. Pawłucki, *Composite differentiable functions*, Duke Math. J. 83 (1996), 607–620.
- [5] L. van den Dries and C. Miller, *Geometric categories and o-minimal structures*, *ibid.* 84 (1996), 497–540.
- [6] G. Glaeser, *Fonctions composées différentiables*, Ann. of Math. 77 (1963), 193–209.
- [7] A. Grothendieck, *Topological Vector Spaces*, Gordon and Breach, 1973.
- [8] B. Koopman and A. Brown, *On the covering of analytic loci by complexes*, Trans. Amer. Math. Soc. 34 (1932), 231–251.
- [9] S. Łojasiewicz, *Sur le problème de la division*, Studia Math. 18 (1959), 87–136, and Rozprawy Mat. 22 (1961).
- [10] —, *Ensembles Semi-analytiques*, I.H.E.S., Bures-sur-Yvette, 1965.
- [11] B. Malgrange, *Ideals of Differentiable Functions*, Oxford Univ. Press, 1966.
- [12] K. J. Nowak, *On the Euler characteristic of the links of a set determined by smooth definable functions*, Ann. Polon. Math. 93 (2008), 231–246.
- [13] —, *Decomposition into special cubes and its application to quasi-subanalytic geometry*, *ibid.* (2009), 65–74.
- [14] —, *Quantifier elimination, valuation property and preparation theorem in quasi-analytic geometry via transformation to normal crossings*, *ibid.* 96 (2009) (2009), 247–282.
- [15] —, *Semicoherent quasi-subanalytic sets and the composite function property*, in preparation.
- [16] —, *Nash quasi-subanalytic sets*, in preparation.

- [17] J.-P. Rolin, P. Speissegger and A. J. Wilkie, *Quasianalytic Denjoy–Carleman classes and o-minimality*, J. Amer. Math. Soc. 16 (2003), 751–777.
- [18] L. Schwartz, *Théorie des Distributions*, Hermann, Paris, 1957–1958.

Krzysztof Jan Nowak
Institute of Mathematics
Jagiellonian University
Łojasiewicza 6
30-348 Kraków, Poland
E-mail: nowak@im.uj.edu.pl

Received June 18, 2010;
received in final form October 2, 2010

(7771)