

Continuity of the fundamental operations on distributions
having a specified wave front set
(with a counterexample by Semyon Alesker)

by

CHRISTIAN BROUDER (Paris), NGUYEN VIET DANG (Lille) and
FRÉDÉRIC HÉLEIN (Paris)

Abstract. The pull-back, push-forward and multiplication of smooth functions can be extended to distributions if their wave front sets satisfy some conditions. Thus, it is natural to investigate the topological properties of these operations between spaces \mathcal{D}'_Γ of distributions having a wave front set included in a given closed cone Γ of the cotangent space. As discovered by S. Alesker, the pull-back is not continuous for the usual topology on \mathcal{D}'_Γ , and the tensor product is not separately continuous. In this paper, a new topology is defined for which the pull-back and push-forward are continuous, and the tensor and convolution products and multiplication of distributions are hypocontinuous.

1. Introduction. The motivation of our work comes from the renormalization of QFT in curved space times; indeed, the question addressed in this paper cannot be avoided in this context and also the technical results of this paper form the core of the proof that perturbative quantum field theories are renormalizable on curved spacetimes [7, 6].

From the work of L. Schwartz, we know that the tensor product of distributions is continuous [16, p. 110] and the product of a distribution by a smooth function is hypocontinuous [16, p. 119] (see Definition 3.1), although it is not jointly continuous [15].

However, in many applications (for instance when considering multiplication of distributions), we cannot work with all distributions and we must consider the subsets \mathcal{D}'_Γ of distributions whose wave front set [3] is included in some closed subsets Γ of $\dot{T}^*\mathbb{R}^n = \{(x; \xi) \in T^*\mathbb{R}^n; \xi \neq 0\}$, where Γ is a *cone* in the sense that $(x; \xi) \in \Gamma$ implies $(x; \lambda\xi) \in \Gamma$ for every $\lambda \in \mathbb{R}_{>0}$.

2010 *Mathematics Subject Classification*: Primary 46F10; Secondary 35A18.

Key words and phrases: microlocal analysis, functional analysis, mathematical physics, renormalization.

Received 11 June 2015; revised 14 February 2016.

Published online 2 May 2016.

If Γ is a cone, then $\Gamma' = \{(x; -\xi); (x; \xi) \in \Gamma\}$. Indeed, the spaces \mathcal{D}'_Γ are widely used in microlocal analysis because wave front set conditions rule the so-called *fundamental operations on distributions*: multiplication, pull-back and push-forward. The tensor product is also a fundamental operation, but it is feasible without any condition.

Hörmander himself, who introduced the concept of a wave front set [11], equipped \mathcal{D}'_Γ with a *pseudo-topology* [11, p. 125], which is not a topology but just a rule describing the convergence of sequences. In particular, when Hörmander writes that the fundamental operations are continuous [12, p. 263], he means “sequentially continuous”. And indeed, Hörmander and his followers proved that, under conditions on the wave front set to be described later, the following operators are sequentially continuous: the pull-back of a distribution by a smooth map [12, Thm. 8.2.4]; the push-forward of a distribution by a proper map [4, p. 528], the tensor product of two distributions [4, p. 511] and the multiplication of two distributions [4, p. 526].

However, sequential continuity was soon found to be too weak for some applications and Duistermaat [8, p. 18] equipped \mathcal{D}'_Γ with a locally convex topology defined in terms of the following seminorms [10, p. 80]:

- (i) All the seminorms on $\mathcal{D}'(\mathbb{R}^n)$ for the weak topology: $\|u\|_\phi = |\langle u, \phi \rangle|$ for all $\phi \in \mathcal{D}(\mathbb{R}^n)$.
- (ii) The seminorms $\|u\|_{N,V,\chi} = \sup_{k \in V} (1 + |k|)^N |\widehat{u\chi}(k)|$, where $N \geq 0$, $\chi \in \mathcal{D}(\mathbb{R}^n)$, and $V \subset \mathbb{R}^n$ is a closed cone with $(\text{supp } \chi \times V) \cap \Gamma = \emptyset$.

These seminorms give \mathcal{D}'_Γ the structure of a locally convex vector space and the corresponding topology is usually called *Hörmander’s topology*. It probably first appeared in the 1970–1971 lecture notes by Duistermaat [8], although the seminorms $\|\cdot\|_{N,V,\chi}$ had already been mentioned by Hörmander [11, p. 128]. The (sequential) convergence in the sense of Hörmander is: a sequence $(u_j) \in \mathcal{D}'_\Gamma$ converges to u in \mathcal{D}'_Γ if and only if $\|u_j - u\|_\phi \rightarrow 0$ for every $\phi \in \mathcal{D}(\mathbb{R}^n)$ and $\|u_j - u\|_{N,V,\chi} \rightarrow 0$ for every $\chi \in \mathcal{D}(\Omega)$, every $N \in \mathbb{N}$ and every closed cone V in \mathbb{R}^n such that $(\text{supp } \chi \times V) \cap \Gamma = \emptyset$. Therefore, it is clear that a sequence converges in the sense of Hörmander if and only if it converges in Hörmander’s topology.

However, for a locally convex space such as \mathcal{D}'_Γ (which is not metrizable), sequential continuity and topological continuity are not equivalent. Therefore, when Duistermaat states, after defining the above topology, that the pull-back [8, p. 19], the push-forward [8, p. 20] and the product of distributions [8, p. 21] are continuous, it is not clear whether he means sequential or topological continuity. When investigating applications to valuation theory [1], Alesker discovered a counterexample proving that the tensor product is not separately continuous and the pull-back is not continuous for Hörmander’s topology. In other words, this topology is too weak to be useful for these questions.

The purpose of the present paper is to describe Alesker’s counterexample and to define a topology for which the fundamental operations have optimal continuity properties: the tensor product is hypocontinuous; the pull-back by a smooth map is continuous; the pull-back by a family of smooth maps depending smoothly on parameters is uniformly continuous; the push-forward by a smooth map is also continuous; the push-forward by a family of smooth maps depending smoothly on parameters is uniformly continuous; and multiplication of distributions and the convolution product are hypocontinuous. Finally, we discuss how the wave front set of distributions on manifolds can be defined in an intrinsic way. In appendices, we prove important technical results concerning the covering of the complement of Γ , the topology of \mathcal{D}'_{\emptyset} and the fact that the additional seminorms used to define the topology of \mathcal{D}'_{Γ} can be taken countably many.

The main applications of these results are to replace technical microlocal proofs by classical topological statements [6, 7].

2. Alesker’s counterexample. Semyon Alesker discovered the following counterexample.

PROPOSITION 2.1. *Let $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ be the projection to the first coordinate. Let $\Gamma = \dot{T}^*\mathbb{R}$, so that $\mathcal{D}'_{\Gamma}(\mathbb{R}) = \mathcal{D}'(\mathbb{R})$ and $f^*\Gamma = \{(x_1, x_2; \xi_1, 0)\}$. Then the map $f^* : \mathcal{D}'_{\Gamma}(\mathbb{R}) \rightarrow \mathcal{D}'_{f^*\Gamma}(\mathbb{R}^2)$ is not topologically continuous for the Hörmander topology.*

Note that the general definition of $f^*\Gamma$ is given in Proposition 5.1.

Proof. Let $\varphi \in \mathcal{D}(\mathbb{R})$ be such that $\varphi|_{[-1,1]} = 1$. Take $\chi = \varphi \otimes \varphi$, $V = \{(\xi_1, \xi_2) \in \mathbb{R}^2; |\xi_1| \leq |\xi_2|\}$ and $N = 0$. The intersection of V with $\{(\xi_1, 0); \xi_1 \neq 0\}$ is empty as $|\xi_1| \leq |\xi_2| = 0$ implies $\xi_1 = \xi_2 = 0$. Therefore, $\|\cdot\|_{N,V,\chi}$ is a seminorm of $\mathcal{D}'_{f^*\Gamma}$ and, if f^* were continuous, it would be possible to bound $\|f^*u\|_{N,V,\chi}$ by $\sup_i |\langle u, f_i \rangle|$ for a finite set of $f_i \in \mathcal{D}(\mathbb{R})$ and every $u \in \mathcal{D}'(\mathbb{R})$.

We are going to show that this is not the case. We have

$$\|f^*u\|_{0,V,\chi} = \sup_{\xi \in V} |\widehat{\varphi u}(\xi_1)| |\widehat{\varphi}(\xi_2)| = \sup_{\xi_1} |\widehat{\varphi u}(\xi_1)| \omega(\xi_1),$$

where $\omega(\xi_1) = \sup_{|\xi_2| \geq |\xi_1|} |\widehat{\varphi}(\xi_2)|$. It is clear that $\omega(\xi_1) > 0$ everywhere since $\widehat{\varphi}$ is a real analytic function. Thus we should show that the map $\mathcal{D}'(\mathbb{R}) \rightarrow \mathbb{R}$ given by $u \mapsto \sup_{\xi \in \mathbb{R}} |\widehat{\varphi u}(\xi)| \omega(\xi)$ is not continuous (for a fixed $\omega > 0$).

If the pull-back were continuous, there would be a finite set χ_1, \dots, χ_t of functions in $\mathcal{D}(\mathbb{R})$ such that

$$\|f^*u\|_{0,V,\chi} \leq \sup_{i=1,\dots,t} |\langle u, \chi_i \rangle|.$$

We can find ξ such that the functions χ_1, \dots, χ_t and $\varphi(x)e^{-ix\xi}$ are linearly

independent. Then there exists $u \in \mathcal{D}'(\mathbb{R})$ such that $\langle u, \chi_i \rangle = 0$ for $i = 1, \dots, t$ and $\widehat{u\varphi}(\xi) = \langle u, \varphi e_\xi \rangle = 1 + 1/\omega(\xi)$, where $e_\xi(x) = e^{-i\xi \cdot x}$. Then $\|f^*u\|_{0,V,\chi} = 1 + \omega(\xi)$, and we reach a contradiction. ■

Thus, the pull-back is not continuous. Moreover, the same example can be considered as an exterior tensor product $u \mapsto u \boxtimes 1$. This shows that the exterior tensor product is not separately continuous for the Hörmander topology.

3. The normal topology and hypocontinuity. We now modify Hörmander’s topology and define what we call the *normal topology* of \mathcal{D}'_Γ . This is a locally convex topology defined by the same seminorms $\|\cdot\|_{N,V,\chi}$ as Hörmander’s topology, but we replace the seminorms $\|\cdot\|_\phi$ of the weak topology of $\mathcal{D}'(\mathbb{R}^n)$ by the seminorms $p_B(u) = \sup_{\phi \in B} |\langle u, \phi \rangle|$ (where B runs over the bounded sets of $\mathcal{D}(\Omega)$) of the strong topology of $\mathcal{D}'(\mathbb{R}^n)$. The functional properties of this topology, like completeness, duality, nuclearity, PLS-property, bornologicity, were investigated in detail [5]. As in the case of standard distributions, several operations will not be jointly continuous but only hypocontinuous. Let us recall

DEFINITION 3.1 ([17, p. 423]). Let E, F and G be topological vector spaces. A bilinear map $f : E \times F \rightarrow G$ is said to be *hypocontinuous* if:

- (i) for every neighborhood W of zero in G and every bounded set $A \subset E$ there is a neighborhood V of zero in F such that $f(A \times V) \subset W$;
- (ii) for every neighborhood W of zero in G and every bounded set $B \subset F$ there is a neighborhood U of zero in E such that $f(U \times B) \subset W$.

If E, F and G are locally convex spaces with topologies defined by the families of seminorms $(p_i)_{i \in I}, (q_j)_{j \in J}$ and $(r_k)_{k \in K}$, respectively, the definition of hypocontinuity can be translated into the following two conditions: (i) for every bounded set A of E and every seminorm r_k , there is a constant M and a finite set of seminorms q_{j_1}, \dots, q_{j_n} (both depending only on k and A) such that

$$(3.1) \quad \forall x \in A, \quad r_k(f(x, y)) \leq M \sup\{q_{j_1}(y), \dots, q_{j_n}(y)\};$$

and (ii) for every bounded set B of F and every seminorm r_k , there is a constant M and a finite set of seminorms p_{i_1}, \dots, p_{i_n} (both depending only on k and B) such that

$$(3.2) \quad \forall y \in B, \quad r_k(f(x, y)) \leq M \sup\{p_{i_1}(x), \dots, p_{i_n}(x)\}.$$

Equivalently [14, p. 155], we can reformulate hypocontinuity using the concept of equicontinuity [13, p. 200], defined as follows:

DEFINITION 3.2. Consider the general context of a locally convex topological vector space E with seminorms $(p_\alpha)_{\alpha \in A}$. Let E^* be its topologi-

cal dual. A set H in E^* is called *equicontinuous* if the family of maps $\ell_v E \ni u \mapsto \langle u, v \rangle \in \mathbb{R}$ is **uniformly** continuous when v runs over H .

Hence f is hypocontinuous if for every bounded set $A \subset E$ and every bounded set $B \subset F$ the sets of maps $\{f_x; x \in A\}$ and $\{f_y; y \in B\}$ are equicontinuous, where $f_x : E \rightarrow G$ and $f_y : F \rightarrow G$ are defined by $f_x(y) = f_y(x) = f(x, y)$.

4. Tensor product of distributions. Let Ω_1 and Ω_2 be open sets in \mathbb{R}^{d_1} and \mathbb{R}^{d_2} , respectively, and $(u, v) \in \mathcal{D}'_{\Gamma_1} \times \mathcal{D}'_{\Gamma_2}$, where $\mathcal{D}'_{\Gamma_1} \subset \mathcal{D}'(\Omega_1)$ and $\mathcal{D}'_{\Gamma_2} \subset \mathcal{D}'(\Omega_2)$. Then the tensor product $u \otimes v$ belongs to $\mathcal{D}'_{\Gamma} \subset \mathcal{D}'(\Omega_1 \times \Omega_2)$ where

$$\begin{aligned} \Gamma &= (\Gamma_1 \times \Gamma_2) \cup ((\Omega_1 \times \{0\}) \times \Gamma_2) \cup (\Gamma_1 \times (\Omega_2 \times \{0\})) \\ &= (\Gamma_1 \cup \{\underline{0}\}_1) \times (\Gamma_2 \cup \{\underline{0}\}_2) \setminus \{(\underline{0}, \underline{0})\}, \end{aligned}$$

$\{\underline{0}\}_1$ means $\Omega_1 \times \{0\}$, $\{\underline{0}\}_2$ means $\Omega_2 \times \{0\}$, and $\{\underline{0}, \underline{0}\}$ means $(\Omega_1 \times \Omega_2) \times \{0, 0\}$. Our goal in this section is to show that the tensor product is hypocontinuous for the normal topology. We denote by $(z; \zeta)$ the coordinates in $T^*(\Omega_1 \times \Omega_2)$, where $z = (x, y)$ with $x \in \Omega_1$ and $y \in \Omega_2$, and $\zeta = (\xi, \eta)$ with $\xi \in \mathbb{R}^{d_1}$ and $\eta \in \mathbb{R}^{d_2}$. We also denote $d = d_1 + d_2$, so that $\zeta \in \mathbb{R}^d$.

LEMMA 4.1. *The seminorms of the strong topology of $\mathcal{D}'(\mathbb{R}^d)$ and the family of seminorms*

$$(4.1) \quad \|t_1 \otimes t_2\|_{N, V, \varphi_1 \otimes \varphi_2} = \sup_{\zeta \in V} (1 + |\zeta|)^N |t_1 \widehat{\varphi_1}(\xi)| |t_2 \widehat{\varphi_2}(\eta)|,$$

where $\zeta = (\xi, \eta)$, $(\varphi_1, \varphi_2) \in \mathcal{D}(\Omega_1) \times \mathcal{D}(\Omega_2)$ and $V \subset \mathbb{R}^d$ are such that $(\text{supp}(\varphi_1 \otimes \varphi_2) \times V) \cap \Gamma = \emptyset$, are a fundamental system of seminorms for the normal topology of \mathcal{D}'_{Γ} .

Proof. We use the following lemma [10, p. 80].

LEMMA 4.2. *Let Ω be an open set in \mathbb{R}^n . If there is a family, indexed by $\alpha \in A$, of $\chi_\alpha \in \mathcal{D}(\Omega)$ and of closed cones $V_\alpha \subset \mathbb{R}^n \setminus \{0\}$ such that $(\text{supp} \chi_\alpha \times V_\alpha) \cap \Gamma = \emptyset$ and*

$$\Gamma^c = \bigcup_{\alpha \in A} \{(x, \xi) \in \dot{T}^* \Omega; \chi_\alpha(x) \neq 0, \xi \in \mathring{V}_\alpha\},$$

then the topology of \mathcal{D}'_{Γ} is already defined by the strong topology of $\mathcal{D}'(\Omega)$ and the seminorms $\|\cdot\|_{N, V_\alpha, \chi_\alpha}$.

It is clear that the family indexed by $\varphi_1 \otimes \varphi_2$ and V with $(\text{supp}(\varphi_1 \otimes \varphi_2) \times V) \cap \Gamma = \emptyset$ satisfies the hypothesis of the lemma. ■

To establish the hypocontinuity of the tensor product, we consider an arbitrary bounded set $B \subset \mathcal{D}'_{\Gamma_1}(\Omega_1)$; according to (3.1), we must show that, for every seminorm r_k of $\mathcal{D}'_{\Gamma}(\Omega_1 \times \Omega_2)$, there is a constant M and a finite

number of seminorms q_j such that $r_k(u \otimes v) \leq M \sup_j q_j(v)$ for all $u \in B$ and $v \in \mathcal{D}'_{\Gamma_2}(\Omega_2)$. By Schwartz' theorem [16, p. 110] we already know that this is true for every seminorm r_k of the strong topology of $\mathcal{D}'_{\Gamma}(\Omega_1 \times \Omega_2)$. It remains to show it for every $\|\cdot\|_{N,V,\varphi_1 \otimes \varphi_2}$. This will be done by first defining a suitable partition of unity on $\Omega_1 \times \Omega_2$ and its corresponding cones. Then, this partition of unity will be used to bound the seminorms by standard microlocal techniques.

LEMMA 4.3. *Let Γ_1, Γ_2 be closed cones in $\dot{T}^*\Omega_1$ and $\dot{T}^*\Omega_2$, respectively. Set $\Gamma = (\Gamma_1 \cup \{\underline{0}\}) \times (\Gamma_2 \cup \{\underline{0}\}) \setminus \{(\underline{0}, \underline{0})\} \subset \dot{T}^*\mathbb{R}^d$. Then for all closed cones $V \subset \mathbb{R}^d$ and $\chi \in \mathcal{D}(\Omega_1 \times \Omega_2)$ such that $(\text{supp } \chi \times V) \cap \Gamma = \emptyset$, there exist a partition of unity $(\psi_{j1} \otimes \psi_{j2})_{j \in J}$ on $\Omega_1 \times \Omega_2$, which is finite on $\text{supp } \chi$, and a family $(W_{j1} \times W_{j2})_{j \in J}$ of closed cones in $(\mathbb{R}^{d_1} \setminus \{0\}) \times (\mathbb{R}^{d_2} \setminus \{0\})$ such that*

$$(4.2) \quad (\text{supp } \psi_{j1} \times W_{j1}^c) \cap \Gamma_1 = (\text{supp } \psi_{j2} \times W_{j2}^c) \cap \Gamma_2 = \emptyset,$$

$$(4.3) \quad V \cap ((W_{j1} \cup \{0\}) \times (W_{j2} \cup \{0\})) = \emptyset$$

if $\text{supp } \chi \cap \text{supp}(\psi_{j1} \otimes \psi_{j2}) \neq \emptyset$.

Proof. We first set some notation. For any $D \in \mathbb{N}$, with the identification $T^*\mathbb{R}^D \simeq \mathbb{R}^D \oplus (\mathbb{R}^D)^*$, we denote by $\underline{\pi} : T^*\mathbb{R}^D \rightarrow \mathbb{R}^D$ and $\overline{\pi} : T^*\mathbb{R}^D \rightarrow (\mathbb{R}^D)^*$ the two projections. We use the distance d_∞ on \mathbb{R}^D (or $(\mathbb{R}^D)^*$) defined by $d_\infty(u, v) := \sup_{1 \leq i \leq D} |u^i - v^i|$. For $u \in \mathbb{R}^D$ and $r \geq 0$ we then set $\overline{B}(u, r) = \{v \in \mathbb{R}^D; d_\infty(u, v) \leq r\}$ and, for any subset $Q \subset \mathbb{R}^D$, $Q_{,r} := \{v \in \mathbb{R}^D; d_\infty(v, Q) \leq r\}$. We note that, for any pair of sets $Q_1 \subset \mathbb{R}^{d_1}$ and $Q_2 \subset \mathbb{R}^{d_2}$, we have $(Q_1 \times Q_2)_{,r} = Q_{1,r} \times Q_{2,r}$ (in particular, $\overline{B}((x, y), r) = \overline{B}(x, r) \times \overline{B}(y, r)$ for $(x, y) \in \Omega_1 \times \Omega_2$). Lastly, for any closed conic subset $W \subset (\mathbb{R}^D)^* \setminus \{0\}$, we set $\overline{W} := W \cup \{0\}$ and $UW := S^{D-1} \cap W$. Similarly if Γ is a conic subset of $T^*\mathbb{R}^D$, we set $U\Gamma = (\mathbb{R}^D \times S^{D-1}) \cap \Gamma$ and $\overline{\Gamma} = \Gamma \cup \underline{0} \subset T^*\mathbb{R}^D$ where $\underline{0}$ is the zero section of $T^*\mathbb{R}^D$.

We will prove that there exists a family $(B_{j1} \times B_{j2})_{j \in J}$ of open balls that covers $\Omega_1 \cap \Omega_2$, which is finite over any compact subset of $\Omega_1 \times \Omega_2$ and in particular on $\text{supp } \chi$ and such that $(\overline{B}_{j1} \times W_{j1}^c) \cap \Gamma_1 = (\overline{B}_{j2} \times W_{j2}^c) \cap \Gamma_2 = \emptyset$, and $V \cap (\overline{W}_{j1} \times \overline{W}_{j2}) = \emptyset$ if $\text{supp } \chi \cap (\overline{B}_{j1} \times \overline{B}_{j2}) \neq \emptyset$. The conclusion of the lemma will then follow by constructing a partition of unity $(\psi_{j1} \otimes \psi_{j2})_{j \in J}$ such that $\text{supp } \psi_{j1} = \overline{B}_{j1}$ and $\text{supp } \psi_{j2} = \overline{B}_{j2}$, for all $j \in J$, by using standard arguments.

STEP 1. If $(\text{supp } \chi \times V) \cap \Gamma = \emptyset$, then there exists $\delta > 0$ such that $d_\infty(\text{supp } \chi \times UV, U\Gamma) \geq 4\delta$. We define $K := (\text{supp } \chi)_{,\delta}$ and note that $d_\infty(K \times UV, U\Gamma) \geq 3\delta$. Without loss of generality, we can assume that δ has been chosen so that $K \subset \Omega_1 \times \Omega_2$. Obviously $\Omega_1 \times \Omega_2$ is covered by $(B((x, y), \delta))_{(x,y) \in \Omega_1 \times \Omega_2}$. Moreover all balls $B((x, y), \delta)$ are contained in K if $(x, y) \in \text{supp } \chi$, and $\text{supp } \chi$ is covered by the subfamily

$(B((x, y), \delta))_{(x,y) \in \text{supp } \chi}$. Since $\text{supp } \chi$ is compact, we can extract a countable family $(B_i)_{i \in I} = (B_{i1} \times B_{i2})_{i \in I}$ of balls which covers $\Omega_1 \times \Omega_2$ and which is finite over $\text{supp } \chi$.

We now set $\gamma := \bar{\pi}(\bar{\pi}^{-1}(K) \cap \Gamma)$ and $U\gamma := \bar{\pi}(\bar{\pi}^{-1}(K) \cap U\Gamma)$ and we estimate the distance of $U\gamma$ to UV :

$$\begin{aligned} d_\infty[U\gamma, UV] &= \inf_{\xi \in \bar{\pi}(\bar{\pi}^{-1}(K) \cap U\Gamma)} \inf_{\eta \in UV} d_\infty(\xi, \eta) \\ &= \inf_{(u, \xi) \in U\Gamma, u \in K} \inf_{(v, \eta) \in K \times UV} d_\infty(\xi, \eta) \\ &= \inf_{(u, \xi) \in U\Gamma, u \in K} \inf_{(v, \eta) \in K \times UV} d_\infty((u, \xi), (v, \eta)), \end{aligned}$$

where the last equality is due to the fact that one can choose $v = u$ in the minimization. We deduce that, by removing the constraint $u \in K$ in the minimization,

$$\begin{aligned} d_\infty[U\gamma, UV] &\geq \inf_{(u, \xi) \in U\Gamma} \inf_{(v, \eta) \in K \times UV} d_\infty((u, \xi), (v, \eta)) \\ &= d_\infty(K \times UV, U\Gamma) \geq 3\delta. \end{aligned}$$

STEP 2. Since γ and V are cones, the previous inequality implies $d_\infty(\xi, V) \geq 2\|\xi\|\delta$ for every $\xi \in \gamma$. For any $i \in I$ such that the ball B_i is centered at a point in $\text{supp } \chi$, the inclusion $\bar{B}_i \subset K$ implies $\bar{\pi}(\bar{\pi}^{-1}(\bar{B}_i) \cap \Gamma) \subset \gamma$. We hence also have

$$(4.4) \quad \forall \xi \in \bar{\pi}(\bar{\pi}^{-1}(\bar{B}_i) \cap \Gamma), \quad d_\infty(\xi, V) \geq 2\|\xi\|\delta.$$

We now set $\bar{W}_{i1} := \{\xi_1 \in (\mathbb{R}^{d_1})^*; d_\infty(\xi_1, \bar{\pi}(\bar{\pi}^{-1}(\bar{B}_{i1}) \cap \Gamma_1)) \leq \|\xi_1\|\delta\}$, $\bar{W}_{i2} := \{\xi_2 \in (\mathbb{R}^{d_2})^*; d_\infty(\xi_2, \bar{\pi}(\bar{\pi}^{-1}(\bar{B}_{i2}) \cap \Gamma_2)) \leq \|\xi_2\|\delta\}$ and $W_{i1} := \bar{W}_{i1} \setminus \{0\}$, $W_{i2} := \bar{W}_{i2} \setminus \{0\}$. By the definition of W_{i1} , $W_{i1}^c \cap \bar{\pi}(\bar{\pi}^{-1}(\bar{B}_{i1}) \cap \Gamma_1) = \emptyset$, which is equivalent to $(\bar{B}_{i1} \times W_{i1}^c) \cap \Gamma_1 = \emptyset$. Similarly $(\bar{B}_{i2} \times W_{i2}^c) \cap \Gamma_2 = \emptyset$.

On the other hand, since

$$\begin{aligned} \bar{\pi}(\bar{\pi}^{-1}(\bar{B}_{i1}) \cap \bar{\Gamma}_1) \times \bar{\pi}(\bar{\pi}^{-1}(\bar{B}_{i2}) \cap \bar{\Gamma}_2) &= \bar{\pi}[\bar{\pi}^{-1}(\bar{B}_{i1} \times \bar{B}_{i2}) \cap (\bar{\Gamma}_1 \times \bar{\Gamma}_2)] \\ &= \bar{\pi}[\bar{\pi}^{-1}(\bar{B}_i) \cap \bar{\Gamma}], \quad \bar{B}_i = \bar{B}_{i1} \times \bar{B}_{i2} \end{aligned}$$

because

$$\begin{aligned} \{\xi_1; \exists(x_1; \xi_1) \in \bar{\Gamma}_1, x_1 \in \bar{B}_{i1}\} \times \{\xi_2; \exists(x_2; \xi_2) \in \bar{\Gamma}_2, x_2 \in \bar{B}_{i2}\} \\ = \{(\xi_1, \xi_2); \exists(x_1, x_2; \xi_1, \xi_2) \in \bar{\Gamma}_1 \times \bar{\Gamma}_2, (x_1, x_2) \in \bar{B}_{i1} \times \bar{B}_{i2}\}, \end{aligned}$$

we also have

$$\begin{aligned} \bar{W}_{i1} \times \bar{W}_{i2} &= \{(\xi_1, \xi_2) \in (\mathbb{R}^d)^*; d_\infty[(\xi_1, \xi_2), \bar{\pi}(\bar{\pi}^{-1}(\bar{B}_i) \cap \Gamma)] \\ &\leq \sup(\|\xi_1\|, \|\xi_2\|)\delta\}. \end{aligned}$$

Hence by (4.4), we deduce that $\bar{W}_{i1} \times \bar{W}_{i2}$ does not meet V . ■

In the rest of the paper, we may identify abusively \mathbb{R}^d and $(\mathbb{R}^d)^*$. We also introduce the notation $e_\zeta(x, y) = e^{i(\xi \cdot x + \eta \cdot y)}$ where $\zeta = (\xi, \eta)$. To estimate

$\|u \otimes v\|_{N,V,\varphi_1 \otimes \varphi_2}$, we use Lemma 4.3 to find a partition of unity $(\psi_{j1} \otimes \psi_{j2})_{j \in J}$ which is finite on $\text{supp}(\varphi_1 \otimes \varphi_2)$ to write

$$\begin{aligned} \widehat{u\varphi_1}(\xi) \widehat{v\varphi_2}(\eta) &= \mathcal{F}(u\varphi_1 \otimes v\varphi_2)(\zeta) = \langle u \otimes v, (\varphi_1 \otimes \varphi_2)e_\zeta \rangle \\ &= \sum_j \langle u \otimes v, (\varphi_1 \psi_{j1} \otimes \varphi_2 \psi_{j2})e_\zeta \rangle = \sum_j \widehat{u\varphi_1 \psi_{j1}}(\xi) \widehat{v\varphi_2 \psi_{j2}}(\eta). \end{aligned}$$

Hence $\|u \otimes v\|_{N,V,\varphi_1 \otimes \varphi_2} \leq \sum_j \|u \otimes v\|_{N,V,\varphi_1 \psi_{j1} \otimes \varphi_2 \psi_{j2}}$, where the sum is finite. By the following lemma, each seminorm on the right hand side is bounded.

LEMMA 4.4. *Let Γ_1, Γ_2 and Γ be closed cones as in the previous lemma, $\psi_1 \in \mathcal{D}(\Omega_1), \psi_2 \in \mathcal{D}(\Omega_2)$ with $(\text{supp}(\psi_1 \otimes \psi_2) \times V) \cap \Gamma = \emptyset$, and closed cones W_1 and W_2 in $\mathbb{R}^{d_1} \setminus \{0\}$ and $\mathbb{R}^{d_2} \setminus \{0\}$ such that*

(4.5) $(W_1 \cup \{0\}) \times (W_2 \cup \{0\}) \cap V = \emptyset,$

(4.6) $(\text{supp } \psi_k \times W_k^c) \cap \Gamma_k = \emptyset \quad \text{for } k = 1, 2.$

Then, for every bounded set $A \subset \mathcal{D}'_{\Gamma_1}$ and every integer N , there are constants m, M_1, M_2 and a bounded set $B \subset \mathcal{D}(K)$, where K is an arbitrary compact neighborhood of $\text{supp } \psi_2$, such that

$$\|t_1 \otimes t_2\|_{N,V,\psi_1 \otimes \psi_2} \leq M_1 \|t_2\|_{N,C_\beta,\psi_2} + M_2 \|t_2\|_{N+m,C_\beta,\psi_2} + p_B(t_2)$$

for all $t_1 \in A$ and $t_2 \in \mathcal{D}'_{\Gamma_2}$, where C_β is any conic neighborhood of W_2 with compact base and p_B is a seminorm of the strong topology of $\mathcal{D}'(\mathbb{R}^{d_2})$.

Proof. We want to calculate

$$\|t_1 \otimes t_2\|_{N,V,\psi_1 \otimes \psi_2} = \sup_{\zeta \in V} (1 + |\zeta|)^N |\mathcal{F}(t_1 \psi_1 \otimes t_2 \psi_2)(\zeta)|.$$

We denote $u = t_1 \psi_1, v = t_2 \psi_2$ and $I = \widehat{u \otimes v}$. From $e_{(\xi,\eta)} = e_\xi \otimes e_\eta$ we find that $I(\xi, \eta) = \langle t, e_{(\xi,\eta)} \rangle = \langle u \otimes v, e_\xi \otimes e_\eta \rangle = \langle u, e_\xi \rangle \langle v, e_\eta \rangle = \widehat{u}(\xi) \widehat{v}(\eta)$. By the shrinking lemma we can slightly enlarge W_1 and W_2 to closed cones having the same properties. Thus, there are two homogeneous functions α and β on \mathbb{R}^{d_1} and \mathbb{R}^{d_2} , respectively, of degree zero, which are smooth except at the origin, nonnegative and bounded by 1, such that:

- (i) $\alpha|_{W_1 \cup \{0\}} = 1$ and $\beta|_{W_2 \cup \{0\}} = 1$;
- (ii) $(\text{supp } \alpha \times \text{supp } \beta) \cap V = \emptyset$;
- (iii) $(\text{supp } \psi_1 \times \text{supp}(1 - \alpha)) \cap \Gamma_1 = \emptyset$;
- (iv) $(\text{supp } \psi_2 \times \text{supp}(1 - \beta)) \cap \Gamma_2 = \emptyset$.

We can write $I = I_1 + I_2 + I_3 + I_4$ where (recalling that $\zeta = (\xi, \eta)$)

$$\begin{aligned} I_1(\zeta) &= \alpha(\xi) \widehat{u}(\xi) \beta(\eta) \widehat{v}(\eta), \\ I_2(\zeta) &= \alpha(\xi) \widehat{u}(\xi) (1 - \beta)(\eta) \widehat{v}(\eta), \\ I_3(\zeta) &= (1 - \alpha)(\xi) \widehat{u}(\xi) \beta(\eta) \widehat{v}(\eta), \\ I_4(\zeta) &= (1 - \alpha)(\xi) \widehat{u}(\xi) (1 - \beta)(\eta) \widehat{v}(\eta). \end{aligned}$$

We have $I_1(\zeta) = 0$ because, by condition (ii), $\alpha(\xi)\beta(\eta) = 0$ for $(\xi, \eta) \in V$. Condition (iii) implies that

$$|(1 - \alpha)(\xi)\widehat{u}(\xi)| \leq \sup_{\xi \in \widehat{C}_\alpha} |t_1\psi_1(\xi)| \leq (1 + |\xi|)^{-N} \|t_1\|_{N, C_\alpha, \psi_1},$$

where $\xi \in C_\alpha = \text{supp}(1 - \alpha)$. This gives, with $C_\beta = \text{supp}(1 - \beta)$,

$$\begin{aligned} |I_4(\zeta)| &\leq (1 + |\xi|)^{-N} (1 + |\eta|)^{-N} \|t_1\|_{N, C_\alpha, \psi_1} \|t_2\|_{N, C_\beta, \psi_2} \\ &\leq (1 + |\zeta|)^{-N} \|t_1\|_{N, C_\alpha, \psi_1} \|t_2\|_{N, C_\beta, \psi_2}, \end{aligned}$$

because $1 + |(\xi, \eta)| \leq 1 + |\xi| + |\eta| \leq (1 + |\xi|)(1 + |\eta|)$. Since the set A is bounded in \mathcal{D}'_{T_1} , there is a constant $M_1 = \sup_{t_1 \in A} \|t_1\|_{N, C_\alpha, \psi_1}$ such that $|I_4(\zeta)| \leq (1 + |\zeta|)^{-N} M_1 \|t_2\|_{N, C_\beta, \psi_2}$.

To estimate I_2 , we use the fact that, $u = t_1\psi_1$ being a compactly supported distribution, there is an integer m such that, for all $t_1 \in A$,

$$|\alpha(\xi)\widehat{u}(\xi)| \leq |\widehat{u}(\xi)| \leq (1 + |\xi|)^m \|\theta^{-m}\widehat{u}\|_{L^\infty}.$$

As for I_4 , we get $|(1 - \beta)(\eta)\widehat{v}(\eta)| \leq (1 + |\eta|)^{-N-m} \|t_2\|_{N+m, C_\beta, \psi_2}$. The set $\overline{\{\zeta \in \text{supp } \alpha \times C_\beta; |\zeta| = 1\} \cap V}$ is **compact** and avoids the set of elements of the form $\zeta = (\xi, 0)$ with $\xi \in \text{supp } \alpha \setminus \{0\}$. Otherwise, we would find some sequence $(\xi_n, \eta_n) \rightarrow (\xi, 0) \in ((\text{supp } \alpha \times \{0\}) \cap V) \subset ((\text{supp } \alpha \times \text{supp } \beta) \cap V)$, which contradicts condition (ii). Let $\epsilon > 0$ be the smallest value of $|\eta|$ in this set. Then, the functions α and β being homogeneous of degree zero, $(\text{supp } \alpha \times C_\beta) \cap V$ is a cone in \mathbb{R}^d and $|\eta|/|\zeta| \geq \epsilon$ for all $\zeta = (\xi, \eta) \in (\text{supp } \alpha \times C_\beta) \cap V$. Thus, $(1 + |\eta|)^{-N-m} \leq \epsilon^{-N-m} (1 + |(\xi, \eta)|)^{-N-m}$ and

$$|I_2(\zeta)| \leq \|\theta^{-m}t_1\widehat{\psi}_1\|_{L^\infty} \|t_2\|_{N+m, C_\beta, \psi_2} \epsilon^{-N-m} (1 + |\zeta|)^{-N} \quad \text{for every } \zeta \in V,$$

because $|\xi| \leq |(\eta, \xi)|$. We now prove an intermediate lemma:

LEMMA 4.5. *Let Ω be an open set in \mathbb{R}^d and B a bounded set in $\mathcal{D}'(\Omega)$. Then for every $\chi \in \mathcal{D}(\Omega)$ there exist an integer M and a constant C (both depending only on B and on an arbitrary relatively compact open neighborhood of $\text{supp } \chi$) such that*

$$\sup_{u \in B} \sup_{\xi \in \mathbb{R}^n} (1 + |\xi|)^{-M} |\widehat{u\chi}(\xi)| < 2^M C \text{Vol}(K) \pi_{M, K}(\chi),$$

where $\pi_{M, K}(\chi) = \sup_{x \in K, |\alpha| \leq M} |\partial^\alpha \chi(x)|$ and $K = \text{supp } \chi$.

Proof. Let Ω_0 be a relatively compact open neighborhood of $K = \text{supp } \chi$. According to Schwartz [16, p. 86], for any bounded set B in $\mathcal{D}'(\Omega)$, there is an integer M (depending only on B and Ω_0) such that every $u \in B$ can be expressed in Ω_0 as $u = \partial^\alpha f_u$ for $|\alpha| \leq M$, where f_u is a continuous function. Moreover, there is a constant C (depending only on B and Ω_0) such that

$|f_u(x)| \leq C$ for all $x \in \Omega_0$ and $u \in B$. Thus,

$$\begin{aligned} \widehat{u\chi}(\xi) &= \int_{\Omega_0} e^{-i\xi \cdot x} \chi(x) \partial^\alpha f_u(x) dx = (-1)^{|\alpha|} \int_{\Omega_0} f_u(x) \partial^\alpha (e^{-i\xi \cdot x} \chi(x)) dx \\ &= (-1)^{|\alpha|} \sum_{\beta \leq \alpha} \binom{\alpha}{\beta} (i\xi)^\beta \int_{\Omega_0} f_u(x) e^{-i\xi \cdot x} \partial^{\beta-\alpha} \chi(x) dx. \end{aligned}$$

By using $|(i\xi)^\beta| \leq (1 + |\xi|)^M$ if $|\beta| \leq M$ we obtain

$$\begin{aligned} (1 + |\xi|)^{-M} |\widehat{u\chi}(\xi)| &\leq \sup_{|\alpha| \leq M} \sum_{\beta \leq \alpha} \binom{\alpha}{\beta} \left| \int_{\Omega_0} f_u(x) e^{-i\xi \cdot x} \partial^{\beta-\alpha} \chi(x) dx \right| \\ &\leq 2^M C \text{Vol}(K) \pi_{M,K}(\chi). \blacksquare \end{aligned}$$

By Lemma 4.5, $\|\theta^{-m} \widehat{t_1 \psi_1}\|_{L^\infty}$ is uniformly bounded for $t_1 \in A$ by a constant $M'_2 = \sup_{t_1 \in A} \|\theta^{-m} t_1 \psi_1\|_{L^\infty}$ with $\theta = 1 + |\xi|$. Therefore, there is a constant $M_2 = M'_2 \epsilon^{-N-m}$ such that, for all $t_1 \in A$ and $t_2 \in \mathcal{D}'_{\Gamma_2}$,

$$|I_2(\zeta)| \leq M_2 \|t_2\|_{N+m, C_\beta, \psi_2}.$$

The term I_3 is treated differently because we want to get the following result: for every bounded set A in \mathcal{D}'_{Γ_1} and every seminorm $\|\cdot\|_{N, V, \chi}$, there is a bounded set $B \subset \mathcal{D}(\Omega_2)$ such that for all $\zeta \in V$,

$$I_3(\zeta) \leq p_B(t_2)(1 + |\zeta|)^{-N} \quad \text{for every } t_2 \in \mathcal{D}'_{\Gamma_2}.$$

This special form of (3.1) is possible because the union of bounded sets is bounded and multiplying a bounded set by a positive constant M yields a bounded set.

We write $I_3(\zeta) = \langle t_2, f_\zeta \rangle$, where

$$f_{(\xi, \eta)}(y) = (1 - \alpha)(\xi) \widehat{u}(\xi) \beta(\eta) \psi_2(y) e_\eta(y);$$

we must show that $B = \{(1 + |\zeta|)^N f_\zeta; \zeta \in V\}$ is a bounded subset of $\mathcal{D}(\Omega_2)$. A subset B of $\mathcal{D}(\Omega_2)$ is bounded if and only if there is a compact set K and, for every integer n , a constant M_n such that $\text{supp } f \subset K$ and $\pi_{n,K}(f) \leq M_n$ for all $f \in B$. All f_ζ are supported on $\text{supp } \psi_2$ and are smooth functions because ψ_2 and e_η are smooth. We have to prove that if t_1 runs over a bounded set in \mathcal{D}'_{Γ_1} , then there are constants M_n such that $\pi_{n,K}(f_\zeta) \leq M_n$ for all $\zeta \in V$, where K is a compact neighborhood of $\text{supp } \psi_2$. We start from

$$\pi_{n,K}(f_{(\xi, \eta)}) = |(1 - \alpha)(\xi) \widehat{u}(\xi) \beta(\eta)| \pi_{n,K}(\psi_2 e_\eta).$$

We notice that $\pi_{n,K}(\psi_2 e_\eta) \leq 2^n \pi_{n,K}(\psi_2) \pi_{n,K}(e_\eta)$ and $\pi_{n,K}(e_\eta) \leq |\eta|^n$. As for the estimate of I_2 , we have

$$|(1 - \alpha)(\xi) \widehat{u}(\xi)| \leq (1 + |\xi|)^{-N-n} \|t_1\|_{N+n, C_\alpha, \psi_1}$$

as $(\text{supp } \varphi_1 \times \text{supp}(1 - \alpha)) \cap \Gamma_1 = \emptyset$, and moreover $(1 + |\xi|)^{-N-n} \leq \epsilon^{-N-n}(1 + |(\xi, \eta)|)^{-N-n}$ for some ϵ as $(\xi, \eta) \in V \cap (\text{supp}(1 - \alpha) \times \text{supp } \beta)$. Therefore

$$\pi_{n,K}(f_\zeta) \leq \|t_1\|_{N+n, C_\alpha, \psi_1} 2^n \pi_{n,K}(\psi_2) \epsilon^{-N-n} (1 + |\zeta|)^{-N},$$

because $|\eta|^n (1 + |(\xi, \eta)|)^{-n} \leq 1$. If t_1 belongs to a bounded subset A of \mathcal{D}'_{Γ_1} , then for each N , $\|t_1\|_{N, C_\alpha, \psi_1}$ is uniformly bounded. Finally,

$$|I_3(\zeta)| \leq p_B(t_2)(1 + |\zeta|)^{-N}. \blacksquare$$

For each j , the conditions of the lemma hold if we set $\psi_1 = \varphi_1 \psi_{j1}$, $\psi_2 = \varphi_2 \psi_{j2}$, $W_1 = W_{j1}$ and $W_2 = W_{j2}$. Thus, for every bounded set A in \mathcal{D}'_{Γ_1} , every $u \in A$ and every $v \in \mathcal{D}'_{\Gamma_2}$ we have

$$\begin{aligned} \|u \otimes v\|_{N, V, \varphi_1 \otimes \varphi_2} &\leq \sum_j \|u \otimes v\|_{N, V, \varphi_1 \psi_{j1} \otimes \varphi_2 \psi_{j2}} \\ &\leq \sum_j [M_{1j} \|v\|_{N, C_{\beta_j}, \varphi_2 \psi_{j2}} + M_2 \|v\|_{N+m, C_{\beta_j}, \varphi_2 \psi_{j2}} + p_{B_j}(v)]. \end{aligned}$$

Since the sum over j is finite, the family of maps $u \times v \mapsto u \otimes v$, where $u \in A$, is equicontinuous for any bounded set $A \subset \mathcal{D}'_{\Gamma_1}$. By the symmetry of the problem, we can prove similarly that the family of maps $u \times v \mapsto u \otimes v$, where $v \in B$, is equicontinuous for any bounded set $B \subset \mathcal{D}'_{\Gamma_2}$. Finally, we have proved

THEOREM 4.6. *Let $\Omega_1 \subset \mathbb{R}^{d_1}$ and $\Omega_2 \subset \mathbb{R}^{d_2}$ be open sets, $\Gamma_1 \in \dot{T}^* \Omega_1$ and $\Gamma_2 \in \dot{T}^* \Omega_2$ be closed cones, and set*

$$\Gamma = (\Gamma_1 \times \Gamma_2) \cup ((\Omega_1 \times \{0\}) \times \Gamma_2) \cup (\Gamma_1 \times (\Omega_2 \times \{0\})).$$

Then the tensor product $(u, v) \mapsto u \otimes v$ is hypocontinuous from $\mathcal{D}'_{\Gamma_1} \times \mathcal{D}'_{\Gamma_2}$ to \mathcal{D}'_Γ in the normal topology.

5. The pull-back. The purpose of this section is to prove

PROPOSITION 5.1. *Let $\Omega_1 \subset \mathbb{R}^{d_1}$ and $\Omega_2 \subset \mathbb{R}^{d_2}$ be open sets and Γ a closed cone in $\dot{T}^* \Omega_2$. Let $f : \Omega_1 \rightarrow \Omega_2$ be a smooth map such that $N_f \cap \Gamma = \emptyset$, where $N_f = \{(f(x); \eta) \in \Omega_2 \times \mathbb{R}^{d_2}; \eta \circ df_x = 0\}$, and define $f^* \Gamma = \{(x; \eta \circ df_x); (f(x); \eta) \in \Gamma\}$, where*

$$\begin{aligned} \eta \circ df_x &:= \sum_{j=1}^{d_2} \eta_j d(y^j \circ f)_x = \sum_{j=1}^{d_2} \eta_j dy^j \circ df_x \\ &= \sum_{j=1}^{d_2} \eta_j df_x^j = \sum_{j=1}^{d_2} \sum_{i=1}^{d_1} \eta_j \frac{\partial f^j}{\partial x^i} dx^i. \end{aligned}$$

Then the pull-back operation $f^ : \mathcal{D}'_\Gamma(\Omega_2) \rightarrow \mathcal{D}'_{f^* \Gamma}(\Omega_1)$ is continuous for the normal topology.*

We will show this by proving that $\langle f^*u, v \rangle$ is continuous for every v in an equicontinuous set. Before doing so, we characterize the equicontinuous sets of the normal topology, which is of independent interest.

5.1. Equicontinuous subsets. Let Ω be open in \mathbb{R}^d and Γ be a closed cone in $\dot{T}^*\Omega$. We define the open cone $\Lambda = \{(x; \xi) \in \dot{T}^*\Omega; (x; -\xi) \notin \Gamma\}$ and consider the space $\mathcal{E}'_\Lambda(\Omega)$ of compactly supported distributions $v \in \mathcal{E}'(\Omega)$ such that $\text{WF}(v) \subset \Lambda$.

According to Definition 3.2, a set H is equicontinuous in $\mathcal{E}'_\Lambda(\Omega)$ (which is the strong dual of $\mathcal{D}'_\Gamma(\Omega)$ [5]) if and only if there are a finite number of seminorms $\|\cdot\|_{N_1, V_1, \chi_1}, \dots, \|\cdot\|_{N_k, V_k, \chi_k}$ of $\mathcal{D}'_\Gamma(\Omega)$, a bounded subset B_0 of $\mathcal{D}(\Omega)$ and a constant M such that

$$(5.1) \quad |\langle u, v \rangle| \leq M \sup\{\|u\|_{N_1, V_1, \chi_1}, \dots, \|u\|_{N_k, V_k, \chi_k}, p_{B_0}(u)\}$$

for all $u \in \mathcal{D}'_\Gamma(\Omega)$ and $v \in H$. There is only one seminorm p_{B_0} because these seminorms are saturated [13, p. 107] in $\mathcal{D}'(\Omega)$ with the strong topology. The following theorem will be useful to prove the continuity of linear maps [13, p. 200]:

THEOREM 5.2. *If E is a locally convex space and $f : E \rightarrow \mathcal{D}'_\Gamma(\Omega)$ is a linear map, then f is continuous if and only if, for every equicontinuous subset H of $\mathcal{E}'_\Lambda(\Omega)$, the seminorm $p_H : E \rightarrow \mathbb{R}$ defined by $p_H(x) = \sup_{v \in H} |\langle f(x), v \rangle|$ is continuous.*

The equicontinuous sets of $\mathcal{E}'_\Lambda(\Omega)$ are known:

LEMMA 5.3. *A subset B of $\mathcal{E}'_\Lambda(\Omega)$ is equicontinuous if and only if there are:*

- (i) *a compact set $K \subset \Omega$ containing the supports of all elements of B ;*
- (ii) *a closed cone $\Xi \subset \Lambda$ such that $B \subset \mathcal{D}'_\Xi(\Omega)$, B is bounded in $\mathcal{D}'_\Xi(\Omega)$ and $\underline{\pi}(\Xi) \subset K$.*

Proof. We first prove that every such B is equicontinuous. We showed in [5] that $\mathcal{E}'_\Lambda(\Omega)$ is the inductive limit of the spaces $E_\ell = \{v \in \mathcal{E}'_\Lambda(\Omega); \text{supp } v \subset L_\ell, \text{WF}(v) \subset \Lambda_\ell\}$, where the compact sets L_ℓ exhaust Ω and the closed cones Λ_ℓ exhaust Λ . Thus, there is an integer ℓ such that $\Xi \subset \Lambda_\ell$ and $B \subset E_\ell$. The inclusion of Ξ in Λ_ℓ implies that every seminorm $\|\cdot\|_{N, V, \chi}$ of E_ℓ is also a seminorm of $\mathcal{D}'_\Xi(\Omega)$ because $\text{supp } \chi \times V$ does not meet Ξ if it does not meet Λ_ℓ . Thus, B is bounded in E_ℓ , and [5, (8)] gives

$$\sup_{v \in B} |\langle u, v \rangle| \leq \sum_j (p_{B_j}(u) + \|u\|_{m+n+1, V_j, \chi_j} C I_n^{n+1} + \|u\|_{n, V_j, \chi_j} M_{n, W_j, \chi_j} I_n^{2n}),$$

which can be converted to the equicontinuity condition (5.1).

To show the converse, we denote by B the set of all $v \in \mathcal{E}'_\Lambda(\Omega)$ that satisfy (5.1). Then, by following [5, proof of Prop. 7], we find that the supports of

all elements of B are included in a compact set $K = \bigcup_j \text{supp } \chi_j \cup K'$, where K' is a compact set containing the supports of all $f \in B_0$. Moreover, the wave front sets of all elements of B are contained in $\Xi = \bigcup_j \text{supp } \chi_j \times (-V_j)$.

It remains to show that B is bounded in $\mathcal{D}'_{\Xi}(\Omega)$ for the normal topology. We first notice that if $\text{supp } \chi \times (-V) \subset \Xi$, then $\|\cdot\|_{N,V,\chi}$ is a continuous seminorm of the strong dual $\mathcal{E}'_{\Xi^c}(\Omega)$ of $\mathcal{D}'_{\Xi}(\Omega)$. Indeed, it was shown in [5, proof of Prop. 7] that we have $\|u\|_{N,V,\chi} = \sup_{\xi \in V} |\langle u, f_{\xi} \rangle|$, where $f_{\xi}(x) = (1 + |\xi|)^N \chi(x) e^{-i\xi \cdot x}$ and the set $\{f_{\xi}; \xi \in V\}$ is bounded in $\mathcal{D}'_{\Xi}(\Omega)$. If B' is a bounded set in $\mathcal{E}'_{\Xi^c}(\Omega)$, the continuous seminorms $\|u\|_{N,V,\chi}$ and $p_{B_0}(u)$ of $\mathcal{E}'_{\Xi^c}(\Omega)$ appearing on the right hand side of (5.1) are bounded over B' . Thus, for any bounded set B' in $\mathcal{E}'_{\Xi^c}(\Omega)$, taking $u \in B'$ and taking the sup in (5.1) over $u \in B'$ shows that $\sup_{u \in B', v \in B} |\langle u, v \rangle|$ is bounded and B is a bounded subset of $\mathcal{D}'_{\Xi}(\Omega)$ when $\mathcal{D}'_{\Xi}(\Omega)$ is equipped with the strong $\beta(\mathcal{D}'_{\Xi}, \mathcal{E}'_{\Xi^c})$ topology. It is shown in [5, Theorem 33] that the bounded sets of $\mathcal{D}'_{\Gamma}(\Omega)$ coincide for the strong and the normal topologies. Thus, B is bounded for the normal topology. ■

We obtain the following characterization of continuous linear maps:

THEOREM 5.4. *Let E be a locally convex space, Ω an open subset of \mathbb{R}^d and Γ a closed cone in $\dot{T}^*\Omega$. A linear map $f : E \rightarrow \mathcal{D}'_{\Gamma}(\Omega)$ is continuous if and only if every map $f_B : E \rightarrow \mathbb{R}$ defined by $f_B(x) = \sup_{v \in B} |\langle f(x), v \rangle|$ is continuous, where B is equicontinuous in $\mathcal{E}'_{\Lambda}(\Omega)$ with $\Lambda = \Gamma^c$.*

The equicontinuous sets of $\mathcal{E}'_{\Lambda}(\Omega)$ intervene also because the duality pairing enjoys a sort of hypocontinuity where, for $\mathcal{E}'_{\Lambda}(\Omega)$, the bounded sets are replaced by the equicontinuous ones:

THEOREM 5.5 ([14, p. 157]). *Let the duality pairing $\mathcal{D}'_{\Gamma}(\Omega) \times \mathcal{E}'_{\Lambda}(\Omega) \rightarrow \mathbb{K}$ be defined by $u \times v \mapsto f(u, v) = \langle u, v \rangle$. Then, for every bounded subset A of $\mathcal{D}'_{\Gamma}(\Omega)$ and every equicontinuous subset B of $\mathcal{E}'_{\Lambda}(\Omega)$ the sets of maps $\{f_u; u \in A\}$ and $\{f_v; v \in B\}$ are equicontinuous.*

5.2. Continuity of pull-back

Strategy of the proof. Let Ω_1 and Ω_2 be open sets in \mathbb{R}^{d_1} and \mathbb{R}^{d_2} , respectively. Let $f : \Omega_1 \rightarrow \Omega_2$ be a smooth map and Γ be a closed cone in $\dot{T}^*\Omega_2$. We want to show that the pull-back $f^* : \mathcal{D}'_{\Gamma}(\Omega_2) \rightarrow \mathcal{D}'_{f^*\Gamma}(\Omega_1)$ is continuous for the normal topology. According to Theorem 5.4, the pull-back is continuous if and only if, for every equicontinuous set $B \subset \mathcal{E}'_{\Lambda}(\Omega_2)$ (where $\Lambda = (f^*\Gamma)^c$) the family of maps $(\rho_v)_{v \in B}$ defined by $\rho_v : u \mapsto \langle f^*u, v \rangle$ is equicontinuous, which implies that $\sup_{v \in B} |\langle f^*u, v \rangle|$ is continuous in u . By Lemma 5.3, there is a compact set $K \subset \Omega_1$ and a closed cone $\Xi \subset (f^*\Gamma^c)$ such that $\text{supp } v \subset K$ and $\text{WF}(v) \subset \Xi$ for all $v \in B$. Choose $\chi \in \mathcal{D}(\Omega_1)$ such that $\chi|_K = 1$. If $(\varphi_i)_{i \in I}$ is a partition of unity on Ω_2 , we can write

$\langle f^*u, v \rangle = \sum_i \langle f^*(u\varphi_i), v\chi \rangle$. The image of $\text{supp } \chi$ by f being compact [2, p. 19], only a finite number of terms of this sum are nonzero and the family ρ_v is equicontinuous if and only if, for every $\varphi \in \mathcal{D}(\Omega_2)$, the family of maps $u \mapsto \langle f^*(u\varphi), v\chi \rangle$ is equicontinuous.

Stationary phase and Schwartz kernels. In order to calculate the pairing between $f^*(u\varphi)$ and v , we first notice that, when u is a locally integrable function, then $u\varphi(y) = \mathcal{F}^{-1}(\widehat{u\varphi})(y) = (2\pi)^{-d_2} \int_{\mathbb{R}^{d_2}} d\eta e^{i\eta \cdot y} \widehat{u\varphi}(\eta)$, so that $f^*(u\varphi)(x) = (2\pi)^{-d_2} \int_{\mathbb{R}^{d_2}} d\eta e^{i\eta \cdot f(x)} \widehat{u\varphi}(\eta)$ and

$$\begin{aligned} \langle f^*(u\varphi), \chi v \rangle &= \frac{1}{(2\pi)^{d_2}} \int_{\Omega_1} \int_{\mathbb{R}^{d_2}} \chi(x)v(x)e^{i\eta \cdot f(x)} \widehat{u\varphi}(\eta) dx d\eta \\ &= \frac{1}{(2\pi)^{d_2}} \int_{\mathbb{R}^{d_2}} \int_{\Omega_1} \int_{\mathbb{R}^{d_2}} \chi(x)v(x)e^{i\eta \cdot f(x)} e^{-iy \cdot \eta} u(y)\varphi(y) dy dx d\eta. \end{aligned}$$

This definition can be extended to any distribution $u \in \mathcal{D}'_\Gamma$ as

$$(5.2) \quad \langle f^*(u\varphi), \chi v \rangle = \frac{1}{(2\pi)^d} \int_{\mathbb{R}^d} u(y)v(x)I(x, y) dx dy,$$

where $d = d_1 + d_2$ and $I(x, y) = \int_{\mathbb{R}^{d_2}} e^{i\eta \cdot (f(x) - y)} \varphi(y)\chi(x) d\eta$. The duality pairing can also be written as $\langle f^*(u\varphi), v\chi \rangle = \langle v \otimes u, I \rangle$. Note that $I(x, y) = (2\pi)^{-d_2} \chi(x)\varphi(y) \int d\eta e^{i\eta \cdot (f(x) - y)}$ is an oscillatory integral [12] with symbol $\chi(x)\varphi(y)$ and phase $\eta \cdot (f(x) - y)$ where $\eta \cdot (f(x) - y)$ is homogeneous of degree 1 with respect to η , and for all $\eta \neq 0, d(\eta \cdot (f(x) - y)) \neq 0$. Therefore, $I \in \mathcal{D}'(\Omega_1 \times \Omega_2)$ is the Schwartz kernel of the bilinear continuous map $(u, v) \mapsto \langle f^*(u\varphi), v\chi \rangle$.

Proof of Proposition 5.1. By Theorem 4.6, the map $(v, u) \mapsto v \otimes u$ is hypocontinuous from $\mathcal{D}'_\Xi \times \mathcal{D}'_\Gamma$ to $\mathcal{D}'_{\Gamma_\otimes}$ where $\Gamma_\otimes = \Xi \times \Gamma \cup (\Omega_1 \times \{0\}) \times \Gamma \cup \Xi \times (\Omega_2 \times \{0\})$. Let Λ_\otimes be the open cone Γ_\otimes^c . Therefore by Theorem 5.5, the family of duality pairings $\mathcal{D}'_{\Gamma_\otimes} \ni u \otimes v \mapsto \langle u \otimes v, r \rangle$ is equicontinuous from $\mathcal{D}'_{\Gamma_\otimes}$ to \mathbb{K} uniformly in $r \in B'$ for every equicontinuous set B' of $\mathcal{E}'_{\Lambda_\otimes}$. In particular, if B' contains only the element I , then the map $v \otimes u \mapsto \langle v \otimes u, I \rangle$ is continuous since I is compactly supported in $\text{supp } \chi \times \text{supp } \varphi$ and its wave front set is contained in Λ_\otimes . Thus, if $\text{WF}(I) \subset \Lambda_\otimes$, then the map $(v, u) \mapsto \langle v \otimes u, I \rangle$ is hypocontinuous by the next lemma.

LEMMA 5.6. *The composition of a hypocontinuous map and a continuous linear map is hypocontinuous.*

Proof. Let $f : E \times F \rightarrow G$ be a hypocontinuous map and $g : G \rightarrow H$ a continuous linear map. The map $g \circ f$ is hypocontinuous if and only if, for every bounded set $B \subset F$ and every neighborhood W of zero in H , there is a neighborhood U of zero in E such that $(g \circ f)(U \times B) \subset W$ (with

a similar condition for $(g \circ f)(A \times V)$). By the continuity of g , there is a neighborhood Z of zero in G such that $g(Z) \subset W$. By the hypocontinuity of f , there is a neighborhood U of zero in E such that $f(U \times B) \subset Z$. Thus, $(g \circ f)(U \times B) \subset g(Z) \subset W$. ■

Therefore, the map $(v, u) \mapsto \langle f^*(u\varphi), \chi v \rangle$ is hypocontinuous; by item (i) of Definition 3.1, this implies that the family of maps $\rho_v : u \mapsto \langle f^*(u\varphi), \chi v \rangle$ with $v \in B$ is equicontinuous. It just remains to check that $\text{WF}(I) \subset \Lambda_\otimes$, i.e. $\text{WF}(I)'$ does not meet Γ_\otimes . We have $\text{WF}(I) \subset \{(x, f(x); -\eta \circ d_x f, \eta) ; x \in \text{supp } \chi\}$ [12, p. 260]. Recall $\Xi \subset (f^* \Gamma)^c = \{(x; -\eta \circ df_x) ; (f(x); \eta) \notin \Gamma\}$. By definition of Γ_\otimes the following three conditions have to be satisfied:

- $\Xi \times \Gamma \cap \text{WF}(I)' = \emptyset$ because it is the set of points $(x, f(x); -\eta \circ d_x f, \eta)$ such that $(f(x); \eta) \notin \Gamma$ by definition of Ξ and $(f(x); \eta) \in \Gamma$ by definition of Γ ;
- $\Xi \times (\Omega_2 \times \{0\}) \cap \text{WF}(I)' = \emptyset$ because we would need $\eta = 0$ whereas $(y, \eta) \in \Gamma$ implies $\eta \neq 0$;
- $(\text{supp } \chi \times \{0\}) \times \Gamma \cap \text{WF}(I)' \subset \{(x, f(x); 0, \eta) ; x \in \text{supp } \chi, \eta \circ df_x = 0, (f(x); \eta) \in \Gamma\}$.

Thus, if $f^* \Gamma \cap N_f = \emptyset$, then $\text{WF}(I)' \cap \Gamma_\otimes = \emptyset$ and the pull-back is continuous.

How to write the pull-back operator in terms of the Schwartz kernel I ? Relationship with the product of distributions. We start from a linear operator $L : \mathcal{D}(\mathbb{R}^{d_2}) \rightarrow \mathcal{D}'(\mathbb{R}^{d_1})$ with Schwartz kernel $K \in \mathcal{D}'(\mathbb{R}^{d_1} \times \mathbb{R}^{d_2})$. Using the standard operations on distributions, we can make sense of the well-known representation formula $Lu = \int_{\mathbb{R}^{d_2}} K(x, y)u(y) dy$ for an operator $L : \mathcal{D}(\mathbb{R}^{d_2}) \rightarrow \mathcal{D}'(\mathbb{R}^{d_1})$ and its kernel $K \in \mathcal{D}'(\mathbb{R}^{d_1} \times \mathbb{R}^{d_2})$. Let $\pi_2 : \mathbb{R}^{d_1} \times \mathbb{R}^{d_2} \ni (x, y) \mapsto y \in \mathbb{R}^{d_2}$ and $\pi_1 : \mathbb{R}^{d_1} \times \mathbb{R}^{d_2} \ni (x, y) \mapsto x \in \mathbb{R}^{d_1}$. Then we define

$$K(x, y)u(y) = K(x, y)\pi_2^*u(x, y) = K(x, y)(1(x) \otimes u(y))$$

where $\pi_2^*u = 1 \otimes u$ and $\int_{\mathbb{R}^{d_2}} dy f(x, y) = \pi_{1*}f(x)$. Therefore

$$(5.3) \quad Lu = \int_{\mathbb{R}^{d_2}} K(x, y)u(y) dy = \pi_{1*}(K(\pi_2^*u)).$$

The interest of the formula $Lu = \pi_{1*}(K(\pi_2^*u))$ is that everything generalizes to oriented manifolds. Replace \mathbb{R}^{d_2} (resp. \mathbb{R}^{d_1}) with a manifold M_2 (resp. M_1) with smooth volume densities $|\omega_2|$ (resp. $|\omega_1|$). The duality pairing is defined as the extension of the usual integration against the volume densities, for instance:

$$\forall (u, \varphi) \in C^\infty(M_1) \times \mathcal{D}(M_1), \quad \langle u, \varphi \rangle_{M_1} = \int_{M_1} (u\varphi)\omega_1.$$

Finally, for the continuous linear map $L : \mathcal{D}(\mathbb{R}^{d_2}) \ni u \mapsto \chi f^*(u\varphi) \in \mathcal{D}'(\mathbb{R}^{d_1})$, we get the formula $Lu = \pi_{1*}(I(\pi_2^*u))$ where

$$I(x, y) = (2\pi)^{-d_2} \chi(x)\varphi(y) \int d\eta e^{i\eta \cdot (f(x)-y)}$$

is the Schwartz kernel of L .

5.3. Pull-back by families of smooth maps. To renormalize quantum field theory in curved spacetimes, it is crucial to pull back by a family of smooth maps. We start with a simple lemma.

LEMMA 5.7. *Let Ω_1, Ω_2, U be open sets in $\mathbb{R}^{d_1}, \mathbb{R}^{d_2}, \mathbb{R}^n$ respectively. For any compact sets $K_1 \subset \Omega_1, K_2 \subset \Omega_2, A \subset U$ and any smooth map $f : \Omega_1 \times U \rightarrow \Omega_2$, the conic set*

$$\Gamma = \{(x, f(x, a); -\eta \circ d_x f(x, a), \eta); (x, a, f(x, a)) \in K_1 \times A \times K_2, \eta \neq 0\}$$

is closed in $\dot{T}^*(\Omega_1 \times \Omega_2)$.

Proof. Let $(x, y; \xi, \eta) \in \bar{\Gamma}$ with $(\xi, \eta) \neq (0, 0)$. Then there is a sequence $(x_k, f(x_k, a_k); -\eta_k \circ d_x f(x_k, a_k), \eta_k) \in \Gamma, (x_k, a_k, f(x_k, a_k)) \in K_1 \times A \times K_2$ which converges to $(x, y; \xi, \eta)$. By compactness of A , we extract a convergent subsequence $a_k \rightarrow a$. By continuity of $d_x f$, we find that $\xi = -\eta \circ d_x f(x, a)$; moreover $\lim_{k \rightarrow \infty} f(x_k, a_k) = f(x, a) \in K_2$ since K_2 is closed; and $\eta \neq 0$ since otherwise $\xi = 0, \eta = 0$. Therefore $(x, y; \xi, \eta) \in \Gamma$ by definition. Finally, $\bar{\Gamma} \subset \Gamma$, hence Γ is closed. ■

PROPOSITION 5.8. *Let Ω_1 be an open set in $\mathbb{R}^{d_1}, \Omega_2$ an open set in \mathbb{R}^{d_2} , and $A \subset U \subset \mathbb{R}^n$ where A is compact and U is open. Let $\chi \in \mathcal{D}(\Omega_1), \varphi \in \mathcal{D}(\Omega_2)$ and $f : \Omega_1 \times U \rightarrow \Omega_2$ a smooth map. Then:*

- (1) *The family $(I_{f(\cdot, a)})_{a \in A}$ of distributions formally defined by*

$$I_{f(\cdot, a)}(x, y) = \chi(x)\varphi(y) \int_{\mathbb{R}^{d_2}} \frac{d\theta}{(2\pi)^{d_2}} e^{i\theta \cdot (f(x, a)-y)}$$

is a bounded set in \mathcal{D}'_Γ , where Γ is the closed cone in $\dot{T}^(\Omega_1 \times \Omega_2)$ defined by*

$$\Gamma = \{(x, f(x, a); -\eta \circ d_x f(x, a), \eta); x \in \text{supp } \chi, f(x, a) \in \text{supp } \varphi, a \in A, \eta \neq 0\}.$$

- (2) *For any open cone Λ containing Γ , $(I_{f(\cdot, a)})_{a \in A}$ is equicontinuous in $\mathcal{E}'_\Lambda(\Omega_1 \times \Omega_2)$.*

In the proof we will use the push-forward Theorem 6.3, whose proof will be given in Section 7.

Proof of Proposition 5.8. From Lemma 5.3 and the fact that $(I_{f(\cdot,a)})_{a \in A}$ is supported in a fixed compact set $\text{supp } \chi \times \text{supp } \varphi$, we deduce that conclusion (2) follows from (1), so it suffices to prove (1).

The conic set Γ is closed by Lemma 5.7. To prove that the family $(I_{f(\cdot,a)})_{a \in A}$ is bounded in \mathcal{D}'_Γ , by [5, Proposition 1] it suffices to check that $\sup_{a \in A} |\langle I_{f(\cdot,a)}, v \rangle| < \infty$ for all $v \in \mathcal{E}'_{\Gamma^c}(\Omega_1 \times \Omega_2)$.

STEP 1. Our goal is to study the map $a \mapsto \int_{\Omega_1 \times \Omega_2} I_f(x, y, a)v(x, y) dx dy$, where

$$I_f(x, y, a) = \chi(x)\varphi(y) \int_{\mathbb{R}^{d_2}} \frac{d\theta}{(2\pi)^{d_2}} e^{i\theta \cdot (f(x,a)-y)}.$$

Let π_{12}, π_3 be projections from $\Omega_1 \times \Omega_2 \times U$ defined by $\pi_{12}(x, y, a) = (x, y)$ and $\pi_3(x, y, a) = a$. Using the dictionary of Subsection 5.2, if v were a test function, then we would find that

$$(5.4) \quad \int_{\Omega_1 \times \Omega_2} I_f(x, y, \cdot)v(x, y) dx dy = \pi_{3*}(I_f \pi_{12}^* v) \in \mathcal{D}'(U).$$

We want to prove that $a \mapsto \int_{\Omega_1 \times \Omega_2} I_f(x, y, a)v(x, y) dx dy$ is smooth in some open neighborhood of A since this will imply that

$$\sup_{a \in A} \left| \int_{\Omega_1 \times \Omega_2} I_f(x, y, a)v(x, y) dx dy \right| = \sup_{a \in A} |\langle I_{f(\cdot,a)}, v \rangle| < \infty.$$

In order to do so, it suffices to prove that the condition $v \in \mathcal{E}'_{\Gamma^c}$ implies that the distributional product $I_f(x, y, a)v(x, y) = I_f(\pi_{12}^* v)(x, y, a)$ makes sense in $\mathcal{D}'(\Omega_1 \times \Omega_2 \times U)$ and the push-forward $\pi_{3*}(I_f \pi_{12}^* v) = \int_{\Omega_1 \times \Omega_2} I_f(x, y, \cdot)v(x, y) dx dy$ has empty wave front set over some open neighborhood of A .

STEP 2. $\text{WF}(I_f)$ is the set of all

$$\begin{pmatrix} x; -\theta \circ d_x f \\ f(x, a); \theta \\ a; -\theta \circ d_a f \end{pmatrix}$$

such that $x \in \text{supp } \varphi$, $f(x, a) \in \text{supp } \chi$, $a \in U$, $\theta \neq 0$. And $\text{WF}(\pi_{12}^* v)$ is the set of all $\begin{pmatrix} x; \xi \\ y; \eta \\ a; 0 \end{pmatrix}$ such that $\begin{pmatrix} x; \xi \\ y; \eta \end{pmatrix} \in \text{WF}(v)$.

Moreover $v \in \mathcal{E}'_{\Gamma^c}$ implies $\text{WF}(v) \cap \Gamma' = \emptyset$ so that $\xi \neq -\eta \circ d_x f$ and

$$\forall \theta, \quad \begin{pmatrix} \xi - \theta \circ d_x f \\ \theta + \eta \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix} \text{ has no solution.}$$

Observe that

$$\text{WF}(I_f) + \text{WF}(\pi_{12}^*v) = \left\{ \begin{pmatrix} x; \xi - \theta \circ d_x f \\ f(x); \theta + \eta \\ a; -\theta \circ d_a f \end{pmatrix}; \theta \in \mathbb{R}^d \setminus \{0\} \right\}$$

implies $(\text{WF}(I_f) + \text{WF}(\pi_{12}^*v)) \cap \underline{0} = \emptyset$ and $(\text{WF}(I_f) \cup \text{WF}(\pi_{12}^*v)) \cap \underline{0} = \emptyset$.

STEP 3. In the last step, we shall prove that the condition $\text{WF}(v) \cap \Gamma' = \emptyset$ actually implies that $\text{WF}(\pi_{3*}(I_f \pi_{12}^*v))$ is empty over some open neighborhood U' of A . The condition $\text{WF}(v) \cap \Gamma' = \emptyset$ implies the existence of some open neighborhood U' of A such that

$$\forall a \in U', \forall (x, f(x, a); \xi, \eta) \in \text{WF}(v), \quad \xi \neq -\eta \circ d_x f(x, a).$$

Since A and $\text{supp } v$ are compact and $A \times \text{WF}(v)$ is closed, we can find $\delta > 0$ such that

$$\forall (a, (x, f(x, a); \xi, \eta)) \in A \times \text{WF}(v), \quad |\xi + \eta \circ d_x f(x, a)| \geq \delta|\eta|.$$

Define

$$U' = \{a \in U; \forall (x, f(x, a); \xi, \eta) \in \text{WF}(v), |\xi + \eta \circ d_x f(x, a)| > (\delta/2)|\eta|\}.$$

The condition $\text{WF}(v) \cap \Gamma' = \emptyset$ ensures that $\pi_{3*}(I_f \pi_{12}^*v)$ is well defined in $\mathcal{D}'_\emptyset(U') = C^\infty(U')$. Hence

$$a \mapsto \langle v, I_{f(\cdot, a)} \rangle = \int_{\Omega_1 \times \Omega_2} I_f(x, y, a)v(x, y) \, dx \, dy$$

is smooth on U' , a fortiori continuous on the compact set A , which means that $\sup_{a \in A} |\langle v, I_{f(\cdot, a)} \rangle| < \infty$. ■

THEOREM 5.9. *Let $\Omega_1 \subset \mathbb{R}^{d_1}$ and $\Omega_2 \subset \mathbb{R}^{d_2}$ be open sets, let $A \subset U \subset \mathbb{R}^n$ where A is compact and U open, and let Γ be a closed cone in $\dot{T}^*\Omega_2$. Let $f : \Omega_1 \times U \rightarrow \Omega_2$ be a smooth map such that $f(\cdot, a)^*\Gamma$ meets the zero section $\underline{0}$ for no $a \in A$, and set $\Theta = \bigcup_{a \in A} f(\cdot, a)^*\Gamma$. Then for all seminorms P_B of $\mathcal{D}'_\Theta(\Omega_1)$ and all $u \in \mathcal{D}'_\Gamma(\Omega_2)$, the family $P_B(f(\cdot, a)^*u)_{a \in A}$ is bounded.*

Proof. We need to prove that $\sup_{(v,a) \in B \times A} |\langle f(\cdot, a)^*u, v \rangle| < \infty$ for any equicontinuous subset B of $\mathcal{E}'_{\Theta^c}(\Omega_1)$. It follows from Lemma 5.3 that there exists a closed cone Ξ such that $\Xi \cap \Theta' = \emptyset$ and $B \subset \mathcal{D}'_\Xi(\Omega_1)$. Set

$$\Gamma_\otimes = (\Xi \times \Gamma) \cup ((\Omega_1 \times \{0\}) \times \Gamma) \cup (\Xi \times (\Omega_2 \times \{0\})).$$

Let Λ_\otimes be the open cone defined by $\Lambda_\otimes = (\Gamma_\otimes)^c$. By Proposition 5.8, we can easily verify as in the proof of Proposition 5.1 that the family $(I_{f(\cdot, a)})_{a \in A}$ is equicontinuous in $\mathcal{E}'_{\Lambda_\otimes}$. Then we prove much as in the proof of the pull-back Proposition 5.1 that the family of maps $\rho_{v,a} : u \mapsto \langle f(\cdot, a)^*(u\varphi), \chi v \rangle$ with $(v, a) \in B \times A$ is equicontinuous where B is equicontinuous in $\mathcal{E}'_{\Theta^c}(\Omega_1)$ and

χ is chosen in such a way that $\chi|_{\text{supp } B} = 1$ and $\varphi|_{f(\text{supp } B)} = 1$. Therefore,

$$\sup_{(v,a) \in B \times A} |\langle f(\cdot, a)^* u, v \rangle| = \sup_{(v,a) \in B \times A} |\langle f(\cdot, a)^*(u\varphi), \chi v \rangle| < \infty,$$

and the result follows from Theorem 5.4. ■

6. Product and push-forward of distributions. Hörmander noticed that the product of distributions u and v can be described as the composition of the tensor product $(u, v) \mapsto u \otimes v$ with the pull-back by the map $f : x \mapsto (x, x)$. If the wave front sets of u and v are contained in Γ_1 and Γ_2 , then the wave front set of $u \otimes v$ is contained in $\Gamma_{\otimes} = (\Gamma_1 \times \Gamma_2) \cup ((\Omega_1 \times \{0\}) \times \Gamma_2) \cup (\Gamma_1 \times (\Omega_2 \times \{0\}))$, and the pull-back is well defined if the set $N_f = \{(x, x; \eta_1, \eta_2); (\eta_1 + \eta_2) \circ dx = 0\}$, which is the conormal bundle of the diagonal $\Delta \subset \mathbb{R}^n \times \mathbb{R}^n$, does not meet Γ , i.e. if there is no point $(x; \eta)$ in Γ_1 such that $(x; -\eta)$ is in Γ_2 . This gives

THEOREM 6.1. *Let $\Omega \subset \mathbb{R}^n$ be an open set and Γ_1, Γ_2 be closed cones in $\dot{T}^*\Omega$ such that $\Gamma_1 \cap \Gamma_2 = \emptyset$. Then the product of distributions is hypocontinuous for the normal topology from $\mathcal{D}'_{\Gamma_1} \times \mathcal{D}'_{\Gamma_2}$ to \mathcal{D}'_{Γ} , where*

$$(6.1) \quad \Gamma = (\Gamma_1 \times_{\Omega} \Gamma_2) \cup ((\Omega \times \{0\}) \times_{\Omega} \Gamma_2) \cup (\Gamma_1 \times_{\Omega} (\Omega \times \{0\})).$$

Proof. The product of distributions is the composition of the hypocontinuous tensor product with the continuous pull-back (see Lemma 5.6). ■

This theorem has the useful corollary:

COROLLARY 6.2. *Let $\Omega \subset \mathbb{R}^n$ be an open set and Γ be a closed cone in $\dot{T}^*\Omega$. Then the product of a smooth map and a distribution is hypocontinuous for the normal topology from $C^\infty(\Omega) \times \mathcal{D}'_{\Gamma}$ to \mathcal{D}'_{Γ} .*

Proof. By Lemma 7.2, $C^\infty(\Omega)$ and \mathcal{D}'_{\emptyset} are topologically isomorphic. Therefore, the corollary follows by applying Theorem 6.1 to $\Gamma_1 = \emptyset$ and $\Gamma_2 = \Gamma$. Equation (6.1) shows that the wave front set of the product is in Γ . ■

6.1. The push-forward as a consequence of the pull-back theorem

THEOREM 6.3. *Let $\Omega_1 \subset \mathbb{R}^{d_1}$ and $\Omega_2 \subset \mathbb{R}^{d_2}$ be two open sets and Γ a closed cone in $\dot{T}^*\Omega_1$. For any smooth map $f : \Omega_1 \rightarrow \Omega_2$ and any closed subset C of Ω_1 such that $f|_C : C \rightarrow \Omega_2$ is proper and $\pi(\Gamma) \subset C$, the map f_* is continuous in the normal topology from $\{u \in \mathcal{D}'_{\Gamma}; \text{supp } u \subset C\}$ to $\mathcal{D}'_{f_*\Gamma}$, where $f_*\Gamma = \{(y; \eta) \in \dot{T}^*\Omega_2; \exists x \in \Omega_1, y = f(x) \text{ and } (x; \eta \circ df_x) \in \Gamma \cup \text{supp } u \times \{0\}\}$.*

Proof. The idea of the proof is to think of a push-forward as the adjoint of a pull-back [8]. For all B equicontinuous in \mathcal{E}'_{Λ} where $\Lambda = f_*\Gamma'^c$, and

all $v \in B$, $\text{supp}(f^*v) \cap C$ is contained in a fixed compact set K since f is proper on C . Hence, for any $\chi \in \mathcal{D}(\mathbb{R}^d)$ such that $\chi|_C = 1$ and f is proper on $\text{supp } \chi$, we should have at least formally $\langle f_*u, v \rangle = \langle u, \chi f^*v \rangle$ if the duality pairings make sense. On the one hand, if $v \in \mathcal{E}'_A(\Omega_2)$ where f^*A does not meet the zero section $\underline{0} \subset T^*\Omega_1$ then the pull-back f^*v would be well defined by the pull-back Proposition 5.1 (which is equivalent to the fact that $N_f \cap A = \emptyset$). On the other hand, the duality pairing $\langle u, \chi(f^*v) \rangle$ is well defined if $f^*A \cap \Gamma' = \emptyset$. Combining both conditions leads to the requirement that $f^*A \cap (\Gamma' \cup \underline{0}) = \emptyset$. But note that:

$$\begin{aligned} (f^*A) \cap (\Gamma' \cup \underline{0}) &= \emptyset \\ \Leftrightarrow \{ (x; \eta \circ df); (f(x); \eta) \in A, (x; \eta \circ df) \in \Gamma' \cup \{0\} \} &= \emptyset \\ \Leftrightarrow [(f(x); \eta) \in A \Rightarrow (x; \eta \circ df) \notin \Gamma' \cup \{0\}] & \\ \Leftrightarrow [(f(x); \eta) \in A' \Rightarrow (x; \eta \circ df) \notin \Gamma \cup \{0\}], & \end{aligned}$$

which is equivalent to the fact that A' does not meet $f_*\Gamma = \{(f(x); \eta); (x; \eta \circ df) \in \Gamma \cup \underline{0}, \eta \neq 0\} \subset T^*\Omega_2$, which is exactly the assumption of our theorem. Therefore, the set of distributions χf^*B is supported in a fixed compact set, bounded in $\mathcal{D}'_{f^*A}(\Omega_1)$ by the pull-back Proposition 5.1 applied to f^* , the duality pairings are well defined, and $\sup_{v \in B} |\langle f_*u, v \rangle| = \sup_{v \in B'} |\langle u, v \rangle|$ where $B' = \chi f^*B$ is equicontinuous in $\mathcal{E}'_{\Gamma'}(\Omega_1)$ (the support of the distribution is compact because f is proper), which means that $\sup_{v \in B'} |\langle u, v \rangle|$ is a continuous seminorm for the normal topology of $\mathcal{D}'_{\Gamma'}(\Omega_1)$. ■

We now prove a parameter version of the push-forward theorem.

THEOREM 6.4. *Let $\Omega_1 \subset \mathbb{R}^{d_1}$ and $\Omega_2 \subset \mathbb{R}^{d_2}$ be open sets, $A \subset U \subset \mathbb{R}^n$, A is compact, U is open and Γ a closed cone in $\dot{T}^*\Omega_1$. For any smooth map $f : \Omega_1 \times U \rightarrow \Omega_2$ and any closed subset C of Ω_1 such that $f : C \times A \rightarrow \Omega_2$ is proper and $\pi(\Gamma) \subset C$, the function $f(\cdot, a)_*$ is uniformly continuous in the normal topology from $\{u \in \mathcal{D}'_{\Gamma}; \text{supp } u \subset C\}$ to \mathcal{D}'_{Ξ} , where $\Xi = \bigcup_{a \in A} f(\cdot, a)_*\Gamma$.*

Proof. We have to check that Ξ is closed over $\dot{T}^*_{f(C \times A)}\Omega_2$. Let $(y; \eta) \in \overline{\Xi} \cap \dot{T}^*_{f(C \times A)}\Omega_2$. Then there exists a sequence $(y_k; \eta_k) \rightarrow (y; \eta)$ such that $(y_k; \eta_k) \in \Xi \cap \dot{T}^*_{f(C \times A)}\Omega_2$. By definition, $y_k = f(x_k, a_k)$ where $(x_k, \eta_k \circ d_x f(x_k, a_k)) \in \Gamma \cup \underline{0}$ and $(x_k, a_k) \in C \times A$. The central observation is that $\{y_k; k \in \mathbb{N}\} \subset f(C \times A)$ and $\{(x_k, a_k); f(x_k, a_k) = y_k, k \in \mathbb{N}\} \subset C \times A$ are compact sets because f is proper on $C \times A$. Then we can extract convergent subsequences $(x_k, a_k) \rightarrow (x, a)$ and $(x, \eta \circ d_x f(x, a)) \in \Gamma \cup \underline{0}$ since $\Gamma \cup \underline{0}$ is closed in $\dot{T}^*\Omega_2$ and $d_x f$ is continuous. By definition of Ξ , this proves that $(y; \eta) \in \Xi$ and we conclude that Ξ is closed.

Now we can repeat the proof of the push-forward theorem except that we use the pull-back theorem with parameters. Let B be equicontinuous in $\mathcal{E}'_{\Xi^c}(\Omega_2)$, hence all elements of B have support contained in some compact set. For every $v \in B$, $\text{supp}(f^*v) \cap C \times A$ is compact, therefore for $\chi = 1$ on $\bigcup_{a \in A} \text{supp}(f(\cdot, a)^*v) \cap C$, the family $B' = \{(\chi f(\cdot, a)^*v); a \in A, v \in B\}$ is equicontinuous in \mathcal{E}'_{Θ} where $\Theta = \Gamma^c$ by the parameter version of the pull-back theorem. Therefore, $u \mapsto \sup_{a \in A} \sup_{v \in B} |\langle f(\cdot, a)_*u, v \rangle| = \sup_{v \in B'} |\langle u, v \rangle|$ is continuous in u since the right hand term is a continuous seminorm for the normal topology of $\mathcal{D}'_{\Gamma}(\Omega_1)$. ■

Convolution product. In the same spirit as for multiplication of distributions, the convolution product $u * v$ can be described as the composition of the tensor product $(u, v) \mapsto u \otimes v$ with the push-forward by the map $\Sigma : (x, y) \mapsto (x + y)$. For a closed subset $X \subset \mathbb{R}^n$, let $\mathcal{D}'_{\Gamma}(X)$ be the set of distributions supported in X with wave front set in Γ .

THEOREM 6.5. *Let Γ_1, Γ_2 be closed conic sets in $\dot{T}^*\mathbb{R}^n$ and X_1, X_2 closed subsets of \mathbb{R}^n such that $\Sigma : X_1 \times X_2 \rightarrow \mathbb{R}^n$ is proper. Then the convolution product of distributions is hypocontinuous from $\mathcal{D}'_{\Gamma_1}(X_1) \times \mathcal{D}'_{\Gamma_2}(X_2)$ to $\mathcal{D}'_{\Gamma}(X_1 + X_2)$ where*

$$(6.2) \quad \Gamma = \{(x + y; \eta); (x; \eta) \in \Gamma_1, (y; \eta) \in \Gamma_2\}.$$

Proof. The convolution product of distributions is the composition of the hypocontinuous tensor product with the continuous push-forward. ■

As an application of the parameter version of the push-forward theorem, we state the coordinate invariant definition of the wave front set, which was proposed by Duistermaat [8, p. 13], correcting a first attempt by Gabor [9]. Its proof is left to the reader.

THEOREM 6.6. *Let $\Omega \subset \mathbb{R}^d$ be an open set, and $u \in \mathcal{D}'(\Omega)$. An element $(x; \xi) \notin \text{WF}(u)$ if and only if for all $f \in C^\infty(\Omega \times \mathbb{R}^n)$ such that $d_x f(x, a_0) = \xi$ for some $a_0 \in \mathbb{R}^n$, there exist neighborhoods A of a_0 and U of x_0 such that for all $\varphi \in \mathcal{D}(U)$, $|\langle u, \varphi e^{i\tau f(\cdot, a)} \rangle| = O(\tau^{-\infty})$ uniformly in some neighborhood of a_0 in A .*

7. Appendix: Technical results. This appendix gathers different useful results. Several of them are folklore results for which we could find no proof in the literature.

7.1. Exhaustion of the complement of Γ . To prove that \mathcal{D}'_{Γ} is nuclear [5], we need to take the additional seminorms in a countable set: the complement $\Gamma^c = \dot{T}^*M \setminus \Gamma$ of any closed cone $\Gamma \subset \dot{T}^*M$ can be exhausted by a countable set of products $U \times V$, where $U \subset M$ is compact and V is a closed conic subset of \mathbb{R}^n .

First we introduce the sphere bundle over M (or unit cotangent bundle) $UT^*M = \{(x; k) \in T^*M; |k| = 1\}$. We then define the set

$$UT_K = \{(x; k) \in \Gamma; x \in K, |k| = 1\} = UT^*M|_K \cap \Gamma,$$

for any compact subset $K \subset M$. Since M can be covered by a countable union of compact sets, we assume without loss of generality that K is covered by a single chart (U, ψ) such that $\psi(K) \subset Q$, where $Q = [-1, 1]^n$ is an n -dimensional cube. Hence we can assume that $M = \mathbb{R}^n$ and $UT^*M = \mathbb{R}^n \times S^{n-1}$. It will be convenient to use the norm d_∞ on \mathbb{R}^n , defined by

$$d_\infty(x, y) := \sup_{1 \leq i \leq n} |x_i - y_i| \quad \text{for } x, y \in \mathbb{R}^n.$$

We denote by $\overline{B}_\infty(x, r) = \{y \in \mathbb{R}^n; d_\infty(x, y) \leq r\}$ the closed ball of radius r for this norm. We also denote the restriction of d_∞ to $S^{n-1} \times S^{n-1}$ by the same letter and, lastly, for $(x; \xi), (y; \eta) \in UT^*\mathbb{R}^n$ we set $d_\infty((x; \xi), (y; \eta)) = \sup(d_\infty(x, y), d_\infty(\xi, \eta))$.

We define cubes centered at rational points in Q : Let $q_j = [-1/2^j, 1/2^j]^n = \overline{B}_\infty(0, 2^{-j})$ and $q_{j,m} = m/2^j + q_j = \overline{B}_\infty(2^{-j}m, 2^{-j})$, where $m \in \mathbb{Z}^n \cap 2^j Q$. In other words, the center of $q_{j,m}$ runs over a hypercubic lattice with coordinates $(2^{-j}m_1, \dots, 2^{-j}m_n)$, where $-2^j \leq m_i \leq 2^j$. Note that, for each nonnegative integer j , the hypercubes $q_{j,m}$ overlap and cover Q :

$$(7.1) \quad Q \subset \bigcup_{m \in \mathbb{Z}^n \cap 2^j Q} q_{j,m}.$$

Denote by $\pi : UT^*\mathbb{R}^n \rightarrow \mathbb{R}^n$ and $\bar{\pi} : UT^*\mathbb{R}^n \rightarrow S^{n-1}$ the projection maps defined by $\pi(x; k) = x$ and $\bar{\pi}(x; k) = k$. We define $F_{j,m} = \pi^{-1}(q_{j,m}) \simeq q_{j,m} \times S^{n-1}$ (see Fig. 1). The set $\bar{\pi}(F_{j,m} \cap UT_K)$ is compact because the projection




Fig. 1. The case $n = 2$, where only one dimension of the cube $[-1, 1]^2$ is shown and the circle S^1 is represented by the vertical segment $[0, 1]$. The large surface is UT_K , the small square around p is the ball $\overline{B}_\infty(p, \delta)$. We see that $q_{j,m}$ contains x and is contained in $\pi(\overline{B}_\infty(p, \delta))$.

$\bar{\pi}$ is continuous and $F_{j,m} \cap U\Gamma_K$ is compact. For any positive integer ℓ , define the compact set $C_{j,m,\ell} = \{\eta \in S^{n-1}; d_\infty(\bar{\pi}(F_{j,m} \cap U\Gamma_K), \eta) \geq 1/\ell\}$. This is the set of points of the sphere which are at least at distance $1/\ell$ from the projection of the slice of $U\Gamma$ inside $F_{j,m}$ (see Fig. 1). We have

$$\bigcup_{\ell > 0} C_{j,m,\ell} = S^{n-1} \setminus \bar{\pi}(F_{j,m} \cap U\Gamma_K).$$

Indeed, by definition, any element of $C_{j,m,\ell}$ is in S^{n-1} and not in $\bar{\pi}(F_{j,m} \cap U\Gamma_K)$; conversely, the compactness of $\bar{\pi}(F_{j,m} \cap U\Gamma_K)$ implies that any point $(x; \xi)$ in $S^{n-1} \setminus \bar{\pi}(F_{j,m} \cap U\Gamma_K)$ is at a finite distance δ from $\bar{\pi}(F_{j,m} \cap U\Gamma_K)$. If we take $\ell > 1/\delta$, we have $(x; \xi) \in C_{j,m,\ell}$. Note that all $C_{j,m,\ell}$ are empty if $\bar{\pi}(F_{j,m} \cap U\Gamma_K) = S^{n-1}$. With this notation we can now state

LEMMA 7.1.

$$(7.2) \quad \bigcup_{j,m,\ell} q_{j,m} \times C_{j,m,\ell} = UT^*M|_K \setminus U\Gamma_K,$$

and, by denoting $V_{j,m,\ell} = \{k \in \mathbb{R}^n \setminus \{0\}; k/|k| \in C_{j,m,\ell}\}$,

$$(7.3) \quad \bigcup_{j,m,\ell} q_{j,m} \times V_{j,m,\ell} = (T^*M \setminus \Gamma)|_K.$$

Proof. We first prove \subset in (7.2). Let $(x; k) \in \bigcup_{j,m,\ell} q_{j,m} \times C_{j,m,\ell}$, so there exist $j \in \mathbb{N}$, $m \in \mathbb{Z}^n \cap 2^j Q$ and $\ell \in \mathbb{N}^*$ such that $(x; k) \in q_{j,m} \times C_{j,m,\ell}$. Hence $(x; k) \in F_{j,m}$ and, by definition of $C_{j,m,\ell}$, $d_\infty(\bar{\pi}(F_{j,m} \cap U\Gamma_K), k) \geq 1/\ell$, which implies that $(x; k) \notin U\Gamma_K$.

Let us prove the reverse inclusion \supset . Let $(x; k) \in UT^*M|_K \setminus U\Gamma_K$. Since this set is open, there exists $\delta > 0$ such that $\bar{B}_\infty((x; k), \delta) \subset UT^*M|_K \setminus U\Gamma_K$. Let $j \in \mathbb{N}^*$ with $2^{-j+1} < \delta$. Because the sets $q_{j,m}$ cover Q (see (7.1)), there is an m such that $x \in q_{j,m}$. Moreover, for all $y \in q_{j,m}$, we have

$$d_\infty(x, y) \leq d_\infty(x, 2^{-j}m) + d_\infty(2^{-j}m, y) \leq 2^{-j} + 2^{-j} < \delta,$$

i.e. $y \in \bar{B}_\infty(x, \delta)$. Hence $q_{j,m} \subset \bar{B}_\infty(x, \delta)$. We deduce that

$$q_{j,m} \times \bar{B}_\infty(k, \delta) \subset \bar{B}_\infty(x, \delta) \times \bar{B}_\infty(k, \delta) = \bar{B}_\infty((x; k), \delta) \subset UT^*M|_K \setminus U\Gamma_K.$$

This means that $(q_{j,m} \times \bar{B}_\infty(k, \delta)) \cap U\Gamma_K = \emptyset$ or equivalently

$$F_{j,m} \cap \bar{\pi}^{-1}(\bar{B}_\infty(k, \delta)) \cap U\Gamma_K = \emptyset.$$

Hence $\bar{B}_\infty(k, \delta) \cap \bar{\pi}(F_{j,m} \cap U\Gamma_K) = \emptyset$, so $d_\infty(k, \bar{\pi}(F_{j,m} \cap U\Gamma_K)) > \delta$. By choosing $\ell \in \mathbb{N}^*$ with $1/\ell \leq \delta$, we deduce that $d_\infty(k, \bar{\pi}(F_{j,m} \cap U\Gamma_K)) > 1/\ell$, i.e. $k \in C_{j,m,\ell}$. Thus $(x; k) \in q_{j,m} \times C_{j,m,\ell}$ and (7.2) is proved.

To prove (7.3), we notice that, because of the conic property of Γ , each $C_{j,m,\ell}$ corresponds to a unique $V_{j,m,\ell}$. ■

Any nonnegative smooth function ψ supported on $[-3/2, 3/2]^n$ and such that $\psi(x) = 1$ for $x \in [-1, 1]^n$ enables us to define scaled and shifted func-

tions $\psi_{j-1,m}(x) = \psi(2^j(x-m))$ supported on $q_{j-1,m}$ and equal to 1 on $q_{j,2m}$. If $C_{j,m,\ell}$ is not empty, we denote by $\alpha_{j,m,\ell} : S^{n-1} \rightarrow \mathbb{R}$ a smooth function supported on $C_{j,m,\ell}$ and equal to 1 on $C_{j,m,\ell+1}$. Note that if $C_{j,m,\ell}$ is a proper subset of S^{n-1} , then it is strictly included in $C_{j,m,\ell+1}$.

7.2. Equivalence of topologies. Grigis and Sjöstrand stated [10, p. 80] that if we have a family χ_α of test functions and closed cones V_α such that $(\text{supp } \chi_\alpha \times V_\alpha) \cap \Gamma = \emptyset$ and $\bigcup_\alpha \{(x, k); \chi_\alpha(x) \neq 0 \text{ and } k \in \check{V}_\alpha\} = \Gamma^c$, then the topology of \mathcal{D}'_Γ is given by the seminorms of the weak topology and the seminorms $\|\cdot\|_{N, V_\alpha, \chi_\alpha}$. By covering M with a countable family of compact sets K_i described in Section 7.1, we see that Lemma 7.1 gives a family of indices $\alpha = (i, j, \ell)$, functions $\chi_{j,m,\ell} = \psi_{j,m}$ and cones $V_{j,m,\ell}$ adapted to K_i such that the conditions of the Grigis–Sjöstrand lemma are satisfied. Therefore, the normal topology is described by the seminorms of the strong topology of $\mathcal{D}'(\Omega)$ and by a countable family of seminorms indexed by (i, j, m, ℓ) .

7.3. Topological equivalence of $C^\infty(X)$ and \mathcal{D}'_\emptyset . As an application of the previous lemma, we show

LEMMA 7.2. *The spaces $C^\infty(X)$ and \mathcal{D}'_\emptyset are topologically isomorphic.*

Proof. The two spaces are identical as vector spaces because a distribution u whose wave front set is empty is smooth everywhere, since its singular support $\text{sing supp}(u) = \pi(\text{WF}(u))$ [12, p. 254] is empty [12, p. 42].

To prove the topological equivalence, we must show that the two inclusions $\mathcal{D}'_\emptyset \hookrightarrow C^\infty(X)$ and $C^\infty(X) \hookrightarrow \mathcal{D}'_\emptyset$ are continuous. Recall that a system of seminorms defining the topology of $C^\infty(X)$ is $\pi_{m,K}$, where m runs over the integers and K runs over the compact subsets of X [16, p. 88]. By a straightforward estimate, we obtain

$$\begin{aligned} \pi_{m,K}(\varphi) &\leq C_n(2\pi)^{-n} \sum_{|\alpha| \leq m} \sup_{k \in \mathbb{R}^n} (1 + |k|^2)^p |k^\alpha \widehat{\varphi} \chi(k)| \\ &\leq C_n(2\pi)^{-n} \binom{m+n}{n} \|\varphi\|_{m+2p, \mathbb{R}^n, \chi}, \end{aligned}$$

where $\chi \in \mathcal{D}(X)$ is equal to one on a compact set whose interior contains K and where we have used $1 + |k|^2 \leq (1 + |k|)^2$, $|k^\alpha| \leq (1 + |k|)^m$ and $\sum_{|\alpha| \leq m} 1 = \binom{m+n}{n}$. Thus, every seminorm of $C^\infty(X)$ is bounded by a seminorm of \mathcal{D}'_\emptyset , and the injection $\mathcal{D}'_\emptyset \hookrightarrow C^\infty(X)$ is continuous.

Conversely, for any closed conic set V and any $\chi \in \mathcal{D}(X)$ we have $\|\varphi\|_{N, V, \chi} \leq \|\varphi\|_{N, \mathbb{R}^n, \chi}$. Thus, it is enough to estimate $\|\varphi\|_{N, \mathbb{R}^n, \chi}$. We also find that, for any integer N and $\alpha = 0$,

$$\sup_{k \in \mathbb{R}^n} (1 + k^2)^N |\widehat{\varphi} \chi(k)| \leq |K| 2^N \pi_{2N, K}(\varphi \chi),$$

where K is the support of φ . Then the inequality $1 + |k| \leq 2(1 + |k|^2)$ and the Leibniz rule give $\|\varphi\|_{N, \mathbb{R}^n, \chi} \leq |K| 8^N \pi_{2N, K}(\chi) \pi_{2N, K}(\varphi)$, so the seminorms $\|\cdot\|_{N, V, \chi}$ are controlled by the seminorms of $C^\infty(X)$. For the seminorms of $\mathcal{D}'(X)$, it is well known that the inclusion $C^\infty(X) \hookrightarrow \mathcal{D}'(X)$ is continuous [16, p. 420] when $\mathcal{D}'(X)$ has its strong topology. Therefore, it is also continuous when $\mathcal{D}'(X)$ is equipped with the weak topology. Thus we have proved that $C^\infty(X)$ and \mathcal{D}'_\emptyset are topologically isomorphic, where \mathcal{D}'_\emptyset can be equipped with the Hörmander or the normal topology. ■

Acknowledgements. We are grateful to Yoann Dabrowski for his generous help in the functional analysis parts of the paper. We thank Camille Laurent-Gengoux for discussions about the geometrical aspects of the pull-back. The research of N. V. Dang was partly supported by the Labex CEMPI (ANR-11-LABX-0007-01).

References

- [1] S. Alesker, *Valuations on manifolds and integral geometry*, *Geom. Funct. Anal.* 20 (2010), 1073–1143.
- [2] G. E. Bredon, *Topology and Geometry*, Springer, New York, 1993.
- [3] Ch. Brouder, N. V. Dang, and F. Hélein, *A smooth introduction to the wavefront set*, *J. Phys. A* 47 (2014), 443001
- [4] J. Chazarain et A. Piriou, *Introduction à la théorie des équations aux dérivées partielles*, Gauthier-Villars, Paris, 1981.
- [5] Y. Dabrowski and Ch. Brouder, *Functional properties of Hörmander’s space of distributions having a specified wavefront set*, *Comm. Math. Phys.* 332 (2014), 1345–1380.
- [6] N. V. Dang, *The extension of distributions on manifolds, a microlocal approach*, *Ann. Henri Poincaré* 17 (2016), 819–859.
- [7] N. V. Dang, *Renormalization of quantum field theory on curved space-times, a causal approach*, Ph.D. thesis, Paris Diderot Univ., 2013; arXiv:1312.5674.
- [8] J. J. Duistermaat, *Fourier Integral Operators*, Birkhäuser, Boston, 1996.
- [9] A. Gabor, *Remarks on the wave front of a distribution*, *Trans. Amer. Math. Soc.* 170 (1972), 239–244.
- [10] A. Grigis and J. Sjöstrand, *Microlocal Analysis for Differential Operators*, Cambridge Univ. Press, London, 1994.
- [11] L. Hörmander, *Fourier integral operators. I*, *Acta Math.* 127 (1971), 79–183.
- [12] L. Hörmander, *The Analysis of Linear Partial Differential Operators I. Distribution Theory and Fourier Analysis*, 2nd ed., Springer, Berlin, 1990.
- [13] J. Horváth, *Topological Vector Spaces and Distributions*, Addison-Wesley, Reading, 1966.
- [14] G. Köthe, *Topological Vector Spaces II*, Springer, New York, 1979.
- [15] J. Kučera and K. McKennon, *Continuity of multiplication of distributors*, *Int. J. Math. Math. Sci.* 4 (1981), 819–822.
- [16] L. Schwartz, *Théorie des distributions*, Hermann, Paris, 1966.
- [17] F. Trèves, *Topological Vector Spaces, Distributions and Kernels*, Dover, New York, 2007.

Christian Brouder
Sorbonne Universités, UPMC Univ. Paris 06
CNRS UMR 7590
and
Muséum National d'Histoire Naturelle, IRD UMR 206
and
Institut de Minéralogie, de Physique des Matériaux
et de Cosmochimie
4 place Jussieu
F-75005 Paris, France

Nguyen Viet Dang
Institut Camille Jordan
Université Claude Bernard Lyon 1
43 boulevard du 11 novembre 1918
69622 Villeurbanne Cedex, France
E-mail: dangnguyenviet20@gmail.com

Frédéric Hélein
Institut de Mathématiques de Jussieu Paris Rive Gauche
Université Denis Diderot Paris 7
Bâtiment Sophie Germain
75205 Paris Cedex 13, France