

Simplest possible locally definable well-orders

by

Peter Holy and Philipp Lücke (Bonn)

Abstract. We study the provable consequences of the existence of a well-order of $H(\kappa^+)$ definable by a Σ_1 -formula over the structure $\langle H(\kappa^+), \in \rangle$ in the case where κ is an uncountable regular cardinal. This is accomplished by constructing partial orders that force the existence of such well-orders while preserving many structural features of the ground model. We will use these constructions to show that the existence of a well-order of $H(\omega_2)$ that is definable over $\langle H(\omega_2), \in \rangle$ by a Σ_1 -formula with parameter ω_1 is consistent with a failure of the GCH at ω_1 . Moreover, we will show that one can achieve this situation also in the presence of a measurable cardinal. In contrast, results of Woodin imply that the existence of such a well-order is incompatible with the existence of infinitely many Woodin cardinals with a measurable cardinal above them all.

1. Introduction. Given an infinite cardinal κ , if the set of all subsets of κ is constructible from some subset z of κ , then there is a well-order of the set $H(\kappa^+)$ of all sets of hereditary cardinality at most κ that is *locally definable* by a Σ_1 -formula with parameter z , i.e. there is such a well-order that is definable over the structure $\langle H(\kappa^+), \in \rangle$ by a Σ_1 -formula ⁽¹⁾ with parameter z . In particular, if the power set of κ is contained in Gödel's constructible universe L , then there is a well-order of $H(\kappa^+)$ that is locally definable by a Σ_1 -formula without parameters. One may view such well-orders as the *simplest possible locally definable well-orders of $H(\kappa^+)$* , because their definition uses no parameters and a short argument shows that no formula that lies lower in the Levy hierarchy defines a well-order over a structure of the form $H(\kappa^+)$ for any infinite cardinal κ . For the sake of completeness, we present

2010 *Mathematics Subject Classification*: 03E35, 03E47, 03E50.

Key words and phrases: definable well-orders, lightface definability, forcing, failures of the GCH, large cardinals.

Received 30 June 2014; revised 22 September 2015 and 30 June 2016.

Published online 2 November 2016.

⁽¹⁾ Note that if \triangleleft is a well-order, then $x \triangleleft y \Leftrightarrow [x \neq y \wedge \neg(y \triangleleft x)]$. Thus any Σ_1 -definable well-order of $H(\kappa^+)$ is in fact Δ_1 -definable. Proposition 1.1 shows that a Σ_1 -formula defining a well-order is not provably equivalent to the Π_1 -formula obtained from the above equivalence.

this argument. Remember that a Σ_1 -formula $\varphi(v_0, \dots, v_{n-1})$ is $\Delta_1^{\text{ZFC}^-}$ if there is a Π_1 -formula $\psi(v_0, \dots, v_{n-1})$ with

$$\text{ZFC}^- \vdash \forall x_0, \dots, x_{n-1} [\varphi(x_0, \dots, x_{n-1}) \leftrightarrow \psi(x_0, \dots, x_{n-1})],$$

where ZFC^- denotes the axioms of ZFC without the Power Set Axiom. Note that every Σ_0 -formula is a $\Delta_1^{\text{ZFC}^-}$ -formula.

PROPOSITION 1.1. *Let κ be an infinite cardinal. Then no well-order of $\mathbb{H}(\kappa^+)$ is definable over $\langle \mathbb{H}(\kappa^+), \in \rangle$ by a $\Delta_1^{\text{ZFC}^-}$ -formula with parameters.*

Proof. Let $\varphi(v_0, v_1, v_2)$ be a $\Delta_1^{\text{ZFC}^-}$ -formula and let $\psi(v_0, v_1, v_2)$ be the corresponding Π_1 -formula. Using ψ , we can construct a Σ_1 -formula $\Phi_0(v)$ such that the axioms of ZFC^- prove that for every set z , the statement $\Phi_0(z)$ is equivalent to the statement that the relation $\{\langle x, y \rangle \mid \varphi(x, y, z)\}$ is not linear. Moreover, we can use φ to construct a Σ_1 -formula $\Phi_1(v)$ such that the axioms of ZFC^- prove that for every set z , the statement $\Phi_1(z)$ is equivalent to the relation $\{\langle x, y \rangle \mid \varphi(x, y, z)\}$ not being well-founded. Set $\Phi(v) \equiv \Phi_0(v) \vee \Phi_1(v)$. By the Σ_1 -Reflection Principle, the axioms of ZFC prove that whenever θ is an infinite cardinal and $z \in \mathbb{H}(\theta^+)$, then $\Phi(z)$ holds in $\langle \mathbb{H}(\theta^+), \in \rangle$ if and only if the set

$$\{\langle x, y \rangle \in \mathbb{H}(\theta^+) \times \mathbb{H}(\theta^+) \mid \langle \mathbb{H}(\theta^+), \in \rangle \models \varphi(x, y, z)\}$$

is not a well-order of $\mathbb{H}(\theta^+)$.

Assume, towards a contradiction, that there is an infinite cardinal κ such that the formula φ and some parameter $z \in \mathbb{H}(\kappa^+)$ define a well-order of $\mathbb{H}(\kappa^+)$ over the structure $\langle \mathbb{H}(\kappa^+), \in \rangle$. By the above remarks, this implies that $\neg\Phi(z)$ holds in $\langle \mathbb{H}(\kappa^+), \in \rangle$. Pick a regular cardinal $\nu > 2^\kappa$ and let G be $\text{Add}(\nu, (\nu^{<\nu})^+)$ -generic over V . Then $\nu = \nu^{<\nu}$ in $V[G]$ and a folklore result (see, for example, [15, Corollary 9.2]) says that no well-order of $\mathbb{H}(\nu^+)^{V[G]}$ is definable over $\langle \mathbb{H}(\nu^+)^{V[G]}, \in \rangle$. In particular, $\Phi(z)$ holds in $\langle \mathbb{H}(\nu^+)^{V[G]}, \in \rangle$, and Σ_1 -reflection implies that it also holds in $\langle \mathbb{H}(\kappa^+)^{V[G]}, \in \rangle$. But $\mathbb{H}(\kappa^+)^{V[G]} = \mathbb{H}(\kappa^+)^V$, a contradiction. ■

A classical theorem of Mansfield shows that the converse of the implication mentioned at the beginning of this section also holds for $\kappa = \omega$, in the sense that the existence of a locally Σ_1 -definable well-order of $\mathbb{H}(\omega_1)$ implies that all subsets of ω are constructible from the parameters of this definition. In particular, the existence of such a well-order implies that CH holds and that there are no measurable cardinals.

THEOREM 1.2 ([17]). *The following statements are equivalent for any $z \subseteq \omega$:*

- (i) *Every subset of ω is an element of $L[z]$.*
- (ii) *There is a well-order of the set $\mathbb{H}(\omega_1)$ that is definable over the structure $\langle \mathbb{H}(\omega_1), \in \rangle$ by a Σ_1 -formula with parameter z .*

- (iii) *There is a well-order of the set ${}^\omega\omega$ of all functions from ω to ω that is definable over $\langle H(\omega_1), \in \rangle$ by a Σ_1 -formula with parameter z .*

In this paper, we are interested in the provable consequences of the existence of locally definable well-orders of $H(\kappa^+)$ of low complexity in the case where κ is an uncountable regular cardinal. In particular, we want to determine the *simplest definition* such that the existence of a well-order definable in this way is consistent together with certain natural set-theoretical assumptions whose negation holds in L . Examples of such assumptions are failures of the GCH at κ or the existence of larger large cardinals above κ . For that purpose, we construct partial orders that force the existence of locally Σ_1 -definable well-orders while preserving many structural features of the ground model.

The starting point of this work is the following result proven in [11]. It can be used to show that many statements are compatible with the existence of locally Σ_1 -definable well-orders if we allow arbitrary parameters in their definitions.

THEOREM 1.3 ([11, Theorem 1.1]). *Let κ be an uncountable cardinal such that $\kappa = \kappa^{<\kappa}$ holds ⁽²⁾ and 2^κ is regular. Then there is a partial order \mathbb{P} with the following properties:*

- (i) \mathbb{P} is $<\kappa$ -closed, and forcing with \mathbb{P} preserves cofinalities less than or equal to 2^κ and the value of 2^κ .
- (ii) *If G is \mathbb{P} -generic over the ground model V , then there is a well-order of $H(\kappa^+)^{V[G]}$ that is definable over the structure $\langle H(\kappa^+)^{V[G]}, \in \rangle$ by a Σ_1 -formula with parameters.*

The parameter used in the definition of this well-order is added by the forcing, and therefore is, in a certain sense, a very complicated object. In this paper, we want to improve the above result by constructing models of set theory possessing locally Σ_1 -definable well-orders that only use *simple* parameters. The first (and in fact main) step towards this goal will be the construction of a locally Σ_1 -definable well-order in a generic extension whose definition only uses parameters that already exist in the ground model. This is achieved (for a somewhat smaller class of cardinals κ) in the next theorem ⁽³⁾. Remember that, given an uncountable cardinal κ , a subset S of κ is *fat stationary* if for every club $C \subseteq \kappa$, the intersection $C \cap S$ contains closed subsets of ordinals of arbitrarily large order-types below κ . Given

⁽²⁾ Note that the assumption $\kappa = \kappa^{<\kappa}$ implies that κ is regular.

⁽³⁾ While the forcing construction in our paper is based on the construction in [11], one could instead base it on the construction that was later provided in [10]. This would eliminate the assumption that 2^κ be regular, and would yield a forcing that preserves all cofinalities (rather than just those less than or equal to 2^κ) in our results below.

regular cardinals $\eta < \kappa$, we let S_η^κ denote the set of all limit ordinals less than κ of cofinality η . The set $S_{<\eta}^\kappa$ is defined analogously.

THEOREM 1.4. *Let κ be an uncountable cardinal such that $\kappa = \kappa^{<\kappa}$, $\eta^{<\eta} < \kappa$ for every $\eta < \kappa$ and 2^κ is regular ⁽⁴⁾. Assume that one of the following statements holds:*

- (a) κ is the successor of a regular cardinal η , and $\langle S_\alpha | \alpha \leq \kappa \rangle$ is a sequence of pairwise disjoint stationary subsets of S_η^κ .
- (b) κ is an inaccessible cardinal, and $\langle S_\alpha | \alpha \leq \kappa \rangle$ is a sequence of pairwise disjoint fat stationary subsets of κ .

Then there is a partial order \mathbb{P} with the following properties:

- (i) \mathbb{P} is $<\kappa$ -distributive, and forcing with \mathbb{P} preserves cofinalities less than or equal to 2^κ and the value of 2^κ .
- (ii) If G is \mathbb{P} -generic over the ground model V , then there is a well-order of $H(\kappa^+)^{V[G]}$ that is definable over $\langle H(\kappa^+)^{V[G]}, \in \rangle$ by a Σ_1 -formula with parameter $\langle S_\alpha | \alpha < \kappa \rangle$.

In the second part of this paper, we use Theorem 1.4 to construct models of set theory containing locally Σ_1 -definable well-orders of some $H(\kappa^+)$ that only use the ordinal κ as a parameter by forcing over certain canonical inner models of set theory. This will allow us to show that the existence of such a well-order is compatible with a failure of the GCH at κ . The following theorem is an example of such a construction.

THEOREM 1.5. *Assume that $V = L$ holds and κ is either the successor of a regular cardinal or an inaccessible cardinal. Let \mathbb{P} be a partial order with the following properties:*

- (a) Forcing with \mathbb{P} preserves cofinalities less than or equal to κ^+ , and preserves fat stationary subsets of κ .
- (b) If G is \mathbb{P} -generic over V , then in $V[G]$, 2^κ is regular, $\kappa = \kappa^{<\kappa}$ and $\eta^{<\eta} < \kappa$ for all $\eta < \kappa$.

*Then there is a \mathbb{P} -name $\dot{\mathbb{Q}}$ for a partial order such that the following statements hold whenever $G * H$ is $(\mathbb{P} * \dot{\mathbb{Q}})$ -generic over V :*

- (i) The partial order $\dot{\mathbb{Q}}^G$ is $<\kappa$ -distributive in $V[G]$.
- (ii) Forcing with $\dot{\mathbb{Q}}^G$ over $V[G]$ preserves all cofinalities less than or equal to $(2^\kappa)^{V[G]}$, and preserves the value of 2^κ .
- (iii) There is a well-ordering of $H(\kappa^+)^{V[G,H]}$ that is definable over the structure $\langle H(\kappa^+)^{V[G,H]}, \in \rangle$ by a Σ_1 -formula with parameter κ .

⁽⁴⁾ Note that the second assumption implies that κ is either an inaccessible cardinal or the successor of a regular cardinal.

For $\mathbb{P} = \text{Add}(\kappa, \kappa^{++})$, Theorem 1.5 shows that a failure of the GCH at an uncountable regular cardinal κ is consistent with the existence of a well-order of $H(\kappa^+)$ that is locally definable by a Σ_1 -formula with parameter κ .

By Mansfield's Theorem 1.2 and the Σ_1 -Reflection Principle, the existence of a well-order of $H(\kappa^+)$ that is definable over $\langle H(\kappa^+), \in \rangle$ by a Σ_1 -formula with parameter $z \in H(\omega_1)$ implies that $\mathcal{P}(\omega) \subseteq L[z]$. In particular, the assumption

$$(\star) \quad \forall z \subseteq \omega \ \exists x \subseteq \omega \ x \notin L[z]$$

implies that there is no such well-order. Note that (\star) holds in all $\text{Add}(\omega, \omega_1)$ -generic extensions and that the partial order $\mathbb{P} = \text{Add}(\omega, \omega_1)$ satisfies requirements (a) and (b) of Theorem 1.5 in L . Since the partial order $\dot{\mathbb{Q}}^G$ from Theorem 1.5 adds no new reals whenever G is \mathbb{P} -generic over L , it preserves (\star) . Thus Theorem 1.5 shows that in this setting, the above forcing construction for $\kappa = \omega_1$ adds a locally definable well-order of $H(\omega_2)$ of the optimal complexity compatible with (\star) , in the sense of providing a Σ_1 -definition with smallest possible parameter.

It is natural to ask whether it is possible to strengthen the above result and force the existence of well-orders of $H(\omega_2)$ that are definable using smaller parameters. By the above remarks, the existence of such a well-order would imply the negation of (\star) . It is not known whether such a well-order can exist outside of models of the form $L[z]$ with $z \subseteq \omega$.

QUESTION 1.6. *Does the existence of a well-order of $H(\omega_2)$ that is definable over $\langle H(\omega_2), \in \rangle$ by a Σ_1 -formula with parameters in $H(\omega_1)$ imply that $\mathcal{P}(\omega_1)$ is constructible from some subset z of ω ?*

In another direction, one can ask whether assumptions like the ones listed in Theorem 1.5 are actually necessary for such constructions. In particular, it is interesting to ask whether the existence of such well-orders is compatible with the existence of larger large cardinals. A modification of the above construction yields the following result that shows that the existence of a well-order of $H(\kappa^+)$ that is definable over $\langle H(\kappa^+), \in \rangle$ by a Σ_1 -formula with parameter κ is consistent with the existence of a measurable cardinal above κ and a failure of the GCH at κ .

THEOREM 1.7. *Assume that U is a normal measure on a cardinal δ , $V = L[U]$ holds and $\kappa < \delta$ is either the successor of a regular cardinal or an inaccessible cardinal. Let $\mathbb{P} \in V_\delta$ be a partial order with properties (a) and (b) listed in Theorem 1.5. Then there is a \mathbb{P} -name $\dot{\mathbb{Q}} \in V_\delta$ for a partial order such that statements (i)–(iii) in Theorem 1.5 hold whenever $G * H$ is $(\mathbb{P} * \dot{\mathbb{Q}})$ -generic over V .*

Note that if $\kappa = \omega_1$ and U witnesses that δ is a measurable cardinal in V , then (\star) holds and the above remarks imply that the well-ordering of $H(\omega_2)$

produced by the above forcing has the optimal complexity that is compatible with the existence of a measurable cardinal. Since the above construction still relies on the assumption that V lies close to some well-behaved inner model, we may ask if it is possible to have such well-orders in the presence of larger large cardinals. It was pointed out to the authors by Daisuke Ikegami that the results of Woodin on the Π_2 -maximality of the \mathbb{P}_{\max} -extension of $L(\mathbb{R})$ (see [14] and [20]) directly imply that the existence of certain large cardinals implies that no well-order of $H(\omega_2)$ is definable over $\langle H(\omega_2), \in \rangle$ by a Σ_1 -formula with parameter ω_1 .

PROPOSITION 1.8. *Assume that there are infinitely many Woodin cardinals with a measurable cardinal above them. If there is a well-order of $H(\omega_2)$ that is definable over $\langle H(\omega_2), \in \rangle$ by a Σ_1 -formula with parameter $z \subseteq \omega_1$, then $z \notin L(\mathbb{R})$.*

Proof. Given $\alpha < \omega_1$, let WO_α denote the set of all $x \in \mathbb{R}$ that code (in some fixed canonical way) a well-ordering of ω of order-type α . Note that the set $\{\langle x, \alpha \rangle \in \mathbb{R} \times \omega_1 \mid x \in WO_\alpha\}$ is definable over $\langle H(\omega_2), \in \rangle$ by a Σ_1 -formula without parameters. Let $\mathcal{L}_{\dot{A}}$ denote the language of set theory extended by a unary predicate symbol A . Then there is a Σ_1 -formula $\varphi_0(v)$ in $\mathcal{L}_{\dot{A}}$ such that the axioms of ZFC prove that

$$\langle H(\omega_2), \in, A \rangle \models \varphi_0(z) \Leftrightarrow z \subseteq \{\alpha < \omega_1 \mid \exists x \in A \ x \in WO_\alpha\}$$

for all $A \subseteq \mathbb{R}$ and $z \in H(\omega_2)$. Moreover, there is a Π_1 -formula $\varphi_1(v)$ in $\mathcal{L}_{\dot{A}}$ such that the axioms of ZFC prove that

$$\langle H(\omega_2), \in, A \rangle \models \varphi_1(z) \Leftrightarrow \{\alpha < \omega_1 \mid \exists x \in A \ x \in WO_\alpha\} \subseteq z$$

for all $A \subseteq \mathbb{R}$ and $z \in H(\omega_2)$.

Fix a Σ_1 -formula $\psi(v_0, v_1, v_2)$ in the language of set theory. Using the formulas constructed above and the arguments used in the proof of Proposition 1.1, we find a Π_2 -sentence Ψ in $\mathcal{L}_{\dot{A}}$ such that the axioms of ZFC prove that whenever $A \subseteq \mathbb{R}$ and $z_A = \{\alpha < \omega_1 \mid \exists x \in A \ x \in WO_\alpha\}$, then the set

$$\{\langle x, y \rangle \in \mathbb{R} \times \mathbb{R} \mid \langle H(\omega_2), \in \rangle \models \psi(x, y, z_A)\}$$

is a well-ordering of \mathbb{R} if and only if $\langle H(\omega_2), \in, A \rangle \models \Psi$.

Assume, towards a contradiction, that there are infinitely many Woodin cardinals with a measurable cardinal above them and there is $z \in \mathcal{P}(\omega_1)^{L(\mathbb{R})}$ such that

$$\{\langle x, y \rangle \in H(\omega_2) \times H(\omega_2) \mid \langle H(\omega_2), \in \rangle \models \psi(x, y, z)\}$$

is a well-ordering of $H(\omega_2)$. Set $A_z = \bigcup_{\alpha \in z} WO_\alpha \in L(\mathbb{R})$. Then $z = z_{A_z}$ and $\langle H(\omega_2), \in, A_z \rangle \models \Psi$. Let G be \mathbb{P}_{\max} -generic over $L(\mathbb{R})$. By [14, Theorem 7.3], we have $\langle H(\omega_2)^{L(\mathbb{R})[G]}, \in, A_z \rangle \models \Psi$, and hence there is a well-order of \mathbb{R} that is definable over $\langle H(\omega_2)^{L(\mathbb{R})[G]}, \in \rangle$ by the formula ψ and the parameter

$z \in L(\mathbb{R})$. By our assumptions and [14, Lemma 2.10], the partial order \mathbb{P}_{\max} is homogeneous ⁽⁵⁾ in $L(\mathbb{R})$. This shows that the set

$$\{\langle x, y \rangle \in \mathbb{R} \times \mathbb{R} \mid \mathbb{1} \Vdash_{\mathbb{P}_{\max}}^{L(\mathbb{R})} \psi(\check{x}, \check{y}, \check{z})\} \in L(\mathbb{R})$$

is a well-order of \mathbb{R} in $L(\mathbb{R})$. By results of Woodin (see [18, Theorem 8.24]), our assumptions imply that AD holds in $L(\mathbb{R})$ and hence $L(\mathbb{R})$ contains no well-orders of \mathbb{R} , a contradiction. ■

Since the existence of a well-order of $H(\omega_2)$ definable over $\langle H(\omega_2), \in \rangle$ by a Σ_1 -formula with parameter ω_1 is consistent with the existence of a measurable cardinal and inconsistent with the existence of infinitely many Woodin cardinals with a measurable cardinal above them, it is natural to ask the following question.

QUESTION 1.9. *What is the weakest large cardinal whose existence implies that no well-order of $H(\omega_2)$ is locally definable by a Σ_1 -formula with parameter ω_1 ?*

This question will be answered in the forthcoming [16], by showing that the existence of a well-ordering of $H(\omega_2)$ that is definable over $\langle H(\omega_2), \in \rangle$ by a Σ_1 -formula with parameter ω_1 is consistent with the existence of a Woodin cardinal and inconsistent with the existence of a Woodin cardinal with a measurable cardinal above it. Moreover, we will use the techniques developed in Section 7 of this paper to show that the existence of such a well-order is compatible with the existence of a Woodin cardinal and a failure of the GCH at ω_1 .

In another direction, the above arguments do not answer the above question for $\kappa > \omega_1$.

QUESTION 1.10. *Given a formula $\varphi(v_0, v_1, v_2)$, do the axioms of ZFC prove that for every supercompact cardinal δ and every regular cardinal $\omega_1 < \kappa < \delta$, the set*

$$\{\langle x, y \rangle \in H(\kappa^+) \times H(\kappa^+) \mid \langle H(\kappa^+), \in \rangle \models \varphi(x, y, \kappa)\}$$

is not a well-ordering of $H(\kappa^+)$?

Next, we consider locally definable well-orders of ${}^\kappa\kappa$. Note that the Σ_1 -Reflection Principle implies that ${}^\kappa\kappa$ is not definable over $\langle H(\kappa^+), \in \rangle$ by a Σ_1 -formula with parameters in $H(\kappa)$, and hence no well-order of this set is definable in this way. In the case of successor cardinals, the following proposition shows that the forcing construction provided by Theorem 1.5

⁽⁵⁾ In the sense that for all conditions $p_0, p_1 \in \mathbb{P}_{\max}$, there are conditions $q_0, q_1 \in \mathbb{P}_{\max}$ such that $q_i \leq_{\mathbb{P}_{\max}} p_i$ and the restriction of \mathbb{P}_{\max} to all conditions below q_0 is isomorphic to the restriction of \mathbb{P}_{\max} to all conditions below q_1 .

adds a locally Π_1 -definable well-order of ${}^\kappa\kappa$ whose definition uses parameters of small cardinality.

PROPOSITION 1.11. *Assume that η is an infinite cardinal, $\kappa = \eta^+$ and there is a well-order of $\mathbf{H}(\kappa^+)$ that is definable over $\langle \mathbf{H}(\kappa^+), \in \rangle$ by a Σ_1 -formula with parameter κ . Then there is a well-order of ${}^\kappa\kappa$ that is definable over $\langle \mathbf{H}(\kappa^+), \in \rangle$ by a Π_1 -formula with parameter η .*

Proof. Fix a Σ_1 -formula $\Phi(v_0, v_1, v_2)$ such that the formula Φ and the parameter κ define a well-order \triangleleft of $\mathbf{H}(\kappa^+)$ over $\langle \mathbf{H}(\kappa^+), \in \rangle$. Let A be the set of all $x \in \mathbf{H}(\kappa^+)$ with $|x| = \kappa$. Then A consists of all $x \in \mathbf{H}(\kappa^+)$ such that there is no surjection from η onto x in $\mathbf{H}(\kappa^+)$. Therefore A is definable over $\langle \mathbf{H}(\kappa^+), \in \rangle$ by a Π_1 -formula $\psi(v_0, v_1)$ with parameter η . If $x \in A$, then κ is the unique ordinal $\alpha < \kappa^+$ such that there is a bijection $b : \alpha \rightarrow x$ in $\mathbf{H}(\kappa^+)$ and for every $\bar{\alpha} < \alpha$ there is a surjection $s : \eta \rightarrow \bar{\alpha}$ in $\mathbf{H}(\kappa^+)$. This shows that there is a Σ_1 -formula $\varphi(v_0, v_1, v_2)$ such that

$$y = \kappa \Leftrightarrow \langle \mathbf{H}(\kappa^+), \in \rangle \models \varphi(x, y, \eta)$$

for all $x, y \in \mathbf{H}(\kappa^+)$ with $x \in A$. Finally, pick a Σ_0 -formula $\phi(v_0, v_1)$ such that $\phi(x, y)$ holds if and only if $x : y \rightarrow y$ is a function.

Define $\blacktriangleleft = \triangleleft \cap ({}^\kappa\kappa \times {}^\kappa\kappa)$. Then \blacktriangleleft is equal to the set of all pairs $\langle x, y \rangle$ in $\mathbf{H}(\kappa^+) \times \mathbf{H}(\kappa^+)$ with

$$\begin{aligned} \langle \mathbf{H}(\kappa^+), \in \rangle \models & x \neq y \wedge \psi(x, \eta) \wedge \psi(y, \eta) \\ & \wedge \forall \alpha [\varphi(x, \alpha, \eta) \rightarrow (\neg \Phi(y, x, \alpha) \wedge \phi(x, \alpha) \wedge \phi(y, \alpha))]. \end{aligned}$$

This shows that the well-ordering \blacktriangleleft of ${}^\kappa\kappa$ is definable over the structure $\langle \mathbf{H}(\kappa^+), \in \rangle$ by a Π_1 -formula with parameter η . ■

Since the set $\{\omega\}$ is definable over $\langle \mathbf{H}(\omega_2), \in \rangle$ by a Σ_0 -formula without parameters, the above proposition shows that the forcing given by Theorem 1.5 produces a locally definable well-ordering of ${}^{\omega_1}\omega_1$ of optimal complexity.

COROLLARY 1.12. *Assume that there is a well-order of $\mathbf{H}(\omega_2)$ that is definable over $\langle \mathbf{H}(\omega_2), \in \rangle$ by a Σ_1 -formula with parameter ω_1 . Then there is a well-order of ${}^{\omega_1}\omega_1$ that is definable over $\langle \mathbf{H}(\omega_2), \in \rangle$ by a Π_1 -formula without parameters. ■*

Finally, motivated by [11, Corollary 1.5], we also consider Bernstein subsets of ${}^\kappa\kappa$. Given an uncountable regular cardinal κ , we equip ${}^\kappa\kappa$ with the topology whose basic open subsets are of the form $N_s = \{x \in {}^\kappa\kappa \mid s \subseteq x\}$ for some function $s : \alpha \rightarrow \kappa$ with $\alpha < \kappa$, and we say that a closed subset of this space is *perfect* if it is homeomorphic to the set ${}^\kappa 2$ equipped with the subspace topology induced by that of ${}^\kappa\kappa$. A *Bernstein subset* of ${}^\kappa\kappa$ is a subset X of ${}^\kappa\kappa$ with the property that X and its complement intersect every perfect subset of ${}^\kappa\kappa$.

By [11, Corollary 1.5], the partial order \mathbb{P} that witnesses Theorem 1.3 introduces a Bernstein subset of ${}^\kappa\kappa$ that is Δ_1 -definable over $\langle \mathbf{H}(\kappa^+), \in \rangle$ with parameters. It is easy to see that the proof of this result, presented in [11, Section 4], also shows that in the forcing extensions produced by the above theorems there are Bernstein subsets of ${}^\kappa\kappa$ with simple definitions.

COROLLARY 1.13.

- (i) *In the situation of Theorem 1.4, forcing with the partial order \mathbb{P} introduces a Bernstein subset of ${}^\kappa\kappa$ that is Δ_1 -definable over $\langle \mathbf{H}(\kappa^+), \in \rangle$ using the parameter $\langle S_\alpha \mid \alpha < \kappa \rangle$.*
- (ii) *In the situation of Theorem 1.5 or Theorem 1.7, forcing with the partial order $\mathbb{P} * \mathbb{Q}$ introduces a Bernstein subset of ${}^\kappa\kappa$ that is Δ_1 -definable over $\langle \mathbf{H}(\kappa^+), \in \rangle$ using the parameter κ . ■*

We now outline the structure of this paper. In Section 2, we start by discussing forcing techniques that allow us to make an arbitrary subset of $\mathbf{H}(\kappa^+)$ definable in a generic extension of the ground model. Section 3 contains the definition of *strongly S -complete forcings* and several observations that will later allow us to show that the forcing constructed in the proof of Theorem 1.4 is $<\kappa$ -distributive and preserves the stationarity of certain subsets of κ . Next, we prove that the generic coding introduced in Section 2 is absolute with respect to strongly S -complete forcings. We continue by constructing the partial order witnessing Theorem 1.4 in Section 5 and by proving the theorem itself in Section 6. Finally, we prove Theorems 1.5 and 1.7 in Section 7 by constructing forcing extensions of canonical inner models that contain simply definable sequences of disjoint fat stationary sets.

2. Almost disjoint coding at uncountable cardinals. In this section, we discuss *almost disjoint coding forcing* (see [3] and [12]) for uncountable cardinals κ that satisfy $\kappa = \kappa^{<\kappa}$. Given such a κ , this forcing technique will allow us to make an arbitrary subset of ${}^\kappa\kappa$ definable by a formula of low complexity in an upwards-absolute way. In order to make this precise, we generalize basic notions of complexity to our uncountable setting.

We equip the set ${}^\kappa\kappa$ with the topology whose basic open sets are of the form $N_s = \{x \in {}^\kappa\kappa \mid s \subseteq x\}$ for some function $s : \alpha \rightarrow \kappa$ with $\alpha < \kappa$. A subset of ${}^\kappa\kappa$ is a Σ_2^0 -subset of ${}^\kappa\kappa$ if it is equal to the union of κ -many closed subsets of ${}^\kappa\kappa$. Note that a subset of ${}^\kappa\kappa$ is closed if and only if it is equal to the set

$$[T] = \{x \in {}^\kappa\kappa \mid \forall \alpha < \kappa \ x \upharpoonright \alpha \in T\}$$

of κ -branches through some subtree $({}^6) T$ of ${}^{<\kappa}\kappa$. In particular, $A \subseteq {}^\kappa\kappa$ is a Σ_2^0 -subset of ${}^\kappa\kappa$ if and only if there is a sequence $\langle T_\alpha \mid \alpha < \kappa \rangle$ of subtrees

(⁶) A nonempty subset $T \subseteq {}^{<\kappa}\kappa$ is a *subtree* of ${}^{<\kappa}\kappa$ if it is closed under initial segments.

of ${}^{<\kappa}\kappa$ with $A = \bigcup_{\alpha < \kappa} [T_\alpha]$. We say that such a sequence of trees *witnesses that A is a Σ_2^0 -subset of ${}^\kappa\kappa$* .

We will now discuss how *almost disjoint coding forcing* at uncountable cardinals κ with $\kappa = \kappa^{<\kappa}$ allows us to make an arbitrary subset of ${}^\kappa\kappa$ Σ_2^0 -definable by a cofinality-preserving forcing.

DEFINITION 2.1. Assume that κ is an uncountable cardinal with $\kappa = \kappa^{<\kappa}$, $A \subseteq {}^\kappa\kappa$ and $\vec{s} = \langle s_\alpha \mid \alpha < \kappa \rangle$ is an enumeration of ${}^{<\kappa}\kappa$ with the property that every element of ${}^{<\kappa}\kappa$ is enumerated κ -many times. We define a partial order $\mathbb{C}_{\vec{s}}(A)$ by the following clauses:

- (i) A condition in $\mathbb{C}_{\vec{s}}(A)$ is a pair $p = \langle t_p, a_p \rangle$ with $t_p : \alpha_p \rightarrow 2$ for some $\alpha_p < \kappa$ and $a_p \in [A]^{<\kappa}$.
- (ii) We have $q \leq_{\mathbb{C}_{\vec{s}}(A)} p$ if and only if $t_p \subseteq t_q$, $a_p \subseteq a_q$ and

$$s_\beta \subseteq x \rightarrow t_q(\beta) = 1$$

for all $x \in a_p$ and $\alpha_p \leq \beta < \alpha_q$.

The following proposition lists the basic properties of this partial order.

PROPOSITION 2.2.

- (i) If $\eta < \kappa$ and $\langle p_\xi \mid \xi < \eta \rangle$ is a descending sequence of conditions in $\mathbb{C}_{\vec{s}}(A)$, then the pair $p = \langle \bigcup_{\xi < \eta} t_{p_\xi}, \bigcup_{\xi < \eta} a_{p_\xi} \rangle$ is a condition in $\mathbb{C}_{\vec{s}}(A)$ with $p \leq_{\mathbb{C}_{\vec{s}}(A)} p_\xi$ for all $\xi < \eta$. In particular, $\mathbb{C}_{\vec{s}}(A)$ is $<\kappa$ -closed with infima.
- (ii) If p and q are conditions in $\mathbb{C}_{\vec{s}}(A)$ with $t_p = t_q$, then p and q are compatible in $\mathbb{C}_{\vec{s}}(A)$. In particular, $\mathbb{C}_{\vec{s}}(A)$ satisfies the κ^+ -chain condition. ■

The coding provided by the above forcing turns out to be much stronger than in the countable setting, because we no longer need to bother about the definability of ${}^\kappa\kappa$ in the ground model (see [9, Section 1]).

There is a sequence $\langle \dot{T}_\alpha \mid \alpha < \kappa \rangle$ of canonical $\mathbb{C}_{\vec{s}}(A)$ -names with the property that whenever G is $\mathbb{C}_{\vec{s}}(A)$ -generic over V and $t_G = \bigcup \{t_p \mid p \in G\} : \kappa \rightarrow 2$, then for every $\alpha < \kappa$,

$$\dot{T}_\alpha^G = \{t \in {}^{<\kappa}\kappa \mid \forall \alpha < \beta < \kappa [t_G(\beta) = 0 \rightarrow s_\beta \not\subseteq t]\}.$$

THEOREM 2.3. *If κ is an uncountable cardinal with $\kappa = \kappa^{<\kappa}$, A is a subset of ${}^\kappa\kappa$ and G is $\mathbb{C}_{\vec{s}}(A)$ -generic over V , then the sequence $\langle \dot{T}_\alpha^G \mid \alpha < \kappa \rangle$ witnesses that A is a Σ_2^0 -subset of ${}^\kappa\kappa$ in $V[G]$.*

Rather than presenting the short proof of this theorem, we will prove an *absoluteness version* of it in Section 4 (see Corollary 4.3) that will imply the above statement. This result will show that the above coding holds true not only in $V[G]$, but is in fact persistent under certain further forcing.

We close this section with a minor observation showing that in a certain sense, the generic coding provided by Theorem 2.3 is optimal.

PROPOSITION 2.4. *Let κ be an uncountable regular cardinal with $\kappa = \kappa^{<\kappa}$. Then there is a subset C of ${}^\kappa\kappa$ with the property that in every generic extension of the ground model by a $<\kappa$ -distributive forcing, the complement of C is not a Σ_2^0 -subset of ${}^\kappa\kappa$.*

Proof. Define \mathcal{C} to be the *club filter* on κ , i.e. \mathcal{C} is the set of all $x \in {}^\kappa\kappa$ such that $\{\alpha < \kappa \mid x(\alpha) = 0\}$ contains a closed unbounded subset of κ . Assume, towards a contradiction, that there is a generic extension $V[G]$ of the ground model V by a $<\kappa$ -distributive forcing \mathbb{P} such the complement of \mathcal{C} is a Σ_2^0 -subset of ${}^\kappa\kappa$ in $V[G]$. Work in $V[G]$. By our assumption, there is a sequence $\langle U_\alpha \mid \alpha < \kappa \rangle$ of open subsets of ${}^\kappa\kappa$ with $\mathcal{C} = \bigcap_{\alpha < \kappa} U_\alpha$. By the $<\kappa$ -distributivity of \mathbb{P} , the set \mathcal{C} is dense in ${}^\kappa\kappa$, and this implies that U_α is open dense for every $\alpha < \kappa$.

Let $h : {}^\kappa\kappa \rightarrow {}^\kappa\kappa$ be the unique function such that $x(\alpha) \geq 2$ implies $h(x)(\alpha) = x(\alpha)$ and $x(\alpha) < 2$ implies $h(x)(\alpha) = 1 - x(\alpha)$ for all $x \in {}^\kappa\kappa$ and $\alpha < \kappa$. Then h is a homeomorphism of ${}^\kappa\kappa$ and $h[\mathcal{C}]$ is also equal to the intersection of κ -many dense open subsets of ${}^\kappa\kappa$. In this situation, a standard argument shows that there are $x \in \mathcal{C} \cap h[\mathcal{C}]$ and closed unbounded subsets C_0 and C_1 of κ in V such that $C_i \subseteq \{\alpha < \kappa \mid x(\alpha) = i\}$ for all $i < 2$. But then $C_0 \cap C_1 = \emptyset$, a contradiction. ■

Note that it is also possible to use [8, Theorem 4.2] to derive a contradiction in the proof of the above proposition.

3. Strongly S -complete forcings. In this section, we define a strengthening of the notion of an S -complete forcing introduced in [19, Chapter V, Definition 1.1]. We will use this property to show that the forcing constructed in the proof of Theorem 1.4 is $<\kappa$ -distributive and preserves the stationarity of certain subsets of κ . This will allow us to generically code information with the help of the sequence $\langle S_\alpha \mid \alpha \leq \kappa \rangle$ of disjoint stationary sets.

DEFINITION 3.1. Assume κ is an uncountable regular cardinal, $S \subseteq \kappa$ and \mathbb{P} is a partial order. We say that \mathbb{P} is *strongly S -complete* if there is a sequence $\vec{D} = \langle D_\alpha \mid \alpha < \kappa \rangle$ of open dense subsets of \mathbb{P} with the property that whenever

- (a) $\theta > \kappa$ is a regular cardinal with $\mathcal{P}(\mathbb{P}) \in H(\theta)$,
- (b) M is an elementary submodel of $H(\theta)$ of cardinality less than κ such that $\vec{D}, \mathbb{P} \in M$ and $\alpha = \kappa \cap M \in S$, and
- (c) $\vec{p} = \langle p_\xi \mid \xi < \eta \rangle$ is a descending sequence of conditions in \mathbb{P} such that $p_\xi \in M$ for every $\xi < \eta$ and $\{\bar{\alpha} < \alpha \mid \exists \xi < \eta \ p_\xi \in D_{\bar{\alpha}}\}$ is unbounded in α ,

then there is a condition p in \mathbb{P} with $p \leq_{\mathbb{P}} p_\xi$ for every $\xi < \eta$.

LEMMA 3.2. *Let κ be an uncountable regular cardinal with $\eta^{<\eta} < \kappa$ for every cardinal $\eta < \kappa$, let S be a fat stationary subset of κ and let \mathbb{P} be a strongly S -complete partial order. Then \mathbb{P} is $<\kappa$ -distributive.*

Proof. We show that \mathbb{P} is $<\eta$ -distributive for every infinite cardinal $\eta \leq \kappa$ by induction on η . Note that limit steps are trivial, for if η is a limit cardinal, then \mathbb{P} is $<\eta$ -distributive iff it is $<\mu$ -distributive for every infinite cardinal $\mu < \eta$. Moreover, any forcing is trivially $<\omega$ -distributive. Similarly, if $\eta < \kappa$ is a singular cardinal such that \mathbb{P} is $<\eta$ -distributive, then \mathbb{P} is in fact $<\eta^+$ -distributive. All of the above hold true for any notion of forcing P . The only nontrivial case in the induction is the following.

Assume $\eta < \kappa$ is a regular cardinal such that \mathbb{P} is $<\eta$ -distributive, let \dot{f} be a \mathbb{P} -name for a function from η to On and let p_0 be a condition in \mathbb{P} . Pick a sequence $\vec{D} = \langle D_\alpha \mid \alpha < \kappa \rangle$ of open dense subsets of \mathbb{P} witnessing that \mathbb{P} is strongly S -complete, and pick a sufficiently large regular cardinal θ .

We inductively construct a continuous \subseteq -increasing chain $\langle M_\beta \mid \beta < \kappa \rangle$ of elementary submodels of $H(\theta)$ of cardinality less than κ and a strictly increasing continuous sequence $\langle \alpha_\beta \mid \beta < \kappa \rangle$ of limit ordinals less than κ such that for every $\beta < \kappa$:

- (i) $\eta, \dot{f}, p_0, \vec{D}, \mathbb{P} \in M_0$.
- (ii) $\alpha_\beta = \kappa \cap M_\beta \in \kappa$.
- (iii) If $\text{cof}(\alpha_\beta) < \eta$, then $\text{cof}(\alpha_\beta) M_\beta \subseteq M_{\beta+1}$.

Then $C = \{\alpha_\beta \mid \beta < \kappa\}$ is a closed unbounded subset of κ , and by the assumption that S is fat stationary, there is a strictly increasing continuous map $b : \eta + 1 \rightarrow \kappa$ such that $\{\alpha_{b(\xi)} \mid \xi \leq \eta\}$ is a closed subset of S of order-type $\eta + 1$. Now we inductively construct a sequence $\langle p_\xi \mid \xi < \eta \rangle$ of conditions in \mathbb{P} and a sequence $\langle t_\xi \mid \xi < \eta \rangle$ such that $p_\xi \in M_{b(\xi+1)}$, $p_{\xi+1} \Vdash " \dot{f} \upharpoonright \check{\xi} = \check{t}_\xi "$ and $p_{\xi+1} \in D_{\alpha_{b(\xi)}}$ for every $\xi < \eta$.

Assume $\langle p_{\bar{\xi}} \mid \bar{\xi} \leq \xi \rangle$ is already constructed. Then p_ξ and $D_{\alpha_{b(\xi)}}$ are both elements of $M_{b(\xi+1)}$, because $\alpha_{b(\xi)} < \alpha_{b(\xi+1)} \subseteq M_{b(\xi+1)}$. Since \mathbb{P} is $<\eta$ -distributive by induction hypothesis, there are $p_{\xi+1} \in D_{\alpha_{b(\xi)}} \cap M_{b(\xi+1)+1} \subseteq M_{b(\xi+2)}$ and t_ξ with $p_{\xi+1} \leq_{\mathbb{P}} p_\xi$ and $p_{\xi+1} \Vdash " \dot{f} \upharpoonright \check{\xi} = \check{t}_\xi "$.

Now, assume $\xi \in \eta \cap \text{Lim}$ and $\langle p_{\bar{\xi}} \mid \bar{\xi} < \xi \rangle$ is already constructed. Then $\{\bar{\alpha} < \alpha_{b(\xi)} \mid \exists \bar{\xi} < \xi \ p_{\bar{\xi}} \in D_{\bar{\alpha}}\}$ is unbounded in $\alpha_{b(\xi)}$. Hence there is a condition q in \mathbb{P} with $q \leq_{\mathbb{P}} p_{\bar{\xi}}$ for every $\bar{\xi} < \xi$. Let $\langle \xi_\mu \mid \mu < \text{cof}(\xi) \rangle$ be a strictly increasing continuous sequence of ordinals that is cofinal in ξ . Since $\text{cof}(\alpha_{b(\xi)}) = \text{cof}(\xi) < \eta$, we can conclude $\text{cof}(\xi) M_{b(\xi)} \subseteq M_{b(\xi)+1} \subseteq M_{b(\xi+1)}$. This shows that the sequence $\langle p_{\xi_\mu} \mid \mu < \text{cof}(\xi) \rangle$ is an element of $M_{b(\xi+1)}$, and by elementarity there is a condition $p_\xi \in M_{b(\xi+1)}$ such that $p_\xi \leq_{\mathbb{P}} p_{\bar{\xi}}$ for every $\bar{\xi} < \xi$.

Since $\{\bar{\alpha} < \alpha_{b(\eta)} \mid \exists \xi < \eta \ p_\xi \in D_{\bar{\alpha}}\}$ is unbounded in $\alpha_{b(\eta)}$, there is a condition $p \in \mathbb{P}$ with $p \leq_{\mathbb{P}} p_\xi$ for every $\xi < \eta$. If we define $f = \bigcup_{\xi < \eta} t_\xi$, then $p \Vdash \text{“}\dot{f} = \check{f}\text{”}$. This shows that \mathbb{P} is η -distributive. ■

We continue with a technical definition that summarizes the important properties that we will deduce from strong S -completeness.

DEFINITION 3.3. Let \mathbb{P} be a partial order, let θ be a regular uncountable cardinal with $\mathcal{P}(\mathbb{P}) \in H(\theta)$ and let M be an elementary submodel of $H(\theta)$ with $\mathbb{P} \in M$. A condition p in \mathbb{P} is *strongly (M, \mathbb{P}) -generic* if whenever D is an open dense subset of \mathbb{P} that is an element of M , then there is a condition $q \in D$ with $q \in M$ and $p \leq_{\mathbb{P}} q$.

The following basic observation will be needed in the proof of the next lemma.

PROPOSITION 3.4. *Let κ be an uncountable regular cardinal and let S be a fat stationary subset of κ . If \mathbb{P} is a $<\kappa$ -closed partial order, then S is a fat stationary subset of κ in every \mathbb{P} -generic extension of V .*

Proof. Let \dot{C} be a name for a club subset of κ , $\gamma < \kappa$ and p_0 be a condition in \mathbb{P} . Then we can inductively construct a descending sequence $\langle p_\alpha \mid \alpha < \kappa \rangle$ of conditions in \mathbb{P} and a sequence $\langle c_\alpha \mid \alpha < \kappa \rangle$ of bounded subsets of κ such that $p_{\alpha+1} \Vdash \text{“}\dot{C} \cap \check{\alpha} = \check{c}_\alpha\text{”}$ for every $\alpha < \kappa$. Then $C = \bigcup_{\alpha < \kappa} c_\alpha$ is a closed unbounded subset of κ and there is an $\alpha_* < \kappa$ such that $C \cap S \cap \alpha_*$ contains a closed subset of order-type γ . Hence $p_{\alpha_*} \leq_{\mathbb{P}} p_0$ forces that the intersection of \dot{C} and S contains a closed subset of order-type γ . ■

We are now ready to show that strongly S -complete forcings contain dense subsets of strongly generic conditions for a great variety of elementary submodels.

LEMMA 3.5. *Assume that*

- (a) κ is an uncountable regular cardinal with $\eta^{<\eta} < \kappa$ for every $\eta < \kappa$,
- (b) S is a fat stationary subset of κ ,
- (c) \mathbb{P} is a partial order that is $<\kappa$ -closed with infima,
- (d) $\dot{\mathbb{Q}}$ is a \mathbb{P} -name for a strongly S -complete partial order,
- (e) $\vec{D} = \langle \dot{D}_\alpha \mid \alpha < \kappa \rangle$ is a sequence of \mathbb{P} -names for open dense subsets of $\dot{\mathbb{Q}}$ such that whenever G is \mathbb{P} -generic over V , then the sequence $\langle \dot{D}_\alpha^G \mid \alpha < \kappa \rangle$ witnesses that $\dot{\mathbb{Q}}^G$ is strongly S -complete in $V[G]$, and
- (f) $\theta > \kappa$ is regular with $\vec{D}, \mathcal{P}(\mathbb{P} * \dot{\mathbb{Q}}) \in H(\theta)$.

Then for every element x of $H(\theta)$ and every regular cardinal $\eta < \kappa$, there is a dense set of conditions $\langle p, \dot{q} \rangle$ in $\mathbb{P} * \dot{\mathbb{Q}}$ such that for some elementary submodel M of $H(\theta)$:

- (i) M has cardinality less than κ , and $x, \mathbb{P}, \dot{\mathbb{Q}} \in M$.

- (ii) $\alpha = \kappa \cap M \in S$ and there is a closed unbounded subset of α of order-type η consisting of elements of S .
- (iii) $\langle p, \dot{q} \rangle$ is strongly $(M, \mathbb{P} * \dot{\mathbb{Q}})$ -generic.
- (iv) The condition p is the infimum of a descending sequence of conditions in $\mathbb{P} \cap M$.

Proof. We start by proving some general facts about the behaviour of elementary submodels of $H(\theta)$ in \mathbb{P} -generic extensions. Let M be an elementary submodel of $H(\theta)$ of size less than κ , with $\vec{D}, \mathbb{P}, \dot{\mathbb{Q}} \in M$ and $\kappa \cap M \in S$, let $\langle \langle p_{\bar{\xi}}, \dot{q}_{\bar{\xi}} \rangle \mid \bar{\xi} < \xi \rangle$ be a descending sequence of conditions in $(\mathbb{P} * \dot{\mathbb{Q}}) \cap M$ of length less than κ , such that every dense subset of $\mathbb{P} * \dot{\mathbb{Q}}$ that is an element of M has some $p_{\bar{\xi}}$ as element, and let p_{ξ} be the infimum of the sequence $\langle p_{\bar{\xi}} \mid \bar{\xi} < \xi \rangle$ in \mathbb{P} . If G is \mathbb{P} -generic over V , then $H(\theta)^{V[G]}$ is a \mathbb{P} -generic extension of $H(\theta)^V$, and we define

$$M[G] = \{ \dot{x}^G \mid \dot{x} \in M \text{ is a } \mathbb{P}\text{-name} \}.$$

The following claims are standard, but we provide their short proofs for the sake of completeness.

CLAIM 1. *If G is \mathbb{P} -generic over V with $p_{\xi} \in G$, then $M[G]$ is an elementary submodel of $H(\theta)^{V[G]}$ with $M \cap \kappa = M[G] \cap \kappa$.*

Proof. Let $\dot{x} \in M$ be a \mathbb{P} -name with $\dot{x}^G \in \kappa$. Then

$$D = \{ \langle p, \dot{q} \rangle \in \mathbb{P} * \dot{\mathbb{Q}} \mid \exists \alpha < \kappa \ p \Vdash \text{“}\dot{x} = \check{\alpha}\text{”} \}$$

is an open dense subset of $\mathbb{P} * \dot{\mathbb{Q}}$ and an element M . Hence $\langle p_{\bar{\xi}}, \dot{q}_{\bar{\xi}} \rangle \in D \cap M$ for some $\bar{\xi} < \xi$, and there is an $\bar{\alpha} \in \kappa \cap M$ with $\dot{x}^G = \bar{\alpha}$.

Assume there are \mathbb{P} -names $\dot{x}_0, \dots, \dot{x}_{n-1} \in M$ such that $\varphi(x, \dot{x}_0^G, \dots, \dot{x}_{n-1}^G)$ holds in $H(\theta)^{V[G]}$ for some formula $\varphi(v_0, \dots, v_n)$ and some $x \in H(\theta)^{V[G]}$. Since $H(\theta)^{V[G]}$ is a \mathbb{P} -generic extension of the model $H(\theta)^V$, the set of all conditions $\langle p, \dot{q} \rangle$ in $\mathbb{P} * \dot{\mathbb{Q}}$ such that either $p \Vdash \text{“}\forall x \neg \varphi(x, \dot{x}_0, \dots, \dot{x}_{n-1})\text{”}$ holds in $H(\theta)^V$ or $p \Vdash \varphi(\dot{x}, \dot{x}_0, \dots, \dot{x}_{n-1})$ holds in $H(\theta)^V$ for some \mathbb{P} -name $\dot{x} \in H(\theta)^V$ is an open dense subset of $\mathbb{P} * \dot{\mathbb{Q}}$, and is an element of M . By our assumptions, this implies that there is a \mathbb{P} -name $\dot{x}_* \in M$ and a $\bar{\xi} < \xi$ such that $p_{\bar{\xi}} \Vdash \varphi(\dot{x}_*, \dot{x}_0, \dots, \dot{x}_{n-1})$ holds in $H(\theta)^V$, and this shows that there is an $x \in M[G]$ such that $\varphi(x, \dot{x}_0^G, \dots, \dot{x}_{n-1}^G)$ holds in $H(\theta)^{V[G]}$. ■

CLAIM 2. *There is a \mathbb{P} -name \dot{q}_{ξ} for a condition in $\dot{\mathbb{Q}}$ such that $\langle p_{\xi}, \dot{q}_{\xi} \rangle$ is a strongly $(M, \mathbb{P} * \dot{\mathbb{Q}})$ -generic condition and $\langle p_{\xi}, \dot{q}_{\xi} \rangle \leq_{\mathbb{P} * \dot{\mathbb{Q}}} \langle p_{\bar{\xi}}, \dot{q}_{\bar{\xi}} \rangle$ for every $\bar{\xi} < \xi$.*

Proof. Let G be \mathbb{P} -generic over V with $p_{\xi} \in G$ and let D be an open dense subset of $\dot{\mathbb{Q}}^G$ that is an element of $M[G]$. Then there is a \mathbb{P} -name

$\dot{D} \in M$ for a dense subset of $\dot{\mathbb{Q}}$ with $D = \dot{D}^G$, and

$$D_* = \{\langle p, \dot{q} \rangle \in \mathbb{P} * \dot{\mathbb{Q}} \mid p \Vdash \text{“}\dot{q} \in \dot{D}\text{”}\}$$

is an open dense subset of $\mathbb{P} * \dot{\mathbb{Q}}$ and an element of M . Hence there is a $\bar{\xi} < \xi$ such that $\langle p_{\bar{\xi}}, \dot{q}_{\bar{\xi}} \rangle$ is an element of $D_* \cap M$, and this yields $\dot{q}_{\bar{\xi}}^G \in D$. We can conclude that for every $\alpha \in \kappa \cap M[G]$ there is $\bar{\xi} < \xi$ with $\dot{q}_{\bar{\xi}}^G$ in \dot{D}_α^G . By our assumptions and the above claim, this shows that there is a condition q_ξ in $\dot{\mathbb{Q}}^G$ such that $q_\xi \leq_{\dot{\mathbb{Q}}^G} \dot{q}_{\bar{\xi}}^G$ holds in $V[G]$ for every $\bar{\xi} < \xi$. These computations yield a \mathbb{P} -name \dot{q}_ξ with the desired properties. ■

Continuing the proof of Lemma 3.5, pick an element x of $H(\theta)$, a condition $\langle p_0, \dot{q}_0 \rangle$ in $\mathbb{P} * \dot{\mathbb{Q}}$ and a regular cardinal $\eta < \kappa$. We inductively construct a continuous \subseteq -increasing chain M_β of elementary submodels of $H(\theta)$ of cardinality less than κ and a strictly increasing continuous sequence $\langle \alpha_\beta \mid \beta < \kappa \rangle$ of limit ordinals less than κ such that, for every $\beta < \kappa$:

- (I) $\eta, p_0, \dot{q}_0, \vec{D}, \mathbb{P}, \dot{\mathbb{Q}} \in M_0$.
- (II) $\alpha_\beta = \kappa \cap M_\beta \in \kappa$.
- (III) $M_\beta \in M_{\beta+1}$.
- (IV) If $\text{cof}(\alpha_\beta) < \eta$, then ${}^{\text{cof}(\alpha_\beta)}M_\beta \subseteq M_{\beta+1}$.

If we define $C = \{\alpha_\beta \mid \beta < \kappa\}$, then there is a continuous strictly increasing function $b : \eta + 1 \rightarrow \kappa$ such that $\{\alpha_{b(\xi)} \mid \xi \leq \eta\}$ is a closed subset of S of order-type $\eta + 1$. We define a decreasing sequence $\langle \langle p_\xi, \dot{q}_\xi \rangle \mid \xi < \eta \rangle$ of conditions in $\mathbb{P} * \dot{\mathbb{Q}}$ such that $\langle p_\xi, \dot{q}_\xi \rangle \in M_{b(\xi+1)}$ and $\langle p_{\xi+1}, \dot{q}_{\xi+1} \rangle$ is an element of every open dense subset of $\mathbb{P} * \dot{\mathbb{Q}}$ that is an element of $M_{b(\xi)}$.

Assume that the sequence $\langle \langle p_{\bar{\xi}}, \dot{q}_{\bar{\xi}} \rangle \mid \bar{\xi} \leq \xi \rangle$ is already constructed. Then Lemma 3.2 and Proposition 3.4 show that the partial order $\mathbb{P} * \dot{\mathbb{Q}}$ is $<\kappa$ -distributive, and this implies that the intersection of all open dense subsets of $\mathbb{P} * \dot{\mathbb{Q}}$ that are elements of $M_{b(\xi)}$ is open dense. Since $p_\xi, \dot{q}_\xi, M_{b(\xi)} \in M_{b(\xi+1)}$ and $M_{b(\xi)}$ has cardinality less than κ in $M_{b(\xi+1)}$, elementarity implies that there is a condition $\langle p_{\xi+1}, \dot{q}_{\xi+1} \rangle$ in $(\mathbb{P} * \dot{\mathbb{Q}}) \cap M_{b(\xi+1)}$ below $\langle p_\xi, \dot{q}_\xi \rangle$ that is an element of every open dense subset of $\mathbb{P} * \dot{\mathbb{Q}}$ that is an element of $M_{b(\xi)}$.

Now assume that $\xi \in \eta \cap \text{Lim}$ and the sequence $\langle \langle p_{\bar{\xi}}, \dot{q}_{\bar{\xi}} \rangle \mid \bar{\xi} < \xi \rangle$ is already constructed. By Claim 2, there is a strongly $(M_{b(\xi)}, \mathbb{P} * \dot{\mathbb{Q}})$ -generic condition r with $r \leq_{\mathbb{P} * \dot{\mathbb{Q}}} \langle p_{\bar{\xi}}, \dot{q}_{\bar{\xi}} \rangle$ for every $\bar{\xi} < \xi$. Pick a sequence $\langle \xi_\mu \mid \mu < \text{cof}(\xi) \rangle$ that is cofinal in ξ . Since $\text{cof}(\alpha_{b(\xi)}) = \text{cof}(\xi) < \eta$, our assumptions imply that the sequence $\langle \langle p_{\xi_\mu}, \dot{q}_{\xi_\mu} \rangle \mid \mu < \text{cof}(\xi) \rangle$ is an element of $M_{b(\xi+1)}$, and since $M_{b(\xi)}$ is an element of $M_{b(\xi+1)}$, elementarity implies that there is a strongly $(M_{b(\xi)}, \mathbb{P} * \dot{\mathbb{Q}})$ -generic condition $\langle p_\xi, \dot{q}_\xi \rangle \in (\mathbb{P} * \dot{\mathbb{Q}}) \cap M_{b(\xi+1)}$ such that $\langle p_\xi, \dot{q}_\xi \rangle \leq_{\mathbb{P} * \dot{\mathbb{Q}}} \langle p_{\bar{\xi}}, \dot{q}_{\bar{\xi}} \rangle$ for every $\bar{\xi} < \xi$.

This completes the construction of $\langle \langle p_\xi, \dot{q}_\xi \mid \xi < \eta \rangle \rangle$. Another application of Claim 2 shows that there is a strongly $(M_{b(\eta)}, \mathbb{P} * \dot{\mathbb{Q}})$ -generic condition $\langle p_\eta, \dot{q}_\eta \rangle$ such that $\langle p_\eta, \dot{q}_\eta \rangle \leq_{\mathbb{P} * \dot{\mathbb{Q}}} \langle p_\xi, \dot{q}_\xi \rangle$ for every $\xi < \eta$ and p_η is the infimum of the sequence $\langle p_\xi \mid \xi < \eta \rangle$ in \mathbb{P} . Moreover, $\{\alpha_{b(\xi)} \mid \xi < \eta\}$ is a club subset of $\alpha_{b(\eta)} \in S$ consisting of elements of S . ■

COROLLARY 3.6. *Assume that $\kappa, S, \mathbb{P}, \dot{\mathbb{Q}}, \vec{D}$ and θ satisfy the properties listed in Lemma 3.5. Then S is a fat stationary subset of κ in every $(\mathbb{P} * \dot{\mathbb{Q}})$ -generic extension of the ground model.*

Proof. Let \dot{C} be a $(\mathbb{P} * \dot{\mathbb{Q}})$ -name for a closed unbounded subset of κ and let $G * H$ be $(\mathbb{P} * \dot{\mathbb{Q}})$ -generic over V . Fix an infinite regular cardinal $\eta < \kappa$ such that either η is uncountable or $\kappa = \omega_1$. By Lemma 3.5, there is an elementary submodel M of $H(\theta)$ containing \dot{C} with $\alpha = \kappa \cap M \in S$, a closed unbounded subset c of α of order-type η contained in S and a strongly $(M, \mathbb{P} * \dot{\mathbb{Q}})$ -generic condition $r \in G$. Then $\alpha \in \text{Lim}(\dot{C}^{G*H}) \cap S$, and if η is uncountable, then $c \cap \dot{C}^{G*H}$ contains a closed subset of order-type η .

If $\kappa > \omega_1$, then this argument shows that $\dot{C}^{G*H} \cap S$ contains a closed subset of order-type $\eta+1$ for every regular cardinal $\eta < \kappa$, and by [1, Lemma 1.2], this implies that S is fat stationary in $V[G, H]$. In the other case, the argument shows that S is a stationary subset of ω_1 in $V[G, H]$ and every such subset is fat by [5]. ■

4. Almost disjoint coding and strongly S-complete forcings.

This section contains the proofs of the absoluteness versions of Theorem 2.3 mentioned in Section 2. Throughout §4, let κ be an uncountable cardinal satisfying $\kappa^{<\kappa} = \kappa$, let $A \subseteq {}^\kappa\kappa$ and let $\langle \dot{T}_\alpha \mid \alpha < \kappa \rangle$ be defined as before the statement of Theorem 2.3.

LEMMA 4.1. *Let $\dot{\mathbb{Q}}$ be a $\mathbb{C}_{\vec{s}}(A)$ -name for a partial order. Assume that for sufficiently large regular cardinals θ and every $x \in H(\theta)$ there is a dense set of conditions $\langle p, \dot{q} \rangle$ in $\mathbb{C}_{\vec{s}}(A) * \dot{\mathbb{Q}}$ that are strongly $(M, \mathbb{C}_{\vec{s}}(A) * \dot{\mathbb{Q}})$ -generic for some elementary submodel M of $H(\theta)$ of cardinality less than κ with $x, \mathbb{C}_{\vec{s}}(A), \dot{\mathbb{Q}} \in M$, $\kappa \cap M \in \kappa$ and $a_p \subseteq M$. Then the sequence $\langle \dot{T}_\alpha^G \mid \alpha < \kappa \rangle$ witnesses that A is a Σ_2^0 -subset of ${}^\kappa\kappa$ in $V[G, H]$ whenever $G * H$ is $(\mathbb{C}_{\vec{s}}(A) * \dot{\mathbb{Q}})$ -generic over V , i.e. $A = \bigcup_{\alpha < \kappa} [\dot{T}_\alpha^G]$ in $V[G, H]$.*

Proof. Fix $x \in A$. Then the set $D = \{p \in \mathbb{C}_{\vec{s}}(A) \mid x \in a_p\}$ is dense, and there is a condition $p \in G$ with $x \in a_p$. Assume, towards a contradiction, that $x \notin [\dot{T}_{\alpha_p}^G]^{V[G]}$. Then there is a $\alpha_p < \beta < \kappa$ with $t_G(\beta) = 0$ and $s_\beta \subseteq x$. In this situation, we can find a $q \in G$ with $q \leq_{\mathbb{C}_{\vec{s}}(A)} p$ and $\beta < \alpha_q$. Then $s_\beta \subseteq x$ implies $t_G(\beta) = t_q(\beta) = 1$, a contradiction. This argument shows that $x \in [\dot{T}_{\alpha_p}^G]^{V[G]} \subseteq [\dot{T}_{\alpha_p}^G]^{V[G, H]}$.

Now, assume that there is an $x \in [\dot{T}_\alpha^G]^{V[G,H]} \setminus A$ for some $\alpha < \kappa$. Pick a $(\mathbb{C}_{\vec{s}}(A) * \dot{\mathbb{Q}})$ -name \dot{x} for an element of ${}^\kappa\kappa$ with $x = \dot{x}^{G*H}$ and a sufficiently large regular cardinal θ . By our assumption, we can find a condition $\langle p_*, \dot{q}_* \rangle$ in $G * H$ such that $\langle p_*, \dot{q}_* \rangle \Vdash "x \in [\dot{T}_\alpha] \setminus \dot{A}"$ and $\langle p_*, \dot{q}_* \rangle$ is strongly $(M, \mathbb{C}_{\vec{s}}(A) * \dot{\mathbb{Q}})$ -generic for some elementary submodel M of $H(\theta)$ of cardinality less than κ with $\alpha, \dot{x}, \mathbb{C}_{\vec{s}}(A), \dot{\mathbb{Q}} \in M$, $\kappa \cap M \in \kappa$ and $a_{p_*} \subseteq M$.

Given $\bar{\alpha} \in \kappa \cap M$, the set

$$D_{\bar{\alpha}} = \{r \in \mathbb{C}_{\vec{s}}(A) * \dot{\mathbb{Q}} \mid r \Vdash "x \notin [\dot{T}_\alpha]" \vee \\ \exists u [\text{lh}(u) = \bar{\alpha} \wedge r \Vdash "\check{u} \subseteq \dot{x} \wedge \check{u} \in \dot{T}_\alpha"]\}$$

is an open dense subset of $\mathbb{C}_{\vec{s}}(A) * \dot{\mathbb{Q}}$ and an element of M . Hence we can find $r_{\bar{\alpha}} \in D_{\bar{\alpha}} \cap M$ with $\langle p_*, \dot{q}_* \rangle \leq_{\mathbb{C}_{\vec{s}}(A)} r_{\bar{\alpha}}$ and $u_{\bar{\alpha}}$ with $\text{lh}(u_{\bar{\alpha}}) = \bar{\alpha}$ and $r_{\bar{\alpha}} \Vdash "\check{u}_{\bar{\alpha}} \subseteq \dot{x} \wedge \check{u}_{\bar{\alpha}} \in \dot{T}_\alpha"$. Define $u = \bigcup \{u_{\bar{\alpha}} \mid \bar{\alpha} < \alpha_{p_\eta}\}$. Then $\langle p_*, \dot{q}_* \rangle \Vdash "\check{u} \subseteq \dot{x}"$ and hence $u \in T_\alpha$.

Next, if $y \in A \cap M$, then the set

$$E_y = \{\langle p, \dot{q} \rangle \in \mathbb{C}_{\vec{s}}(A) * \dot{\mathbb{Q}} \mid y \in a_p \wedge \\ (r \Vdash "x = \check{y}" \vee \exists \delta < \kappa r \Vdash "x \upharpoonright \check{\delta} \neq \check{y} \upharpoonright \check{\delta}")\}$$

is an element of M and an open dense subset of $\mathbb{C}_{\vec{s}}(A) * \dot{\mathbb{Q}}$. This shows that for every $y \in A \cap M$ there is a condition $r_y \in E_y \cap M$ with $\langle p_*, \dot{q}_* \rangle \leq_{\mathbb{C}_{\vec{s}}(A) * \dot{\mathbb{Q}}} r_y$. Having established that $a_{p_*} \subseteq M$, we can conclude that $a_{p_*} = A \cap M$ and $u \not\subseteq y$ for every $y \in A \cap M$.

By our assumptions on \vec{s} , there is an $\alpha_{p_*} \leq \alpha_* < \kappa$ with $u = s_{\alpha_*}$, and we can find a condition \bar{p} in $\mathbb{C}_{\vec{s}}(A)$ with $\bar{p} \leq_{\mathbb{C}_{\vec{s}}(A)} p_*$, $\alpha_* < \alpha_{\bar{p}}$, $a_{\bar{p}} = a_{p_*}$ and $t_{\bar{p}}(\alpha_*) = 0$. Let $\bar{G} * \bar{H}$ be $(\mathbb{C}_{\vec{s}}(A) * \dot{\mathbb{Q}})$ -generic over V with $\langle \bar{p}, \dot{q}_* \rangle \in \bar{G} * \bar{H}$. Then $\dot{x}^{\bar{G}} \in [\dot{T}_\alpha^{\bar{G}}]^{V[\bar{G}, \bar{H}]}$ and hence $u = \dot{x}^{\bar{G}} \upharpoonright \text{lh}(u) \in \dot{T}_\alpha^{\bar{G}}$. But also $\alpha_* \geq \alpha_{p_*} > \alpha$, $s_{\alpha_*} = u$ and $t_{\bar{G}}(\alpha_*) = t_{\bar{p}}(\alpha_*) = 0$, which contradicts the definition of \dot{T}_α . ■

COROLLARY 4.2. *Assume that $\eta^{<\eta} < \kappa$ for every $\eta < \kappa$. If S is a fat stationary subset of κ , $\dot{\mathbb{Q}}$ is a $\mathbb{C}_{\vec{s}}(A)$ -name for a strongly S -complete partial order and $G * H$ is $(\mathbb{C}_{\vec{s}}(A) * \dot{\mathbb{Q}})$ -generic over V , then the sequence $\langle \dot{T}_\alpha^G \mid \alpha < \kappa \rangle$ witnesses that A is a Σ_2^0 -subset of ${}^\kappa\kappa$ in $V[G, H]$.*

Proof. Pick θ satisfying Lemma 3.5(f) and some $x \in H(\theta)$. By Lemma 3.5, there is a dense set of conditions $\langle p, \dot{q} \rangle$ in $\mathbb{C}_{\vec{s}}(A) * \dot{\mathbb{Q}}$ such that statements (i)–(iv) of the lemma hold for some elementary submodel M of $H(\theta)$ of cardinality less than κ with $x \in M$. Since every such condition p is the infimum of a sequence of conditions in $\mathbb{C}_{\vec{s}}(A) \cap M$, Proposition 2.2 shows that $a_p \subseteq M$ for these conditions. We conclude that the assumptions of Lemma 4.1 are satisfied in this case. ■

We close this section with another corollary of Lemma 4.1 that directly implies the statement of Theorem 2.3. Note that, in contrast to the last corollary, the following argument only uses the assumption that κ is an uncountable cardinal with $\kappa = \kappa^{<\kappa}$.

COROLLARY 4.3. *Let $\dot{\mathbb{Q}}$ be a $\mathbb{C}_{\bar{s}}(A)$ -name for a σ -closed, $<\kappa$ -distributive partial order. If $G * H$ is $(\mathbb{C}_{\bar{s}}(A) * \dot{\mathbb{Q}})$ -generic over V , then the sequence $\langle \dot{I}_\alpha^G \mid \alpha < \kappa \rangle$ witnesses that A is a Σ_2^0 -subset of ${}^\kappa\kappa$ in $V[G, H]$.*

Proof. We show that the assumptions of Lemma 4.1 are satisfied. Let θ be a sufficiently large regular cardinal, $x \in H(\theta)$ and r be a condition in $\mathbb{C}_{\bar{s}}(A) * \dot{\mathbb{Q}}$. Since our assumptions imply that the partial order $\mathbb{C}_{\bar{s}}(A) * \dot{\mathbb{Q}}$ is $<\kappa$ -distributive and therefore the intersection of less than κ -many open dense subsets of this partial order is nonempty, we can simultaneously construct a \subseteq -increasing sequence $\langle M_n \mid n < \omega \rangle$ of elementary submodels of $H(\theta)$ of cardinality less than κ and a descending sequence $\langle \langle p_n, \dot{q}_n \rangle \mid n < \omega \rangle$ of conditions in $\mathbb{C}_{\bar{s}}(A) * \dot{\mathbb{Q}}$ below r such that, for every $n < \omega$:

- (i) $r, x, \mathbb{C}_{\bar{s}}(A) * \dot{\mathbb{Q}} \in M_0, \langle p_n, \dot{q}_n \rangle, M_n \in M_{n+1}$ and $\kappa \cap M_n \in \kappa$.
- (ii) $\langle p_n, \dot{q}_n \rangle$ is an element of every open dense subset of $\mathbb{C}_{\bar{s}}(A) * \dot{\mathbb{Q}}$ that is an element of M_n .

Set $M = \bigcup_{n < \omega} M_n$ and let p be the infimum of the sequence $\langle p_n \mid n < \omega \rangle$ in $\mathbb{C}_{\bar{s}}(A)$. By our assumption, there is a $\mathbb{C}_{\bar{s}}(A)$ -name \dot{q} for a condition in $\dot{\mathbb{Q}}$ with $\langle p, \dot{q} \rangle \leq_{\mathbb{C}_{\bar{s}}(A) * \dot{\mathbb{Q}}} \langle p_n, \dot{q}_n \rangle$ for all $n < \omega$. Then $\langle p, \dot{q} \rangle$ is strongly $(M, \mathbb{C}_{\bar{s}}(A) * \dot{\mathbb{Q}})$ -generic with the desired properties. ■

5. The main forcing construction. In this section and the next, we work under the following assumptions:

- (1) κ is an uncountable cardinal with $\kappa = \kappa^{<\kappa}$, and $\eta^{<\eta} < \kappa$ for every $\eta < \kappa$.
- (2) $\lambda = 2^\kappa$ is a regular cardinal.
- (3) $\langle S_\alpha \mid \alpha \leq \kappa \rangle$ is a sequence of disjoint stationary subsets of S_η^κ if κ is the successor of a regular cardinal η , and a sequence of disjoint fat stationary subsets of κ if κ is inaccessible.

We define $S = S_\kappa, \vec{S} = \langle S_\alpha \mid \alpha < \kappa \rangle$ and $\tilde{S}_\alpha = \kappa \setminus S_\alpha$ for every $\alpha < \kappa$.

Note that assumption (1) implies that κ is either an inaccessible cardinal or the successor of a regular cardinal. In the following, we will work with stationary subsets R of κ such that either $\kappa = \eta^+$ and $R \subseteq S_\eta^\kappa$, or κ is inaccessible and R is fat stationary. We write $R^* = R \cup S_{<\eta}^\kappa$ in the first case and $R^* = R$ in the second. By using [1, Lemma 1.2], it is easy to see that R^* is a fat stationary subset of κ in both cases.

Towards the proof of Theorem 1.4, we recursively define a forcing that simultaneously performs the following three tasks ⁽⁷⁾:

- Generically add a sequence $\vec{A} = \langle A_\delta \mid \delta < 2^\kappa \rangle$ of subsets of κ in the generic extension $V[G]$ such that every element of $H(\kappa^+)^{V[G]}$ is coded (in a sense made precise later on) by exactly one A_δ .
- Generically code \vec{A} to ensure that this sequence is definable over $H(\kappa^+)^{V[G]}$ by a Σ_1 -formula using a parameter $y \subseteq \kappa$ that is added by our forcing.
- Ensure that the parameter y is definable in $H(\kappa^+)^{V[G]}$ by a Σ_1 -formula that uses the sequence $\langle S_\alpha \mid \alpha < \kappa \rangle$ as a parameter.

In this situation, we can well-order $H(\kappa^+)^{V[G]}$ in the desired way by identifying each element of $H(\kappa^+)^{V[G]}$ with the unique A_δ coding it. The generic coding used in this construction will be a variation of the *almost disjoint coding forcing* from Section 2 that was introduced in [2, Section 2], and combines the original forcing with iterated club shooting. The additional coding to make the parameter y definable from $\langle S_\alpha \mid \alpha < \kappa \rangle$ will be achieved by further iterated club shooting.

Before we begin the construction of our forcing, we specify a number of notions used in this construction and fix some more assumptions. We start with several notions of coding sets into other sets:

- We let $\prec, \succ : \text{On} \times \text{On} \rightarrow \text{On}$ denote the *Gödel pairing function*.
- We say that $A \subseteq \kappa$ codes an element z of $H(\kappa^+)$ if there is a bijection $b : \kappa \rightarrow \text{tc}(\{z\})$ such that

$$A = \{ \prec 0, \prec \alpha, \beta \succ \mid \alpha, \beta < \kappa, b(\alpha) \in b(\beta) \} \\ \cup \{ \prec 1, \alpha \succ \mid \alpha < \kappa, b(\alpha) \in z \}.$$

Note that z and b are uniquely determined by A .

- Given $x, y \in {}^\kappa \kappa$, we define $x \oplus y \in {}^\kappa \kappa$ by setting, for all $\alpha < \kappa$,

$$(x \oplus y)(\alpha) := \begin{cases} x(\beta) & \text{if } \alpha = \prec 0, \beta \succ, \\ y(\beta) & \text{if } \alpha = \prec 1, \beta \succ, \\ 0 & \text{otherwise.} \end{cases}$$

- Given $\alpha, \beta < \kappa$, we define $c(\alpha, \beta) \in {}^\kappa 2$ by setting, for all $\gamma < \kappa$,

$$c(\alpha, \beta)(\gamma) := \begin{cases} 1 & \text{if } \gamma \in \{ \prec 0, \alpha \succ, \prec 1, \beta \succ \}, \\ 0 & \text{otherwise.} \end{cases}$$

⁽⁷⁾ This extends the construction of the forcing to witness Theorem 1.3, as provided in [11, Section 2], by the additional third task below. Note that this task introduces the additional technical difficulty that the witnessing forcing for Theorem 1.4 cannot be (unlike the witnessing forcing for Theorem 1.3) $< \kappa$ -closed. The final forcing to witness Theorem 1.4 will be a two-step iteration of the forcing described in this section preceded by a preliminary forcing to achieve assumption (5) below. This two-step iteration is described in detail at the end of Section 6.

In addition to our previous assumptions, we also assume that the following objects exist. (Note that we can (and will) achieve (5) by a preparatory almost disjoint coding forcing, using Corollary 4.2.)

- (4) $\vec{w} = \langle w_\gamma \mid \gamma < \lambda \rangle$ is a sequence of pairwise distinct elements of ${}^\kappa 2$.
(5) $\vec{T} = \langle T_\alpha \mid \alpha < \kappa \rangle$ is a sequence of subtrees of ${}^{<\kappa} \kappa$ with the property that

$$(\dagger) \quad \{w_{\bar{\gamma}} \oplus w_\gamma \mid \bar{\gamma} < \gamma < \lambda\} = \bigcup_{\alpha < \kappa} [T_\alpha]^{V[G]}$$

whenever $V[G]$ is a generic extension of the ground model V by a strongly S^* -complete forcing.

Thus, \vec{T} witnesses that $\{w_{\bar{\gamma}} \oplus w_\gamma \mid \bar{\gamma} < \gamma < \lambda\}$ is a Σ_2^0 -subset of ${}^\kappa \kappa$ in every generic extension by a strongly S^* -complete forcing.

In the following, we inductively construct a sequence $\vec{\mathbb{P}}_{\vec{w}} = \langle \mathbb{P}_\gamma \mid \gamma \leq \lambda \rangle$ of partial orders such that \mathbb{P}_δ is a complete subforcing of \mathbb{P}_γ whenever $\delta < \gamma \leq \lambda$. Fix $\gamma \leq \lambda$ and assume that we have constructed \mathbb{P}_δ with that property for every $\delta < \gamma$.

DEFINITION 5.1. We call a tuple

$$p = \langle s_p, t_p, \vec{d}_p, \vec{c}_p, \vec{A}_p \rangle$$

a \mathbb{P}_γ -candidate if, for some ordinals $\beta_p < \kappa$ and $\gamma_p < \min\{\gamma + 1, \lambda\}$:

- (i) $s_p : \beta_p + 1 \rightarrow {}^{<\kappa} 2$.
(ii) $t_p : \beta_p + 1 \rightarrow 2$.
(iii) $\vec{d}_p = \langle d_{p,\alpha} \mid \alpha \leq \beta_p \rangle$ is such that $d_{p,\alpha}$ is a closed subset of $\tilde{S}_\alpha \cap (\beta_p + 1)$ for every $\alpha \leq \beta_p$. We require that $d_{p,\alpha}$ is the empty set if $\alpha = \beta \cdot 7 + i$ with $i < 7$, $\beta = \prec \gamma, \delta \succ$, and one of the following statements holds ⁽⁸⁾:
(a) $i < 2$ and $s_p(\gamma)(\delta) = i$.
(b) $i = 2$ and $\text{lh}(s_p(\gamma)) \leq \delta$.
(c) $2 < i < 5$ and $t_p(\beta) = i - 3$.
(d) $i = 5$ and $s_p(\gamma) \notin T_\delta$.
(e) $i = 6$ and $s_p(\gamma) \in T_\delta$.
(iv) $\vec{c}_p = \langle c_{p,x} \mid x \in a_p \rangle$ is a sequence such that:
(a) a_p is a subset of $\{w_\delta \oplus c(\alpha, i) \mid \delta < \gamma_p, \alpha < \kappa, i < 2\}$ of cardinality less than κ .

⁽⁸⁾ We will shoot clubs through the \tilde{S}_α , but only for specific α , in order to ensure that in the end, certain sets are definable using \vec{S} as a parameter by checking which of the S_α remained stationary. Coding each piece of information both positively and negatively will make sure that the complexity of those definitions is in fact Δ_1 . Note that each \tilde{S}_α is a fat stationary subset of κ .

- (b) If $x \in a_p$, then $c_{p,x}$ is a closed subset of $\beta_p + 1$ and the implication

$$s_p(\alpha) \subseteq x \rightarrow t_p(\alpha) = 1$$

holds for every $\alpha \in c_{p,x}$.

- (v) $\vec{A}_p = \langle \dot{A}_{p,\delta} \mid \delta < \gamma_p \rangle$ is a sequence such that:

- (a) If $\delta < \gamma_p$, then $\dot{A}_{p,\delta}$ is a \mathbb{P}_δ -nice name for a subset of κ (and thus by our assumptions a $\mathbb{P}_{\bar{\delta}}$ -nice name for a subset of κ for every $\delta \leq \bar{\delta} < \gamma$).
- (b) If $\bar{\gamma} < \gamma_p$ and G is $\mathbb{P}_{\bar{\gamma}}$ -generic over the ground model V , then either $|\lambda|^{V[G]} = |\bar{\gamma}|^{V[G]}$ holds ⁽⁹⁾, or in $V[G]$, there is a sequence $\langle y_\delta \mid \delta \leq \bar{\gamma} \rangle$ of pairwise distinct elements of $H(\kappa^+)$ such that $\dot{A}_{p,\delta}^G$ codes y_δ for every $\delta \leq \bar{\gamma}$.

Given a \mathbb{P}_γ -candidate p and $\delta \leq \gamma$, we define $p \upharpoonright \delta$ to be the tuple

$$\langle s_p, t_p, \vec{d}_p, \langle c_{p,x} \mid x \in a_p \upharpoonright \delta \rangle, \vec{A}_p \upharpoonright \min\{\gamma_p, \delta\} \rangle,$$

where $a_p \upharpoonright \delta = a_p \cap \{w_{\bar{\delta}} \oplus c(\alpha, i) \mid \bar{\delta} < \delta, \alpha < \kappa, i < 2\}$. It is straightforward to check that whenever p is a \mathbb{P}_γ -candidate and $\delta \leq \gamma$, then $p \upharpoonright \delta$ is a \mathbb{P}_δ -candidate.

DEFINITION 5.2. A \mathbb{P}_γ -candidate p is a condition in \mathbb{P}_γ if the following statement holds for all $\delta < \gamma_p$, $\alpha < \kappa$ and $i < 2$ with $w_\delta \oplus c(\alpha, i) \in a_p$:

- (vi) If $p \upharpoonright \delta$ is a condition in \mathbb{P}_δ , then

$$p \upharpoonright \delta \Vdash_{\mathbb{P}_\delta} "i = 1 \leftrightarrow \check{\alpha} \in \dot{A}_{p,\delta}" \text{ (10)}.$$

Given conditions p and q in \mathbb{P}_γ , we define $q \leq_{\mathbb{P}_\gamma} p$ to hold if $s_p = s_q \upharpoonright (\beta_p + 1)$, $t_p = t_q \upharpoonright (\beta_p + 1)$, $d_{p,\alpha} = d_{q,\alpha} \upharpoonright (\beta_p + 1)$ for every $\alpha \leq \beta_p$, $a_p \subseteq a_q$, $\vec{A}_p = \vec{A}_q \upharpoonright \gamma_p$ and $c_{p,x} = c_{q,x} \upharpoonright (\beta_q + 1)$ for every $x \in a_p$.

PROPOSITION 5.3. *If p is a condition in \mathbb{P}_γ and $\delta < \gamma$, then $p \upharpoonright \delta$ is a condition in \mathbb{P}_δ . In particular, every condition p in \mathbb{P}_γ is also a condition in \mathbb{P}_{γ_p} .*

Proof. Let $\delta < \gamma$ and assume that $p \upharpoonright \bar{\delta}$ is a condition in $\mathbb{P}_{\bar{\delta}}$ for every $\bar{\delta} < \delta$. Fix $\bar{\delta} < \delta$, $\alpha < \kappa$ and $i < 2$ with $w_{\bar{\delta}} \oplus c(\alpha, i) \in a_{p \upharpoonright \bar{\delta}}$. Then $(p \upharpoonright \delta) \upharpoonright \bar{\delta} = p \upharpoonright \bar{\delta}$ is a condition in $\mathbb{P}_{\bar{\delta}}$ and $a_{p \upharpoonright \delta} = a_p \upharpoonright \delta \subseteq a_p$. Since p is a condition in \mathbb{P}_γ , this

⁽⁹⁾ We will show later that this case never occurs (see Corollary 5.12).

⁽¹⁰⁾ The idea behind this construction is that the set a_p collects information about the interpretations of names in \vec{A}_p that is already decided by the condition p . This will allow us to use the almost disjoint coding part of the forcing (see clause (iv)(b)) to add a subset of κ that in the end codes $\bigcup_{p \in G} a_p$ and thus also $\bigcup_{p \in G} \vec{A}_p$ whenever G is \mathbb{P}_λ -generic. In clause (iii), we simultaneously work on making this subset of κ lightface definable.

implies $\bar{\delta} < \gamma_p$ and

$$(p \upharpoonright \bar{\delta}) \upharpoonright \bar{\delta} \Vdash_{\mathbb{P}_{\bar{\delta}}} "i = 1 \leftrightarrow \check{\alpha} \in \dot{A}_{p, \bar{\delta}}".$$

We can conclude that $p \upharpoonright \bar{\delta}$ is a condition in $\mathbb{P}_{\bar{\delta}}$. ■

The following statement is a direct consequence of the above definitions.

PROPOSITION 5.4. *If p is a condition in \mathbb{P}_{γ} and \vec{A} is a sequence of length smaller than $\min\{\gamma + 1, \lambda\}$ such that $\vec{A}_p \subseteq \vec{A}$ and \vec{A} satisfies the statements in Definition 5.1(v), then $\langle s_p, t_p, \vec{d}_p, \vec{c}_p, \vec{A} \rangle$ is a condition in \mathbb{P}_{γ} that is stronger than p . ■*

PROPOSITION 5.5. *If $\bar{\gamma} < \min\{\gamma + 1, \lambda\}$, then the set of all conditions p in \mathbb{P}_{γ} with $\gamma_p \geq \bar{\gamma}$ is dense in \mathbb{P}_{γ} .*

Proof. Fix a condition p in \mathbb{P}_{γ} with $\gamma_p < \bar{\gamma}$. Since $\bar{\gamma} < \lambda = 2^{\kappa}$, we can recursively construct a sequence \vec{A} of length $\bar{\gamma}$ that satisfies the statements in Definition 5.1(v). By Proposition 5.4, the resulting tuple $\langle s_p, t_p, \vec{d}_p, \vec{c}_p, \vec{A} \rangle$ is a condition in \mathbb{P}_{γ} that is stronger than p . ■

LEMMA 5.6. *If $\delta < \gamma$, then \mathbb{P}_{δ} is a complete subforcing of \mathbb{P}_{γ} .*

Proof. Every condition in \mathbb{P}_{δ} is a condition in \mathbb{P}_{γ} , $\leq_{\mathbb{P}_{\delta}} = \leq_{\mathbb{P}_{\gamma}} \upharpoonright (\mathbb{P}_{\delta} \times \mathbb{P}_{\delta})$, and if q is a condition in \mathbb{P}_{δ} and p is a condition in \mathbb{P}_{γ} with $p \leq_{\mathbb{P}_{\gamma}} q$, then Proposition 5.3 shows that $p \upharpoonright \delta$ is a condition in \mathbb{P}_{δ} and it is easy to check that $p \upharpoonright \delta \leq_{\mathbb{P}_{\delta}} q$. Hence it suffices to show that every maximal antichain in \mathbb{P}_{δ} is maximal in \mathbb{P}_{γ} .

Fix a maximal antichain \mathcal{A} of \mathbb{P}_{δ} and a condition p_0 in \mathbb{P}_{γ} . By Proposition 5.5, there is a condition p with $p \leq_{\mathbb{P}_{\gamma}} p_0$ and $\gamma_p \geq \delta$. Proposition 5.3 implies that $p \upharpoonright \delta$ is a condition in \mathbb{P}_{δ} . Hence we find a condition q in \mathbb{P}_{δ} and $r \in \mathcal{A}$ with $q \leq_{\mathbb{P}_{\delta}} p \upharpoonright \delta, r$. Then $\gamma_q = \delta$. Define p^* to be the tuple

$$\langle s_q, t_q, \vec{d}_q, \langle c_{p,x} \mid x \in a_p \setminus a_q \rangle \cup \langle c_{q,x} \mid x \in a_q \rangle, \vec{A}_p \rangle.$$

Then p^* is a \mathbb{P}_{γ} -candidate with $\gamma_{p^*} = \gamma_p$. Fix $\bar{\delta} < \gamma$, $\alpha < \kappa$ and $i < 2$ such that $p^* \upharpoonright \bar{\delta}$ is a condition in $\mathbb{P}_{\bar{\delta}}$ and $x = w_{\bar{\delta}} \oplus c(\alpha, i) \in a_p \cup a_q$. If $x \in a_q$, then $\bar{\delta} < \delta \leq \gamma_{p^*}$ and $\vec{A}_q = \vec{A}_p \upharpoonright \bar{\delta}$ implies that $p^* \upharpoonright \bar{\delta} \leq_{\mathbb{P}_{\bar{\delta}}} q \upharpoonright \bar{\delta}$. Hence

$$p^* \upharpoonright \bar{\delta} \Vdash_{\mathbb{P}_{\bar{\delta}}} "i = 1 \leftrightarrow \check{\alpha} \in \dot{A}_{p, \bar{\delta}}"$$

in this case. Now assume that $x \in a_p \setminus a_q$. Since $q \leq_{\mathbb{P}_{\delta}} p \upharpoonright \delta$, we have $p^* \upharpoonright \bar{\delta} \leq_{\mathbb{P}_{\bar{\delta}}} p \upharpoonright \bar{\delta}$ and this implies that the above forcing statement also holds in this case. Therefore p^* is a condition in \mathbb{P}_{γ} and our construction ensures that $p^* \leq_{\mathbb{P}_{\gamma}} p, q$. Hence \mathcal{A} is a maximal antichain in \mathbb{P}_{γ} . ■

This completes the construction of the sequence $\vec{\mathbb{P}}_{\vec{w}}$ of partial orders.

PROPOSITION 5.7. *Let $\gamma \leq \lambda$, $\bar{\lambda} < \lambda$ and $\langle p_\alpha \mid \alpha < \bar{\lambda} \rangle$ be a sequence of conditions in \mathbb{P}_γ such that $\vec{A}_{p_\alpha} \subseteq \vec{A}_{p_\beta}$ for all $\alpha < \beta < \bar{\lambda}$. Then $\vec{A} = \bigcup \{ \vec{A}_{p_\alpha} \mid \alpha < \bar{\lambda} \}$ satisfies the statements in Definition 5.1(v). ■*

PROPOSITION 5.8. *If $\bar{\beta} < \kappa$ and $\gamma \leq \lambda$, then the set of all conditions q in \mathbb{P}_γ with $\beta_q \geq \bar{\beta}$ is dense in \mathbb{P}_γ . In particular, if $\gamma \leq \lambda$ and G is \mathbb{P}_γ -generic over V , then $\kappa = \sup \{ \beta_p \mid p \in G \}$.*

Proof. Fix a condition p in \mathbb{P}_γ with $\beta_p < \bar{\beta}$ and define q to be the tuple $\langle s_p \cup \langle (\alpha, \emptyset) \mid \beta_p \leq \alpha \leq \bar{\beta} \rangle, t_p \cup \langle (\alpha, 1) \mid \beta_p \leq \alpha \leq \bar{\beta} \rangle, \langle d_{q,\alpha} \mid \alpha \leq \bar{\beta} \rangle, \vec{c}_p, \vec{A}_p \rangle$ with $d_{q,\alpha} = d_{p,\alpha}$ for $\alpha \leq \beta_p$ and $d_{q,\alpha} = \emptyset$ for all $\beta_p < \alpha \leq \bar{\beta}$. Then it is easy to see that q is a condition in \mathbb{P}_γ with $q \leq_{\mathbb{P}_\gamma} p$ and $\beta_q = \bar{\beta}$. ■

LEMMA 5.9. *If $\gamma \leq \lambda$, then \mathbb{P}_γ is strongly S^* -complete. Moreover, if p is a condition in \mathbb{P}_γ and $\zeta \leq \beta_p$ is such that $d_{p,\zeta}$ is required to be the empty set by one of the statements in Definition 5.1(iii), then the partial order of conditions in \mathbb{P}_γ below p is strongly S^* -complete.*

Proof. Define $\vec{D} = \langle D_\alpha \mid \alpha < \kappa \rangle$ by setting $D_\alpha = \{ q \in \mathbb{P}_\gamma \mid \beta_q \geq \alpha \}$ for every $\alpha < \kappa$. Each D_α is open dense in \mathbb{P}_γ by Proposition 5.8. We show that \vec{D} witnesses that \mathbb{P}_γ is strongly S^* -complete.

Assume $\theta > \kappa$ is a regular cardinal with $\mathcal{P}(\mathbb{P}_\gamma) \in H(\theta)$, M is an elementary substructure of $H(\theta)$ of cardinality less than κ with $\beta = \sup(M \cap \kappa) \in S^*$ and $\vec{D}, \mathbb{P}_\gamma \in M$, and $\vec{p} = \langle p_\xi \mid \xi < \eta \rangle \subseteq M$ is a descending sequence of conditions in \mathbb{P}_γ such that $\{ \alpha < \beta \mid \exists \xi < \eta \ p_\xi \in D_\alpha \}$ is unbounded in β . This implies that $\beta = \sup_{\xi < \eta} \beta_{p_\xi} \in S^*$. We define a tuple

$$p_* = \langle s, t, \langle d_\alpha \mid \alpha \leq \beta \rangle, \langle c_x \mid x \in a \rangle, \vec{A} \rangle$$

by setting

$$\begin{aligned} s &= \{ \langle \beta, \emptyset \rangle \} \cup \bigcup \{ s_{p_\xi} \mid \xi < \eta \}, \\ t &= \{ \langle \beta, 1 \rangle \} \cup \bigcup \{ t_{p_\xi} \mid \xi < \eta \}, \\ \bar{d}_\alpha &= \bigcup \{ d_{p_\xi, \alpha} \mid \xi < \eta, \alpha \leq \beta_{p_\xi} \} \text{ for every } \alpha < \beta, \\ d_\alpha &= \begin{cases} \bar{d}_\alpha \cup \{ \beta \} & \text{if } \alpha < \beta \text{ and } \bar{d}_\alpha \neq \emptyset, \\ \emptyset & \text{if either } \alpha = \beta, \text{ or } \alpha < \beta \text{ and } \bar{d}_\alpha = \emptyset, \end{cases} \\ a &= \bigcup \{ a_{p_\xi} \mid \xi < \eta \}, \\ c_x &= \{ \beta \} \cup \bigcup \{ c_{p_\xi, x} \mid \xi < \eta, x \in a_{p_\xi} \} \text{ for each } x \in a, \\ \vec{A} &= \bigcup \{ \vec{A}_{p_\xi} \mid \xi < \eta \}. \end{aligned}$$

By Proposition 5.7, \vec{A} satisfies the statements in Definition 5.1(v). Since $\beta \in S^* \subseteq \bar{S}_\alpha$ for every $\alpha < \beta$, we conclude that p_* is a \mathbb{P}_γ -candidate. Fix

$\delta < \gamma$, $\nu < \kappa$ and $i < 2$ with $x = w_\delta \oplus c(\nu, i) \in a$. Then there is $\xi < \eta$ with $x \in a_{p_\xi}$ and hence $\delta < \gamma_{p_\xi} \leq \gamma_{p_*}$. If $p_* \upharpoonright \delta$ is a condition in \mathbb{P}_δ , then $p_* \upharpoonright \delta \leq_{\mathbb{P}_\delta} p_\xi \upharpoonright \delta$ and hence this forces that α is an element of $\dot{A}_{p_*, \delta}$ if and only if $i = 1$. Therefore p_* is a condition in \mathbb{P}_γ and our construction ensures that $p_* \leq_{\mathbb{P}_\gamma} p_\xi$ for every $\xi < \eta$.

The proof of the second statement is similar, noting that if p and ζ are as in its hypothesis, then $S_\zeta^* \subseteq \tilde{S}_\alpha$ for all $\zeta \neq \alpha < \kappa$ and $d_{q, \zeta} = \emptyset$ for every condition q in \mathbb{P}_γ below p . Thus, if θ , M , β and \vec{p} satisfy the above statements with S^* replaced by S_ζ^* , then we can use the same construction to obtain a condition p_* witnessing strong S_ζ^* -completeness. ■

If $\kappa = \eta^+$ is a successor cardinal, then the assumption $S_\alpha \subseteq S_\eta^\kappa$ can be used to show that \mathbb{P}_γ is in fact $< \eta$ -closed.

LEMMA 5.10. *If $\gamma < \lambda$ and $p \in \mathbb{P}_\gamma$ is a condition with $\gamma_p = \gamma$, then the forcing \mathbb{P}_γ satisfies the κ^+ -cc below p .*

Proof. This is a standard Δ -system argument, using the assumption that $\kappa = \kappa^{< \kappa}$ and that whenever $q \leq p$ in \mathbb{P}_γ , we have $\vec{A}_q = \vec{A}_p$. ■

LEMMA 5.11. *If q is a condition in \mathbb{P}_λ and \mathcal{D} is a collection of less than λ -many open dense subsets of \mathbb{P}_λ , then there is a condition p in \mathbb{P}_λ such that $p \leq_{\mathbb{P}_\lambda} q$ and $D \cap \mathbb{P}_{\gamma_p}$ is dense below p in \mathbb{P}_{γ_p} for every $D \in \mathcal{D}$.*

Proof. We start by proving the following claim. Iterated application of it will yield the statement of the lemma.

CLAIM. *Let q_0 be a condition in \mathbb{P}_λ and D be an open dense subset of \mathbb{P}_λ . Then there is a condition q_0^* in \mathbb{P}_λ such that $q_0^* = \langle s_{q_0}, t_{q_0}, \vec{d}_{q_0}, \vec{c}_{q_0}, \vec{A}_{q_0^*} \rangle \leq_{\mathbb{P}_\lambda} q_0$ and $D \cap \mathbb{P}_{\gamma_{q_0^*}}$ is dense below q_0^* in $\mathbb{P}_{\gamma_{q_0^*}}$.*

Proof. We inductively construct a sequence $\langle q_\alpha \mid 0 < \alpha < \theta \rangle$ of incompatible conditions below q_0 in \mathbb{P}_λ with $0 < \theta \leq \kappa^+$ and $\vec{A}_{q_{\bar{\alpha}}} \subseteq \vec{A}_{q_\alpha}$ for all $\bar{\alpha} < \alpha < \theta$: Assume that $\langle q_{\bar{\alpha}} \mid 0 < \bar{\alpha} < \alpha \rangle$ is already constructed. If there is $p_\alpha \in D$ such that $p_\alpha \leq_{\mathbb{P}_\lambda} \langle s_{q_0}, t_{q_0}, \vec{d}_{q_0}, \vec{c}_{q_0}, \bigcup_{\bar{\alpha} < \alpha} \vec{A}_{p_{\bar{\alpha}}} \rangle$ and p_α and $q_{\bar{\alpha}}$ are incompatible in \mathbb{P}_λ for all $0 < \bar{\alpha} < \alpha$, then we set $q_\alpha = p_\alpha$ and we continue our construction. Otherwise, we stop and set $\theta = \alpha$.

Define $\vec{A} = \bigcup_{\alpha < \theta} \vec{A}_{q_\alpha}$ and $q_\alpha^* = \langle s_{q_\alpha}, t_{q_\alpha}, \vec{d}_{q_\alpha}, \vec{c}_{q_\alpha}, \vec{A} \rangle$ for all $\alpha < \theta$. Given $\alpha < \theta$, Proposition 5.7 shows that q_α^* is a condition in $\mathbb{P}_{\gamma_{q_0^*}}$ below q_0^* and q_α . In particular, $\mathcal{A} = \{q_\alpha^* \mid 0 < \alpha < \theta\}$ is an antichain in $\mathbb{P}_{\gamma_{q_0^*}}$ below q_0^* . By Lemma 5.10, this means that the above construction has stopped at stage $\theta < \kappa^+$, because no suitable condition p_θ could be found. This implies that \mathcal{A} is a *maximal* antichain in $\mathbb{P}_{\gamma_{q_0^*}}$ below q_0^* .

Pick a condition p in $\mathbb{P}_{\gamma_{q_0^*}}$ below q_0^* . Then there is $0 < \alpha < \theta$ and a condition r in $\mathbb{P}_{\gamma_{q_0^*}}$ with $r \leq_{\mathbb{P}_{\gamma_{q_0^*}}} p, q_\alpha^*$. Since q_α^* is an element of D , we get $r \in D$. This shows that the condition q_0^* has the desired properties. ■

Let $\langle D_\alpha \mid \alpha < \bar{\lambda} \rangle$ be an enumeration of \mathcal{D} such that $\bar{\lambda} < \lambda$ is a limit ordinal. By the above claim and Proposition 5.7, we can construct a decreasing sequence $\langle q_\alpha \mid \alpha \leq \bar{\lambda} \rangle$ of conditions in \mathbb{P}_λ such that $q = q_0$, $q_\alpha = \langle s_q, t_q, \vec{d}_q, \vec{c}_q, \vec{A}_{q_\alpha} \rangle$ for all $\alpha \leq \bar{\lambda}$, and $D_\alpha \cap \mathbb{P}_{\gamma_{q_{\alpha+1}}}$ is dense below $q_{\alpha+1}$ in $\mathbb{P}_{\gamma_{q_{\alpha+1}}}$ for all $\alpha < \bar{\lambda}$.

Pick a condition r in $\mathbb{P}_{\gamma_{q_{\bar{\lambda}}}}$ below $q_{\bar{\lambda}}$ and $\alpha < \bar{\lambda}$. Then our construction ensures $\vec{A}_r = \vec{A}_{q_{\bar{\lambda}}}$ and $r \upharpoonright \gamma_{q_{\alpha+1}} \leq_{\mathbb{P}_{\gamma_{q_{\alpha+1}}}} q_{\bar{\lambda}} \upharpoonright \gamma_{q_{\alpha+1}} = q_{\alpha+1}$. This allows us to find a condition $\bar{r}_\alpha \in D_\alpha$ with $\bar{r}_\alpha \leq_{\mathbb{P}_{\gamma_{q_{\alpha+1}}}} r \upharpoonright \gamma_{q_{\alpha+1}}$. We define $\vec{c} = \langle c_x \mid x \in a_r \cup a_{\bar{r}_\alpha} \rangle$ by letting $c_x = c_{\bar{r}_\alpha, x}$ if $x \in a_{\bar{r}_\alpha}$, and letting $c_x = c_{r, x}$ otherwise. Then $r_\alpha = \langle s_{\bar{r}_\alpha}, t_{\bar{r}_\alpha}, \vec{d}_{\bar{r}_\alpha}, \vec{c}, \vec{A}_r \rangle$ is a $\mathbb{P}_{\gamma_{q_{\bar{\lambda}}}}$ -candidate with $\bar{r}_\alpha = r_\alpha \upharpoonright \gamma_{q_{\alpha+1}}$. Moreover, if $\delta < \gamma_{q_{\bar{\lambda}}}$ and $r_\alpha \upharpoonright \delta$ is a condition in \mathbb{P}_δ , then this condition is stronger than $r \upharpoonright \delta$. We can conclude that r_α is actually a condition in $\mathbb{P}_{\gamma_{q_{\bar{\lambda}}}}$ that is a common extension of r and \bar{r}_α , and an element of D_α . This shows that $p = q_{\bar{\lambda}}$ has the desired properties. ■

COROLLARY 5.12. *Forcing with \mathbb{P}_λ preserves all cofinalities less than or equal to λ .*

Proof. By Lemmas 3.2 and 5.9, forcing with \mathbb{P}_λ preserves cofinalities less than or equal to κ . Let $\gamma \leq \lambda$ be a limit ordinal with $\text{cof}(\gamma) > \kappa$ and let ν be a regular cardinal with $\kappa \leq \nu < \text{cof}(\gamma)$. Assume, towards a contradiction, that there is $q \in \mathbb{P}_\lambda$ and a \mathbb{P}_λ -name \dot{c} with $q \Vdash_{\mathbb{P}_\lambda} \text{“}\dot{c} : \dot{\nu} \rightarrow \dot{\gamma} \text{ is cofinal”}$. Given $\alpha < \nu$, define

$$D_\alpha = \{p \in \mathbb{P}_\lambda \mid \exists \beta < \gamma \ p \Vdash_{\mathbb{P}_\lambda} \text{“}\dot{c}(\dot{\alpha}) = \dot{\beta}\text{”}\}.$$

Let G be \mathbb{P}_λ -generic over V . By Lemma 5.11, there is $p \in G$ such that $p \leq_{\mathbb{P}_\lambda} q$ and $D_\alpha \cap \mathbb{P}_{\gamma_p}$ is dense below p in \mathbb{P}_{γ_p} for every $\alpha < \nu$. By Lemma 5.10, \mathbb{P}_{γ_p} satisfies the κ^+ -cc below p . Therefore we can define $c : \nu \rightarrow \gamma$ in V by setting

$$c(\alpha) = \sup\{\beta + 1 \mid \exists r \in \mathbb{P}_{\gamma_p} [r \leq_{\mathbb{P}_{\gamma_p}} p \wedge r \Vdash_{\mathbb{P}_\lambda} \text{“}\dot{c}(\dot{\alpha}) = \dot{\beta}\text{”}]\}$$

for every $\alpha < \nu$. Pick $\alpha < \nu$. By Lemma 5.6, $\vec{G} = G \cap \mathbb{P}_{\gamma_p}$ is \mathbb{P}_{γ_p} -generic over V . Since $p \in \vec{G}$, the above computations show that there is an $r \in D_\alpha \cap \vec{G}$. If $\beta < \gamma$ witnesses that r is an element of D_α , then $\dot{c}^G(\alpha) = \beta < c(\alpha)$. This shows that the range of c is unbounded in γ , a contradiction. ■

COROLLARY 5.13. *Let G be \mathbb{P}_λ -generic over V and A be a subset of κ in $V[G]$. Then there is a $\gamma < \lambda$ such that $A = \dot{A}^{G \cap \mathbb{P}_\gamma}$ for some \mathbb{P}_γ -name \dot{A} for a subset of κ .*

Proof. Let \dot{A}_0 be a \mathbb{P}_λ -name for a subset of κ with $A = \dot{A}_0^G$, and given $\alpha < \kappa$, let D_α be the open dense subset of \mathbb{P}_λ consisting of all conditions in \mathbb{P}_λ that decide the statement “ $\check{\alpha} \in \dot{A}_0$ ”. By Lemma 5.11, there is a $p \in G$ such that $D_\alpha \cap \mathbb{P}_{\gamma_p}$ is dense below p for every $\alpha < \kappa$. Define

$$\dot{A} = \{ \langle \check{\alpha}, r \rangle \mid \alpha < \kappa, r \in D_\alpha \cap \mathbb{P}_{\gamma_p}, r \leq_{\mathbb{P}_\lambda} p, r \Vdash_{\mathbb{P}_\lambda} \text{“}\check{\alpha} \in \dot{A}_0\text{”} \}.$$

Then \dot{A} is a \mathbb{P}_{γ_p} -name for a subset of κ , and we can use Lemma 5.6 to conclude that $A = \dot{A}^G = \dot{A}^{G \cap \mathbb{P}_{\gamma_p}}$. ■

6. The proof of Theorem 1.4. We are now ready to show how the forcing constructed in the last section can be used to produce a locally Σ_1 -definable well-order of $H(\kappa^+)$ using only the sequence \vec{S} as a parameter.

LEMMA 6.1. *If G is \mathbb{P}_λ -generic over V and y is an element of $H(\kappa^+)^{V[G]}$, then there is a unique ordinal $\delta < \lambda$ such that for some $p \in G$ with $\delta < \gamma_p$ the set $\dot{A}_{p,\delta}^G$ codes y .*

Proof. By Corollary 5.13, there is a $\gamma < \lambda$ and a \mathbb{P}_γ -name \dot{y} such that $y = \dot{y}^{G \cap \mathbb{P}_\gamma}$. Fix a condition p in \mathbb{P}_λ with $\gamma_p \geq \gamma$. Let \dot{A} be a \mathbb{P}_{γ_p} -name for a subset of κ such that the following statements hold whenever H is \mathbb{P}_{γ_p} -generic over V with $p \in H$ and $\dot{y}^H \in H(\kappa^+)^{V[G]}$:

- If there is no $\delta < \gamma_p$ such that $\dot{A}_{p,\delta}^H$ codes \dot{y}^H , then \dot{A}^H codes \dot{y}^H .
- Otherwise, \dot{A}^H codes an element of $H(\kappa^+)^V$ that is not coded by some $\dot{A}_{p,\delta}^H$ with $\delta < \gamma_p$ (note that Corollary 5.12 implies that such an element always exists).

Define $\vec{A} = \vec{A}_p \cup \{ \langle \gamma_p, \dot{A} \rangle \}$. Then \vec{A} satisfies the statements in Definition 5.1(v) and $\langle s_p, t_p, \vec{q}_p, \vec{c}_p, \vec{A} \rangle$ is a condition in \mathbb{P}_λ below p . This density argument shows that there are $q \in G$ and $\delta < \gamma_q$ such that $\gamma_q > \gamma$ and $\dot{A}_{q,\delta}^{G \cap \mathbb{P}_{\gamma_q}} = \dot{A}_{q,\delta}^G$ codes $\dot{y}^{G \cap \mathbb{P}_{\gamma_q}} = \dot{y}^G$.

Now assume, towards a contradiction, that there are $\delta_0 < \delta_1 < \lambda$ and $p_0, p_1 \in G$ such that both \dot{A}_{p_0,δ_0}^G and \dot{A}_{p_1,δ_1}^G code y . Pick $p \in G$ with $p \leq_{\mathbb{P}_\lambda} p_0, p_1$. Then $\vec{G} = G \cap \mathbb{P}_{\delta_1}$ is \mathbb{P}_{δ_1} -generic over V , and Corollary 5.12 implies $|\delta_1|^{V[G]} < |\lambda|^{V[G]}$. Thus $\dot{A}_{p,\delta_0}^{\vec{G}} = \dot{A}_{p_0,\delta_0}^G$ and $\dot{A}_{p,\delta_1}^{\vec{G}} = \dot{A}_{p_1,\delta_1}^G$ code the same element of $H(\kappa^+)^{V[\vec{G}]}$, contradicting Definition 5.1(v) for the condition p . ■

COROLLARY 6.2. *Forcing with \mathbb{P}_λ preserves the value of 2^κ .* ■

If G is \mathbb{P}_λ -generic over V , then we define

$$D(G) = \{ w_\delta \oplus c(\alpha, i) \mid i < 2, \exists p \in G [\delta < \gamma_p \wedge (i = 1 \leftrightarrow \alpha \in \dot{A}_{p,\delta}^G)] \}.$$

PROPOSITION 6.3. *If G is \mathbb{P}_λ -generic over V and $x = w_\delta \oplus c(\alpha, i) \in D(G)$, then there is a condition $p \in G$ with $x \in a_p$. In particular, $D(G) = \bigcup_{p \in G} a_p$.*

Proof. Assume $x = w_\delta \oplus c(\alpha, i) \in D(G)$. Then there is a condition $q \in G$ such that $\delta < \gamma_q$ and $q \upharpoonright \delta \Vdash_{\mathbb{P}_\delta} "i = 1 \leftrightarrow \check{\alpha} \in \dot{A}_{q,\delta}"$. We may assume that $x \notin a_q$. Fix $p_0 \in \mathbb{P}_\lambda$ with $p_0 \leq_{\mathbb{P}_\lambda} q$ and $x \notin a_{p_0}$. If we define

$$p = \langle s_{p_0}, t_{p_0}, \vec{d}_{p_0}, \{\langle x, \emptyset \rangle\} \cup \langle c_{p_0, y} \mid y \in a_{p_0} \rangle, \vec{A}_{p_0} \rangle,$$

then the above assumptions imply that p is a condition in \mathbb{P}_λ that is stronger than p_0 . Hence the set of all conditions p in \mathbb{P}_λ with $x \in a_p$ is dense below $q \in G$.

The second statement is immediate from the first and the definition of \mathbb{P}_λ . ■

PROPOSITION 6.4. *If G is \mathbb{P}_λ -generic over V and $x \in D(G)$, then*

$$\kappa = \sup\{\sup(c_{p,x}) \mid p \in G, x \in a_p\}$$

and for every $\alpha < \kappa$, either $\kappa = \sup\{\sup(d_{p,\alpha}) \mid p \in G\}$ or $d_{p,\alpha} = \emptyset$ for every $p \in G$ with $\alpha \leq \beta_p$. Moreover, the latter case occurs if and only if there is a condition $p \in G$ such that $\alpha \leq \beta_p$ and $d_{p,\alpha}$ is required to be the empty set by one of the statements in Definition 5.1(iii).

Proof. Fix a condition q in \mathbb{P}_λ with $x \in a_q$, and $\beta \in S$ with $\beta_q < \beta < \kappa$. Moreover, if there is an $\alpha \leq \beta_q$ such that $d_{q,\alpha}$ is not required to be the empty set by one of the statements in Definition 5.1(iii), then we also fix such an α . We define p to be the tuple

$$\langle s, t, \langle d_\zeta \mid \zeta \leq \beta \rangle, \langle c_x \mid x \in a_q \rangle, \vec{A}_q \rangle$$

with

$$\begin{aligned} s &= s_q \cup \langle \langle \xi, \emptyset \rangle \mid \beta_q < \xi \leq \beta \rangle, \\ t &= t_q \cup \langle \langle \xi, 1 \rangle \mid \beta_q < \xi \leq \beta \rangle, \\ c_x &= c_{q,x} \cup (\beta_q, \beta] \text{ for all } x \in a_q, \\ d_\zeta &= \begin{cases} d_{q,\zeta} & \text{if } \zeta \leq \beta_q \text{ with } \alpha \neq \zeta, \\ d_{q,\alpha} \cup \{\beta\} & \text{if } \alpha = \zeta, \\ \emptyset & \text{if } \beta_q < \zeta \leq \beta. \end{cases} \end{aligned}$$

Then p is a condition in \mathbb{P}_λ with $p \leq_{\mathbb{P}_\lambda} q$, $\beta_p = \beta$, $\sup(c_{p,x}) = \beta$ and $\sup(d_{p,\alpha}) = \beta$. Together with Proposition 6.3, this implies all but the backwards direction of the last statement of the assertion. To see that this direction holds as well, note that if $p \in G$ and $d_{p,\alpha}$ is required to be the empty set by one of the statements in Definition 5.1(iii) and q is a condition in \mathbb{P}_λ below p , then $d_{q,\alpha}$ is required to be empty by that same statement, and hence $d_{q,\alpha} = \emptyset$ for every $q \in G$ with $\alpha \leq \beta_q$. ■

We fix \mathbb{P}_λ -names \dot{s} and \dot{t} in V such that $\dot{s}^H = \bigcup\{s_p \mid p \in H\} : \kappa \rightarrow <^{\kappa} 2$ and $\dot{t}^H = \bigcup\{t_p \mid p \in H\} : \kappa \rightarrow 2$ hold whenever H is \mathbb{P}_λ -generic over V .

LEMMA 6.5. *If G is \mathbb{P}_λ -generic over V , then $D(G)$ is definable over the structure $\langle H(\kappa^+)^{V[G]}, \in \rangle$ by a Σ_1 -formula with parameters \dot{s}^G and \dot{t}^G .*

Proof. Let G be \mathbb{P}_λ -generic over V . The statement of the lemma will be a consequence of the following two claims. The first claim is a direct consequence of the definition of \mathbb{P}_λ and Proposition 6.4.

CLAIM 1. *If $x \in D(G)$, then $C_G^x = \bigcup \{c_{p,x} \mid p \in G, x \in a_p\}$ is a club subset of κ such that the implication*

$$(\ddagger) \quad \dot{s}^G(\alpha) \subseteq x \rightarrow \dot{t}^G(\alpha) = 1$$

holds for all $\alpha \in C_G^x$. ■

CLAIM 2. *Assume that $x \in (\kappa^2)^{V[G]}$ is such that (\ddagger) holds for every element α of some club subset C of κ . Then x is an element of $D(G)$.*

Proof. Note that the following proof will be similar to that of Lemma 4.1. Let \dot{a} be the canonical \mathbb{P}_λ -name such that $\dot{a}^H = \bigcup \{a_p \mid p \in H\}$ whenever H is \mathbb{P}_λ -generic over V . Assume, towards a contradiction, that x is not an element of $\dot{a}^G = D(G)$. Then we can find $q \in G$ and \mathbb{P}_λ -names \dot{C} and \dot{x} such that $x = \dot{x}^G$ and

$$q \Vdash_{\mathbb{P}_\lambda} \text{“} \dot{x} \in \check{\kappa}^2 \setminus \dot{a} \wedge \dot{C} \subseteq \check{\kappa} \text{ club} \wedge \forall \alpha \in \dot{C} [\dot{s}(\alpha) \subseteq \dot{x} \rightarrow \dot{t}(\alpha) = 1]\text{”}.$$

Fix a condition p_0 in \mathbb{P}_λ below q . Let M be a countable elementary substructure of $\langle H(\theta), \in \rangle$ for some large, regular θ with the property that $\beta = \sup(M \cap \kappa) \in S^*$ and $\mathbb{P}_\lambda, p_0, q, \dot{a}, \dot{C}, \dot{x} \in M$. Pick a decreasing sequence of conditions $\langle p_n \mid n < \omega \rangle \subseteq M$ so that $p_n \in D$ for some n whenever $D \in M$ is a dense subset of \mathbb{P}_λ . By the genericity of the p_n and the fact that forcing with \mathbb{P}_λ preserves the regularity of κ , we have $\beta = \sup_{n < \omega} \beta_{p_n}$ and (using Lemmas 3.2 and 5.9) there is $u : \beta \rightarrow 2$ such that for every $n < \omega$ there is an $m \geq n$ with $p_m \Vdash_{\mathbb{P}_\lambda} \text{“} \dot{x} \upharpoonright \check{\beta}_{p_n} = \check{u} \upharpoonright \check{\beta}_{p_n}\text{”}$ and $y \upharpoonright \beta_{p_n} \neq u \upharpoonright \beta_{p_n}$ for all $y \in a_{p_n}$. Define

$$p = \langle s, t, \langle d_\alpha \mid \alpha \leq \beta \rangle, \langle c_y \mid y \in a \rangle, \vec{A} \rangle$$

by setting

$$\begin{aligned} s &= \{ \langle \beta, u \rangle \} \cup \bigcup_{n < \omega} s_{p_n}, \\ t &= \{ \langle \beta, 0 \rangle \} \cup \bigcup_{n < \omega} t_{p_n}, \\ \bar{d}_\alpha &= \bigcup \{ d_{p_n, \alpha} \mid n < \omega, \alpha \leq \beta_{p_n} \} \quad \text{for every } \alpha < \beta, \\ d_\alpha &= \begin{cases} \bar{d}_\alpha \cup \{ \beta \} & \text{if } \alpha < \beta \text{ and } \bar{d}_\alpha \neq \emptyset, \\ \emptyset & \text{if either } \alpha = \beta, \text{ or } \alpha < \beta \text{ and } \bar{d}_\alpha = \emptyset, \end{cases} \end{aligned}$$

$$\begin{aligned}
a &= \bigcup_{n < \omega} a_{p_n}, \\
c_y &= \{\beta\} \cup \bigcup \{c_{p_\alpha, y} \mid \alpha < \eta, y \in a_{p_\alpha}\} \quad \text{for every } y \in a, \\
\vec{A} &= \bigcup_{n < \omega} \vec{A}_{p_n}.
\end{aligned}$$

Since $\beta \in S^*$ and $u \not\subseteq y$ for every $y \in a$, we can deduce that p is a condition in \mathbb{P}_λ that is stronger than p_0 . This construction ensures

$$p \Vdash_{\mathbb{P}_\lambda} \text{“}\check{\beta} \in \dot{C} \wedge \dot{s}(\check{\beta}) = \check{s}(\check{\beta}) \subseteq \dot{x} \wedge \dot{t}(\check{\beta}) = 0\text{”},$$

a contradiction. Hence we conclude that $x \in \dot{a}^G$. ■

By Claims 1–2 we can see that

$$D(G) = \{x \in (\kappa^2)^{V[G]} \mid \exists C \subseteq \kappa \text{ club } \forall \alpha \in C [\dot{s}^G(\alpha) \subseteq x \rightarrow \dot{t}^G(\alpha) = 1]\}.$$

This yields a Σ_1 -definition of $D(G)$ over $\langle \mathbb{H}(\kappa^+)^{V[G]}, \in \rangle$ using the parameters \dot{s}^G and \dot{t}^G . ■

LEMMA 6.6. *Let $\alpha < \kappa$ and G be \mathbb{P}_λ -generic over V . Then S_α is a stationary subset of κ in $V[G]$ if and only if there is a $p \in G$ such that $d_{p, \alpha}$ is required to be the empty set by one of the statements in Definition 5.1(iii).*

Proof. By Lemma 5.9, the partial order of conditions in \mathbb{P}_λ below p is strongly S_α^* -complete whenever $d_{p, \alpha}$ is required to be the empty set by one of the statements in Definition 5.1(iii). Thus, if there is such a condition p in G , then Corollary 3.6 shows that forcing with \mathbb{P}_λ preserves the fat stationarity of S_α^* . In the other case, forcing with \mathbb{P}_λ destroys the stationarity of S_α , because Proposition 6.4 shows that $\bigcup \{d_{p, \alpha} \mid p \in G, \alpha \leq \beta_p\}$ is a closed unbounded subset of \tilde{S}_α in $V[G]$. ■

LEMMA 6.7. *Let G be \mathbb{P}_λ -generic over V . Then the sets \dot{s}^G , \dot{t}^G and \vec{T} are Δ_1 -definable in $\langle \mathbb{H}(\kappa^+)^{V[G]}, \in \rangle$ using the sequence \vec{S} as a parameter.*

Proof. From Definition 5.1(iii) and Lemma 6.6, it follows that the following equivalences hold for all $\beta, \gamma, \delta < \kappa$ with $\beta = \prec \gamma, \delta \succ$:

$$\begin{aligned}
\dot{s}^G(\gamma)(\delta) = 0 &\Leftrightarrow S_{\beta \cdot 7} \text{ is stationary} \\
&\Leftrightarrow S_{\beta \cdot 7+1} \text{ is not stationary} \wedge S_{\beta \cdot 7+2} \text{ is not stationary,} \\
\dot{s}^G(\gamma)(\delta) = 1 &\Leftrightarrow S_{\beta \cdot 7+1} \text{ is stationary} \\
&\Leftrightarrow S_{\beta \cdot 7} \text{ is not stationary} \wedge S_{\beta \cdot 7+2} \text{ is not stationary,} \\
\dot{t}^G(\beta) = 0 &\Leftrightarrow S_{\beta \cdot 7+3} \text{ is stationary} \Leftrightarrow S_{\beta \cdot 7+4} \text{ is not stationary,} \\
\dot{t}^G(\beta) = 1 &\Leftrightarrow S_{\beta \cdot 7+4} \text{ is stationary} \Leftrightarrow S_{\beta \cdot 7+3} \text{ is not stationary,} \\
\dot{s}^G(\gamma) \notin T_\delta &\Leftrightarrow S_{\beta \cdot 7+5} \text{ is stationary} \Leftrightarrow S_{\beta \cdot 7+6} \text{ is not stationary,} \\
\dot{s}^G(\gamma) \in T_\delta &\Leftrightarrow S_{\beta \cdot 7+6} \text{ is stationary} \Leftrightarrow S_{\beta \cdot 7+5} \text{ is not stationary.}
\end{aligned}$$

These equivalences yield Δ_1 -definitions of \dot{s}^G , \dot{t}^G and \vec{T} in $\mathsf{H}(\kappa^+)^{\mathsf{V}[G]}$ that only use the sequence \vec{S} as a parameter. ■

LEMMA 6.8. *Let G be \mathbb{P}_λ -generic over V . Then there is a well-order of $\mathsf{H}(\kappa^+)^{\mathsf{V}[G]}$ that is definable over $\langle \mathsf{H}(\kappa^+)^{\mathsf{V}[G]}, \in \rangle$ by a Σ_1 -formula with parameter \vec{S} .*

Proof. Define $W = \{w_\delta \mid \delta < \lambda\}$. Then (\dagger) implies that W and \vec{w} are both definable over $\langle \mathsf{H}(\kappa^+)^{\mathsf{V}[G]}, \in \rangle$ by Σ_1 -formulas that use the sequence \vec{T} of subtrees of ${}^{<\kappa}\kappa$ as a parameter.

CLAIM 1. *If $p \in G$ and $\delta < \gamma_p$, then*

$$\dot{A}_{p,\delta}^G = \{\alpha < \kappa \mid w_\delta \oplus c(\alpha, 1) \in D(G)\} = \{\alpha < \kappa \mid w_\delta \oplus c(\alpha, 0) \notin D(G)\}.$$

Proof. By the definition of $D(G)$, we have

$$\alpha \in \dot{A}_{p,\delta}^G \Leftrightarrow \exists q \in G [\delta < \gamma_q \wedge \alpha \in \dot{A}_{q,\delta}^G] \Leftrightarrow w_\delta \oplus c(\alpha, 1) \in D(G)$$

and

$$\alpha \notin \dot{A}_{p,\delta}^G \Leftrightarrow \exists q \in G [\delta < \gamma_q \wedge \alpha \notin \dot{A}_{q,\delta}^G] \Leftrightarrow w_\delta \oplus c(\alpha, 0) \in D(G). \blacksquare$$

Working in $\mathsf{V}[G]$, we define P to be the set of all pairs $\langle z, v \rangle$ such that $z \in \mathsf{H}(\kappa^+)^{\mathsf{V}[G]}$, $v \in W$ and there is a subset A of κ coding z that satisfies

$$[\alpha \in A \rightarrow v \oplus c(\alpha, 1) \in D(G)] \wedge [\alpha \notin A \rightarrow v \oplus c(\alpha, 0) \in D(G)].$$

CLAIM 2. *Let $z \in \mathsf{H}(\kappa^+)^{\mathsf{V}[G]}$ and let δ_z be the unique ordinal (given by Lemma 6.1) such that $\delta_z < \gamma_p$ and \dot{A}_{p,δ_z}^G codes z for some $p \in G$. Then w_{δ_z} is the unique element of W with $\langle z, w_{\delta_z} \rangle \in P$.*

Proof. By Claim 1, the subset \dot{A}_{p,δ_z}^G of κ witnesses that $\langle z, w_{\delta_z} \rangle$ is an element of P . Now assume, towards a contradiction, that there is $\delta < \lambda$ with $\delta \neq \delta_z$ and $\langle z, w_\delta \rangle \in P$. Let $A \subseteq \kappa$ witness that $\langle z, w_\delta \rangle \in P$. By Claim 1, $A = \dot{A}_{q,\delta}^G$ for some $q \in G$ with $\bar{\gamma} = \max\{\delta, \delta_z\} < \gamma_q$. If we set $\bar{G} = G \cap \mathbb{P}_{\bar{\gamma}}$, then Corollary 5.12 implies $|\bar{\gamma}|^{\mathsf{V}[\bar{G}]} < |\lambda|^{\mathsf{V}[\bar{G}]}$, and the subsets $\dot{A}_{q,\delta}^{\bar{G}} = \dot{A}_{q,\delta}^G$ and $\dot{A}_{q,\delta_z}^{\bar{G}} = \dot{A}_{q,\delta_z}^G$ code the same element of $\mathsf{H}(\kappa^+)^{\mathsf{V}[\bar{G}]}$. This contradicts Definition 5.1(v). ■

Let $\prec_{\vec{w}}$ denote the well-order on W induced by its enumeration \vec{w} . Define \prec_* to be the set of all pairs $\langle z, \bar{z} \rangle$ in $\mathsf{H}(\kappa^+)$ such that

$$\exists v, \bar{v} \in W [\langle z, v \rangle \in P \wedge \langle \bar{z}, \bar{v} \rangle \in P \wedge v \prec_{\vec{w}} \bar{v}].$$

Lemma 6.5 implies that P is Σ_1 -definable over $\mathsf{H}(\kappa^+)^{\mathsf{V}[G]}$ with parameters \vec{T} , \dot{s}^G and \dot{t}^G . Thus the assumptions made at the beginning of Section 5 imply that \prec_* is Σ_1 -definable over $\mathsf{H}(\kappa^+)^{\mathsf{V}[G]}$ with parameters \vec{T} , \dot{s}^G and \dot{t}^G . Lemma 6.7 shows that each of these parameters is itself definable in $\mathsf{H}(\kappa^+)^{\mathsf{V}[G]}$ by a Σ_1 -formula with parameter \vec{S} . In particular, the relation \prec_* is definable over $\mathsf{H}(\kappa^+)^{\mathsf{V}[G]}$ by a Σ_1 -formula with parameter \vec{S} .

Given $z_0, z_1 \in \mathbb{H}(\kappa^+)^{V[G]}$ and $\delta_0, \delta_1 < \lambda$ such that δ_i is the unique ordinal with the property that $\delta_i < \gamma_p$ and \dot{A}_{p, δ_i}^G codes z_i for some $p \in G$, we have $z_0 \prec_* z_1$ if and only if $\delta_0 < \delta_1$. This shows that \prec_* is a well-order of $\mathbb{H}(\kappa^+)$. ■

Proof of Theorem 1.4. Let κ and $\langle S_\alpha \mid \alpha \leq \kappa \rangle$ be as in the statement of the theorem and $\lambda = 2^\kappa$. Fix an injective sequence $\vec{w} = \langle w_\gamma \mid \gamma < \lambda \rangle$ of elements of ${}^\kappa 2$ and define $A = \{w_\delta \oplus w_\gamma \mid \delta < \gamma < \lambda\}$. Fix an enumeration \vec{s} of ${}^{<\kappa} \kappa$ as in Definition 2.1. Let $C_{\vec{s}}(A)$ be the notion of forcing corresponding to A given by Definition 2.1. Since forcing with $C_{\vec{s}}(A)$ preserves our assumptions on κ and Corollary 4.2 shows that all assumptions listed at the beginning of Section 5 hold in all $C_{\vec{s}}(A)$ -generic extensions of the ground model V , there is a canonical $C_{\vec{s}}(A)$ -name \dot{Q} with the property that $\dot{Q}^G = \mathbb{P}_\lambda$ whenever G is $C_{\vec{s}}(A)$ -generic over V and $\mathbb{P}_{\vec{w}}^{V[G]} = \langle \mathbb{P}_\gamma \mid \gamma \leq \lambda \rangle$ is the corresponding sequence of partial orders constructed in $V[G]$ with respect to \vec{w} .

Then the combination of Proposition 2.2, Lemma 5.9 and Corollaries 5.12 and 6.2 implies that $\mathbb{P} = C_{\vec{s}}(A) * \dot{Q}$ is $<\kappa$ -distributive, and that forcing with \mathbb{P} preserves all cofinalities less than or equal to λ and the value of 2^κ . If $G * H$ is $(C_{\vec{s}}(A) * \dot{Q})$ -generic over V , then Lemma 6.8 implies that there is a well-order of $\mathbb{H}(\kappa^+)^{V[G*H]}$ that is definable over $\langle \mathbb{H}(\kappa^+)^{V[G*H]}, \in \rangle$ by a Σ_1 -formula with parameter \vec{S} . ■

7. Σ_1 -definable sequences of disjoint fat stationary sets. This section contains the proofs of Theorems 1.5 and 1.7. We start with a definition that will allow us to prove both theorems using the same techniques.

DEFINITION 7.1. Let κ be an uncountable regular cardinal. We say that a tuple $\langle \delta, \theta, \nu, \triangleleft \rangle$ is *suitable for κ* if $\theta > \kappa$ is a regular cardinal, $\delta \geq \theta$ is a strong limit cardinal, $\nu \leq \kappa$ is an ordinal, \triangleleft is a well-ordering of $\mathbb{H}(\theta)$ of order-type θ , and the following statements hold:

- (i) The set $I(\triangleleft) = \{\{x \mid x \triangleleft y\} \mid y \in \mathbb{H}(\theta)\}$ of all proper initial segments of \triangleleft is definable over $\langle \mathbb{H}(\theta), \in \rangle$ by a Σ_1 -formula with parameter ν .
- (ii) If \mathbb{P} is a partial order of cardinality less than δ with the property that forcing with \mathbb{P} preserves cofinalities less than or equal to κ^+ and G is \mathbb{P} -generic over V , then $\mathbb{H}(\kappa^+)^V$ is definable over $\langle \mathbb{H}(\kappa^+)^{V[G]}, \in \rangle$ by a Σ_1 -formula with parameters κ and ν .
- (iii) There is a closed unbounded subset of $[\mathbb{H}(\theta)]^{<\kappa}$ consisting of elementary submodels M of $\mathbb{H}(\theta)$ with $\pi[I(\triangleleft) \cap M] \subseteq I(\triangleleft)$, where $\pi : M \rightarrow N$ denotes the corresponding transitive collapse.

The following proposition shows that the last statement listed in the above definition follows from the first statement in the case where $\nu < \kappa$.

PROPOSITION 7.2. *Let $\kappa < \theta$ be uncountable regular cardinals and let \triangleleft be a well-ordering of $\mathbf{H}(\theta)$. Assume that there is $\nu < \kappa$ such that the set $I(\triangleleft)$ of all proper initial segments of \triangleleft is definable over $\langle \mathbf{H}(\theta), \in \rangle$ by a Σ_1 -formula with parameter ν . If M is an elementary substructure of $\mathbf{H}(\theta)$ of cardinality less than κ with $\kappa, \nu \in M$, $M \cap \kappa \in \kappa$ and $\pi : M \rightarrow N$ denotes the transitive collapse of M , then $\pi[I(\triangleleft) \cap M] \subseteq I(\triangleleft)$.*

Proof. Fix a Σ_0 -formula $\varphi(v_0, v_1, v_2)$ such that

$$I(\triangleleft) = \{A \in \mathbf{H}(\theta) \mid \exists x \in \mathbf{H}(\theta) \varphi(A, x, \nu)\},$$

and pick $A \in I(\triangleleft) \cap M$. By elementarity, there is $X \in M$ such that $\varphi(A, X, \nu)$ holds. Since $\pi(\nu) = \nu$ and $N \subseteq \mathbf{H}(\theta)$, we can apply Σ_0 -absoluteness to conclude that $\varphi(\pi(A), \pi(X), \nu)$ holds and $\pi(A)$ is an element of $I(\triangleleft)$. ■

In the following, we present two settings in which the above requirements are satisfied. We start by showing that in \mathbf{L} there is a suitable tuple for every uncountable regular cardinal.

LEMMA 7.3. *Assume that $\mathbf{V} = \mathbf{L}$. If $\theta > \kappa$ is a regular cardinal, $\delta \geq \theta$ is a strong limit cardinal and \triangleleft denotes the restriction of the canonical well-ordering of \mathbf{L} to $\mathbf{H}(\theta)$, then the tuple $\langle \delta, \theta, 0, \triangleleft \rangle$ is suitable for κ .*

Proof. The set $I(\triangleleft)$ consists of all $A \in \mathbf{H}(\theta)$ with the property that there is an $\alpha < \theta$ with $A \in \mathbf{L}_\alpha$ and $\langle \mathbf{L}_\alpha, \in \rangle \models$ “ A is an initial segment of $\triangleleft_{\mathbf{L}}$ ”. This shows that $I(\triangleleft)$ is definable over $\langle \mathbf{H}(\theta), \in \rangle$ by a Σ_1 -formula without parameters.

Let \mathbb{P} be a partial order with the property that forcing with \mathbb{P} preserves cofinalities less than or equal to κ^+ , and let G be \mathbb{P} -generic over \mathbf{V} . Then $\mathbf{H}(\kappa^+)^{\mathbf{V}}$ is equal to $\mathbf{L}_{(\kappa^+)^{\mathbf{V}[G]}}$. This shows that the set $\mathbf{H}(\kappa^+)^{\mathbf{V}}$ is definable over $\langle \mathbf{H}(\kappa^+)^{\mathbf{V}[G]}, \in \rangle$ by a Σ_1 -formula without parameters. ■

Next, we present a setting in which suitable tuples exist for all uncountable regular cardinals below a measurable cardinal. The reasonings make use of some basic properties of the *Dodd–Jensen core model* \mathbf{K}^{DJ} (see [4]). The following observation will allow us to show that the third clause of Definition 7.1 is satisfied in this setting.

PROPOSITION 7.4. *Assume that $\mathbf{V} = \mathbf{L}[E]$ is an extender model in the sense of [21]. If κ is an uncountable regular cardinal, \triangleleft is the restriction of the canonical well-ordering of $\mathbf{L}[E]$ to $\mathbf{H}(\kappa^+)$ and $I(\triangleleft)$ is the set of all proper initial segments of \triangleleft , then there is a closed unbounded subset of $[\mathbf{H}(\kappa^+)]^{<\kappa}$ consisting of elementary submodels M of $\mathbf{H}(\kappa^+)$ with $\pi[I(\triangleleft) \cap M] \subseteq I(\triangleleft)$, where $\pi : M \rightarrow N$ denotes the corresponding transitive collapse.*

Proof. By [6, Theorem 8], we know that *Local Club Condensation* holds and we can use [7, Theorem 88] to find a closed unbounded subset of $[H(\kappa^+)]^{<\kappa}$ consisting of elementary submodels of $H(\kappa^+)$ with the desired properties. ■

In the proof of the following lemma, we use the presentation of the core model given in [13]. We only consider premice over the empty set. Therefore we omit the index D (as used in [13]) in these arguments.

LEMMA 7.5. *Assume that U is a normal measure on a cardinal δ and that $V = L[U]$. If $\kappa < \delta$ is an uncountable regular cardinal and \triangleleft denotes the restriction of the canonical well-ordering of the Dodd–Jensen core model K^{DJ} to $H(\kappa^+)^{K^{DJ}}$, then the tuple $\langle \delta, \kappa^+, \kappa, \triangleleft \rangle$ is suitable for κ .*

Proof. Let K abbreviate K^{DJ} . Then the results of [4] show that $H(\delta) \subseteq K$ and \triangleleft is a well-ordering of $H(\kappa^+)$ of order-type κ^+ . Let $\mathbb{P} \in H(\delta)$ be a (possibly trivial) partial order with the property that forcing with \mathbb{P} preserves cofinalities less than or equal to κ^+ , and let G be \mathbb{P} -generic over V . Then $H(\kappa^+)^{V[G]} = H(\kappa^+)^{K[G]}$ and the results of [13] show that $K = (K^{DJ})^{K[G]}$. By [13, Theorem 2.7], the set of all *mice* in $H(\kappa^+)^{V[G]}$ is definable over the structure $\langle H(\kappa^+)^{V[G]}, \in \rangle$ by a Σ_1 -formula with parameter κ . Since $V = L[U]$ and $H(\delta)^V \subseteq K$, a standard argument using elementary substructures of $H(\delta^+)^V$ of cardinality κ shows that $H(\kappa^+)^V$ is equal to the union of all *low parts* $\text{lp}(M)$ of mice M (see [13, Section 1]) in $H(\kappa^+)^{V[G]}$, and we can conclude that $H(\kappa^+)^V$ is definable over $\langle H(\kappa^+)^{V[G]}, \in \rangle$ by a Σ_1 -formula with parameter κ . Moreover, given $A \in H(\kappa^+)^{V[G]}$, we can use [13, Theorems 2.10 and 3.4] to see that A is an element of $I(\triangleleft)$ if and only if there is a mouse $M = J_\alpha[F] \in H(\kappa^+)^{V[G]}$ such that $A \in \text{lp}(M)$ and $\langle M, \in \rangle \models$ “ A is a proper initial segment of $\langle \cdot, J[F] \rangle$ ”. This shows that $I(\triangleleft)$ is definable over $\langle H(\kappa^+)^{V[G]}, \in \rangle$ by a Σ_1 -formula with parameter κ . Finally, K is an extender model and therefore Proposition 7.4 implies that the third clause of Definition 7.1 is satisfied in this setting. ■

In the subsequent paper mentioned after Question 1.9, we will show that if M_1 exists, δ is the unique Woodin cardinal in M_1 and \triangleleft is the canonical well-ordering of $H(\omega_2)$ in M_1 , then the tuple $\langle \delta, \omega_2, \omega_1, \triangleleft \rangle$ is suitable for ω_1 in M_1 . With the help of the techniques developed in this section, we will use this result in that paper to show that the existence of a well-ordering of $H(\omega_2)$ that is locally definable by a Σ_1 -formula with parameter ω_1 is consistent with the existence of a Woodin cardinal and a failure of the GCH at ω_1 .

Next, we show how the concept of κ -suitable tuples can be combined with our previous forcing constructions to obtain well-orders of $H(\kappa^+)$ whose Σ_1 -definition only uses the cardinal κ as a parameter. We start by proving some direct consequences of suitability.

LEMMA 7.6. *Let κ be an uncountable regular cardinal and let $\langle \delta, \theta, \nu, \triangleleft \rangle$ be suitable for κ . Define $\triangleleft_\kappa = \triangleleft \upharpoonright (\mathbf{H}(\kappa) \times \mathbf{H}(\kappa))$.*

- (i) *The order \triangleleft is definable over $\langle \mathbf{H}(\theta), \in \rangle$ by a Σ_1 -formula with parameter ν .*
- (ii) *The set $\mathbf{H}(\kappa)$ is an element of $I(\triangleleft)$ and κ is \triangleleft_κ -cofinal in $\mathbf{H}(\kappa)$.*
- (iii) *The well-order $\langle \mathbf{H}(\kappa), \triangleleft_\kappa \rangle$ has order-type κ and $\kappa = \kappa^{<\kappa}$ holds.*
- (iv) *The set $\{\mathbf{H}(\kappa)\}$ is definable over $\langle \mathbf{H}(\theta), \in \rangle$ by a Σ_1 -formula with parameters κ and ν .*
- (v) *The set $\{\triangleleft_\kappa\}$ is definable over $\langle \mathbf{H}(\theta), \in \rangle$ by a Σ_1 -formula with parameters κ and ν .*
- (vi) *If κ is the successor of a regular cardinal η , then the set $\{S_\eta^\kappa\}$ is definable over $\langle \mathbf{H}(\theta), \in \rangle$ by a Σ_1 -formula with parameters κ and ν .*

Proof. (i) Given $x, y \in \mathbf{H}(\theta)$, we have $x \triangleleft y$ if and only if there are $A, B \in I(\triangleleft)$ with $x \in A$ and $y \in B \setminus A$. By our assumptions on $I(\triangleleft)$, this yields the first statement of the lemma.

(ii) Fix $x, y \in \mathbf{H}(\theta)$ with $x \triangleleft y$ and $y \in \mathbf{H}(\kappa)$. Since \triangleleft has order-type θ , there is $\lambda < \theta$ with $y \triangleleft \lambda$. Pick $A, B \in I(\triangleleft)$ with $y \in A$ and $\lambda \in B \setminus A$. Choose an elementary submodel M of $\mathbf{H}(\theta)$ contained in the closed unbounded set described in Definition 7.1(iii) with $\text{tc}(\{y\}) \cup \{A, B, \lambda, \kappa, \nu\} \subseteq M$, and let $\pi : M \rightarrow N$ denote the corresponding transitive collapse. Then $\pi(A), \pi(B) \in I(\triangleleft)$, $y = \pi(y) \in \pi(A) \in I(\triangleleft)$ and hence $x \in \pi(A) \subseteq N \subseteq \mathbf{H}(\kappa)$. Moreover, $y = \pi(y) \in \pi(B) \setminus \pi(A)$, $\pi(\lambda) \in \pi(B)$ and hence $y \triangleleft \pi(\lambda) < \kappa$.

(iii) The second statement of the lemma implies that the well-order $\langle \mathbf{H}(\kappa), \triangleleft_\kappa \rangle$ has order-type at least κ . Assume toward a contradiction that there is $A \in I(\triangleleft)$, a bijection $b : \kappa \rightarrow A$ and $y \in \mathbf{H}(\kappa)$ with $A = \{x \in \mathbf{H}(\kappa) \mid x \triangleleft y\}$. Choose an elementary submodel M of $\mathbf{H}(\theta)$ contained in the closed unbounded set described in Definition 7.1(iii) with $\text{tc}(\{y\}) \cup \{A, b, \kappa, \nu\} \subseteq M$, and let $\pi : M \rightarrow N$ denote the corresponding transitive collapse. Then we have $\pi(A), \pi(A \cup \{y\}) \in I(\triangleleft)$, $\pi(A) \cup \{y\} = \pi(A \cup \{y\})$, which shows that $A = \pi(A)$. But elementarity implies that there is a bijection between $\pi(A)$ and $\pi(\kappa) < \kappa$, a contradiction. In particular, this shows that $\mathbf{H}(\kappa)$ has cardinality κ and hence we also get $\kappa = \kappa^{<\kappa}$.

(iv) By the second part of the lemma, $\mathbf{H}(\kappa)$ is the unique element M of $\mathbf{H}(\theta)$ such that $M \in I(\triangleleft)$, $M \cap \kappa = \kappa$ and κ is \triangleleft -cofinal in M . By Definition 7.1(i) and the first part of the lemma, this yields the assertion.

(v) This follows from our assumptions and the above statements.

(vi) Since the sets S_η^κ and $\{\eta\}$ are definable over $\langle \mathbf{H}(\kappa), \in \rangle$, this follows directly from (iv). ■

The next result shows how we can use suitable tuples to replace sequences of fat stationary subsets of κ by the cardinal κ as the parameter in a Σ_1 -definition of a well-ordering of $\mathbf{H}(\kappa^+)$.

THEOREM 7.7. *Let κ be an uncountable regular cardinal and let $\langle \delta, \theta, \nu, \triangleleft \rangle$ be suitable for κ . Then there is a sequence $\langle S_\alpha \mid \alpha \leq \kappa \rangle$ of disjoint fat stationary subsets of κ with the property that the set $\{ \langle S_\alpha \mid \alpha < \kappa \rangle \}$ is definable over $\langle H(\kappa^+)^{V[G]}, \in \rangle$ by a Σ_1 -formula with parameters κ and ν whenever \mathbb{P} is a partial order of cardinality less than δ such that forcing with \mathbb{P} preserves cofinalities less than or equal to κ^+ and G is \mathbb{P} -generic over V .*

Proof. Define $F : \kappa \cap \text{Lim} \rightarrow \kappa$ to be the unique function such that the following statements hold for all $\alpha \in \kappa \cap \text{Lim}$:

- (i) Assume that there is a triple $\langle \gamma, \lambda, C \rangle$ such that $\gamma, \lambda < \alpha$ and $C \subseteq \alpha \cap \text{Lim}$ is a club in α with the property that, for every closed bounded subset c of C of order-type λ , there is $\bar{\alpha} \in c$ with $F(\bar{\alpha}) \neq \gamma$. Then $F(\alpha) = \gamma_0$, where $\langle \gamma_0, \lambda_0, C_0 \rangle$ denotes the \triangleleft -minimal triple with this property.
- (ii) If there is no such triple, then $F(\alpha) = \alpha$.

By the Recursion Theorem, F is definable over $\langle H(\theta), \in \rangle$ by a formula with parameters κ and ν .

CLAIM 1. *Given $\beta < \kappa$, the set $F^{-1}\{\beta\}$ is a fat stationary subset of κ .*

Proof. Assume, towards a contradiction, that the statement of the claim fails. Then there is a triple $\langle \beta, \lambda, C \rangle$ such that $\beta, \lambda < \kappa$ and $C \subseteq \kappa \cap \text{Lim}$ is a club subset of κ with the property that, for every closed bounded subset c of C of order-type λ , there is an $\alpha \in c$ with $F(\alpha) \neq \beta$. Let $\langle \beta_0, \lambda_0, C_0 \rangle$ denote the \triangleleft -minimal triple with these properties, and set

$$A = \{x \mid x \triangleleft \langle \beta_0, \lambda_0, C_0 \rangle\} \cup \{\langle \beta_0, \lambda_0, C_0 \rangle\} \in I(\triangleleft).$$

Since $\langle \delta, \theta, \nu, \triangleleft \rangle$ is suitable for κ , we can find a monotone enumeration $\langle \alpha_\xi \mid \xi < \kappa \rangle$ of a club subset of κ and a continuous ascending sequence $\langle M_\xi \mid \xi < \kappa \rangle$ of elementary submodels of $H(\theta)$ of size less than κ such that, for all $\xi < \kappa$:

- (1) $\alpha_\xi = \kappa \cap M_\xi$ and $\beta_0, \lambda_0, \kappa, \nu, A, C_0 \in M_\xi$.
- (2) M_ξ is contained in the club described in Definition 7.1(iii), thus if $\pi_\xi : M_\xi \rightarrow N_\xi$ denotes the transitive collapse of M_ξ , then we have $\pi[I(\triangleleft) \cap M] \subseteq I(\triangleleft)$.

Fix $\xi < \kappa$. Then $\beta_0, \lambda_0 < \alpha_\xi = \pi_\xi(\kappa) \in \text{Lim}$, $C_0 \cap \alpha_\xi = \pi_\xi(C_0) \subseteq \alpha_\xi \cap \text{Lim}$ is a club in α_ξ and $\alpha_\xi \in C_0$. Pick a closed bounded subset c of $C_0 \cap \alpha_\xi$ of order-type λ . Then c is a closed bounded subset of C_0 , and there is $\bar{\alpha} \in c$ with $F(\bar{\alpha}) \neq \beta_0$. This shows that the triple $\langle \beta_0, \lambda_0, C_0 \cap \kappa_\alpha \rangle$ satisfies the assumption in (i) with respect to α_ξ , and hence $F(\alpha_\xi) < \alpha_\xi$ by elementarity of M_ξ . Next, pick $\langle \gamma, \rho, D \rangle \triangleleft \langle \beta_0, \lambda_0, C_0 \cap \kappa_\alpha \rangle$, if it exists, such that $\gamma, \rho < \alpha_\xi$ and $D \subseteq \alpha_\xi \cap \text{Lim}$ is a club in α_ξ . Since (2) implies that $\pi_\xi(A) \in I(\triangleleft)$ and $\langle \gamma, \rho, D \rangle \in \pi_\xi(A) \subseteq N_\xi$, there is $\bar{D} \subseteq \kappa \cap \text{Lim}$ club in κ such that $\bar{D} \in M_\xi$,

$D = \pi_\xi(\bar{D}) = \bar{D} \cap \alpha_\xi$ and $\langle \gamma, \rho, D \rangle \triangleleft \langle \beta_0, \lambda_0, C_0 \rangle$. In this situation, the \triangleleft -minimality of $\langle \beta_0, \gamma_0, C_0 \rangle$ and elementarity imply that there is a closed bounded subset d of \bar{D} of order-type ρ such that $d \in M_\xi$ and $F(\alpha) = \gamma$ for all $\alpha \in d$. Then $\sup(d) < \alpha_\xi$ and d is a closed bounded subset of D of order-type ρ . These computations show that $\langle \beta_0, \lambda_0, C_0 \cap \kappa_\alpha \rangle$ is the \triangleleft -minimal triple that satisfies the assumption in (i) with respect to α_ξ , and we conclude that $F(\alpha_\xi) = \beta_0$.

Set $c = \{\alpha_\xi \mid \xi < \lambda_0\}$. We have just shown that c is a closed bounded subset of C_0 with $F(\alpha) = \beta_0$ for all $\alpha \in c$. This contradicts the choice of $\langle \beta_0, \lambda_0, C_0 \rangle$. ■

Set $S_\kappa = F^{-1}\{0\}$ and $S_\beta = F^{-1}\{1 + \beta\}$ for all $\beta < \kappa$. By Claim 1, $\{S_\beta \mid \beta \leq \kappa\}$ consists of pairwise disjoint fat stationary subsets of κ .

CLAIM 2. *The set $\{\langle S_\beta \mid \beta < \kappa \rangle\}$ is definable over $\langle H(\theta), \in \rangle$ by a Σ_1 -formula with parameters κ and ν .*

Proof. By Lemma 7.6, the set $\{\langle H(\kappa), \in, \triangleleft_\kappa \rangle\}$ is definable over the structure $\langle H(\theta), \in \rangle$ by a Σ_1 -formula with parameters κ and ν . Since $\langle H(\kappa), \in, \triangleleft_\kappa \rangle$ is a model of $ZFC^{\dot{A}}$ (the canonical extension of the axioms of ZFC^- to the language of set theory extended by a new predicate symbol \dot{A} , that includes all instances of Replacement and Separation for formulas in the extended language), we can use the Recursion Theorem within this structure in order to show that F is definable over $\{\langle H(\kappa), \in, \triangleleft_\kappa \rangle\}$ by a formula without parameters. In combination, these observations yield the conclusion. ■

Now, let \mathbb{P} be a partial order of cardinality less than δ such that forcing with \mathbb{P} preserves cofinalities less than or equal to κ^+ , and let G be \mathbb{P} -generic over V . By Definition 7.1(ii), $H(\kappa^+)^V$ is definable over $\langle H(\kappa^+)^{V[G]}, \in \rangle$ by a Σ_1 -formula with parameters κ and ν . By Claim 2 and Σ_1 -absoluteness, $\{\langle S_\beta \mid \beta < \kappa \rangle\}$ is definable over $\langle H(\kappa^+)^V, \in \rangle$ by a Σ_1 -formula with parameters κ and ν . Together, these statements imply that $\{\langle S_\beta \mid \beta < \kappa \rangle\}$ is definable over $\langle H(\kappa^+)^{V[G]}, \in \rangle$ by a Σ_1 -formula with parameters κ and ν . This completes the proof Theorem 7.7. ■

COROLLARY 7.8. *Let η be an infinite regular cardinal and let $\langle \delta, \theta, \nu, \triangleleft \rangle$ be suitable for $\kappa = \eta^+$. Then there is a sequence $\langle S_\alpha \mid \alpha \leq \kappa \rangle$ of pairwise disjoint stationary subsets of S_η^κ with the property that $\{\langle S_\alpha \mid \alpha < \kappa \rangle\}$ is definable over $\langle H(\kappa^+)^{V[G]}, \in \rangle$ by a Σ_1 -formula with parameters κ and ν whenever \mathbb{P} is a partial order of cardinality less than δ such that forcing with \mathbb{P} preserves cofinalities less than or equal to κ^+ and G is \mathbb{P} -generic over V .*

Proof. Let $\langle \bar{S}_\alpha \mid \alpha \leq \kappa \rangle$ be the sequence of pairwise disjoint fat stationary subsets of κ produced by Theorem 7.7. Given $\alpha \leq \kappa$, set $S_\alpha = \bar{S}_\alpha \cap S_\eta^\kappa$. Then S_α is a stationary subset of κ for each $\alpha \leq \kappa$.

Let \mathbb{P} be a partial order of cardinality less than δ such that forcing with \mathbb{P} preserves cofinalities less than or equal to κ^+ , and let G be \mathbb{P} -generic over V . By Lemma 7.6(vi) and Σ_1 -reflection, the set $\{S_\eta^\kappa\}$ is definable over $\langle H(\kappa^+)^{V[G]}, \in \rangle$ by a Σ_1 -formula with parameters κ and ν . Since Theorem 7.7 shows that the set $\{\vec{S}_\alpha \mid \alpha < \kappa\}$ is definable in the same way, this yields the statement of the corollary. ■

COROLLARY 7.9. *Assume that κ is either the successor of a regular cardinal or an inaccessible cardinal. Let $\langle \delta, \theta, \nu, \triangleleft \rangle$ be suitable for κ and let \mathbb{P} be a partial order of cardinality less than δ with the following properties:*

- (a) *Forcing with \mathbb{P} preserves cofinalities less than or equal to κ^+ , and preserves fat stationary subsets of κ .*
- (b) *If G is \mathbb{P} -generic over V , then 2^κ is regular, $\kappa = \kappa^{<\kappa}$ and $\eta^{<\eta} < \kappa$ for all $\eta < \kappa$ in $V[G]$.*

*Then there is a \mathbb{P} -name $\dot{\mathbb{Q}}$ for a partial order such that the following statements hold whenever $G * H$ is $(\mathbb{P} * \dot{\mathbb{Q}})$ -generic over V :*

- (i) *The partial order $\dot{\mathbb{Q}}^G$ is $<\kappa$ -distributive in $V[G]$.*
- (ii) *Forcing with $\dot{\mathbb{Q}}^G$ over $V[G]$ preserves all cofinalities less than or equal to $(2^\kappa)^{V[G]}$, and preserves the value of 2^κ .*
- (iii) *There is a well-ordering of $H(\kappa^+)^{V[G,H]}$ that is definable over the structure $\langle H(\kappa^+)^{V[G,H]}, \in \rangle$ by a Σ_1 -formula with parameters κ and ν .*

Proof. If κ is the successor of a regular cardinal η , let $\vec{S} = \langle S_\alpha \mid \alpha \leq \kappa \rangle$ denote the sequence of pairwise disjoint stationary subsets of S_η^κ produced by Corollary 7.8. If κ is an inaccessible cardinal, then we let \vec{S} denote the sequence of fat stationary subsets of κ provided by Theorem 7.7. In either case, by (a), \vec{S} is a sequence of either fat stationary subsets of κ or stationary subsets of S_η^κ respectively in every \mathbb{P} -generic extension of the ground model. Together with (b), this shows that κ and \vec{S} satisfy the requirements of Theorem 1.4 in every \mathbb{P} -generic extension of the ground model. Let $\dot{\mathbb{Q}}$ be a \mathbb{P} -name for the partial order given by Theorem 1.4. Since \mathbb{P} has cardinality less than δ , and δ is a strong limit cardinal, the construction of the forcing in the proof of Theorem 1.4 shows that we can find such a name with $\mathbb{P} * \dot{\mathbb{Q}}$ also having cardinality less than δ .

Let $G * H$ be $(\mathbb{P} * \dot{\mathbb{Q}})$ -generic over V . By Theorem 1.4, $\dot{\mathbb{Q}}^G$ is $<\kappa$ -distributive in $V[G]$, forcing with $\dot{\mathbb{Q}}^G$ over $V[G]$ preserves all cofinalities less than or equal to $(2^\kappa)^{V[G]}$ and preserves the value of 2^κ , and there is a well-ordering \triangleleft of $H(\kappa^+)^{V[G,H]}$ that is definable over $\langle H(\kappa^+)^{V[G,H]}, \in \rangle$ by a Σ_1 -formula with parameter $\langle S_\alpha \mid \alpha < \kappa \rangle$. Since Theorem 7.7 shows that $\{\langle S_\alpha \mid \alpha < \kappa \rangle\}$ is definable over $\langle H(\kappa^+)^{V[G,H]}, \in \rangle$ by a Σ_1 -formula with pa-

rameters κ and ν , we conclude that the well-order \blacktriangleleft is also definable over $\langle \mathbb{H}(\kappa^+)^{V[G,H]}, \in \rangle$ by a Σ_1 -formula with parameters κ and ν . ■

Proof of Theorem 1.5. Let \mathbb{P} be a partial order with properties (a) and (b) of Corollary 7.9. Pick a strong limit cardinal δ with $|\mathbb{P}| < \delta$. By Lemma 7.3, there is a well-order \triangleleft of $\mathbb{H}(\kappa^+)$ such that the tuple $\langle \delta, \kappa^+, 0, \triangleleft \rangle$ is suitable for κ . Let \dot{Q} be the \mathbb{P} -name for a partial order given by Corollary 7.9. Then the corollary implies that statements (i)–(iii) of the theorem hold. ■

Proof of Theorem 1.7. Let $\mathbb{P} \in V_\delta$ be a partial order with properties (a) and (b) of Corollary 7.9. By Lemma 7.5, there is a well-order \triangleleft of $\mathbb{H}(\kappa^+)$ such that the tuple $\langle \delta, \kappa^+, \kappa, \triangleleft \rangle$ is suitable for κ . Let \dot{Q} be the \mathbb{P} -name for a partial order given by Corollary 7.9 and let $G * H$ be $(\mathbb{P} * \dot{Q})$ -generic over V . Then the corollary implies that the partial order \dot{Q}^G is $<\kappa$ -distributive in $V[G]$, forcing with \dot{Q}^G over $V[G]$ preserves all cofinalities less than or equal to $(2^\kappa)^{V[G]}$ and preserves the value of 2^κ , and there is a well-ordering of $\mathbb{H}(\kappa^+)^{V[G,H]}$ that is definable over $\langle \mathbb{H}(\kappa^+)^{V[G,H]}, \in \rangle$ by a Σ_1 -formula with parameter κ . ■

Acknowledgements. During the preparation of the paper the first author was partially supported by the EPSRC Project EP/J005630/1. During the revision of the paper both authors were partially supported by the DFG through Projekt LU2020/1-1. Both authors would like to thank the organizers of the 2013 *ESI Workshop on Forcing and Large Cardinals* in Vienna, Austria, during which the results of this paper were mostly obtained. We would also like to thank the ESI itself for its hospitality during our stay. Moreover, we would like to thank Daisuke Ikegami, Peter Koepke, Ralf Schindler and David Schrittesser for valuable discussions. Finally, we thank the anonymous referee for the careful reading of the paper and numerous useful comments.

References

- [1] U. Abraham and S. Shelah, *Forcing closed unbounded sets*, J. Symbolic Logic 48 (1983), 643–657.
- [2] D. Asperó, P. Holy, and P. Lücke, *Forcing lightface definable well-orders without the GCH*, Ann. Pure Appl. Logic 166 (2015), 553–582.
- [3] A. Beller, R. Jensen, and P. Welch, *Coding the Universe*, London Math. Soc. Lecture Note Ser. 47, Cambridge Univ. Press, Cambridge, 1982.
- [4] A. Dodd and R. Jensen, *The core model*, Ann. Math. Logic 20 (1981), 43–75.
- [5] H. Friedman, *On closed sets of ordinals*, Proc. Amer. Math. Soc. 43 (1974), 190–192.
- [6] S.-D. Friedman and P. Holy, *Condensation and large cardinals*, Fund. Math. 215 (2011), 133–166.
- [7] S.-D. Friedman and P. Holy, *A quasi-lower bound on the consistency strength of PFA*, Trans. Amer. Math. Soc. 366 (2014), 4021–4065.

- [8] A. Halko and S. Shelah, *On strong measure zero subsets of ${}^{\kappa}2$* , Fund. Math. 170 (2001), 219–229.
- [9] L. Harrington, *Long projective wellorderings*, Ann. Math. Logic 12 (1977), 1–24.
- [10] P. Holy, *Σ_1 -wellorders without collapsing*, Arch. Math. Logic 54 (2015), 453–462.
- [11] P. Holy and P. Lücke, *Locally Σ_1 -definable well-orders of $H(\kappa^+)$* , Fund. Math. 226 (2014), 221–236.
- [12] R. B. Jensen and R. M. Solovay, *Some applications of almost disjoint sets*, in: Mathematical Logic and Foundations of Set Theory (Jerusalem, 1968), North-Holland, Amsterdam, 1970, 84–104.
- [13] P. Koepke, *Some applications of short core models*, Ann. Pure Appl. Logic 37 (1988), 179–204.
- [14] P. B. Larson, *Forcing over models of determinacy*, in: Handbook of Set Theory, Vol. 3, Springer, Dordrecht, 2010, 2121–2177.
- [15] P. Lücke, *Σ_1^1 -definability at uncountable regular cardinals*, J. Symbolic Logic 77 (2012), 1011–1046.
- [16] P. Lücke, R. Schindler, and P. Schlicht, *$\Sigma_1(\kappa)$ -definable subsets of $H(\kappa^+)$* , submitted.
- [17] R. Mansfield, *The non-existence of Σ_2^1 well-orderings of the Cantor set*, Fund. Math. 86 (1975), 279–282.
- [18] I. Neeman, *Determinacy in $L(\mathbb{R})$* , in: Handbook of Set Theory, Vol. 3, Springer, Dordrecht, 2010, 1877–1950.
- [19] S. Shelah, *Proper and Improper Forcing*, 2nd ed., Perspec. Math. Logic, Springer, Berlin, 1998.
- [20] W. H. Woodin, *The Axiom of Determinacy, Forcing Axioms, and the Nonstationary Ideal*, de Gruyter Ser. Logic Appl. 1, de Gruyter, Berlin, 1999.
- [21] M. Zeman, *Inner Models and Large Cardinals*, de Gruyter Ser. Logic Appl. 5, de Gruyter, Berlin, 2002.

Peter Holy, Philipp Lücke
Mathematisches Institut
Rheinische Friedrich-Wilhelms-Universität Bonn
Endenicher Allee 60
53115 Bonn, Germany
E-mail: pholy@math.uni-bonn.de
pluecke@uni-bonn.de

