

## Two inequalities between cardinal invariants

by

Dilip Raghavan (Singapore) and Saharon Shelah (Jerusalem)

**Abstract.** We prove two ZFC inequalities between cardinal invariants. The first inequality involves cardinal invariants associated with an analytic P-ideal, in particular the ideal of subsets of  $\omega$  of asymptotic density 0. We obtain an upper bound on the  $*$ -covering number, sometimes also called the weak covering number, of this ideal by proving that  $\text{cov}^*(\mathcal{Z}_0) \leq \mathfrak{d}$ . Next, we investigate the relationship between the bounding and splitting numbers at regular uncountable cardinals. We prove that, in sharp contrast to the case when  $\kappa = \omega$ , if  $\kappa$  is any regular uncountable cardinal, then  $\mathfrak{s}_\kappa \leq \mathfrak{b}_\kappa$ .

**1. Introduction.** Cardinal invariants associated with analytic P-ideals and their quotients are becoming increasingly well studied. Several cardinal invariants have been defined and investigated for quotients of the form  $\mathcal{P}(\omega)/\mathcal{I}$ , where  $\mathcal{I}$  is some definable ideal, guided by analogy with the familiar case of the quotient  $\mathcal{P}(\omega)/\text{FIN}$ . The most interesting among these have been the cases where  $\mathcal{I}$  is either  $F_\sigma$  or an analytic P-ideal. Recall that an ideal  $\mathcal{I}$  on  $\omega$  is called a *P-ideal* if for every collection  $\{a_n : n \in \omega\} \subset \mathcal{I}$ , there exists  $a \in \mathcal{I}$  such that  $\forall n \in \omega [a_n \subset^* a]$ . Here  $a \subset^* b$  means that  $a \setminus b$  is finite. When  $\mathcal{I}$  is a tall ideal, it is possible to associate some cardinals with  $\mathcal{I}$  that it would not necessarily make sense to do with FIN. Recall that an ideal  $\mathcal{I}$  on  $\omega$  is *tall* if it is *proper*, meaning  $\omega \notin \mathcal{I}$ , it is *non-principal*, meaning that  $[\omega]^{<\omega} \subset \mathcal{I}$ , and it has the property that  $\forall a \in [\omega]^\omega \exists b \in [a]^\omega [b \in \mathcal{I}]$ .

DEFINITION 1.1. Let  $\mathcal{I}$  be a tall P-ideal on  $\omega$ . Define

$$\begin{aligned} \text{add}^*(\mathcal{I}) &= \min\{|\mathcal{F}| : \mathcal{F} \subset \mathcal{I} \wedge \forall b \in \mathcal{I} \exists a \in \mathcal{F} [a \not\subset^* b]\}, \\ \text{cov}^*(\mathcal{I}) &= \min\{|\mathcal{F}| : \mathcal{F} \subset \mathcal{I} \wedge \forall a \in [\omega]^\omega \exists b \in \mathcal{F} [|a \cap b| = \omega]\}, \end{aligned}$$

---

2010 *Mathematics Subject Classification*: 03E17, 03E55, 03E05, 03E20.

*Key words and phrases*: asymptotic density, cardinal invariants, dominating number, weakly compact cardinal.

Received 22 April 2016; revised 21 July 2016.

Published online 1 December 2016.

$$\begin{aligned} \text{cof}^*(\mathcal{I}) &= \min\{|\mathcal{F}| : \mathcal{F} \subset \mathcal{I} \wedge \forall b \in \mathcal{I} \exists a \in \mathcal{F} [b \subset^* a]\}, \\ \text{non}^*(\mathcal{I}) &= \min\{|\mathcal{F}| : \mathcal{F} \subset [\omega]^\omega \wedge \forall b \in \mathcal{I} \exists a \in \mathcal{F} [|a \cap b| < \omega]\}. \end{aligned}$$

Cardinals of this kind were first considered by Brendle and Shelah [4] and by Bartoszyński [1]. Brendle and Shelah [4] referred to  $\text{add}^*(\mathcal{I})$  as  $\mathfrak{p}(\mathcal{I}^*)$  and to  $\text{cov}^*(\mathcal{I})$  as  $\pi\mathfrak{p}(\mathcal{I}^*)$ , where  $\mathcal{I}^* = \{\omega \setminus a : a \in \mathcal{I}\}$  is the *dual filter* to  $\mathcal{I}$ . The present terminology was popularized by Hernández-Hernández and Hrušák [9] who carried out a detailed investigation of these invariants for tall analytic P-ideals. Their choice of terminology was motivated by analogy with the following definitions which make sense for any ideal whatsoever.

DEFINITION 1.2. Let  $\mathcal{I}$  be any ideal on a set  $X$ . Define

$$\begin{aligned} \text{add}(\mathcal{I}) &= \min\left\{|\mathcal{F}| : \mathcal{F} \subset \mathcal{I} \wedge \bigcup \mathcal{F} \notin \mathcal{I}\right\}, \\ \text{cov}(\mathcal{I}) &= \min\left\{|\mathcal{F}| : \mathcal{F} \subset \mathcal{I} \wedge \bigcup \mathcal{F} = X\right\}, \\ \text{cof}(\mathcal{I}) &= \min\{|\mathcal{F}| : \mathcal{F} \subset \mathcal{I} \wedge \forall B \in \mathcal{I} \exists A \in \mathcal{F} [B \subset A]\}, \\ \text{non}(\mathcal{I}) &= \{ |Y| : Y \subset X \wedge Y \notin \mathcal{I} \}. \end{aligned}$$

It is possible to associate with each tall ideal  $\mathcal{I}$  on  $\omega$  an ideal  $\hat{\mathcal{I}}$  on  $\mathcal{P}(\omega)$  which is generated by Borel subsets of  $\mathcal{P}(\omega)$  in a natural way. For each  $a \in \mathcal{P}(\omega)$ , let  $\hat{a} = \{b \subset \omega : |a \cap b| = \omega\}$ . This is a  $G_\delta$  subset of  $\mathcal{P}(\omega)$ . If  $\mathcal{I}$  is a tall ideal on  $\omega$ , then  $\hat{\mathcal{I}} = \{X \subset \mathcal{P}(\omega) : \exists a \in \mathcal{I} [X \subset \hat{a}]\}$  is an ideal on  $\mathcal{P}(\omega)$  generated by Borel sets. Moreover  $\mathcal{I}$  is a P-ideal iff  $\hat{\mathcal{I}}$  is a  $\sigma$ -ideal. Now the invariants from Definition 1.2 associated with  $\hat{\mathcal{I}}$  correspond exactly with the  $*$ -invariants from Definition 1.1 associated with  $\mathcal{I}$ . It can be shown (see [9, Proposition 1.2]) that  $\text{add}(\hat{\mathcal{I}}) = \text{add}^*(\mathcal{I})$ ,  $\text{cov}(\hat{\mathcal{I}}) = \text{cov}^*(\mathcal{I})$ ,  $\text{cof}(\hat{\mathcal{I}}) = \text{cof}^*(\mathcal{I})$ ,  $\text{non}(\hat{\mathcal{I}}) = \text{non}^*(\mathcal{I})$ .

One of the main tools used in [9] for analyzing the  $*$ -invariants of tall analytic P-ideals is the Katětov order. Let  $\mathcal{I}$  and  $\mathcal{J}$  be ideals on  $\omega$ . Recall that  $\mathcal{I}$  is *Katětov below*  $\mathcal{J}$ , written  $\mathcal{I} \leq_K \mathcal{J}$ , if there is a function  $f : \omega \rightarrow \omega$  such that  $\forall a \in \mathcal{I} [f^{-1}(a) \in \mathcal{J}]$ . The significance of this ordering lies in the fact that  $\mathcal{I} \leq_K \mathcal{J}$  implies both that  $\text{cov}^*(\mathcal{I}) \geq \text{cov}^*(\mathcal{J})$  and that  $\text{non}^*(\mathcal{I}) \leq \text{non}^*(\mathcal{J})$  (see [9, Proposition 3.1]). The Tukey ordering is also relevant here. We say that  $\langle \mathcal{I}, \subset^* \rangle$  is *Tukey below*  $\langle \mathcal{J}, \subset^* \rangle$ , and we write  $\mathcal{I} \leq_T^* \mathcal{J}$ , if there is a map  $\varphi : \mathcal{I} \rightarrow \mathcal{J}$  such that if  $X \subset \mathcal{I}$  is any set that does not have an upper bound in the partial order  $\langle \mathcal{I}, \subset^* \rangle$ , then  $\varphi''X$  does not have an upper bound in the partial order  $\langle \mathcal{J}, \subset^* \rangle$ . If  $\mathcal{I} \leq_T^* \mathcal{J}$ , then  $\text{add}^*(\mathcal{I}) \geq \text{add}^*(\mathcal{J})$  and  $\text{cof}^*(\mathcal{I}) \leq \text{cof}^*(\mathcal{J})$  (see [9, Proposition 2.1]). The ideal  $\mathcal{Z}_0$  of sets of asymptotic density 0 is a critical object of study in [9].

DEFINITION 1.3. A set  $A \subset \omega$  is said to have *asymptotic density 0* if  $\lim_{n \rightarrow \infty} |A \cap n|/n = 0$ . Set  $\mathcal{Z}_0 = \{A \subset \omega : \lim_{n \rightarrow \infty} |A \cap n|/n = 0\}$ .

$\mathcal{Z}_0$  is easily seen to be a tall  $F_{\sigma\delta}$  P-ideal. It is pointed out in [9] that  $\text{add}^*(\mathcal{Z}_0) = \text{add}(\mathcal{N})$  and  $\text{cof}^*(\mathcal{Z}_0) = \text{cof}(\mathcal{N})$ , where  $\mathcal{N}$  is the ideal of subsets of  $\mathbb{R}$  that have Lebesgue measure 0. This follows from earlier work of Todorčević [15] and Fremlin [7] on the Tukey order  $\leq_T^*$ . Hernández-Hernández and Hrušák [9] prove that

$$\mathcal{ED}_{\text{fin}} \leq_K \mathcal{I}_{1/n} \leq_K \text{tr}(\mathcal{N}) \leq_K \mathcal{Z}_0.$$

Here  $\mathcal{ED}_{\text{fin}}$  is the ideal on  $\omega \times \omega$  generated by graphs of functions that lie below the diagonal; more formally, if  $\Delta = \{\langle n, i \rangle \in \omega \times \omega : i \leq n\}$ , then  $\mathcal{ED}_{\text{fin}} = \{A \subset \Delta : \exists m \forall n [|\{i : \langle n, i \rangle \in A\}| \leq m]\}$ . This is an  $F_\sigma$ -ideal. Next,  $\mathcal{I}_{1/n}$  is the ideal of summable sets: a set  $A \subset \omega$  is called *summable* if  $\sum_{n \in A} 1/(n+1) < \infty$ , and  $\mathcal{I}_{1/n} = \{A \subset \omega : A \text{ is summable}\}$ . This is an  $F_\sigma$  P-ideal. And finally,  $\text{tr}(\mathcal{N})$  is the trace of the null ideal: for a set  $A \subset 2^{<\omega}$ , we let  $G(A) = \{f \in 2^\omega : \exists^\infty n \in \omega [f \upharpoonright n \in A]\}$ ; then  $\text{tr}(\mathcal{N}) = \{A \subset 2^{<\omega} : \mu(G(A)) = 0\}$ , where  $\mu$  is the Lebesgue measure on  $2^\omega$ .

The upshot of all these Katětov reductions is that  $\text{add}(\mathcal{N}) \leq \text{cov}^*(\mathcal{Z}_0) \leq \text{non}(\mathcal{M})$  and dually  $\text{cov}(\mathcal{M}) \leq \text{non}^*(\mathcal{Z}_0) \leq \text{cof}(\mathcal{N})$ , where  $\mathcal{M}$  is the ideal of meager subsets of  $\mathbb{R}$ . Hernández-Hernández and Hrušák [9] further prove the inequalities

$$\min\{\text{cov}(\mathcal{N}), \mathfrak{b}\} \leq \text{cov}^*(\mathcal{Z}_0) \leq \max\{\mathfrak{b}, \text{non}(\mathcal{N})\}$$

and the dual inequalities

$$\min\{\mathfrak{d}, \text{cov}(\mathcal{N})\} \leq \text{non}^*(\mathcal{Z}_0) \leq \max\{\mathfrak{d}, \text{non}(\mathcal{N})\}.$$

Here  $\mathfrak{b}$  is the least size of an unbounded family in  $\omega^\omega$  with respect to the ordering of eventual domination, and  $\mathfrak{d}$  is the least size of a cofinal family. It is also proved in [9] that  $\mathcal{Z}_0$  is Katětov minimal among all density ideals.

Given these results the following question naturally suggests itself.

QUESTION 1.4 ([9, Question 3.23(a)]). *Is  $\text{cov}^*(\mathcal{Z}_0) \leq \mathfrak{d}$ ?*

Apart from the intrinsic interest in locating  $\text{cov}^*(\mathcal{Z}_0)$  in relation to the cardinals in Cichoń’s diagram, Question 1.4 also has a motivation coming from forcing theory. Recall the following definition.

DEFINITION 1.5. Let  $\mathbf{V}$  be any ground model and  $\mathbb{P} \in \mathbf{V}$  be a notion of forcing. Let  $\mathcal{I} \in \mathbf{V}$  be an ideal on  $\omega$ . We say that  $\mathbb{P}$  *diagonalizes*  $\mathbf{V} \cap \mathcal{I}$  if there exists  $\dot{A} \in \mathbf{V}^{\mathbb{P}}$  such that  $\Vdash_{\mathbb{P}} \dot{A} \in [\omega]^\omega$  and for each  $X \in \mathbf{V} \cap \mathcal{I}$ ,  $\Vdash_{\mathbb{P}} |X \cap \dot{A}| < \omega$ .

If  $\mathcal{I}$  is a definable tall ideal and if  $\mathbb{P} \in \mathbf{V}$  diagonalizes  $\mathbf{V} \cap \mathcal{I}$ , then  $\mathbb{P}$  tends to push  $\text{cov}^*(\mathcal{I})$  up. A classical theorem of Laflamme [11] says that any  $F_\sigma$  ideal can be diagonalized by a proper  $\omega^\omega$ -bounding forcing. When combined with standard preservation theorems and bookkeeping arguments, Laflamme’s result enables the construction of a model where  $\text{cov}^*(\mathcal{I}) > \mathfrak{d}$

for every tall  $F_\sigma$  ideal  $\mathcal{I}$ . Laflamme’s result has led to speculation about whether all tall  $F_{\sigma\delta}$  P-ideals, which arguably constitute the second nicest class of definable ideals after the  $F_\sigma$  ideals, could also be diagonalized by a proper  $\omega^\omega$ -bounding forcing.

QUESTION 1.6. *Suppose  $\mathcal{I} \in \mathbf{V}$  is an  $F_{\sigma\delta}$  P-ideal. Does there exist a proper  $\omega^\omega$ -bounding  $\mathbb{P} \in \mathbf{V}$  which diagonalizes  $\mathbf{V} \cap \mathcal{I}$ ? Is it consistent that  $\text{cov}^*(\mathcal{I}) > \mathfrak{d}$  for all tall  $F_{\sigma\delta}$  P-ideals  $\mathcal{I}$ ?*

It has long been known that one cannot hope for anything like this if one moves up one level to consider  $F_{\sigma\delta\sigma}$  ideals. The ideal  $\text{FIN} \times \text{FIN}$ , which is defined to be  $\{a \subset \omega \times \omega : \{n \in \omega : \{m \in \omega : \langle n, m \rangle \in a\} \text{ is infinite}\} \text{ is finite}\}$ , is an  $F_{\sigma\delta\sigma}$  ideal and any  $\mathbb{P}$  that diagonalizes it must add a dominating real. Indeed, diagonalizing  $\text{FIN} \times \text{FIN}$  is easily seen to be equivalent to adding a dominating real.

In Section 2 we give a positive answer to Question 1.4, along with a negative answer to Question 1.6, by proving the following theorem.

THEOREM 1.7.  $\text{cov}^*(\mathcal{Z}_0) \leq \mathfrak{d}$ .

The proof will show that if a proper forcing  $\mathbb{P} \in \mathbf{V}$  diagonalizes  $\mathbf{V} \cap \text{cov}^*(\mathcal{Z}_0)$ , then  $\mathbb{P}$  necessarily adds an unbounded real over  $\mathbf{V}$ . So we get a negative answer to both parts of Question 1.6 (see Corollary 2.10). Our proof also dualizes to show that  $\mathfrak{b} \leq \text{non}^*(\mathcal{Z}_0)$ .

Section 3 deals with cardinal invariants on uncountable cardinals.

DEFINITION 1.8. Let  $\kappa > \omega$  be a regular cardinal. For  $f, g \in \kappa^\kappa$ ,  $f <^* g$  means that  $|\{\alpha < \kappa : g(\alpha) \leq f(\alpha)\}| < \kappa$ . A set  $F \subset \kappa^\kappa$  is said to be *unbounded* if there does not exist  $g \in \kappa^\kappa$  such that  $\forall f \in F [f <^* g]$ . A set  $F \subset \kappa^\kappa$  is said to be *dominating* if  $\forall f \in \kappa^\kappa \exists g \in F [f <^* g]$ .

For  $a, b \in \mathcal{P}(\kappa)$ , we write  $a \subset^* b$  to mean that  $|a \setminus b| < \kappa$ . Since  $\kappa$  is regular, this is equivalent to saying that  $\exists \delta < \kappa [(a \setminus \delta) \subset b]$ . For  $a, b \in \mathcal{P}(\kappa)$  we say that  $a$  *splits*  $b$  if both  $b \cap a$  and  $b \cap (\kappa \setminus a)$  have cardinality  $\kappa$ . A family  $F \subset \mathcal{P}(\kappa)$  is called a *splitting family* if  $\forall b \in [\kappa]^\kappa \exists a \in F [a \text{ splits } b]$ .

We define the cardinal invariants  $\mathfrak{b}_\kappa$ ,  $\mathfrak{d}_\kappa$ , and  $\mathfrak{s}_\kappa$  as follows:

$$\begin{aligned} \mathfrak{b}_\kappa &= \min\{|F| : F \subset \kappa^\kappa \wedge F \text{ is unbounded}\}; \\ \mathfrak{d}_\kappa &= \min\{|F| : F \subset \kappa^\kappa \wedge F \text{ is dominating}\}; \\ \mathfrak{s}_\kappa &= \min\{|F| : F \subset \mathcal{P}(\kappa) \wedge F \text{ is a splitting family}\}. \end{aligned}$$

These are of course direct analogues of the cardinals  $\mathfrak{b}$ ,  $\mathfrak{d}$ , and  $\mathfrak{s}$  that play an important role in the theory of cardinal characteristics on  $\omega$ . Historically, one of the first works to investigate some of these higher analogues in depth was the paper [5] by Cummings and Shelah. They showed that for a regular  $\kappa > \omega$ ,  $\kappa^+ \leq \text{cf}(\mathfrak{b}_\kappa) = \mathfrak{b}_\kappa \leq \text{cf}(\mathfrak{d}_\kappa) \leq 2^\kappa$ . They also proved that these are

essentially the only restrictions on  $\mathfrak{b}_\kappa$  and  $\mathfrak{d}_\kappa$  that are provable in ZFC. In this sense,  $\mathfrak{b}_\kappa$  and  $\mathfrak{d}_\kappa$  behave in exactly the same way as  $\mathfrak{b}$  and  $\mathfrak{d}$ .

While the results of Cummings and Shelah [5] do not involve any large cardinals, it has lately become clear that obtaining consistency results on cardinal invariants at large cardinals is easier than obtaining the same consistency results at accessible cardinals. For example, a recent work of Garti and Shelah [8] proves the consistency of  $\mathfrak{u}_\kappa < 2^\kappa$  at a supercompact cardinal  $\kappa$ , based on earlier methods introduced by Džamonja and Shelah [6]. Here  $\mathfrak{u}_\kappa$  is the smallest number of sets needed to generate a uniform ultrafilter on  $\kappa$ . On the other hand, it is completely open whether this situation is consistent at  $\kappa = \omega_1$ .

It is a classical result of Shelah [12] that  $\mathfrak{b} < \mathfrak{s}$  is consistent. This result had a lot of impact on the study of cardinal invariants on  $\omega$ . It was the first published application of creature forcing, a method that has subsequently become indispensable to many consistency results on cardinal invariants on  $\omega$ . The importance of this result is one of the motivations for posing the following question.

**QUESTION 1.9.** *Is it consistent to have a regular uncountable cardinal  $\kappa$  such that  $\mathfrak{b}_\kappa < \mathfrak{s}_\kappa$ ?*

The cardinal  $\mathfrak{s}_\kappa$  has been investigated by Kamo [10], Suzuki [14], and Zapletal [16]. Suzuki [14] proved that  $\mathfrak{s}_\kappa$  is small for most regular uncountable cardinals. He showed that for a regular  $\kappa > \omega$ ,  $\mathfrak{s}_\kappa \geq \kappa$  iff  $\kappa$  is strongly inaccessible and  $\mathfrak{s}_\kappa \geq \kappa^+$  iff  $\kappa$  is weakly compact. It is easy to see that for a regular  $\kappa$ ,  $\kappa^+ \leq \mathfrak{b}_\kappa$ . Consequently,  $\mathfrak{b}_\kappa < \mathfrak{s}_\kappa$  implies  $\mathfrak{s}_\kappa \geq \kappa^{++}$ , and if  $\kappa$  is not weakly compact, then  $\mathfrak{s}_\kappa < \kappa^+ \leq \mathfrak{b}_\kappa$ . The main result of Zapletal [16] is that if it is consistent to have a regular uncountable cardinal  $\kappa$  such that  $\mathfrak{s}_\kappa \geq \kappa^{++}$ , then it is also consistent that there is a  $\kappa$  with  $o(\kappa) \geq \kappa^{++}$ . In particular, any positive answer to Question 1.9 would have had to start with a substantial large cardinal hypothesis. On the other hand, Kamo [10] proved that it is consistent relative to a supercompact cardinal that  $\mathfrak{s}_\kappa \geq \kappa^{++}$  at a supercompact  $\kappa$ . It was perhaps hoped that a positive answer to Question 1.9 would lead to new techniques for forcing at uncountable cardinals  $\kappa$ , at least when  $\kappa$  is supercompact, and help generate further results like the consistency of  $\mathfrak{b}_\kappa < \mathfrak{a}_\kappa$ . However, we will prove that Question 1.9 has a negative answer.

**THEOREM 1.10.** *For any regular uncountable cardinal  $\kappa$ ,  $\mathfrak{s}_\kappa \leq \mathfrak{b}_\kappa$ .*

It should be noted that this is not the first time that a significant difference has been observed in the behavior of cardinal characteristics between  $\omega$  and larger regular cardinals. Blass, Hyttinen, and Zhang [3] proved that  $\mathfrak{d}_\kappa = \kappa^+$  implies  $\mathfrak{a}_\kappa = \kappa^+$  for all regular uncountable cardinals  $\kappa$ , while the question of whether  $\mathfrak{d} = \omega_1$  implies  $\mathfrak{a} = \omega_1$  is a long-standing unresolved problem.

**2. A bound for  $\text{cov}^*(\mathcal{Z}_0)$ .** Theorem 1.7 is proved in this section.

DEFINITION 2.1. An *interval partition* is a strictly increasing function  $I = \langle i_n : n \in \omega \rangle \in \omega^\omega$  such that  $i_0 = 0$ . If  $I$  is an interval partition and  $n \in \omega$ , then  $I_n$  is the  $n$ th interval of the partition. In other words,  $I_n = [i_n, i_{n+1}) = \{k \in \omega : i_n \leq k < i_{n+1}\}$ .

LEMMA 2.2. *Let  $I$  be an interval partition. Let  $A \subset \omega$  be such that for each  $l \geq 0$ , there exists  $N \in \omega$  such that for each  $n \geq N$ :*

- (1)  $|A \cap I_n|/|I_n| \leq 2^{-l}$ ;
- (2)  $\forall i, j \in A \cap I_n [i \neq j \Rightarrow |i - j| > 2^{l-1}]$ .

Then  $A$  has density 0.

*Proof.* Fix  $l \geq 0$ . Using the given hypotheses, fix  $N \in \omega$  such that:

- (3)  $N > 0$  and  $i_N \geq 2^{l+1}$ ;
- (4) for each  $n \geq N$ :
  - (a)  $|A \cap I_n|/|I_n| \leq 2^{-l-2}$ ;
  - (b)  $\forall i, j \in A \cap I_n [i \neq j \Rightarrow |i - j| > 2^{l+1}]$ .

Find  $L \in \omega$  with  $M = i_L \geq 2^{l+2}i_N$ . We will show that for each  $k \geq M$ ,  $|A \cap k|/k \leq 2^{-l}$ . This will suffice to prove that  $A \in \mathcal{Z}_0$ . To this end, we first show that for each  $m \in \omega$ , if  $i_m \geq M$ , then  $|A \cap i_m|/i_m \leq 2^{-l-1}$ . Fix any such  $m$ . Note that  $m > 0$ ,  $i_N > 0$ , and  $i_m \geq 2^{l+2}i_N > i_N$ . Set  $Z = [i_N, i_m) = \bigcup_{N \leq k \leq m-1} I_k$ . For each  $N \leq k \leq m-1$ ,  $|A \cap I_k|/|I_k| \leq 2^{-l-2}$ , and so  $|A \cap Z|/|Z| \leq 2^{-l-2}$ . Therefore,

$$\frac{|A \cap i_m|}{i_m} = \frac{|A \cap i_N|}{i_m} + \frac{|A \cap Z|}{i_m} \leq \frac{i_N}{i_m} + \frac{|A \cap Z|}{|Z|}.$$

Since  $i_N/i_m \leq 2^{-l-2}$ , we get  $|A \cap i_m|/i_m \leq 2^{-l-2} + 2^{-l-2} = 2^{-l-1}$ , as needed.

Next, if  $k \geq M$ , then there exists  $m \in \omega$  such that  $i_m \geq M$  and  $k \in I_m$ . Thus it suffices to prove that for all  $m \in \omega$ , if  $i_m \geq M$ , then for all  $k \in I_m$ ,  $|A \cap k|/k \leq 2^{-l}$ . Fix any such  $m$ . If  $I_m \cap A = 0$ , then for any  $k \in I_m$ ,  $A \cap k = A \cap i_m$ , and so  $|A \cap k|/k \leq |A \cap k|/i_m = |A \cap i_m|/i_m \leq 2^{-l-1} \leq 2^{-l}$ . Thus we may assume that  $I_m \cap A \neq 0$ . Let  $\{a_1, \dots, a_p\}$  enumerate  $I_m \cap A$  in increasing order. Fix any  $k \in I_m$ . If  $k \leq a_1$ , then  $A \cap i_m = A \cap k$ , and so once again  $|A \cap k|/k \leq |A \cap k|/i_m = |A \cap i_m|/i_m \leq 2^{-l-1} \leq 2^{-l}$ . We may assume that  $a_1 < k$ . Set  $q = \max\{1 \leq q \leq p : a_q < k\}$  and note that  $A \cap k \subset (A \cap a_1) \cup \{a_1, \dots, a_q\}$ . By the remarks above,  $|A \cap a_1|/a_1 \leq 2^{-l-1}$ , and so  $|A \cap a_1|/k \leq |A \cap a_1|/a_1 \leq 2^{-l-1}$ . Clause (4)(b) implies that  $a_q - a_1 \geq (q-1)2^{l+1}$ , and clause (3) implies that  $2^{l+1} < a_1$ . It follows that  $q/k \leq 2^{-l-1}$ . Therefore  $|A \cap k|/k \leq |A \cap a_1|/k + q/k \leq 2^{-l-1} + 2^{-l-1} = 2^{-l}$ . ■

The next lemma describes an iterative procedure for dividing a finite set in half by taking every other element.

LEMMA 2.3. *Let  $l$  be a member of  $\omega$  greater than 0 and let  $X \subset \omega$  with  $|X| = 2^l$ . Then there exists a sequence  $\{A_\sigma : \sigma \in 2^{\leq l}\}$  such that:*

- (1)  $\forall m \leq l$   $[\bigcup_{\sigma \in 2^m} A_\sigma = X \wedge \forall \sigma, \tau \in 2^m [\sigma \neq \tau \Rightarrow A_\sigma \cap A_\tau = 0]]$ ;
- (2)  $\forall \sigma \in 2^{\leq l}$   $[|A_\sigma| = 2^{l-|\sigma|}]$  and  $\forall \sigma, \tau \in 2^{\leq l}$   $[\sigma \subset \tau \Rightarrow A_\tau \subset A_\sigma]$ ;
- (3) for each  $\sigma \in 2^{\leq l}$ ,  $\forall i, j \in A_\sigma$   $[i \neq j \Rightarrow |i - j| > 2^{|\sigma|-1}]$ .

*Proof.* Build the  $A_\sigma$  by induction on  $m \leq l$ . When  $m = 0$ , set  $A_\emptyset = X$ . Clause (3) is satisfied because  $\forall i, j \in X$   $[i \neq j \Rightarrow |i - j| \geq 1 > 1/2]$ . Now suppose  $m < l$  and that  $A_\sigma$  is given for every  $\sigma \in 2^m$ . Fix any  $\sigma \in 2^m$ . Then  $|A_\sigma| = 2^{l-m}$ . Note that  $2^{l-m} \geq 2$ . Let  $\phi : 2^{l-m} \rightarrow A_\sigma$  be the order isomorphism. Set  $E = \{x < 2^{l-m} : x \text{ is even}\}$  and  $O = \{x < 2^{l-m} : x \text{ is odd}\}$ . As  $2^{l-m} \geq 2$ ,  $|E| = |O| = 2^{l-m-1}$  and  $E \cup O = 2^{l-m}$ . Define  $A_{\sigma \frown \langle 0 \rangle} = \phi''E \subset A_\sigma$  and  $A_{\sigma \frown \langle 1 \rangle} = \phi''O \subset A_\sigma$ . It is easy to verify that the  $A_{\sigma \frown \langle i \rangle}$  satisfy (1)–(3) for  $\sigma \in 2^m$  and  $i \in 2$ . ■

The next lemma is a variation on a well-known characterization of the cardinal  $\mathfrak{d}$ , which may be found, for example, in [2]. We include a proof here for completeness.

LEMMA 2.4. *There is a family  $D$  of interval partitions such that:*

- (1)  $|D| \leq \mathfrak{d}$ ;
- (2) for each  $I \in D$  and for each  $n \in \omega$ , there exists  $l_n \in \omega$  such that  $l_n > 0$ ,  $l_n \geq n$ , and  $|I_n| = 2^{l_n}$ ;
- (3) for any interval partition  $J$  there exists  $I \in D$  such that  $\forall^\infty n \in \omega$   $\exists k > n$   $[J_k \subset I_n]$ .

*Proof.* Let  $F \subset \omega^\omega$  be a dominating family with  $|F| = \mathfrak{d}$ . Define an interval partition  $I_f$  as follows. First,  $i_{f,0} = 0$ . Given  $i_{f,n} \in \omega$ , let

$$M = \max(\{f(x) : x \leq i_{f,n} + 1\} \cup \{i_{f,n} + 1\}).$$

Find  $l_n \in \omega$  such that  $l_n > 0$ ,  $l_n \geq n$ , and  $2^{l_n} + i_{f,n} > M$ . Define  $i_{f,n+1} = 2^{l_n} + i_{f,n}$ . This completes the definition of the interval partition  $I_f$ .

Define  $D = \{I_f : f \in F\}$ . It is clear that  $|D| \leq |F| = \mathfrak{d}$ . And the definition of  $I_f$  ensures that for each  $n \in \omega$ ,  $|I_{f,n}| = 2^{l_n}$  for some  $l_n \in \omega$  with  $l_n > 0$  and  $l_n \geq n$ .

Now suppose that  $J$  is any interval partition. Define  $g \in \omega^\omega$  as follows: start with  $g(0) = 0$ . For any  $n \in \omega$ , given  $g(n) \in \omega$ , define  $g(n+1)$  as follows. Set  $x = g(n) + 1$  and let  $m_x$  be the unique  $m \in \omega$  such that  $x \in J_m$ . Let  $m = \max\{m_x, n\}$  and define  $g(n+1) = j_{m+2}$ . Note that  $g(n) < g(n) + 1 = x < j_{(m_x+2)} \leq j_{(m+2)} = g(n+1)$ . Thus  $g$  is strictly increasing. Now using the fact that  $F$  is dominating, find  $f \in F$  and  $N \in \omega$

such that  $\forall n \geq N [f(n) \geq g(n)]$ . Fix any  $n \geq N$ . Note that  $i_{f,n} + 1 > i_{f,n} \geq n \geq N$ . So  $g(i_{f,n} + 1) \leq f(i_{f,n} + 1)$ . Also, since  $g$  is strictly increasing,  $i_{f,n} \leq g(i_{f,n})$ . By the definition of  $g(i_{f,n} + 1)$ , there exists  $m \in \omega$  such that  $n \leq i_{f,n} \leq m$  and  $i_{f,n} \leq g(i_{f,n}) < g(i_{f,n}) + 1 < j_{m+1} < j_{m+2} = g(i_{f,n} + 1) \leq f(i_{f,n} + 1) < i_{f,n+1}$ . Thus  $J_{m+1} \subset I_{f,n}$ . Since  $m + 1 > n$ , we have shown that  $\forall n \geq N \exists k > n [J_k \subset I_{f,n}]$ , as needed. ■

**DEFINITION 2.5.** Let  $J$  be an interval partition such that for each  $n \in \omega$  there exists  $l_n \in \omega$  such that  $l_n > 0$ ,  $l_n \geq n$ , and  $|J_n| = 2^{l_n}$ . Applying Lemma 2.3, fix a sequence  $\bar{A} = \langle A_{n,\sigma} : n \in \omega \wedge \sigma \in 2^{\leq l_n} \rangle$  such that for each  $n \in \omega$ , the sequence  $\{A_{n,\sigma} : \sigma \in 2^{\leq l_n}\}$  satisfies (1)–(3) of Lemma 2.3 with  $l := l_n$  and  $X := J_n$ . Define  $\mathcal{F}_{J,\bar{A}}$  to be the collection of all functions  $f \in \omega^\omega$  such that for each  $n \in \omega$  and  $l < l_n$ , there exists  $\sigma \in 2^{l+1}$  such that  $f^{-1}(\{l\}) \cap J_n = A_{n,\sigma}$ , and there exists  $\tau \in 2^{l_n}$  such that  $f^{-1}(\{l_n\}) \cap J_n = A_{n,\tau}$ .

Observe that if  $f \in \mathcal{F}_{J,\bar{A}}$ , then  $f(k) \leq l_n$  for all  $n \in \omega$  and  $k \in J_n$ . Also, for any  $n, l \in \omega$ ,

$$\frac{|\{k \in J_n : f(k) \geq l\}|}{|J_n|} \leq 2^{-l},$$

and for any  $i, j \in \{k \in J_n : f(k) \geq l\}$ , if  $i \neq j$ , then  $|i - j| > 2^{l-1}$ . Moreover for any  $f \in \mathcal{F}_{J,\bar{A}}$ ,  $n \in \omega$ , and  $l \leq l_n$ , there is  $\sigma_{f,n,l} \in 2^l$  such that  $A_{n,\sigma_{f,n,l}} = \{k \in J_n : f(k) \geq l\}$ .

The next lemma is the crux of the construction. We define by induction on  $\omega_1$  a sequence of functions in such a way that any large enough finite set of these functions “covers” some infinite set of intervals in a suitable sense.

**LEMMA 2.6.** *Let  $J$  and  $\bar{A}$  be as in Definition 2.5. There exists a sequence of functions  $\langle f_{J,\bar{A},\alpha} : \alpha < \omega_1 \rangle$  such that:*

- (1)  $f_{J,\bar{A},\alpha} \in \mathcal{F}_{J,\bar{A}}$  for all  $\alpha < \omega_1$ ;
- (2) for each triple  $\langle i, m, F \rangle$  such that  $i, m \in \omega$ ,  $m \leq 2^i$ , and  $F \in [\omega_1]^m$ , there exists  $B_{i,m,F} \in [(\omega \setminus i)^\omega]$  such that

$$\forall \alpha, \beta \in F \forall n \in B_{i,m,F} [\alpha \neq \beta \Rightarrow \sigma_{f_{J,\bar{A},\alpha},n,i} \neq \sigma_{f_{J,\bar{A},\beta},n,i}].$$

*Proof.* We write  $\sigma_{\alpha,n,i}$  instead of  $\sigma_{f_{J,\bar{A},\alpha},n,i}$  and  $f_\alpha$  instead of  $f_{J,\bar{A},\alpha}$  to simplify the notation. We first define  $B_{i,0,\emptyset} = \omega \setminus i$  for all  $i \in \omega$ . Fix  $\alpha < \omega_1$ . Suppose that  $f_\xi$  has been defined for all  $\xi < \alpha$ . Suppose also that  $B_{i,m,F}$  has been defined for all triples  $\langle i, m, F \rangle$  such that  $i, m \in \omega$ ,  $m \leq 2^i$ , and  $F \in [\alpha]^m$ . Let  $\{\langle i_n, m_n, F_n \rangle : n \in \omega\}$  be a 1-1 enumeration of all triples  $\langle i, m, F \rangle$  such that  $i, m \in \omega$ ,  $m < 2^i$ , and  $F \in [\alpha]^m$ . Let  $B_n$  denote  $B_{i_n,m_n,F_n}$ . Find a sequence  $\langle C_n : n \in \omega \rangle$  such that  $C_n \in [B_n]^\omega$  and  $\forall n < m < \omega [C_n \cap C_m = \emptyset]$ . For each  $\langle i, m, F \rangle$  with  $i, m \in \omega$ ,  $m < 2^i$ , and  $F \in [\alpha]^m$ , define  $B_{i,m+1,F \cup \{\alpha\}} = C_n$ , where  $n$  is the unique  $n \in \omega$  such that  $\langle i_n, m_n, F_n \rangle = \langle i, m, F \rangle$ . Note that  $B_{i,m+1,F \cup \{\alpha\}} \in [(\omega \setminus i)^\omega]$ .

We now define  $f_\alpha \in \mathcal{F}_{J, \bar{A}}$ . Fix  $k \in \omega$ . Suppose first that  $k \in \bigcup_{n \in \omega} C_n$ . Let  $n$  be the unique  $n \in \omega$  such that  $k \in C_n$ . Since  $i_n \leq k \leq l_k$ ,  $\sigma_{\xi, k, i_n}$  is defined and belongs to  $2^{i_n}$  for each  $\xi \in F_n$ . Hence  $\{\sigma_{\xi, k, i_n} : \xi \in F_n\}$  is a subset of  $2^{i_n}$  with cardinality less than  $2^{i_n}$ . So choose  $\eta \in 2^{l_k}$  such that  $\eta \upharpoonright i_n \notin \{\sigma_{\xi, k, i_n} : \xi \in F_n\}$ . If  $k \notin \bigcup_{n \in \omega} C_n$ , then choose  $\eta \in 2^{l_k}$  to be arbitrary. In either case, for each  $l < l_k$ , define  $\tau_l = (\eta \upharpoonright l) \wedge \langle 1 - \eta(l) \rangle \in 2^{l+1}$  and define  $\tau_{l_k} = \eta$ . It is clear that for all  $l < l' \leq l_k$ ,  $A_{k, \tau_l} \cap A_{k, \tau_{l'}} = 0$  and that  $\bigcup_{l \leq l_k} A_{k, \tau_l} = J_k$ . Define  $f''_\alpha A_{k, \tau_l} = \{l\}$  for each  $l \leq l_k$ . This completes the definition of  $f_\alpha$ .

In the case when  $k \in C_n$  for some  $n \in \omega$ , we have  $\sigma_{\alpha, k, i_n} = \eta \upharpoonright i_n$ . One sees that  $f_\alpha \in \mathcal{F}_{J, \bar{A}}$  and  $\sigma_{\alpha, k, i_n} = \sigma_{\xi, k, i_n}$  for each  $n \in \omega$ ,  $\xi \in F_n$ , and  $k \in C_n$ .

Now to verify clause (2) after stage  $\alpha$  of the construction, suppose that  $\langle i, m, F \rangle$  is a triple such that  $i, m \in \omega$ ,  $m \leq 2^i$ , and  $F \in [\alpha + 1]^m$ . If  $F \in [\alpha]^m$ , then (2) holds by the induction hypothesis. So assume that  $\alpha \in F$  and let  $G = F \setminus \{\alpha\}$ . Let  $n$  be the unique element of  $\omega$  such that  $\langle i_n, m_n, F_n \rangle = \langle i, m - 1, G \rangle$ . Then  $B_{i, m, F} = C_n \subset B_n = B_{i, m-1, G}$ . Take  $\xi, \zeta \in F$  and  $k \in B_{i, m, F}$ . Suppose  $\xi < \zeta$ . If  $\xi, \zeta \in G$ , then since  $k \in B_{i, m-1, G}$ , we have  $\sigma_{\xi, k, i} \neq \sigma_{\zeta, k, i}$  by the induction hypothesis. If  $\zeta = \alpha$ , then  $\sigma_{\zeta, k, i} = \sigma_{\alpha, k, i} = \sigma_{\alpha, k, i_n} \neq \sigma_{\xi, k, i_n} = \sigma_{\xi, k, i}$ . This finishes the construction. ■

Note that if  $m = 2^i$  and  $F \in [\omega_1]^m$ , then for any  $k \in B_{i, m, F}$ ,  $\{\sigma_{\alpha, k, i} : \alpha \in F\}$  is a subset of  $2^i$  of size  $2^i$ . Therefore  $\bigcup_{\alpha \in F} A_{k, \sigma_{\alpha, k, i}} = \bigcup_{\sigma \in 2^i} A_{k, \sigma} = J_k$ . This consequence of Lemma 2.6 will be used below.

**DEFINITION 2.7.** Let  $D$  be a family of interval partitions as in Lemma 2.4. For each  $J \in D$  fix a sequence  $\bar{A}_J = \langle A_{J, k, \sigma} : k \in \omega \wedge \sigma \in 2^{\leq l_{J, k}} \rangle$  as in Definition 2.5. Use Lemma 2.6 to fix a sequence  $\langle f_{J, \bar{A}_J, \alpha} : \alpha < \omega_1 \rangle$  of functions satisfying (1) and (2) of the lemma. We will write  $f_{J, \alpha}$  instead of  $f_{J, \bar{A}_J, \alpha}$ . For any tuple  $\langle I, J, \alpha, l \rangle \in D \times D \times \omega_1 \times \omega$ , let  $Z_{I, J, \alpha, l} = \bigcup_{k \in I_l} \{x \in J_k : f_{J, \alpha}(x) \geq l\}$ . For each triple  $\langle I, J, \alpha \rangle \in D \times D \times \omega_1$  define  $Z_{I, J, \alpha} = \bigcup_{l \in \omega} Z_{I, J, \alpha, l} \subset \omega$ . Note that the family  $\{Z_{I, J, \alpha} : \langle I, J, \alpha \rangle \in D \times D \times \omega_1\}$  has size at most  $\mathfrak{d}$ .

**LEMMA 2.8.**  $Z_{I, J, \alpha} \in \mathcal{Z}_0$  for every triple  $\langle I, J, \alpha \rangle \in D \times D \times \omega_1$ .

*Proof.* We apply Lemma 2.2 with  $J$  as  $I$  and  $Z_{I, J, \alpha}$  as  $A$ . Fix  $l \geq 0$ . Let  $N = i_l$  and let  $k \geq N$  be given. Let  $l^*$  be the unique member of  $\omega$  such that  $k \in I_{l^*}$ . Note that  $l \leq l^*$ . Now  $Z_{I, J, \alpha} \cap J_k = \{x \in J_k : f_{J, \alpha}(x) \geq l^*\}$ . Since  $f_{J, \alpha} \in \mathcal{F}_{J, \bar{A}_J}$ , we have  $|Z_{I, J, \alpha} \cap J_k|/|J_k| \leq 2^{-l^*} \leq 2^{-l}$  and for any  $i, j \in Z_{I, J, \alpha} \cap J_k$ , if  $i \neq j$ , then  $|i - j| > 2^{l^*-1} \geq 2^{l-1}$ , exactly as needed. ■

**LEMMA 2.9.** Fix  $J \in D$  and  $\alpha \in \omega_1$ . Suppose  $A \subset \omega$  and suppose that  $A \cap Z_{I, J, \alpha}$  is finite for every  $I \in D$ . Then there exists  $i \in \omega$  such that  $f''_{J, \alpha} A \subset i$ .

*Proof.* Suppose not. Then for every  $i \in \omega$ , there exists  $x \in A$  such that  $f_{J,\alpha}(x) \geq i$ . Define an interval partition  $K$  as follows. First, set  $k_0 = 0$ . Given  $k_n$ , let  $U = f''_{J,\alpha}(\bigcup_{k < k_n} J_k)$ . Let  $i = \max(U \cup \{n\}) + 1$ . Find  $x \in A$  such that  $f_{J,\alpha}(x) \geq i$ . Let  $k^* \in \omega$  be such that  $x \in J_{k^*}$ . Note that  $k_n \leq k^*$ . Set  $k_{n+1} = k^* + 1$ . This finishes the construction of  $K$ .

By construction we have  $\forall n \in \omega \exists k \in K_n \exists x \in J_k \cap A [f_{J,\alpha}(x) > n]$ . Find  $I \in D$  and  $N \in \omega$  such that  $\forall l \geq N \exists n > l [K_n \subset I_l]$ . It follows that  $\forall l \geq N [Z_{I,J,\alpha,l} \cap A \neq \emptyset]$ . This contradicts the hypothesis that  $Z_{I,J,\alpha} \cap A$  is finite. ■

*Proof of Theorem 1.7.* The family  $\{Z_{I,J,\alpha} : \langle I, J, \alpha \rangle \in D \times D \times \omega_1\}$  is a subset of  $\mathcal{Z}_0$  of cardinality at most  $\mathfrak{d}$ . Suppose for a contradiction that  $A \in [\omega]^\omega$  and that  $A \cap Z_{I,J,\alpha}$  is finite for every  $\langle I, J, \alpha \rangle \in D \times D \times \omega_1$ . Fix  $J \in D$  such that  $\forall^\infty k \in \omega [A \cap J_k \neq \emptyset]$ . By Lemma 2.9,  $\forall \alpha < \omega_1 \exists i_\alpha \in \omega [f''_{J,\alpha} A \subset i_\alpha]$ . There exist  $i \in \omega$  and  $S \in [\omega_1]^{\omega_1}$  such that  $\forall \alpha \in S [i = i_\alpha]$ . Let  $m = 2^i$  and  $F \in [S]^m$ . By the remark following Lemma 2.6, there exists  $B_{J,\bar{A}_J,i,m,F} \in [\omega]^\omega$  such that for every  $k \in B_{J,\bar{A}_J,i,m,F}$ , we have  $\bigcup_{\alpha \in F} A_{J,k,\sigma_{J,\bar{A}_J,\alpha,k,i}} = J_k$ , where  $\{x \in J_k : f_{J,\alpha}(x) \geq i\} = A_{J,k,\sigma_{J,\bar{A}_J,\alpha,k,i}}$ . It follows that  $A \cap J_k = \emptyset$  for each  $k \in B_{J,\bar{A}_J,i,m,F}$ . However, this contradicts the choice of  $J$ . ■

The proof of Theorem 1.7 shows that if  $\mathbb{P} \in \mathbf{V}$  is an  $\omega_1$ -preserving forcing and if there is a set  $A \in \mathbf{V}^\mathbb{P} \cap [\omega]^\omega$  which is almost disjoint from every member of  $\{Z_{I,J,\alpha} : \langle I, J, \alpha \rangle \in D \times D \times \omega_1\}$ , then  $D$  no longer satisfies clause (3) of Lemma 2.4 in  $\mathbf{V}^\mathbb{P}$ . Hence  $\mathbb{P}$  necessarily adds an unbounded real.

**COROLLARY 2.10.** *Let  $\mathbf{V}$  be any ground model and let  $E \in \mathbf{V}$  be a dominating family of minimal size. If  $\mathbb{P} \in \mathbf{V}$  preserves  $\omega_1$  and diagonalizes  $\mathcal{Z}_0 \cap \mathbf{V}$ , then  $E$  is no longer a dominating family in  $\mathbf{V}^\mathbb{P}$ .*

It is fairly straightforward to dualize the proof of Theorem 1.7 to show that  $\mathfrak{b} \leq \text{non}^*(\mathcal{Z}_0)$ . The proof of Theorem 1.7 can also be adapted to cover several density ideals (see [9, Definition 1.6] for the exact definition of a density ideal). By the results in [9],  $\text{cov}^*(\mathcal{Z}_0) \leq \text{cov}^*(\mathcal{I})$  for every density ideal  $\mathcal{I}$ .

Solecki and Todorcevic [13] have introduced the more general notion of a density-like ideal.

**DEFINITION 2.11.** Let  $\varphi : 2^\omega \rightarrow [0, \infty]$  be a lower semicontinuous submeasure. Let  $\mathcal{I} = \text{Exh}(\varphi) = \{X \subset \omega : \lim_{n \rightarrow \infty} \varphi(X \setminus n) = 0\}$ . Then  $\mathcal{I}$  is said to be *density-like* if for every  $\epsilon > 0$  there exists  $\delta > 0$  such that for every sequence  $\langle F_n : n \in \omega \rangle$  of finite sets, if  $\forall n \in \omega [\varphi(F_n) < \delta]$ , then there exists  $a \in [\omega]^\omega$  such that  $\varphi(\bigcup_{n \in a} F_n) < \epsilon$ .

It would be of interest to see whether a bound on  $\text{cov}^*(\mathcal{I})$  similar to that in Theorem 1.7 can be proved for all density-like ideals  $\mathcal{I}$ .

**3. The bounding and splitting numbers at uncountable cardinals.** Theorem 1.10, which says that  $\mathfrak{s}_\kappa \leq \mathfrak{b}_\kappa$  for every regular uncountable cardinal  $\kappa$ , is proved in this section. We begin with an observation, due to Suzuki [14], that for a regular uncountable  $\kappa$ ,  $\mathfrak{s}_\kappa > \kappa$  implies that  $\kappa$  is weakly compact. Though we will not need this observation, we include a proof for completeness. Its converse is also true and was noted by Suzuki.

LEMMA 3.1 (Suzuki). *Let  $\kappa > \omega$  be a regular cardinal. If  $\mathfrak{s}_\kappa > \kappa$ , then  $\kappa$  is weakly compact.*

*Proof.* We show that  $\kappa \rightarrow (\kappa)_2^2$ . Let  $c : [\kappa]^2 \rightarrow 2$  be a coloring. For each  $\alpha < \kappa$  and  $i \in 2$ , let  $K_{\alpha,i} = \{\beta > \alpha : c(\{\alpha, \beta\}) = i\}$ . Since  $\{K_{\alpha,0} : \alpha < \kappa\}$  is not a splitting family, there exists  $x \in [\kappa]^\kappa$  such that  $\forall \alpha < \kappa \exists i_\alpha \in 2 [x \subset^* K_{\alpha,i_\alpha}]$ . Find  $y \in [x]^\kappa$  and  $i \in 2$  such that  $\forall \gamma \in y [i_\gamma = i]$ . Define a sequence  $\langle \gamma_\alpha : \alpha < \kappa \rangle \subset y$  by induction on  $\alpha < \kappa$  as follows. Fix  $\alpha < \kappa$  and suppose that  $\langle \gamma_\xi : \xi < \alpha \rangle \subset y$  is given. For each  $\xi < \alpha$ , there exists  $\delta_\xi < \kappa$  such that  $x \setminus \delta_\xi \subset K_{\gamma_\xi,i}$ . Set  $\delta = \sup\{\delta_\xi : \xi < \alpha\} < \kappa$  and  $\gamma_\alpha = \min(y \setminus \delta)$ . It is clear that for each  $\xi < \alpha$ ,  $\gamma_\xi < \gamma_\alpha$  and  $c(\{\gamma_\xi, \gamma_\alpha\}) = i$ . Therefore  $\{\gamma_\alpha : \alpha < \kappa\}$  is a homogeneous set of cardinality  $\kappa$  for the color  $i$ . ■

DEFINITION 3.2. Let  $\kappa > \omega$  be regular and suppose that  $\kappa^+ < \mathfrak{s}_\kappa$ . Let  $\lambda$  be a cardinal such that  $\kappa < \lambda < \mathfrak{s}_\kappa$ . Fix a sufficiently large regular cardinal  $\theta$  ( $\theta = (2^{2^{\mathfrak{s}_\kappa}})^+$  will suffice). Let  $M \prec H(\theta)$  be such that  $\lambda \subset M$  and  $|M| = \lambda$ . We are going to define a filter  $D$  on  $\kappa$ , an equivalence relation  $\sim_D$  on  $M \cap \kappa^\kappa$ , and an ordering  $<_D$  on the  $\sim_D$ -equivalence classes as follows.  $M \cap \mathcal{P}(\kappa)$  is not a splitting family. So there exists  $A_* \in [\kappa]^\kappa$  such that for all  $x \in M \cap \mathcal{P}(\kappa)$  either  $A_* \subset^* \kappa \setminus x$  or  $A_* \subset^* x$ . Define  $D$  to be  $\{x \in \mathcal{P}(\kappa) : A_* \subset^* x\}$ . For any  $f, g \in M \cap \kappa^\kappa$ , define  $f \sim_D g$  iff  $\{\alpha < \kappa : f(\alpha) = g(\alpha)\} \in D$ . This is an equivalence relation on  $M \cap \kappa^\kappa$ . For  $f \in M \cap \kappa^\kappa$ , let  $[f]_D = \{g \in M \cap \kappa^\kappa : f \sim_D g\}$ . For  $f, g \in M \cap \kappa^\kappa$ , define  $[f]_D \sim_D [g]_D$  iff  $\{\alpha < \kappa : f(\alpha) < g(\alpha)\} \in D$ . Let  $L = \{[f]_D : f \in M \cap \kappa^\kappa\}$ .

We also fix some notation. Let  $i : \kappa \rightarrow \kappa$  be the identity function on  $\kappa$ . For each  $\alpha \in \kappa$ , let  $c_\alpha : \kappa \rightarrow \kappa$  be the function such that  $c_\alpha(\beta) = \alpha$  for all  $\beta \in \kappa$ . Note that  $i, c_\alpha \in M \cap \kappa^\kappa$  for all  $\alpha \in \kappa$ .

Observe that  $D$  is a  $\kappa$ -complete filter on  $\kappa$ . Also, if  $X \in D$ , then  $|X| = \kappa$ . Next we note that for any  $\delta < \kappa$ ,  $\kappa \setminus \delta \in D$ . Finally, if  $0 < \delta < \kappa$  and  $\langle X_\alpha : \alpha < \delta \rangle \in M$  is a partition of  $\kappa$ , then there is a unique  $\alpha < \delta$  such that  $X_\alpha \in D$ . To see this note that  $\delta \subset \kappa \subset \lambda \subset M$ , and so  $\{X_\alpha : \alpha < \delta\} \subset M$ . For each  $\alpha < \delta$  there exists  $i_\alpha \in 2$  such that  $A_* \subset^* X_\alpha^{i_\alpha}$ , where  $X_\alpha^0 = X_\alpha$  and  $X_\alpha^1 = \kappa \setminus X_\alpha$ . Thus  $\{X_\alpha^{i_\alpha} : \alpha < \delta\} \subset D$ , and by the  $\kappa$ -completeness of  $D$ ,  $X = \bigcap \{X_\alpha^{i_\alpha} : \alpha < \delta\} \in D$ . So  $|X| = \kappa$ . By hypothesis, if  $\alpha < \beta < \delta$ , then  $X_\alpha \cap X_\beta = 0$ , and also  $\bigcap \{X_\alpha^1 : \alpha < \delta\} = 0$ . It follows that  $i_\alpha = 0$  for some unique  $\alpha < \delta$ .

LEMMA 3.3. *The structure  $\langle L, <_D \rangle$  is a linear order. Moreover  $\{[c_\alpha]_D : \alpha < \kappa\}$  has a least upper bound in  $L$ .*

*Proof.* The relation  $<_D$  is transitive because  $D$  is a filter. Given  $f, g \in M \cap \kappa^\kappa$ , the sets  $\{\alpha < \kappa : f(\alpha) = g(\alpha)\}$ ,  $\{\alpha < \kappa : f(\alpha) < g(\alpha)\}$ , and  $\{\alpha < \kappa : g(\alpha) < f(\alpha)\}$  all belong to  $M$  and they partition  $\kappa$ . So by the remarks above, exactly one of them belongs to  $D$ , whence exactly one of  $[f]_D = [g]_D$ ,  $[f]_D \sim_D [g]_D$ , or  $[g]_D \sim_D [f]_D$  holds.

For the “moreover” statement, note that  $[i]_D \in L$  and is an upper bound of  $\{[c_\alpha]_D : \alpha < \kappa\}$ . If there is no least upper bound, then we can get a sequence  $\langle f_n : n \in \omega \rangle \subset M \cap \kappa^\kappa$  such that for each  $n \in \omega$ ,  $[f_n]_D \in L$  is an upper bound of  $\{[c_\alpha]_D : \alpha < \kappa\}$  and  $[f_{n+1}]_D \sim_D [f_n]_D$ . Thus for each  $n \in \omega$ ,  $X_n = \{\beta < \kappa : f_{n+1}(\beta) < f_n(\beta)\} \in D$ . By  $\kappa$ -completeness of  $D$ ,  $X = \bigcap_{n \in \omega} X_n \in D$ , and so  $X \neq \emptyset$ . Choosing  $\beta \in X$ , we get an infinite descending sequence of ordinals  $f_0(\beta) > f_1(\beta) > \dots$ , which is a contradiction. ■

Of course the argument of Lemma 3.3 shows that  $\langle L, <_D \rangle$  is a well-order. But we will not need this in what follows. One can also take the reduced power of the structure  $\langle M, \in \rangle$  with respect to the filter  $D$ . The above argument shows that this structure will be well-founded. It is also possible to argue for Theorem 1.10 in terms of the resulting embedding, which may not be elementary. Similar ideas were used by Zapletal [16] to prove that  $\mathfrak{s}_\kappa > \kappa^+$  has large consistency strength. The proof we give below avoids dealing with the reduced power of  $\langle M, \in \rangle$ .

DEFINITION 3.4. Fix a function  $f_* \in M \cap \kappa^\kappa$  such that  $[f_*]_D \in L$  is a least upper bound of  $\{[c_\alpha]_D : \alpha < \kappa\}$ .

LEMMA 3.5. *If  $C \in M$  is a club in  $\kappa$ , then  $f_*^{-1}(C) \in D$ .*

*Proof.* Suppose for a contradiction that  $f_*^{-1}(C) \notin D$ . Since  $f_*, C, \kappa \in M$ , both  $f_*^{-1}(C)$  and  $\kappa \setminus f_*^{-1}(C) = f_*^{-1}(\kappa \setminus C)$  belong to  $M$  and partition  $\kappa$ . Therefore,  $X_0 = f_*^{-1}(\kappa \setminus C) \in D$ . Next since  $[c_0]_D \sim_D [f_*]_D$ , we have  $X_1 = \{\beta < \kappa : 0 < f_*(\beta)\} \in D$ . Thus  $X = X_0 \cap X_1 \in M \cap D$ . Define a function  $f : \kappa \rightarrow \kappa$  as follows. For any  $\alpha \in \kappa$ , if  $\alpha \in X$ , then let  $f(\alpha) = \sup(C \cap f_*(\alpha))$ , and let  $f(\alpha) = 0$  otherwise. If  $\alpha \in X$ , then since  $f_*(\alpha) \notin C$ , we have  $f(\alpha) = \sup(C \cap f_*(\alpha)) < f_*(\alpha)$ . It is clear that  $f \in M$ . Thus  $[f]_D \in L$  and  $[f]_D \sim_D [f_*]_D$ . On the other hand, consider any  $\alpha < \kappa$ . Fix  $\delta \in C$  with  $\delta > \alpha$ . Since  $[c_\delta]_D \sim_D [f_*]_D$ , we have  $Y = \{\beta < \kappa : \delta < f_*(\beta)\} \in D$ . So  $Z = X \cap Y \in D$  and  $\forall \beta \in Z [c_\alpha(\beta) = \alpha < f(\beta)]$ , whence  $[c_\alpha]_D \sim_D [f]_D$ . However, this contradicts the choice of  $f_*$ . ■

LEMMA 3.6.  *$M \cap \kappa^\kappa$  is bounded.*

*Proof.* First note that  $f_*'' A_*$  is an unbounded subset of  $\kappa$ . This is because for any  $\alpha < \kappa$ ,  $\{\beta < \kappa : \alpha < f_*(\beta)\} \in D$ , whence  $A_* \subset^* \{\beta < \kappa : \alpha < f_*(\beta)\}$ .

So we can find  $\beta \in A_*$  with  $\alpha < f_*(\beta)$  because  $|A_*| = \kappa$ . Next if  $C \in M$  is any club in  $\kappa$ , then by Lemma 3.5,  $A_* \subset^* f_*^{-1}(C)$ . It follows that  $f''A_* \subset^* C$ . Now since  $\kappa$  is regular,  $\text{otp}(f''A_*) = \kappa$ . Let  $g : \kappa \rightarrow f''A_*$  be the unique order isomorphism. Define  $h : \kappa \rightarrow \kappa$  by  $h(\alpha) = g(\alpha + 1)$  for each  $\alpha \in \kappa$ .

We claim that  $h$  bounds  $M \cap \kappa^\kappa$ . Indeed, let  $f \in M \cap \kappa^\kappa$ . Then  $C_f = \{\alpha < \kappa : \alpha \text{ is closed under } f\} \in M$  is a club in  $\kappa$ . Therefore there exists  $\delta < \kappa$  such that  $(f''A_*) \setminus \delta \subset C_f$ . Consider any  $\alpha < \kappa$  with  $\alpha \geq \delta$ . Then  $h(\alpha) = g(\alpha + 1) \in f''A_*$ , and since  $g$  is order preserving,  $\delta \leq \alpha < \alpha + 1 \leq g(\alpha + 1)$ . Therefore  $g(\alpha + 1) \in C_f$ , and since  $\alpha < g(\alpha + 1)$ , we obtain  $f(\alpha) < g(\alpha + 1) = h(\alpha)$ . Thus we have shown that  $\forall \alpha < \kappa [\alpha \geq \delta \Rightarrow f(\alpha) < h(\alpha)]$ , as required. ■

*Proof of Theorem 1.10.* Suppose for a contradiction that  $\mathfrak{b}_\kappa < \mathfrak{s}_\kappa$ . Define  $\lambda = \mathfrak{b}_\kappa$ . By the results in [5],  $\kappa < \lambda < \mathfrak{s}_\kappa$ . Let  $\{f_\xi : \xi < \lambda\} \subset \kappa^\kappa$  be an unbounded family of size  $\lambda$ . Let  $M \prec H(\theta)$  be such that  $|M| = \lambda$  and  $\lambda \cup \{f_\xi : \xi < \lambda\} \subset M$ . Applying Lemma 3.6, we deduce that  $\{f_\xi : \xi < \lambda\} \subset M \cap \kappa^\kappa$  is bounded, which is a contradiction. ■

We do not know whether the above proof can be dualized to show that  $\mathfrak{d}_\kappa \leq \mathfrak{r}_\kappa$  for all regular uncountable  $\kappa$ . The invariant  $\mathfrak{d}_\kappa$  has already been defined in 1.8, and  $\mathfrak{r}_\kappa$  is the dual of the splitting number, known as the *reaping number* for  $\kappa$ . A set  $a \in \mathcal{P}(\kappa)$  is said to *reap* a family  $\mathcal{F} \subset [\kappa]^\kappa$  if  $a$  splits each  $b \in \mathcal{F}$ . A family  $\mathcal{F}$  is *unreaped* if there is no  $a \in \mathcal{P}(\kappa)$  reaping  $\mathcal{F}$ . Now  $\mathfrak{r}_\kappa$  is defined as follows:

$$\mathfrak{r}_\kappa = \min\{|\mathcal{F}| : \mathcal{F} \subset [\kappa]^\kappa \wedge \mathcal{F} \text{ is unreaped}\}.$$

The proof of Theorem 1.10 given in this section does not provide much information about how  $\mathfrak{r}_\kappa$  and  $\mathfrak{d}_\kappa$  are related.

**4. Some questions.** It is unknown to what extent the bound on  $\text{cov}^*(\mathcal{Z}_0)$  given by Theorem 1.7 can be improved.

QUESTION 4.1. *Is  $\text{cov}^*(\mathcal{Z}_0) \leq \mathfrak{b}$ ?*

This is essentially equivalent to asking whether there is a proper forcing that diagonalizes  $\mathbf{V} \cap \mathcal{Z}_0$  while preserving all unbounded families in  $\mathbf{V}$ .

The following is an outstanding open problem about cardinal invariants above the continuum.

QUESTION 4.2 <sup>(1)</sup>. *Is it consistent to have a regular uncountable cardinal  $\kappa$  for which  $\mathfrak{b}_\kappa < \mathfrak{a}_\kappa$ ?*

Finally, we ask whether the inequality  $\mathfrak{s}_\kappa \leq \mathfrak{b}_\kappa$  can be dualized in ZFC.

---

<sup>(1)</sup> The authors have recently proved the consistency of  $\mathfrak{b}_\kappa < \mathfrak{a}_\kappa$  for a supercompact cardinal  $\kappa$  assuming the consistency of another supercompact cardinal  $\sigma > \kappa$ .

QUESTION 4.3. *Is it consistent to have a regular uncountable cardinal  $\kappa$  for which  $\mathfrak{r}_\kappa < \mathfrak{d}_\kappa$ ?*

**Acknowledgments.** The first author was partially supported by National University of Singapore research grant number R-146-000-211-112. Both authors were partially supported by European Research Council grant 338821. This is paper 1060 on Shelah's list.

### References

- [1] T. Bartoszyński, *Invariants of measure and category*, in: Handbook of Set Theory, Vol. 1, Springer, Dordrecht, 2010, 491–555.
- [2] A. Blass, *Combinatorial cardinal characteristics of the continuum*, in: Handbook of Set Theory, Vol. 1, Springer, Dordrecht, 2010, 395–489.
- [3] A. Blass, T. Hyttinen, and Y. Zhang, *Mad families and their neighbours*, preprint.
- [4] J. Brendle and S. Shelah, *Ultrafilters on  $\omega$ —their ideals and their cardinal characteristics*, Trans. Amer. Math. Soc. 351 (1999), 2643–2674.
- [5] J. Cummings and S. Shelah, *Cardinal invariants above the continuum*, Ann. Pure Appl. Logic 75 (1995), 251–268.
- [6] M. Džamonja and S. Shelah, *Universal graphs at the successor of a singular cardinal*, J. Symbolic Logic 68 (2003), 366–388.
- [7] D. H. Fremlin, *Measure Theory. Vol. 5. Set-Theoretic Measure Theory*, Torres Fremlin, Colchester, 2008.
- [8] S. Garti and S. Shelah, *Partition calculus and cardinal invariants*, J. Math. Soc. Japan 66 (2014), 425–434.
- [9] F. Hernández-Hernández and M. Hrušák, *Cardinal invariants of analytic  $P$ -ideals*, Canad. J. Math. 59 (2007), 575–595.
- [10] S. Kamo, *Splitting numbers on uncountable regular cardinals*, preprint.
- [11] C. Laflamme, *Zapping small filters*, Proc. Amer. Math. Soc. 114 (1992), 535–544.
- [12] S. Shelah, *On cardinal invariants of the continuum*, in: Axiomatic Set Theory (Boulder, CO, 1983), Contemp. Math. 31, Amer. Math. Soc., Providence, RI, 1984, 183–207.
- [13] S. Solecki and S. Todorcevic, *Avoiding families and Tukey functions on the nowhere-dense ideal*, J. Inst. Math. Jussieu 10 (2011), 405–435.
- [14] T. Suzuki, *About splitting numbers*, Proc. Japan Acad. Ser. A Math. Sci. 74 (1998), 33–35.
- [15] S. Todorcevic, *Analytic gaps*, Fund. Math. 150 (1996), 55–66.
- [16] J. Zapletal, *Splitting number at uncountable cardinals*, J. Symbolic Logic 62 (1997), 35–42.

Dilip Raghavan  
 Department of Mathematics  
 National University of Singapore  
 Singapore 119076  
 E-mail: raghavan@math.nus.edu.sg  
 URL: <http://www.math.nus.edu.sg/~raghavan/>

Saharon Shelah  
 Institute of Mathematics  
 The Hebrew University  
 Jerusalem 9190401, Israel  
 E-mail: shelah@math.huji.ac.il  
 URL: <http://shelah.logic.at/>