

Almost disjoint refinements and mixing reals

by

Barnabás Farkas (Vienna), Yurii Khomskii (Hamburg)
and Zoltán Vidnyánszky (Budapest)

Abstract. We investigate families of subsets of ω with almost disjoint refinements in the classical case as well as with respect to given ideals on ω .

We prove the following generalization of a result due to J. Brendle: If $V \subseteq W$ are transitive models, $\omega_1^W \subseteq V$, $\mathcal{P}(\omega) \cap V \neq \mathcal{P}(\omega) \cap W$, and \mathcal{J} is an analytic or coanalytic ideal coded in V , then there is an \mathcal{J} -almost disjoint refinement of $\mathcal{J}^+ \cap V$ in W .

We study the existence of perfect \mathcal{J} -almost disjoint families, and the existence of \mathcal{J} -almost disjoint refinements in which any two distinct sets have finite intersection.

We introduce the notion of mixing real (motivated by the construction of an almost disjoint refinement of $[\omega]^\omega \cap V$ after adding a Cohen real to V) and discuss logical implications between the existence of mixing reals in forcing extensions and classical properties of forcing notions.

1. Introduction. Let us begin with our motivations which led us to work on almost disjoint refinements and their generalizations. First of all, the following easy fact seems to be somewhat surprising (see also Proposition 1.9):

FACT 1.1. *If $\mathcal{H} \subseteq [\omega]^\omega$ ($= \{X \subseteq \omega : |X| = \omega\}$) is of size $< \mathfrak{c}$, then \mathcal{H} has an almost disjoint refinement $\{A_H : H \in \mathcal{H}\}$, that is, (i) $A_H \in [H]^\omega$ for every $H \in \mathcal{H}$, and (ii) $|A_H \cap A_K| < \omega$ for every $H \neq K$ from \mathcal{H} .*

The following theorem due to B. Balcar and P. Vojtáš is probably the most well-known general result on the existence of almost disjoint refinements.

THEOREM 1.2 (see [BaV80]). *Every ultrafilter on ω has an almost disjoint refinement.*

2010 *Mathematics Subject Classification:* 03E05, 03E15, 03E35.

Key words and phrases: analytic ideal, coanalytic ideal, almost disjoint family, almost disjoint refinement, Mansfield–Solovay theorem, mixing real, meager ideal.

Received 19 October 2015; revised 9 January 2017 and 19 July 2017.

Published online 30 March 2018.

B. Balcar and T. Pazák, and independently J. Brendle, proved the following theorem:

THEOREM 1.3 (see [BaP10], [Sk08]). *Assume that $V \subseteq W$ are transitive models and $\mathcal{P}(\omega) \cap V \neq \mathcal{P}(\omega) \cap W$. Then $[\omega]^\omega \cap V$ has an almost disjoint refinement in W (where by transitive model we mean a transitive model of a “large enough” finite fragment of ZFC).*

One of our main results is a generalization of this theorem in the context of “nice” ideals on ω , that is, we change the notion of *smallness* in the setting above by replacing *finite* with *element of an ideal \mathcal{J}* .

In order to formulate our generalization and to give a setting for other related results, we have to introduce some notation and appropriate versions of classical notions.

Let \mathcal{J} be an ideal on a countably infinite set X . We always assume that $[X]^{<\omega} = \{Y \subseteq X : |Y| < \omega\} \subseteq \mathcal{J}$ and $X \notin \mathcal{J}$. Denote by $\mathcal{J}^+ = \mathcal{P}(X) \setminus \mathcal{J}$ the family of \mathcal{J} -positive sets, and by $\mathcal{J}^* = \{X \setminus A : A \in \mathcal{J}\}$ the *dual filter* of \mathcal{J} . If $Y \in \mathcal{J}^+$ then let $\mathcal{J}|Y = \{A \in \mathcal{J} : A \subseteq Y\} = \{B \cap Y : B \in \mathcal{J}\}$ be the *restriction* of \mathcal{J} to Y (which is an ideal on Y). If X is clear from the context, then the ideal of finite subsets of X will be denoted by Fin .

DEFINITION 1.4. We say that a non-empty family $\mathcal{A} \subseteq \mathcal{I}^+$ is \mathcal{I} -almost disjoint (\mathcal{I} -AD) if $A \cap B \in \mathcal{I}$ for any distinct $A, B \in \mathcal{A}$. A family $\mathcal{A} \subseteq \mathcal{J}^+$ is $(\mathcal{J}, \text{Fin})$ -AD if $|A \cap B| < \omega$ for any distinct $A, B \in \mathcal{A}$.

DEFINITION 1.5. Let $\mathcal{H} \subseteq \mathcal{J}^+$. We say that a family $\mathcal{A} = \{A_H : H \in \mathcal{H}\}$ is an \mathcal{J} -AD refinement (\mathcal{J} -ADR) of \mathcal{H} if (i) $A_H \subseteq H$, $A_H \in \mathcal{J}^+$ for every H , and (ii) $A_{H_0} \cap A_{H_1} \in \mathcal{J}$ for any distinct $H_0, H_1 \in \mathcal{H}$ (in particular, \mathcal{A} is an \mathcal{J} -AD family). If $\mathcal{J} = \text{Fin}$ we simply say *AD refinement* (ADR).

We say that a family $\mathcal{A} = \{A_H : H \in \mathcal{H}\}$ is an $(\mathcal{J}, \text{Fin})$ -AD refinement $((\mathcal{J}, \text{Fin})$ -ADR) of \mathcal{H} if (i) holds and (ii)' $|A_{H_0} \cap A_{H_1}| < \omega$ for any distinct $H_0, H_1 \in \mathcal{H}$.

Notice that an ideal on a countably infinite X can be regarded as a subset of the Polish space $2^X \simeq 2^\omega$ by using a bijection between X and ω . Thus, it makes sense to talk about Borel, analytic, etc. ideals and about certain descriptive properties of ideals, such as the Baire property or meagerness (it is easy to see that these properties do not depend on the choice of the bijection). In the past three decades the study of certain definable (e.g. Borel, analytic, coanalytic, etc.) ideals has become a central topic in set theory. It turned out that they play an important role in combinatorial set theory, and in the theory of cardinal invariants of the continuum as well as the theory of forcing (see e.g. [Ma91], [So99], [F00], [Hr11] and many other publications).

Now we can formulate our generalization of Theorem 1.3:

THEOREM 1.6. *Assume that $V \subseteq W$ are transitive models, $\omega_1^W \subseteq V$, $\mathcal{P}(\omega) \cap V \neq \mathcal{P}(\omega) \cap W$, and \mathcal{J} is an analytic or coanalytic ideal coded in V . Then there is an \mathcal{J} -ADR of $\mathcal{J}^+ \cap V$ in W .*

We say that an ideal \mathcal{J} on X (where $|X| = \omega$) is *everywhere meager* if $\mathcal{J} \upharpoonright Y$ is meager in $\mathcal{P}(Y)$ for every $Y \in \mathcal{J}^+$. In particular, analytic and coanalytic ideals are everywhere meager because their restrictions are also analytic and coanalytic, respectively, hence have the Baire property, and we can apply the following well-known characterization theorem (due to Sierpiński (1) \leftrightarrow (2), and Talagrand (2) \leftrightarrow (3); for the proofs see e.g. [BrJ95, Thms. 4.1.1–4.1.2]).

THEOREM 1.7. *Let \mathcal{J} be an ideal on ω . Then the following are equivalent: (1) \mathcal{J} has the Baire property, (2) \mathcal{J} is meager, and (3) there is a partition $\{P_n : n \in \omega\}$ of ω into finite sets such that $\{n \in \omega : P_n \subseteq A\}$ is finite for each $A \in \mathcal{J}$.*

From now on, when working with partitions of a set, we always assume that every element of the partition is non-empty. From this theorem we can also deduce the following corollary:

COROLLARY 1.8. *If \mathcal{J} is a meager ideal, then there is a perfect $(\mathcal{J}, \text{Fin})$ -AD family. In particular, if \mathcal{J} is everywhere meager, then there are perfect $(\mathcal{J}, \text{Fin})$ -AD families on every $X \in \mathcal{J}^+$.*

Proof. It is easy to define a perfect AD family \mathcal{A} on ω (e.g. consider the branches of $2^{<\omega}$ in $\mathcal{P}(2^{<\omega})$). Fix a partition $(P_n)_{n \in \omega}$ of ω into finite sets such that $\{n \in \omega : P_n \subseteq A\}$ is finite for every $A \in \mathcal{J}$. For each $A \in \mathcal{A}$ let $A' = \bigcup\{P_n : n \in A\} \in \mathcal{J}^+$, and let $\mathcal{A}' = \{A' : A \in \mathcal{A}\}$. Then $|A' \cap B'| < \omega$ for any distinct $A, B \in \mathcal{A}$, hence \mathcal{A}' is an $(\mathcal{J}, \text{Fin})$ -AD family. The function $\mathcal{P}(\omega) \rightarrow \mathcal{P}(\omega)$, $A \mapsto A'$ is injective and continuous, hence \mathcal{A}' is perfect. ■

Concerning the reverse implications in Corollary 1.8, in Section 5 we prove the following:

- (a) The existence of a perfect $(\mathcal{J}, \text{Fin})$ -AD family does not imply that \mathcal{J} is meager (Example 5.1). Moreover, if $\mathfrak{b} = \mathfrak{c}$ then there is a non-meager ideal \mathcal{J} such that there are perfect $(\mathcal{J}, \text{Fin})$ -AD families on every $X \in \mathcal{J}^+$ (Theorem 5.2). Here \mathfrak{c} stands for the continuum and \mathfrak{b} for the *bounding number*, that is, $\mathfrak{b} = \min\{|F| : F \subseteq \omega^\omega \text{ is } \leq^* \text{-unbounded}\}$ where $f \leq^* g$ iff the set $\{n \in \omega : f(n) > g(n)\}$ is finite.
- (b) There is an ideal \mathcal{J} such that every \mathcal{J} -AD family is countable but \mathcal{J} is *nowhere maximal*, that is, $\mathcal{J} \upharpoonright X$ is not a prime ideal for any $X \in \mathcal{J}^+$ (in particular, there are infinite \mathcal{J} -AD families); and it is independent from ZFC whether such an ideal can be chosen as Σ_2^1 (Proposition 5.3).

Corollary 1.8 has an easy but important application. Clearly, if \mathcal{J} is an ideal on ω then there is a family (e.g. \mathcal{J}^+) of size \mathfrak{c} which does not have any

J-ADR's. Conversely, we have the following very special case of results from [BHM75] and [BaSV81]:

PROPOSITION 1.9. *If \mathcal{J} is an everywhere meager ideal and $\mathcal{H} \in [\mathcal{J}^+]^{<\mathfrak{c}}$, then \mathcal{H} has an J-ADR.*

Proof. Let $\mathcal{H} = \{H_\alpha : \alpha < \kappa\}$. Applying Corollary 1.8, we can fix an J-AD family $\mathcal{A} = \{A_\xi : \xi < \kappa^+\}$ on H_0 and for every $\beta < \kappa$ let $T_\beta = \{\xi < \kappa^+ : H_\beta \cap A_\xi \in \mathcal{J}^+\}$; furthermore let $R = \{\beta < \kappa : |T_\beta| \leq \kappa\}$ (we know that $0 \notin R$). By induction on $\alpha \in \kappa \setminus R$ we can pick a

$$\xi_\alpha \in T_\alpha \setminus \left(\bigcup_{\beta \in R} T_\beta \cup \{\xi_{\alpha'} : \alpha' \in \alpha \setminus R\} \right)$$

because $|T_\alpha| = \kappa^+$ and $|\bigcup\{T_\beta : \beta \in R\}| \leq \kappa$, and let $E_\alpha = H_\alpha \cap A_{\xi_\alpha} \in \mathcal{J}^+$. Then the family $\{E_\alpha : \alpha \in \kappa \setminus R\}$ is an J-ADR of $\{H_\alpha : \alpha \in \kappa \setminus R\}$. We can continue the procedure on $\{H_\beta : \beta \in R\}$ because $E_\alpha \cap H_\beta \in \mathcal{J}$ for every $\alpha \in \kappa \setminus R$ and $\beta \in R$. ■

This proposition motivates the following:

QUESTION 1.10. *Let \mathcal{J} be an everywhere meager ideal and $\mathcal{H} \in [\mathcal{J}^+]^{<\mathfrak{c}}$. Does \mathcal{H} have an (J, Fin)-ADR?*

In Section 6, we answer this question (at least consistently): Assume MA_κ and let \mathcal{J} be an everywhere meager ideal; then every $\mathcal{H} \in [\mathcal{J}^+]^{\leq \kappa}$ has an (J, Fin)-ADR (Theorem 6.1).

We also define new notions of mixing and injective mixing reals, and investigate connections between adding (injective) mixing reals and classical properties of forcing notions (such as adding Cohen/random/splitting/dominating reals and the Laver/Sacks properties).

DEFINITION 1.11. Let \mathbb{P} be a forcing notion. We say that an $f \in \omega^\omega \cap V^{\mathbb{P}}$ is a *mixing real* over V if $|f[X] \cap Y| = \omega$ for any $X, Y \in [\omega]^\omega \cap V$. If f is one-to-one, then we call it an *injective mixing real* or *mixing injection*.

Mixing reals can be seen as “infinite splitting partitions” (see Proposition 7.1): There is a mixing real in $V^{\mathbb{P}}$ (over V) iff there is a partition $(Y_n)_{n \in \omega} \in V^{\mathbb{P}}$ of ω into infinite sets such that $\forall X \in [\omega]^\omega \cap V \forall n |X \cap Y_n| = \omega$. Recall that a set $S \subseteq \omega$ is a *splitting real* over V if $|X \cap S| = |X \setminus S| = \omega$ for every $X \in [\omega]^\omega \cap V$; in other words, $P = \{S, \omega \setminus S\}$ is a partition of ω such that $\forall X \in [\omega]^\omega \cap V \forall Y \in P |X \cap Y| = \omega$. Therefore, mixing reals are strong variants of splitting reals.

Why are mixing reals relevant to almost disjoint refinements? Fix an AD family $\mathcal{A} = \{A_\alpha : \alpha < \mathfrak{c}\}$ in V , and let $\{X_\alpha : \alpha < \mathfrak{c}\}$ be an enumeration of $[\omega]^\omega$ in V . If $f \in \omega^\omega \cap V^{\mathbb{P}}$ is a mixing injection over V , then the family $\{f[A_\alpha] \cap X_\alpha : \alpha < \mathfrak{c}\} \in V^{\mathbb{P}}$ is an ADR of $[\omega]^\omega \cap V$.

In Proposition 7.2 we prove the following. Let \mathbb{P} be a forcing notion.

- (i) If \mathbb{P} adds random reals, then it adds mixing reals.
- (ii) If \mathbb{P} adds dominating reals, then it adds mixing reals.
- (iii) If \mathbb{P} adds Cohen reals, then it adds mixing injections.
- (iv) If \mathbb{P} adds mixing injections, then it adds unbounded reals.
- (v) If \mathbb{P} has the Laver property, then it does not add injective mixing reals.

Our paper is organized as follows. In Section 2 we recall some notation and classical results of descriptive set theory we will need later.

The next two sections are focused on descriptive aspects of nice ideals and almost disjoint refinements. In Section 3 we present a plethora of examples of Borel and projective ideals on ω . In Section 4 we prove Theorem 1.6 by modifying Brendle's proof of Theorem 1.3.

The next two sections contain combinatorial results. In Section 5 we study the reverse implications in Corollary 1.8. In Section 6 we answer (consistently) Question 1.10.

In Section 7 we study the notions of mixing and injective mixing reals. In this section we will heavily use standard facts about forcing notions; for the details see [BrJ95].

Finally, in Section 8, we list some open questions concerning our results.

2. Descriptive set theory and ideals. As usual, $\Sigma_\alpha^0, \Pi_\alpha^0$ will stand for the α th level of the Borel hierarchy while we denote by Σ_n^1, Π_n^1 the levels of the projective hierarchy. If r is a real, the appropriate relativized versions are denoted by $\Sigma_\alpha^0(r), \Pi_\alpha^0(r)$, etc. For the ambiguous classes we write Δ_α^i and $\Delta_\alpha^i(r)$.

Suppose that \mathcal{J} is an ideal on a set X . As mentioned before, if X is countable, then we can talk about complexity of ideals: \mathcal{J} is $F_\sigma, \Sigma_\alpha^0, \Pi_n^1$, etc. if $\mathcal{J} \subseteq \mathcal{P}(X) \simeq 2^X$ is an $F_\sigma, \Sigma_\alpha^0, \Pi_n^1$, etc. set in the usual compact Polish topology on 2^X . If we fix a bijection between ω and X , we can define the collection of $\Sigma_\alpha^0(r), \Pi_\alpha^0(r)$, etc. subsets of 2^X as well. If $X = \omega^n$ and $\Delta = \{(n, m) \in \omega^2 : m \leq n\}, [\omega]^n, 2^{<\omega}, \omega^{<\omega}, \mathbb{Q}$ ($= \{\text{rational numbers}\}$) then we will always assume that the bijection is the usual, recursive one.

For example, $\text{Fin} = [\omega]^{<\omega}$ is an F_σ ideal, $\mathcal{Z} = \{A \subseteq \omega : |A \cap n|/n \rightarrow 0\}$ is $F_{\sigma\delta}$, and $\text{Conv} = \{A \subseteq \mathbb{Q} \cap [0, 1] : A \text{ has only finitely many accumulation points}\}$ is $F_{\sigma\delta\sigma}$, etc. (see more examples in Section 3). Similarly, we can associate descriptive complexity to any $\mathcal{X} \subseteq \mathcal{P}(\omega)$, and we can also talk about the Baire property and measurability of subsets of $\mathcal{P}(\omega)$. Clearly, if $Y \in \mathcal{J}^+$ then $\mathcal{J} \upharpoonright Y$ belongs to the same Borel or projective class in $\mathcal{P}(Y)$ as \mathcal{J} in $\mathcal{P}(\omega)$ (simply because $\mathcal{J} \upharpoonright Y$ is a continuous preimage of \mathcal{J}).

For a family $\mathcal{H} \subseteq \mathcal{P}(X)$ we will denote by $\text{id}(\mathcal{H})$ the ideal generated by the sets in \mathcal{H} . We say that an ideal \mathcal{J} on a countably infinite set X is

- *tall* if every infinite subset of X contains an infinite element of \mathcal{J} ;
- a *P-ideal* if for every sequence $A_n \in \mathcal{J}$ ($n \in \omega$), there is an $A \in \mathcal{J}$ such that $A_n \subseteq^* A$, that is, $|A_n \setminus A| < \omega$ for every n .

We will need the following two fundamental results of descriptive set theory (see e.g. [J03]):

THEOREM 2.1 (Shoenfield Absoluteness Theorem). *If $V \subseteq W$ are transitive models, $\omega_1^W \subseteq V$, and $r \in \omega^\omega \cap V$, then $\Sigma_2^1(r)$ formulas are absolute between V and W .*

COROLLARY 2.2. *If $X \subseteq \mathcal{P}(\omega)$ is an analytic or coanalytic set in the parameter $r \in \omega^\omega$, then the statement “ X is an ideal” is absolute for transitive models $V \subseteq W$ with $\omega_1^W \subseteq V$ and $r \in V$.*

Proof. Let $\varphi(x, r)$ be a $\Sigma_1^1(r)$ or $\Pi_1^1(r)$ definition of X ($r \in \omega^\omega$). Then the statement “ X is an ideal” is the conjunction of the following formulas: (i) $\forall a \in \text{Fin } \varphi(a, r)$, (ii) $\forall x, y$ ($x \not\subseteq y$ or $\neg\varphi(y, r)$ or $\varphi(x, r)$), and (iii) $\forall x, y$ ($\neg\varphi(x, r)$ or $\neg\varphi(y, r)$ or $\varphi(x \cup y, r)$). In particular, “ X is an ideal” is $\Pi_2^1(r)$ and hence we can apply the Shoenfield Absoluteness Theorem. ■

THEOREM 2.3 (Mansfield–Solovay Theorem). *If $A \not\subseteq L[r]$ is a $\Sigma_2^1(r)$ set, then A contains a perfect subset.*

Other than the notions and results above, we will use descriptive-set-theoretic tools such as Γ -completeness, Γ -hardness, etc. which can all be found in [K95].

Let $\text{Tree} = \{T \subseteq \omega^{<\omega} : T \text{ is a tree}\}$ be the usual Polish space of all trees on ω (a closed subset of $\mathcal{P}(\omega^{<\omega})$) and as usual, denote by $[T] = \{x \in \omega^\omega : \forall n \ x \upharpoonright n \in T\}$ the *body* of T , i.e. the set of all branches of T .

3. Examples of Borel and projective ideals. There are many classical examples of Borel ideals. Here we present some of those that have easily understandable definitions, and the reader can see that these examples are motivated by a wide variety of backgrounds. For the important role of these ideals, especially in characterization results, see [Hr11].

Some F_σ ideals

Summable ideals. Let $h : \omega \rightarrow [0, \infty)$ be a function such that $\sum_{n \in \omega} h(n) = \infty$. The *summable ideal associated to h* is

$$\mathcal{I}_h = \left\{ A \subseteq \omega : \sum_{n \in A} h(n) < \infty \right\}.$$

It is easy to see that a summable ideal \mathcal{I}_h is tall iff $\lim_{n \rightarrow \infty} h(n) = 0$, and that summable ideals are F_σ P-ideals. The *classical summable ideal* is $\mathcal{J}_{1/n} = \mathcal{J}_h$ where $h(n) = 1/(n+1)$, or $h(0) = 1$ and $h(n) = 1/n$ if $n > 0$. We know that

there are tall F_σ P-ideals which are not summable ideals: Farah's example [F00, Example 1.11.1] is the following:

$$\mathcal{J}_F = \left\{ A \subseteq \omega : \sum_{n < \omega} \frac{\min \{n, |A \cap [2^n, 2^{n+1})|\}}{n^2} < \infty \right\}.$$

The *eventually different ideals* are

$$\mathcal{ED} = \left\{ A \subseteq \omega \times \omega : \limsup_{n \rightarrow \infty} |(A)_n| < \infty \right\}$$

where $(A)_n = \{k \in \omega : (n, k) \in A\}$, and $\mathcal{ED}_{\text{fin}} = \mathcal{ED} \upharpoonright \Delta$ where $\Delta = \{(n, m) \in \omega \times \omega : m \leq n\}$. Note that \mathcal{ED} and $\mathcal{ED}_{\text{fin}}$ are not P-ideals.

The *van der Waerden ideal* is

$$\mathcal{W} = \{A \subseteq \omega : A \text{ does not contain arbitrarily long arithmetic progressions}\}.$$

Van der Waerden's well-known theorem says that \mathcal{W} is a proper ideal. Again, \mathcal{W} is not a P-ideal. For some set-theoretic results about this ideal see e.g. [F109] and [F110].

The *random graph ideal* is

$$\text{Ran} = \text{id}(\{\text{homogeneous subsets of the random graph}\})$$

where the *random graph* (ω, E) , $E \subseteq [\omega]^2$, is up to isomorphism uniquely determined by the following property: If $A, B \in [\omega]^{<\omega}$ are non-empty and disjoint, then there is an $n \in \omega \setminus (A \cup B)$ such that $\{\{n, a\} : a \in A\} \subseteq E$ and $\{\{n, b\} : b \in B\} \cap E = \emptyset$. A set $H \subseteq \omega$ is *(E-)homogeneous* if $[H]^2 \subseteq E$ or $[H]^2 \cap E = \emptyset$. The ideal Ran is not a P-ideal.

The *ideal of graphs with finite chromatic number* is

$$\mathcal{G}_{\text{fc}} = \{E \subseteq [\omega]^2 : \chi(\omega, E) < \omega\}.$$

It is not a P-ideal.

Solecki's ideal. Let $\text{CO}(2^\omega)$ be the family of clopen subsets of 2^ω (it is easy to see that $|\text{CO}(2^\omega)| = \omega$), and let $\Omega = \{A \in \text{CO}(2^\omega) : \lambda(A) = 1/2\}$ where λ is the usual product measure on 2^ω . The ideal \mathcal{S} on Ω is generated by $\{I_x : x \in 2^\omega\}$ where $I_x = \{A \in \Omega : x \in A\}$. It is not a P-ideal.

Some $F_{\sigma\delta}$ ideals

Density ideals. Let $(P_n)_{n \in \omega}$ be a sequence of pairwise disjoint finite subsets of ω and let $\vec{\mu} = (\mu_n)_{n \in \omega}$ be a sequence of measures concentrated on P_n such that $\limsup_{n \rightarrow \infty} \mu_n(\omega) > 0$. The *density ideal generated by $\vec{\mu}$* is

$$\mathcal{Z}_{\vec{\mu}} = \left\{ A \subseteq \omega : \lim_{n \rightarrow \infty} \mu_n(A) = 0 \right\}.$$

A density ideal $\mathcal{Z}_{\vec{\mu}}$ is tall iff $\max\{\mu_n(\{i\}) : i \in P_n\} \xrightarrow{n \rightarrow \infty} 0$, and density ideals are $F_{\sigma\delta}$ P-ideals. The *density zero ideal*

$$\mathcal{Z} = \left\{ A \subseteq \omega : \lim_{n \rightarrow \infty} |A \cap n|/n = 0 \right\}$$

is a tall density ideal. Indeed, let $P_n = [2^n, 2^{n+1})$ and $\mu_n(A) = |A \cap P_n|/2^n$. Then it is easy to see that $J_{1/n} \subsetneq \mathcal{Z}$, and Szemerédi's famous theorem implies that $\mathcal{W} \subseteq \mathcal{Z}$ (see [Sz75]). The stronger statement $\mathcal{W} \subseteq J_{1/n}$ is a still open Erdős prize problem.

The ideal of nowhere dense subsets of the rationals is

$$\text{Nwd} = \{A \subseteq \mathbb{Q} : \text{int}(\overline{A}) = \emptyset\}$$

where $\text{int}(\cdot)$ stands for the interior operation on subsets of the reals, and \overline{A} is the closure of A in \mathbb{R} . Nwd is not a P-ideal.

The trace ideal of the null ideal. Let \mathcal{N} be the σ -ideal of subsets of 2^ω with measure zero (with respect to the usual product measure). The G_δ -closure of a set $A \subseteq 2^{<\omega}$ is $[A]_\delta = \{x \in 2^\omega : \exists^\infty n \ x \upharpoonright n \in A\}$, a G_δ subset of 2^ω . The trace of \mathcal{N} is defined by

$$\text{tr}(\mathcal{N}) = \{A \subseteq 2^{<\omega} : [A]_\delta \in \mathcal{N}\}.$$

It is a tall $F_{\sigma\delta}$ P-ideal.

Some tall $F_{\sigma\delta\sigma}$ (non-P-)ideals. The ideal Conv is generated by those infinite subsets of $\mathbb{Q} \cap [0, 1]$ which are convergent in $[0, 1]$, in other words

$$\text{Conv} = \{A \subseteq \mathbb{Q} \cap [0, 1] : |\text{accumulation points of } A \text{ (in } \mathbb{R})| < \omega\}.$$

Another example is the Fubini product of Fin by itself:

$$\text{Fin} \otimes \text{Fin} = \{A \subseteq \omega \times \omega : \forall^\infty n \in \omega \ |(A)_n| < \omega\}.$$

Some non-tall ideals. Here we mention an important F_σ ideal:

$$\text{Fin} \otimes \{\emptyset\} = \{A \subseteq \omega \times \omega : \forall^\infty n \in \omega \ (A)_n = \emptyset\},$$

and its $F_{\sigma\delta}$ brother (a density ideal)

$$\{\emptyset\} \otimes \text{Fin} = \{A \subseteq \omega \times \omega : \forall n \in \omega \ |(A)_n| < \omega\}.$$

Applying the Baire Category Theorem, it is easy to see that there are no G_δ (i.e. $\underline{\Pi}_2^0$) ideals and we already presented many F_σ (i.e. $\underline{\Sigma}_2^0$) ideals. In general, we have Borel ideals at arbitrarily high levels of the Borel hierarchy:

THEOREM 3.1 (see [C85] and [C88]). *There are $\underline{\Sigma}_\alpha^0$ - and $\underline{\Pi}_\alpha^0$ -complete ideals for every $\alpha \geq 3$.*

For ideals on the ambiguous levels of the Borel hierarchy see [E94].

We also present some (co)analytic examples.

THEOREM 3.2 (see [Z90, p. 321]). *For every $x \in \omega^\omega$ let $I_x = \{s \in \omega^{<\omega} : x \upharpoonright |s| \not\leq s\}$ where \leq is the coordinatewise ordering on every ω^n . Then the ideal on $\omega^{<\omega}$ generated by $\{I_x : x \in \omega^\omega\}$ is $\underline{\Sigma}_1^1$ -complete.*

THEOREM 3.3. *The ideal of graphs without infinite complete subgraphs,*

$$\mathcal{G}_c = \{E \subseteq [\omega]^2 : \forall X \in [\omega]^\omega \ [X]^2 \not\subseteq E\},$$

is a $\underline{\Pi}_1^1$ -complete (in $\mathcal{P}([\omega]^2)$), tall, non-P-ideal.

Proof. Tallness is trivial. If we define $E_n = \{\{k, m\} : k \leq n, m \neq k\} \in \mathcal{G}_c$ and $E_n \subseteq^* E \subseteq [\omega]^2$ for every n , then E contains a complete subgraph (see also [Me09]), hence \mathcal{G}_c is not a P-ideal.

Let $\text{WF} = \{T \in \text{Tree} : [T] = \emptyset\}$ be the \mathbb{Q}_1^1 -complete set of well-founded trees. Furthermore, let Tree' be the family of those trees T such that (i) every $t \in T$ is strictly increasing, and (ii) if $\{t \in T : n \in \text{ran}(t)\} \neq \emptyset$ then it has a \subseteq -minimal element ($n \in \omega$). Then it is not hard to see that Tree' is also closed in $\mathcal{P}(\omega^{<\omega})$, hence Polish. Finally, let $\text{WF}' = \{T \in \text{Tree}' : [T] = \emptyset\}$; clearly, it is also \mathbb{Q}_1^1 .

We will construct Wadge reductions $\text{WF} \leq_W \text{WF}' \leq_W \mathcal{G}_c$.

$\text{WF} \leq_W \text{WF}'$: Fix an order preserving isomorphism j between $\omega^{<\omega}$ and a $T_0 \in \text{Tree}'$. More precisely, for a $t = (k_0, k_1, \dots, k_{m-1}) \in \omega^{<\omega}$ let $j(t) = (p_{k_0}^1, p_{k_0}^1 p_{k_1}^2, \dots, p_{k_0}^1 p_{k_1}^2 \dots p_{k_{m-1}}^m)$ where p_i denotes the i th prime number. Then j is one-to-one, order preserving, and $T_0 = j[\omega^{<\omega}]$ is a tree containing strictly increasing sequences. To show that T_0 satisfies (ii), assume that $n \in \text{ran}(j(t))$ for some $n \in \omega$ and $t \in \omega^{<\omega}$. Then, by the definition of j , $n = p_{k_0}^1 p_{k_1}^2 \dots p_{k_{m-1}}^m$ where $s = (k_0, k_1, \dots, k_{m-1}) \leq t$, and if $n \in \text{ran}(j(t'))$ for some $t' \in \omega^{<\omega}$ then $s \leq t'$, hence $j(s)$ is \subseteq -minimal in $\{h \in T_0 : n \in \text{ran}(h)\}$.

The map $\text{Tree} \rightarrow \text{Tree}'$, $T \mapsto j[T]$ is a continuous reduction of WF to WF' . Continuity is trivial, as is the fact that $[T] = \emptyset$ iff $[j[T]] \neq \emptyset$, in other words, $T \in \text{WF}$ iff $j[T] \in \text{WF}'$.

$\text{WF}' \leq_W \mathcal{G}_c$: For every $T \in \text{Tree}'$ let $E_T = \bigcup\{[\text{ran}(t)]^2 : t \in T\}$. We show that the function $T \mapsto E_T$ is continuous. If $u, v \in [[\omega]^2]^{<\omega}$ are disjoint then it is easy to see that the preimage of the basic clopen set $[u, v] = \{E \subseteq [\omega]^2 : u \subseteq E, v \cap E = \emptyset\} \subseteq \mathcal{P}([\omega]^2)$ is

$$\{T \in \text{Tree}' : (\forall \{x, y\} \in u \exists t \in T \ x, y \in \text{ran}(t)) \\ \text{and } (\forall t \in T \ v \cap [\text{ran}(t)]^2 = \emptyset)\}.$$

Although, as the collection of the sets satisfying the second part of the condition is a countable intersection of clopen sets, this set seems to be closed (and it is enough to prove that \mathcal{G}_c is \mathbb{Q}_1^1 -complete), actually, it is open in Tree' : Let $m = \max(\cup v) + 1$. Then the set $\{T \in \text{Tree}' : \forall t \in T \ v \cap [\text{ran}(t)]^2 = \emptyset\}$ is the intersection of Tree' and the clopen (in $\mathcal{P}(\omega^{<\omega})$) set

$$[\emptyset, \{t \in m^{<m} : t \text{ is strictly increasing and } v \cap [\text{ran}(t)]^2 \neq \emptyset\}].$$

The function $T \mapsto E_T$ is a reduction of WF' to \mathcal{G}_c : Clearly, if $T \in \text{Tree}'$ and $x \in [T]$ then $X = \text{ran}(x) \in [\omega]^\omega$ shows that $E_T \notin \mathcal{G}_c$ (i.e. $[X]^2 \subseteq E$). Conversely, if $[X]^2 \subseteq E_T$ and $X = \{k_0 < k_1 < \dots\}$, then for every n there is a $t_n \in T$ such that $k_n, k_{n+1} \in \text{ran}(t_n)$; we can assume that t_n is minimal in $\{s \in T : k_{n+1} \in \text{ran}(s)\}$. This implies that $t_0 \subseteq t_1 \subseteq t_2 \subseteq \dots$ is an infinite chain in T . ■

In the following example, we show that a seemingly “very” Π_2^1 definition can also give us a $\underline{\Pi}_1^1$ -complete ideal.

THEOREM 3.4. *The ideal*

$\mathcal{J}_0 = \{A \subseteq \omega \times \omega : \forall X, Y \in [\omega]^\omega \exists X' \in [X]^\omega \exists Y' \in [Y]^\omega A \cap (X' \times Y') = \emptyset\}$
is a $\underline{\Pi}_1^1$ -complete (in $\mathcal{P}(\omega \times \omega)$), tall, non-P-ideal.

Proof. Tallness is trivial because injective partial functions from ω to ω belong to \mathcal{J}_0 . The failure of the P property is also easy: Consider the sets $n \times \omega \in \mathcal{J}$. If for some A we have $n \times \omega \subseteq^* A$ for every n then every vertical section of A is cofinite, and such a set is clearly \mathcal{J}_0 -positive.

Now we show that \mathcal{J}_0 is $\underline{\Pi}_1^1$, for which the next Claim is clearly enough. For $X, Y \in [\omega]^\omega$ define $T^\uparrow(X, Y) = \{(n, k) \in X \times Y : n < k\}$ and $T^\downarrow(X, Y) = \{(n, k) \in X \times Y : n > k\}$.

CLAIM. *$A \in \mathcal{J}_0$ iff for every infinite X and Y the set A does not contain $T^\uparrow(X, Y)$ or $T^\downarrow(X, Y)$.*

Proof of Claim. The “only if” part is trivial. Conversely, assume that $A \notin \mathcal{J}_0$, i.e. there are $X, Y \in [\omega]^\omega$ such that $A \cap (X' \times Y') \neq \emptyset$ for all $X' \in [X]^\omega$ and $Y' \in [Y]^\omega$. Fix increasing enumerations $X = \{x_0 < x_1 < \dots\}$ and $Y = \{y_0 < y_1 < \dots\}$. By shrinking the sets X and Y , we can assume that $x_0 < y_0 < x_1 < y_1 < \dots$, in particular $X \cap Y = \emptyset$. Consider the following coloring $c : [\omega]^2 \rightarrow 2 \times 2$: for $m < n$ let $c(m, n) = (\chi_A(x_m, y_n), \chi_A(x_n, y_m))$ where $\chi_A(x, y) = 1$ iff $(x, y) \in A$.

By Ramsey’s theorem, there is an infinite homogeneous subset $S \subseteq \omega$. Let $S = Z \cup W$ be a partition into infinite subsets such that the elements of Z and W alternate in S . Then the elements of the sets $X' = \{x_m : m \in Z\}$ and $Y' = \{y_n : n \in W\}$ alternate in ω as well.

Note that S cannot be homogeneous in color $(0, 0)$, since otherwise $A \cap (X' \times Y') = \emptyset$ would hold. Similarly, if S is homogeneous in color $(1, 1)$ then $X' \times Y' \subset A$ and we are done. Now suppose that S is homogeneous in color $(1, 0)$ (for $(0, 1)$ the same argument works). If $x_m \in X'$, $y_n \in Y'$ and $x_m < y_n$ then $m < n$ because $Z \cap W = \emptyset$. Hence by the homogeneity of S we can conclude $(x_m, y_n) \in A$, so $T^\uparrow(X', Y') \subseteq A$. ■

Now we show that \mathcal{I}_0 is $\underline{\Pi}_1^1$ -complete. We will use the fact [K95, 27.B] that the set

$$S = \{C \in \mathcal{K}(2^\omega) : \forall x \in C \forall^\infty n \in \omega x(n) = 0\}$$

is $\underline{\Pi}_1^1$ -complete, where $\mathcal{K}(2^\omega)$ stands for the family of compact subsets of 2^ω equipped with the Hausdorff metric, i.e. with the Vietoris topology, we know that $\mathcal{K}(2^\omega)$ is a compact Polish space.

To finish the proof, we will define a Borel map $\mathcal{K}(2^\omega) \rightarrow \mathcal{P}(\omega \times \omega)$, $C \mapsto A_C$, such that $C \in S$ iff $A_C \in \mathcal{I}_0$. Fix an enumeration $\{s_m : m \in \omega\}$

of $2^{<\omega}$, for every $s \in 2^{<\omega}$ define $[s] = \{x \in 2^\omega : s \subseteq x\}$ (a basic clopen subset of 2^ω), and let

$$A_C = \{(m, n) : |s_m| > n, s_m(n) = 1, \text{ and } [s_m] \cap C \neq \emptyset\}.$$

For $C \in S$ we show that $A_C \in \mathcal{J}_0$. Let $X, Y \in [\omega]^\omega$ be arbitrary. If the set $\{m \in X : [s_m] \cap C = \emptyset\}$ is infinite then we are done, since

$$A_C \cap (\{m \in X : [s_m] \cap C = \emptyset\} \times Y) = \emptyset.$$

Otherwise, using the compactness of C we can choose $\{m_0 < m_1 < \dots\} = X' \in [X]^\omega$ and a convergent sequence $(x_i)_{i \in \omega}$ such that $x_i \in [s_{m_i}] \cap C$ for every i . If $x_i \rightarrow x$ then $x \in C \in S$ so $x(n) = 0$ for every $n \geq n_0$ for some n_0 . If $n \in Y \setminus n_0$ then for every large enough i we have $n < |s_{m_i}|$ and $s_{m_i}(n) = x(n) = 0$, hence the section $\{m : (m, n) \in A_C \cap (X' \times Y)\}$ is finite. On the other hand, for a fixed m , if $|s_m| \leq n$ then $(m, n) \notin A_C$, therefore the section $\{n : (m, n) \in A_C \cap (X' \times Y)\}$ is also finite. By an easy induction, one can define an $X'' \in [X']^\omega$ and a $Y'' \in [Y]^\omega$ such that $A_C \cap (X'' \times Y'') = \emptyset$.

Now we show that if $C \notin S$ then $A_C \notin \mathcal{I}_0$. Let $x \in C$ be such that $Y = \{n : x(n) = 1\}$ is infinite and let $X = \{m : x \in [s_m]\}$. Now clearly, if $(m, n) \in X \times Y$ then $(m, n) \in A_C$ if and only if $n < |s_m|$. In particular, for every $n \in Y$ the set $\{m \in X : (m, n) \notin A_C\}$ is finite, and this clearly implies that the rectangle $X \times Y$ witnesses that $A_C \notin \mathcal{J}_0$. ■

REMARK 3.5. One can give an alternative proof of Theorem 3.3 by constructing a Borel reduction of the set C to \mathcal{G}_c .

THEOREM 3.6. *There exist Σ_n^1 - and Π_n^1 -complete tall ideals for every $n \geq 1$.*

Proof. First we will construct Σ_n^1 -complete ideals. Let \mathcal{J} be a tall Borel ideal, \mathcal{A} be a perfect \mathcal{J} -AD family, and let \mathcal{A}_n be a Σ_n^1 -complete subset of the Polish space \mathcal{A} . Define $\mathcal{J}_n = \text{id}(\mathcal{J} \cup \mathcal{A}_n)$, i.e. \mathcal{J}_n is the ideal generated by $\mathcal{J} \cup \mathcal{A}_n$. Then \mathcal{J}_n is a tall proper (because \mathcal{A}_n is infinite) ideal. Moreover, \mathcal{J}_n is Σ_n^1 because

$$\mathcal{J}_n = \{X \subseteq \omega : \exists k \in \omega \exists (A_i)_{i < k} \in \mathcal{A}_n^k \ X \setminus (A_0 \cup A_1 \cup \dots \cup A_{k-1}) \in \mathcal{J}\}.$$

In order to see that \mathcal{J}_n is Σ_n^1 -complete, we remark that if B is a Σ_n^1 set in a Polish space \mathcal{X} , then it can be reduced to \mathcal{A}_n with a continuous map $f : \mathcal{X} \rightarrow \mathcal{A} \subseteq \mathcal{P}(\omega)$; furthermore, applying the trivial observation that $\mathcal{A}_n = \mathcal{J}_n \cap \mathcal{A}$, we conclude that this map is in fact a reduction of B to \mathcal{J}_n as well.

Now we proceed to Π_n^1 ideals. We can assume that $n > 1$ (see the last two theorems above). Again, there exists a Π_n^1 -complete set $\mathcal{B}_n \subseteq \mathcal{A}$. The previous argument shows that the ideal $\mathcal{J}'_n = \text{id}(\mathcal{J} \cup \mathcal{B}_n)$ is Π_n^1 -hard, so it is enough to prove that \mathcal{J}'_n is Π_n^1 . To see this, just notice that since \mathcal{A} is an

\mathcal{J} -AD-family, if $\mathcal{J}_0 = \text{id}(\mathcal{J} \cup \mathcal{A})$ then

$$X \in \mathcal{J}_0 \setminus \mathcal{J}'_n \text{ iff } X \in \mathcal{J}_0 \text{ and } \exists A \in \mathcal{A} \setminus \mathcal{B}_n \ A \cap X \in \mathcal{J}^+.$$

This implies, as \mathcal{J}_0 is clearly $\underline{\Sigma}_1^1$, that $\mathcal{J}_0 \setminus \mathcal{J}'_n$ is a $\underline{\Sigma}_n^1$ set, and hence \mathcal{J}'_n is $\underline{\Pi}_n^1$ (here we have used $\mathcal{J}'_n \subseteq \mathcal{J}_0$ and $n > 1$). ■

The idea of the above proof can be used to construct $\underline{\Sigma}_\alpha^0$ -complete ideals for $\alpha \geq 3$ as well.

4. Proof of Theorem 1.6. Applying Corollary 1.8, we can fix perfect \mathcal{J} -AD families \mathcal{A}_X on every $X \in \mathcal{J}^+$. The statement “ \mathcal{A}_X is an \mathcal{J} -AD family” is (at most) $\underline{\Pi}_2^1$, hence absolute because if $\mathcal{A}_X = [T]$ is coded by a perfect tree $T \in \text{Tree}_2 = \{T \subseteq 2^{<\omega} : T \text{ is a tree}\}$ then

$$“\mathcal{A}_X \text{ is an } \mathcal{J}\text{-AD family}” \equiv \forall x, y \in [T] \ (x \in \mathcal{J}^+ \text{ and } (x = y \text{ or } x \cap y \in \mathcal{J})),$$

where of course we are working on 2^ω and $(x \cap y)(n) = x(n) \cdot y(n)$ for all n .

For any $X, Y \in \mathcal{J}^+$ let $B(X, Y) = \{A \in \mathcal{A}_X : A \cap Y \in \mathcal{J}^+\}$. Then it is a continuous preimage of \mathcal{J}^+ (under $\mathcal{A}_X \rightarrow \mathcal{P}(\omega)$, $A \mapsto A \cap Y$), hence if \mathcal{J} is analytic then $B(X, Y)$ is coanalytic, and similarly if \mathcal{J} is coanalytic then $B(X, Y)$ is analytic.

Let $\kappa = |\mathfrak{c}^V|^W$ and fix an enumeration $\{X_\alpha : \alpha < \kappa\}$ of the set $\mathcal{J}^+ \cap V$ in W . Working in W , we will construct the desired \mathcal{J} -AD refinement $\{A_\alpha : \alpha < \kappa\}$ with $A_\alpha \subseteq X_\alpha$ by recursion on κ . During this process, we will also define a sequence $(B_\alpha)_{\alpha < \kappa}$ in \mathcal{J}^+ .

Assume that $\{A_\xi : \xi < \alpha\}$ and $(B_\xi)_{\xi < \alpha}$ are found. Let γ_α be minimal such that $B(X_{\gamma_\alpha}, X_\alpha)$ contains a perfect set. This property, namely, that an analytic or coanalytic set $H \subseteq \mathcal{P}(\omega)$ contains a perfect set, is absolute because if H is analytic then “ H contains a perfect subset” iff “ H is uncountable” is of the form “ $\forall f \in \mathcal{P}(\omega)^\omega \exists x (x \in H \text{ and } x \notin \text{ran}(f))$ ”, hence it is $\underline{\Pi}_2^1$; and if H is coanalytic then “ H contains a perfect set” is of the form “ $\exists T \in \text{Tree}_2 (T \text{ is perfect and } \forall x \in [T] \ x \in H)$ ”, hence it is $\underline{\Sigma}_2^1$. In particular, $\gamma_\alpha \leq \alpha$. We also know that if C is a perfect set coded in V , then in W it contains κ many new elements: we know it holds for 2^ω e.g. because of the group structure on it, and we can compute new elements of C along a homeomorphism between C and 2^ω fixed in V . Let

$$B_\alpha \in B(X_{\gamma_\alpha}, X_\alpha) \setminus (V \cup \{B_\xi : \xi < \alpha\})$$

be arbitrary, and set $A_\alpha = X_\alpha \cap B_\alpha \in \mathcal{J}^+$. We claim that $\{A_\alpha : \alpha < \kappa\}$ is an \mathcal{J} -AD family (it is clearly a refinement of $\mathcal{J}^+ \cap V$). Let $\alpha, \beta < \kappa$, $\alpha \neq \beta$.

If $\gamma_\alpha = \gamma_\beta = \gamma$ then $B_\alpha, B_\beta \in \mathcal{A}_{X_\gamma}$ are distinct, and hence $A_\alpha \cap A_\beta \subseteq B_\alpha \cap B_\beta \in \mathcal{J}$ (actually, we can assume that it is finite).

If $\gamma_\alpha < \gamma_\beta$, then because of the minimality of γ_β , we know that $B(X_{\gamma_\alpha}, X_\beta)$ does not contain perfect subsets. It is enough to see that $B(X_{\gamma_\alpha}, X_\beta)$ is the

same set in V and W , i.e. if $\psi(x, r)$ is a $\Sigma_1^1(r)$ or $\Pi_1^1(r)$ definition of this set then $\forall x \in W (\psi(x, r) \rightarrow x \in V)$. Why? Because then $B_\alpha \notin B(X_{\gamma_\alpha}, X_\beta)$ but $B_\alpha \in \mathcal{A}_{X_{\gamma_\alpha}}$, hence $A_\alpha \cap A_\beta \subseteq B_\alpha \cap X_\beta \in \mathcal{J}$.

The set $K := B(X_{\gamma_\alpha}, X_\beta)$ is analytic or coanalytic and does not contain perfect subsets (in V or in W). Applying the Mansfield–Solovay Theorem, we find that $K \subseteq L[r]$ ($r \in V$). We also know that $(L[r])^V \cap \mathcal{P}(\omega) = (L[r])^W \cap \mathcal{P}(\omega)$ because $\omega_1^W \subseteq V$, hence $K^V = K^W$.

REMARK 4.1. It is natural to ask the following: Assume that $V \subseteq W$ are transitive models, W contains new reals, and let C be a perfect set coded in W . Does C contain at least $|c^V|^W$ many new elements in W ? In other words: Does $|C^W \setminus V|^W \geq |c^V|^W$ hold? Surprisingly, the answer is no! Moreover, there may be a perfect set of ground model reals in the extension [VW98].

REMARK 4.2. What can we say about possible generalizations of Theorem 1.6? For example, can we weaken the condition on the complexity of the ideal? In general, this statement is false. Let $\varphi(x)$ be a Σ_2^1 definition of a Σ_2^1 (i.e. Δ_2^1) prime P-ideal \mathcal{J} in L . (How to construct such an ideal? Using a Δ_2^1 -good well-order \leq on $\mathcal{P}(\omega)$, by the most natural recursion, at every stage extending our family with a \leq -minimal element which can be added without generating $\mathcal{P}(\omega)$ and also with a \leq -minimal pseudounion of the previous elements, avoiding universal quantification by applying goodness, we obtain such an ideal.) We cannot expect that $\varphi(x)$ defines an ideal in general but we can talk about the *generated* ideal: $x \in \mathcal{J}$ iff “ $\exists y \in \mathcal{J} x \subseteq y$ ”, which is Σ_2^1 too. If r is a Sacks real over L , then \mathcal{J} is still a prime P-ideal in $L[r]$ (see [BrJ95, Lemma 7.3.48]), hence $\mathcal{J}^+ \cap L$ does not have any \mathcal{J} -ADR’s in $L[r]$.

5. On the existence of perfect $(\mathcal{J}, \text{Fin})$ -AD families. First of all, we show that the reverse implication in the first part of Corollary 1.8 does not hold.

EXAMPLE 5.1. The assumption that there is a perfect $(\mathcal{J}, \text{Fin})$ -AD family does not imply that \mathcal{J} is meager: Fix a prime ideal \mathcal{J} on ω . For every partition $P = (P_n)_{n \in \omega}$ of ω into finite sets, fix an $X_P \in [\omega]^\omega$ such that $A_P = \bigcup \{P_n : n \in X_P\} \in \mathcal{J}$ (notice that \mathcal{J} cannot be meager); and let the ideal \mathcal{J} on $2^{<\omega}$ be generated by the sets of the form $A'_P = \bigcup \{2^k : k \in A_P\}$.

Clearly, the family $\{\{f \upharpoonright n : n \in \omega\} : f \in 2^{<\omega}\}$ of branches of $2^{<\omega}$ is a perfect AD family. We show that $\{f \upharpoonright n : n \in \omega\} \in \mathcal{J}^+$. Notice that $\{\text{dom}(s) : s \in A'_P\} = A_P \in \mathcal{J}$ for every P . Thus, a set of the form $B_f = \{f \upharpoonright n : n \in \omega\}$ cannot be an element of the ideal because $\{\text{dom}(s) : s \in B_f\} = \omega$.

To see that \mathcal{J} is not meager, assume the contrary; then by Theorem 1.7 there exists a partition $Q = (Q_n)_{n \in \omega}$ of $2^{<\omega}$ into finite sets such that $\{n \in \omega : Q_n \subseteq A\}$ is finite for every $A \in \mathcal{J}$. Then there is a partition $P = (P_n)_{n \in \omega}$ of ω

into finite sets such that for every n there is an m with $Q_m \subseteq \bigcup\{2^k : k \in P_n\}$. We know that $A'_P \in \mathcal{J}$, a contradiction because A'_P contains infinitely many Q_m 's.

What can we say if there are perfect $(\mathcal{J}, \text{Fin})$ -AD families on every $X \in \mathcal{J}^+$? In this case we have only consistent counterexamples.

THEOREM 5.2. *Assume that $\mathfrak{b} = \mathfrak{c}$. Then there is a non-meager ideal \mathcal{J} on ω such that there are perfect $(\mathcal{J}, \text{Fin})$ -AD families on every $X \in \mathcal{J}^+$.*

Proof. Let $[\omega]^\omega = \{X_\alpha : \alpha < \mathfrak{c}\}$ and $\{\text{partitions of } \omega \text{ into finite sets}\} = \{P_\alpha = (P_n^\alpha)_{n \in \omega} : \alpha < \mathfrak{c}\}$ be enumerations. We will construct the desired ideal \mathcal{J} as an increasing union $\bigcup\{\mathcal{J}_\alpha : \alpha < \mathfrak{c}\}$ of ideals by recursion on $\alpha < \mathfrak{c}$. At the α th stage we will make sure that

- (i) \mathcal{J}_α is generated by $|\alpha|$ many elements;
- (ii) P_α cannot witness that \mathcal{J}_α is meager;
- (iii) either X_α belongs to \mathcal{J}_α , or there is a perfect $(\mathcal{J}_\alpha, \text{Fin})$ -AD family on X_α ;
- (iv) we do not destroy the $(\mathcal{J}_\beta, \text{Fin})$ -AD families we may have constructed at previous stages.

Let $\mathcal{J}_0 = \text{Fin}$ and fix a perfect AD family \mathcal{A}_0 on X_0 . At stage $\alpha > 0$ we already have the ideals \mathcal{J}_β for every $\beta < \alpha$; let $\mathcal{J}_{<\alpha} = \bigcup\{\mathcal{J}_\beta : \beta < \alpha\}$. We also have perfect $(\mathcal{J}_{<\alpha}, \text{Fin})$ -AD families \mathcal{A}_β on $X_\beta \in \mathcal{J}_{<\alpha}^+$ for certain $\beta \in D_\alpha \subseteq \alpha$.

If we can add X_α to $\mathcal{J}_{<\alpha}$, that is, $\mathcal{A}_\beta \cap \text{id}(\mathcal{J}_{<\alpha} \cup \{X_\alpha\}) = \emptyset$ for every $\beta \in D_\alpha$, then let $\mathcal{J}'_\alpha = \text{id}(\mathcal{J}_{<\alpha} \cup \{X_\alpha\})$ and $D'_\alpha = D_\alpha$.

Suppose that we cannot add X_α to $\mathcal{J}_{<\alpha}$, that is, $\mathcal{A}_\beta \cap \text{id}(\mathcal{J}_{<\alpha} \cup \{X_\alpha\}) \neq \emptyset$ for some $\beta \in D_\alpha$. Since $\mathcal{J}_{<\alpha}$ is generated by $< \mathfrak{b} = \mathfrak{c}$ many sets, it is an everywhere meager ideal (see [Sm77] or [Bl10, Thm. 9.10]). We can apply Corollary 1.8 to obtain a perfect $(\mathcal{J}_{<\alpha}, \text{Fin})$ -AD family \mathcal{A}_α on X_α ; let $\mathcal{J}'_\alpha = \mathcal{J}_{<\alpha}$, and let $D'_\alpha = D_\alpha \cup \{\alpha\}$.

Fix a partition $Q = (Q_n)_{n \in \omega}$ of ω into finite sets such that $\{n \in \omega : Q_n \subseteq A\}$ is finite for every $A \in \mathcal{J}'_\alpha$ (we know that \mathcal{J}'_α is meager).

CLAIM. *There exist partitions $Q_{\beta,B} = (Q_n^{\beta,B})_{n \in \omega}$ for every $\beta \in D'_\alpha$ and $B \in \mathcal{J}'_\alpha$ such that $A \cap Q_n^{\beta,B} \setminus B \neq \emptyset$ for every $\beta \in D'_\alpha$, $A \in \mathcal{A}_\beta$, $B \in \mathcal{J}'_\alpha$, and $n \in \omega$.*

Proof of Claim. Let $\beta \in D'_\alpha$ and $B \in \mathcal{J}'_\alpha$. We know that \mathcal{A}_β is compact as a subset of $\mathcal{P}(\omega)$. Basic open sets in $\mathcal{P}(\omega)$ are of the form $[s, t] = \{A \subseteq \omega : s \cap A = \emptyset \text{ and } t \subseteq A\}$ for disjoint, finite $s, t \subseteq \omega$. Then $\mathcal{A}_\beta \subseteq \bigcup\{[\emptyset, \{n\}] : n \in \omega \setminus B\}$ because $A \setminus B$ is infinite for every $A \in \mathcal{A}_\beta$. Therefore $\mathcal{A}_\beta \subseteq \bigcup\{[\emptyset, \{n\}] : n \in N_0 \setminus B\}$ for an $N_0 \in \omega$, in particular, $A \cap N_0 \setminus B \neq \emptyset$ for every $A \in \mathcal{A}_\beta$. Let $Q_0^{\beta,B} = [0, N_0)$. We can proceed by the same argument: $\mathcal{A}_\beta \subseteq \bigcup\{[\emptyset, \{n\}] : n \in [N_0, \omega) \setminus B\}$, hence there is an $N_1 > N_0$ such that

$\mathcal{A}_\beta \subseteq \bigcup\{[\emptyset, \{n\}] : n \in [N_0, N_1] \setminus B\}$, in other words, $A \cap [N_0, N_1] \setminus B \neq \emptyset$ for every $A \in \mathcal{A}_\beta$. Let $Q_1^{\beta, B} = [N_0, N_1)$ etc. ■

Now we have the family $\mathcal{Q} = \{P_\alpha\} \cup \{Q\} \cup \{Q_{\beta, B} : \beta \in D'_\alpha, B \in \mathcal{C}_\alpha\}$ of partitions where $\mathcal{C}_\alpha \subseteq \mathcal{J}'_\alpha$ is a cofinal family, $|\mathcal{C}_\alpha| \leq \max\{|\alpha|, \omega\}$. Since $|\mathcal{Q}| < \mathfrak{c} = \mathfrak{b}$, there is a partition $R = (R_m)_{m \in \omega}$ which dominates all of these partitions, that is, $\forall P = (P_n)_{n \in \omega} \in \mathcal{Q} \forall^\infty m \exists n P_n \subseteq R_m$ [Bl10, Thm. 2.10]. Let $Y = \bigcup\{R_{2n} : n \in \omega\}$ and $\mathcal{J}_\alpha = \text{id}(\mathcal{J}'_\alpha \cup \{Y\})$.

Then (i) is clearly satisfied. In order to see (ii) notice that since the partition R_m was dominating and $P_\alpha \in \mathcal{Q}$, for almost every m there exists an n with $P_n^\alpha \subseteq R_{2m}$. Condition (iii) is also clear if $X_\alpha \in \mathcal{J}'_\alpha$.

If $X_\alpha \notin \mathcal{J}'_\alpha$ then by definition $\alpha \in D'_\alpha$, so to see (iii) and (iv) we have to show that for every $\beta \in D'_\alpha$ the family \mathcal{A}_β is not just an $(\mathcal{J}'_\alpha, \text{Fin})$ -AD family, but also an $(\mathcal{J}_\alpha, \text{Fin})$ -AD family. In other words, it is enough to check that for every $A \in \mathcal{A}_\beta$ and $B \in \mathcal{J}'_\alpha$ we have $A \setminus (B \cup Y) \neq \emptyset$. Fix such A and B ; we can assume that $B \in \mathcal{C}_\alpha$. Then for almost every m , there is an n_m such that $Q_{n_m}^{\beta, B} \subseteq R_{2m+1}$, and by the Claim we know that $A \cap Q_{n_m}^{\beta, B} \setminus B \neq \emptyset$. Therefore, $A \setminus (B \cup Y)$ is infinite, hence $\mathcal{A}_\beta \cap \mathcal{J}_\alpha = \emptyset$ for every $\beta \in D'_\alpha$. ■

What can we say about ideals on the second level of the projective hierarchy? Do there always exist perfect or at least uncountable $(\mathcal{J}, \text{Fin})$ -AD families? If all Σ_2^1 and Π_2^1 sets have the Baire property, then of course yes, because then Σ_2^1 and Π_2^1 ideals are meager and we can apply Corollary 1.8. On the other hand, if \mathcal{J} is a Σ_2^1 (i.e. Δ_2^1) prime ideal (e.g. in L) then every \mathcal{J} -AD family is a singleton.

Similarly, we can construct a Σ_2^1 ideal \mathcal{J} in L such that there are infinite \mathcal{J} -AD families but all of them are countable: copy the above ideal \mathcal{J} to the elements of a partition $\{P_n : n \in \omega\} \subseteq [\omega]^\omega$ of ω , and let \mathcal{J} be the generated ideal.

This last example is very artificial, in the sense that this ideal is constructed from maximal ideals in a very “obvious” way, many of its restrictions are prime ideals. However, we can construct even more peculiar ideals:

PROPOSITION 5.3. *Suppose that there exists a Δ_n^1 prime ideal on ω for some n . Then there exists a Δ_n^1 ideal \mathcal{J} such that \mathcal{J} is nowhere maximal but every \mathcal{J} -AD family is countable. In particular, there exists such a Δ_2^1 ideal in L .*

Proof. Let \mathcal{U} be an ultrafilter and define $\mu : \mathcal{P}(\omega) \rightarrow [0, 1]$ as $\mu(A) = \lim_{\mathcal{U}} |A \cap n|/n$ where $\lim_{\mathcal{U}}$ stands for the \mathcal{U} -limit operation on sequences in topological spaces, that is, $\lim_{\mathcal{U}}(a_n) = a$ iff $\{\overline{n \in \omega : a_n \in V}\} \in \mathcal{U}$ for every neighborhood V of a . It is easy to see that if $\{a_n : n \in \omega\}$ is compact, then $\lim_{\mathcal{U}}(a_n)_{n \in \omega}$ exists, in particular, μ is defined on every $A \in \mathcal{P}(\omega)$. It is also straightforward to show that μ is a finitely additive non-atomic probability

measure on $\mathcal{P}(\omega)$, that is, $\mu(\emptyset) = 0$, $\mu(A \cup B) = \mu(A) + \mu(B)$ if $A \cap B = \emptyset$, $\mu(\omega) = 1$, and if $\mu(X) = \varepsilon > 0$ then for every $\delta \in (0, \varepsilon)$ there is a $Y_\delta \subseteq X$ with $\mu(Y_\delta) = \delta$.

Let $\mathcal{J} = \{A \subseteq \omega : \mu(A) = 0\}$. Then \mathcal{J} is an ideal; it is nowhere maximal because μ is non-atomic (in particular, there are infinite \mathcal{J} -AD families). We show that every \mathcal{J} -AD family is countable. If there was an uncountable \mathcal{J} -AD family \mathcal{A} , then $\mathcal{A}_n = \{A \in \mathcal{A} : \mu(A) > 1/n\}$ would be uncountable for some $n \in \omega$, and therefore among any n elements of \mathcal{A}_n there would be two with \mathcal{J} -positive intersection.

Notice that if \mathcal{U} is Δ_n^1 ($n \geq 2$) then \mathcal{J} is also Δ_n^1 because $A \in \mathcal{J}$ iff $\forall k \in \omega \{n \in \omega : |A \cap n|/n < 2^{-k}\} \in \mathcal{U}$, and the function $A \mapsto \{n \in \omega : |A \cap n|/n < 2^{-k}\}$ is continuous (for every k). ■

6. On $(\mathcal{J}, \text{Fin})$ -ADR's. In this section, we study Question 1.10.

THEOREM 6.1. *Assume MA_κ and let \mathcal{J} be an everywhere meager ideal. Then every $\mathcal{H} \in [\mathcal{J}^+]^{\leq \kappa}$ has an $(\mathcal{J}, \text{Fin})$ -ADR.*

Proof. Let $\mathcal{H} = \{H_\alpha : \alpha < \kappa\}$ be an enumeration. Define $p \in \mathbb{P} = \mathbb{P}(\mathcal{H})$ iff p is a function, $\text{dom}(p) \in [\kappa]^{< \omega}$, and $p(\alpha) \in [H_\alpha]^{< \omega}$ for every $\alpha \in \text{dom}(p)$; also declare $p \leq q$ iff $\text{dom}(p) \supseteq \text{dom}(q)$, $\forall \alpha \in \text{dom}(q) p(\alpha) \supseteq q(\alpha)$, and $\forall \{\alpha, \beta\} \in [\text{dom}(q)]^2 p(\alpha) \cap p(\beta) = q(\alpha) \cap q(\beta)$.

Then \mathbb{P} is a poset. First of all, we show that \mathbb{P} has the ccc. Let $\{p_\xi : \xi < \omega_1\} \subseteq \mathbb{P}$. Then $\{\text{dom}(p_\xi) : \xi < \omega_1\} \subseteq [\kappa]^{< \omega}$. We can assume that this family forms a Δ -system, and $\text{dom}(p_\xi) = D_\xi \cup R$. There are at most ω many functions $R \rightarrow \text{Fin}$, hence we can also assume that there is a $q \in \mathbb{P}$ such that $p_\xi \upharpoonright R = q$ for every $\xi < \omega_1$. Clearly, $p_\xi \cup p_\zeta \in \mathbb{P}$ and $p_\xi \cup p_\zeta \leq p_\xi$ for all $\xi, \zeta < \omega_1$.

It is easy to see that for every $\alpha < \kappa$ the set $D_\alpha = \{p \in \mathbb{P} : \alpha \in \text{dom}(p)\}$ is dense in \mathbb{P} . If G is a $\{D_\alpha : \alpha < \kappa\}$ -generic filter, then let $F_G : \kappa \rightarrow \mathcal{P}(\omega)$, $F_G(\alpha) = \bigcup \{p(\alpha) : p \in G\}$. Clearly, $F_G(\alpha) \subseteq H_\alpha$ for every α .

We show that $F_G(\alpha) \cap F_G(\beta)$ is finite for any distinct $\alpha, \beta < \kappa$. Let $p \in D_\alpha \cap G$, $q \in D_\beta \cap G$, and $r \in G$ be a common lower bound of them. It is easy to see that $F_G(\alpha) \cap F_G(\beta) = r(\alpha) \cap r(\beta)$.

If somehow we can make sure that $F_G(\alpha) \in \mathcal{J}^+$, then we are done because $\{F_G(\alpha) : \alpha < \kappa\}$ will be an $(\mathcal{J}, \text{Fin})$ -ADR of \mathcal{H} . We show that if G is (V, \mathbb{P}) -generic then $F_G(\alpha)$ is a Cohen real in $\mathcal{P}(H_\alpha)$ over V . This is enough because then $F_G(\alpha) \notin \mathcal{J} \upharpoonright H_\alpha$ (we know that $\mathcal{J} \upharpoonright H_\alpha$ is meager), and to show that $V[F_G(\alpha)] \models F_G(\alpha) \notin \mathcal{J} \upharpoonright H_\alpha$, it is enough to use countably many dense sets. Why? For every α we can fix a countable family $\mathcal{C}_\alpha = \{C_n^\alpha : n \in \omega\}$ of closed nowhere dense subsets of $\mathcal{P}(H_\alpha)$ which covers $\mathcal{J} \upharpoonright H_\alpha$, and hence we have countably many dense subsets of the Cohen forcing such that if a filter is generic for this family then the generic real is not covered by any element

of \mathcal{C}_α . More precisely, we have to translate these dense subsets of the Cohen forcing to dense subsets in \mathbb{P} , which can be done by applying the (inverse of the) projection $\mathbb{P} \rightarrow \mathbb{C}(H_\alpha)$ defined below.

Fix an $\alpha < \kappa$, let $\mathbb{C}(H_\alpha) = \{s : s \text{ is a finite partial function from } H_\alpha \text{ to } 2\}$ where $s \leq t$ iff $s \supseteq t$ (then $\mathbb{C}(H_\alpha)$ adds a Cohen subset of H_α over V), and define the map $e = e_\alpha : \mathbb{P} \rightarrow \mathbb{C}(H_\alpha)$ as follows:

- (i) $\text{dom}(e(p)) = \bigcup \{p(\beta) \cap H_\alpha : \beta \in \text{dom}(p)\}$;
- (ii) $e(p)(n) = 1$ iff $n \in p(\alpha)$.

We show that e is a *projection* (see e.g. [A10, p. 335]), that is,

- (1) e is order preserving, onto, and $e(\emptyset) = \emptyset$;
- (2) $\forall p \in \mathbb{P} \forall s \in \mathbb{C}(H_\alpha) (s \leq e(p) \rightarrow \exists p' \leq p e(p') = s)$.

Clearly, $e(\emptyset) = \emptyset$. Assume that $p \leq q$. Then clearly $\text{dom}(e(p)) \supseteq \text{dom}(e(q))$. If $n \in \text{dom}(e(q))$ and $n \in q(\alpha) \subseteq p(\alpha)$ then $e(q)(n) = e(p)(n) = 1$; if $n \in \text{dom}(e(p))$ and $n \in q(\beta) \setminus q(\alpha)$ for some $\beta \neq \alpha$ then, as $p(\alpha) \cap p(\beta) = q(\alpha) \cap q(\beta)$, we have $n \in p(\beta) \setminus p(\alpha)$ and hence $e(q)(n) = e(p)(n) = 0$. This shows that e is indeed order preserving.

To show that e is onto, we have to assume that $H_\alpha \subseteq \bigcup \{H_\beta : \beta \neq \alpha\}$ (and without loss of generality we can do so by extending \mathcal{H} to be a cover of ω and adding ω as an element to \mathcal{H}). For an $s \in \mathbb{C}(H_\alpha)$ define $p \in \mathbb{P}$ as follows: Fix a finite $D \subseteq \kappa$ containing α such that $\text{dom}(s) \subseteq \bigcup \{H_\beta : \beta \in D\}$, let $\text{dom}(p) = D$, and define $p(\alpha) = s^{-1}(1)$ and $p(\beta) = \{n \in H_\beta \cap H_\alpha : s(n) = 0\}$. Then $e(p) = s$.

To show that e satisfies (2), fix a $p \in \mathbb{P}$, an $s \in \mathbb{C}(H_\alpha)$, and assume that $s \leq e(p)$. Define $p' \in \mathbb{P}$ as follows: For every $n \in J = (s \setminus e(p))^{-1}(0)$ pick a $\gamma_n \in \kappa \setminus \{\alpha\}$ such that $n \in H_{\gamma_n}$. Let $\text{dom}(p') = \text{dom}(p) \cup \{\gamma_n : n \in J\}$ and define $p'(\alpha) = p(\alpha) \cup s^{-1}(1)$, and if $\beta \in \text{dom}(p') \setminus \{\alpha\}$ then $p'(\beta) = p(\beta) \cup \{n \in J : \beta = \gamma_n\}$. It is straightforward to see that $p' \in \mathbb{P}$, $p' \leq p$, and $e(p') = s$.

We know that if G is (V, \mathbb{P}) -generic then $e[G]$ generates a (V, \mathbb{C}) -generic filter G' . Notice that the Cohen real defined from G' is $F_G(\alpha)$, so we are done. ■

Unfortunately, at this moment, we do not know whether we really needed Martin's Axiom in the previous theorem or whether the theorem holds in ZFC. We show that if we attempt to construct a counterexample, that is, say, a tall Borel ideal \mathcal{J} and a family $\mathcal{H} \in [\mathcal{J}^+]^{<\mathfrak{c}}$ without an $(\mathcal{J}, \text{Fin})$ -ADR, we have to be careful. Let us define the following cardinal invariants of tall ideals on ω : The *star-additivity* of \mathcal{J} is

$$\text{add}^*(\mathcal{J}) = \min\{|\mathcal{X}| : \mathcal{X} \subseteq \mathcal{J} \text{ and } \nexists A \in \mathcal{J} \forall X \in \mathcal{X} X \subseteq^* A\},$$

the *Fodor number* of \mathcal{J} is

$$F(\mathcal{J}) = \min\{|\mathcal{H}| : \mathcal{H} \subseteq \mathcal{J}^+ \text{ has no } \mathcal{J}\text{-ADR}\},$$

and the *star-Fodor number* of \mathcal{J} is

$$F^*(\mathcal{J}) = \min\{|\mathcal{H}| : \mathcal{H} \subseteq \mathcal{J}^+ \text{ has no } (\mathcal{J}, \text{Fin})\text{-ADR}\}.$$

Clearly, \mathcal{J} is a P-ideal iff $\text{add}^*(\mathcal{J}) > \omega$. Proposition 1.9 says that $F(\mathcal{J}) = \mathfrak{c}$ whenever \mathcal{J} is everywhere meager; and clearly $F^*(\mathcal{J}) \leq F(\mathcal{J})$.

FACT 6.2. *If $\text{add}^*(\mathcal{J}) < F(\mathcal{J})$ then $\text{add}^*(\mathcal{J}) < F^*(\mathcal{J})$. If $\text{add}^*(\mathcal{J}) = F(\mathcal{J})$ then $F(\mathcal{J}) = F^*(\mathcal{J})$.*

Proof. Assume that $\mathcal{H} = \{H_\alpha : \alpha < \kappa\} \subseteq \mathcal{J}^+$ where $\kappa = \text{add}^*(\mathcal{J}) < F(\mathcal{J})$. First fix an \mathcal{J} -ADR $\{A_\alpha : \alpha < \kappa\}$ of \mathcal{H} ($A_\alpha \subseteq H_\alpha$). Then for every $\alpha < \kappa$ fix a $B_\alpha \in \mathcal{J}$ such that $A_\alpha \cap A_\beta \subseteq^* B_\alpha$ for every $\beta < \alpha$, and let $A'_\alpha = A_\alpha \setminus B_\alpha$. Then $\{A'_\alpha : \alpha < \kappa\}$ is an $(\mathcal{J}, \text{Fin})$ -ADR of \mathcal{H} . The second statement can be proved by the same argument. ■

In particular, if \mathcal{J} is an everywhere meager P-ideal and $F^*(\mathcal{J}) < \mathfrak{c}$, then $F^*(\mathcal{J}) < F(\mathcal{J})$, hence $\text{add}^*(\mathcal{J}) < F(\mathcal{J})$ and so $\omega_1 \leq \text{add}^*(\mathcal{J}) < F^*(\mathcal{J}) < \mathfrak{c}$, therefore $\mathfrak{c} \geq \omega_3$.

7. Mixing reals. In this section, we study two closely related properties of forcing notions, one of which is slightly stronger than “[ω] $^\omega \cap V$ has an ADR in $V^\mathbb{P}$ ”.

First of all, let us recall the following definition from the introduction: Let \mathbb{P} be a forcing notion. We say that an $f \in \omega^\omega \cap V^\mathbb{P}$ is a *mixing real* over V if $|f[X] \cap Y| = \omega$ for all $X, Y \in [\omega]^\omega \cap V$. (Clearly, it is enough to require that $f[X] \cap Y \neq \emptyset$ for all $X, Y \in [\omega]^\omega \cap V$.) If such an f is one-to-one, then we call it an *injective mixing real* or a *mixing injection*.

We already mentioned that mixing reals can be seen as “infinite splitting partitions” (see the proof below), and that if \mathbb{P} adds a mixing injection then it is easy to find an ADR of $[\omega]^\omega \cap V$ in $V^\mathbb{P}$.

PROPOSITION 7.1. *Let \mathbb{P} be a forcing notion. Then the following are equivalent:*

- (i) *There is a mixing real $f \in \omega^\omega \cap V^\mathbb{P}$ over V .*
- (ii) *There is an $f \in \omega^\omega \cap V^\mathbb{P}$ such that $f[X] = \omega$ for all $X \in [\omega]^\omega \cap V$.*
- (iii) *There is a partition $(Y_n)_{n \in \omega}$ of ω into infinite sets in $V^\mathbb{P}$ such that $\forall X \in [\omega]^\omega \cap V \forall n |X \cap Y_n| = \omega$.*
- (iii)' *There is a partition $(Y_n)_{n \in \omega}$ of ω into infinite sets in $V^\mathbb{P}$ such that $\forall X \in [\omega]^\omega \cap V \forall n X \cap Y_n \neq \emptyset$.*

Proof. (ii)→(i) and (iii)↔(iii)' are trivial. (ii)↔(iii)' is justified by letting $Y_n = f^{-1}(n)$ (and vice versa). To verify (i)→(ii), fix a partition $(C_n)_{n \in \omega}$ of

ω into infinite sets in V and let $g : \omega \rightarrow \omega$, $g \upharpoonright C_n \equiv n$. If f is a mixing real over V , then $h = g \circ f$ has the required property. ■

PROPOSITION 7.2. *Let \mathbb{P} be a forcing notion.*

- (i) *If \mathbb{P} adds random reals then it adds mixing reals.*
- (ii) *If \mathbb{P} adds dominating reals then it adds mixing reals.*
- (iii) *If \mathbb{P} adds Cohen reals then it adds mixing injections.*
- (iv) *If \mathbb{P} adds mixing injections then it adds unbounded reals.*
- (v) *If \mathbb{P} has the Laver property then it does not add injective mixing reals.*

Proof. (i) Let λ be the usual probability measure on ω^ω , that is, λ is uniquely determined by the values $\lambda([s]) = 2^{-s(0)-s(1)-\dots-s(n-1)-n}$ where $s : n \rightarrow \omega$ and $[s] = \{f \in \omega^\omega : s \subseteq f\}$. If $\mathcal{N}_\lambda = \{A \subseteq \omega^\omega : \lambda(A) = 0\}$, then it is well-known that $\text{Borel}(\omega^\omega)/\mathcal{N}_\lambda$ is forcing equivalent to the random forcing. It is enough to see that the set $A_{X,Y} = \{f \in \omega^\omega : |f[X] \cap Y| < \omega\}$ is a null set in ω^ω for all $X, Y \in [\omega]^\omega$: $A_{X,Y} = \bigcup_{n \in \omega} \{f \in \omega^\omega : f[X] \cap Y \subseteq n\}$ and if $X = \{x_k : k \in \omega\}$ and $n \in \omega$ then $\{f : f[X] \cap Y \subseteq n\} = \{f : \forall k f(x_k) \in n \cup (\omega \setminus Y)\}$. Clearly, $\sum \{2^{-m-1} : m \in n \cup (\omega \setminus Y)\} = \varepsilon < 1$ and hence $\lambda(\{f : f[X] \cap Y \subseteq n\}) \leq \lim_{k \rightarrow \infty} \varepsilon^k = 0$.

(ii) Trivial modification of the proof of the fact (see e.g. [Hb12, Fact 20.1]) that adding a dominating real implies adding a splitting real works here as well: adding a dominating real is equivalent to adding a dominating partition $(P_n)_{n \in \omega}$ of ω into finite sets [Bl10, Thm. 2.10], that is, for every partition $(Q_m)_{m \in \omega} \in V$ of ω into finite sets, $\forall^\infty n \exists m Q_m \subseteq P_n$. Now any infinite partition of ω consisting of unions of infinitely many P_n 's satisfies (iii) from Proposition 7.1.

(iii) We can talk about *injective Cohen reals*. Simply consider the forcing notion (Inj, \supseteq) where $\text{Inj} = \{s \in \omega^{<\omega} : s \text{ is one-to-one}\}$, or the forcing notion $(\text{Borel}(\text{INJ}) \setminus \mathcal{M}(\text{INJ}), \subseteq)$ where $\text{INJ} = \{f \in \omega^\omega : f \text{ is one-to-one}\}$ is a nowhere dense closed subset of ω^ω and $\mathcal{M}(\text{INJ})$ is the meager ideal on this Polish space. It is not difficult to see that these forcing notions are forcing equivalent to the Cohen forcing (moreover, INJ is homeomorphic to ω^ω).

If c is an injective Cohen real over V , then c is mixing: For any X, Y in $[\omega]^\omega$, the set $A'_{X,Y} = A_{X,Y} \cap \text{INJ} = \bigcup_{n \in \omega} \{f \in \text{INJ} : f[X] \cap Y \subseteq n\}$ is meager because $\{f \in \text{INJ} : f[X] \cap Y \subseteq n\}$ is closed and nowhere dense in INJ .

(iv) Let $f \in \text{INJ} \cap V^{\mathbb{P}}$ be a mixing injection and assume on the contrary that there is a strictly increasing $g \in \omega^\omega \cap V$ such that $f, f^{-1} < g$ (where of course $f^{-1} < g$ means that $f^{-1}(k) < g(k)$ for every $k \in \text{ran}(f)$).

We define $X = \{x_k : k \in \omega\}$, $Y = \{y_k : k \in \omega\} \in [\omega]^\omega$ in V as follows: $x_0 = 0$, $y_0 = g(0)$, $x_n = \max\{g(y_k) : k < n\}$, and $y_n = g(x_n)$. Suppose that

$f(x_k) = y_l$ for some $k, l \in \omega$. If $k \leq l$ then

$$f(x_k) < g(x_k) = g\left(\max_{m < k} g(y_m)\right) \leq g\left(\max_{m < l} g(y_m)\right) = x_l < g(x_l) = y_l,$$

a contradiction. If $k > l$ then

$$x_k = f^{-1}(y_l) < g(y_l) \leq \max\{g(y_m) : m < k\} = x_k,$$

which is again impossible. Thus, $f[X] \cap Y = \emptyset$, so f cannot be a mixing injection.

(v) Fix a sequence $(a_n)_{n \in \omega} \in \omega^\omega \cap V$ satisfying $a_{n+1} - a_n > (n+2)2^{n+1}$ and $a_0 > 1$. Assume that $p \Vdash \dot{f} \in \text{INJ}$. Let \dot{g} be a \mathbb{P} -name for a function on ω such that $p \Vdash \dot{g}(n) = \dot{f} \cap (a_n \times a_n) = \{(k, l) \in a_n \times a_n : f(k) = l\}$ for every n (in particular, $p \Vdash \text{“}\dot{g}(n) \text{ is an injective partial function from } a_n \text{ to } a_n\text{”}$). Then $p \Vdash \dot{g} \in \prod_{n \in \omega} \mathcal{P}(a_n \times a_n)$, hence, applying the Laver property of our forcing notion to the name \dot{g} for a function from ω to $[\omega \times \omega]^{<\omega}$, we find a $q \leq p$ and a “slalom” $S : \omega \rightarrow [[\omega \times \omega]^{<\omega}]^{<\omega}$ in V which catches \dot{g} , that is, $S(n) \subseteq \mathcal{P}(a_n \times a_n)$, $|S(n)| \leq 2^n$, and $q \Vdash \dot{g}(n) \in S(n)$ for every n . Without loss of generality we can assume that all elements of $S(n)$ are injective partial functions $a_n \rightarrow a_n$.

Working in V , we will define sets $X = \{x_n : n \in \omega\}, Y = \{y_n : n \in \omega\}$ in $[\omega]^\omega$ by recursion on n so that $q \Vdash \dot{f}[X] \cap Y = \emptyset$.

Let $x_0 \in a_0$ be arbitrary. We know that there is a $y_0 \in a_0$ such that $(x_0, y_0) \notin \bigcup S(0)$ (a function cannot cover $\{(x_0, k) : k < a_0\}$).

Assume that we already have $X_n = \{x_k : k \leq n\}$ and $Y_n = \{y_k : k \leq n\}$ such that $(X_n \times Y_n) \cap \bigcup_{k \leq n} \bigcup S(k) = \emptyset$. There is an $x_{n+1} \in a_{n+1} \setminus a_n$ such that


$$\{s(x_{n+1}) : s \in S(n+1), x_{n+1} \in \text{dom}(s)\} \cap Y_n = \emptyset.$$

Why? If for every $m \in a_{n+1} \setminus a_n$ there is an $s_m \in S(n+1)$ such that $s_m(m) \in Y_n$ then there is a set $H \in [a_{n+1} \setminus a_n]^{n+2}$ such that $s_m = s$ does not depend on $m \in H$ (because $|a_{n+1} \setminus a_n| > (n+2)2^{n+1}$ and $|S(n+1)| \leq 2^{n+1}$). But it would mean that $H \subseteq \text{dom}(s)$ and $|s[H]| \leq |Y_n| = n+1$, which is a contradiction because s is injective.

We also want to fix a $y_{n+1} \in a_{n+1} \setminus a_n$ such that $y_{n+1} \neq s(x_k)$ for any $k \leq n+1$, $s \in S(n+1)$ if $x_k \in \text{dom}(s)$. The set of forbidden values is of size at most $2^{n+1}(n+2)$ hence there is such a y_{n+1} . ■

In the diagram below, we summarize the logical implications between classical properties of forcing notions and the ones we defined above. We will show that arrows without a $*$ are strict (i.e. not equivalences), and that there are no other implications between these properties. The question mark means that we do not know whether the implication holds (but the reverse implication is false). Of course, \mathbb{C} stands for the Cohen forcing, \mathbb{B} is the

random forcing, and to keep the diagram small, we did not put “ \mathbb{P} adds ...” and “ \mathbb{P} has the ...” before the properties we deal with.



The non-trivial non-implications in the diagram are the following:

- \neg Laver prop. $\not\rightarrow$ splitting reals ⁽¹⁾. The infinitely equal forcing $\mathbb{E}\mathbb{E}$ is ω^ω -bounding, preserves P-points (hence cannot add splitting reals), and $\Vdash_{\mathbb{E}\mathbb{E}}$ “ $2^\omega \cap V$ is a null set” [BrJ95, Lemmas 7.4.13–7.4.15]. Further, $\mathbb{E}\mathbb{E}$ cannot have the Laver property because otherwise it would have the Sacks property as well but then it could not force $2^\omega \cap V$ to be of measure zero (this follows from e.g. [BrJ95, Thm. 2.3.12]).
- unbounded reals $\not\rightarrow$ splitting reals: The Miller forcing [BrJ95, 7.3.E].
- spl. reals $\not\rightarrow$ \neg Sacks prop.: The Silver forcing adds splitting reals [Hb12, Lemma 2.3] and it is straightforward to show that it satisfies the Sacks property.

We list the remaining questions in the next section.

8. Related questions. We already presented Σ_n^1 - and Π_n^1 -complete ideals but our construction was pretty artificial.

QUESTION 8.1. *Can we define “natural” Σ_n^1 - and Π_n^1 -complete ideals?*

QUESTION 8.2. *Assume that V, W and \mathcal{J} are as in Theorem 1.6. Does there exist an $(\mathcal{J}, \text{Fin})$ -ADR of $\mathcal{J}^+ \cap V$ in W ? Or at least an \mathcal{J} -ADR $\{A_X : X \in \mathcal{J}^+ \cap V\} \in W$ such that for any distinct $X, Y \in \mathcal{J}^+ \cap V$ (using the notation from the proof of Theorem 1.6) there is a $B_{X,Y} \in \mathcal{J} \cap V$ such that $A_X \cap A_Y \subseteq B_{X,Y}$?*

QUESTION 8.3. *Does there exist a non-meager ideal \mathcal{J} (in ZFC) such that there are perfect $(\mathcal{J}, \text{Fin})$ -AD families on every $X \in \mathcal{J}^+$?*

⁽¹⁾ Notice that showing “ \neg Sacks prop. $\not\rightarrow$ splitting reals” is easier as the Miller forcing is a natural counterexample (for details see [BrJ95, 7.3.E]).

In Proposition 5.3, assuming that there is a \mathcal{A}_2^1 ultrafilter, we constructed a \mathcal{A}_2^1 ideal \mathcal{J} such that every \mathcal{J} -AD family is countable but \mathcal{J} is nowhere maximal.

QUESTION 8.4. *Is it consistent that there are no \mathcal{A}_2^1 ultrafilters but there is a \mathcal{A}_2^1 ideal \mathcal{J} such that every \mathcal{J} -AD family is countable but \mathcal{J} is nowhere maximal?*

A remark to Question 8.4: We know [BrJ95, Thm. 9.3.9(2)] that if there are no dominating reals over $L[r]$ for any $r \in \omega^\omega$, then there is a Σ_2^1 unbounded hence non-meager filter. If every \mathcal{A}_2^1 set is Lebesgue measurable or has the Baire property, then there are no \mathcal{A}_2^1 ultrafilters. For instance, these conditions hold in the Cohen and random models over $V = L$ [BrJ95, Thm. 9.2.1]. In these models a non-meager Σ_2^1 ideal \mathcal{J} must be nowhere maximal (otherwise a restriction of \mathcal{J} would be a Δ_2^1 prime ideal). It would be interesting to know the possible sizes of \mathcal{J} -AD families in these models.

QUESTION 8.5. *Is it consistent that for some (tall) Borel (P -)ideal \mathcal{J} a family $\mathcal{H} \in [\mathcal{J}^+]^{<\mathfrak{c}}$ does not have an $(\mathcal{J}, \text{Fin})$ -ADR (i.e. $F^*(\mathcal{J}) < \mathfrak{c}$)?*

QUESTION 8.6. *Does adding mixing injections imply adding Cohen reals?*

QUESTION 8.7. *Does the Sacks property of a forcing notion imply that it does not add mixing reals?*

Proposition 7.1 motivates the following notion: Let $n \geq 2$. We say that a forcing notion adds an n -splitting partition if there is a partition $(Y_k)_{k < n}$ of ω into infinite sets in $V^\mathbb{P}$ such that $|X \cap Y_k| = \omega$ for every $X \in [\omega]^\omega \cap V$ and $k < n$. In particular, adding 2-splitting partitions is the same as adding splitting reals, and adding ω -splitting (infinite splitting) partitions is equivalent to adding mixing reals.

It is easy to see that if \mathbb{P} adds a splitting real then the n -stage iteration of \mathbb{P} adds a 2^n -splitting partition. In fact, splitting reals and n -splitting partitions cannot be separated in terms of cardinal invariants. Let us denote by \mathfrak{s}_n ($2 \leq n < \omega$) the least size of a family \mathcal{S}_n of partitions of ω into n many infinite sets such that

$$(*) \quad \forall X \in [\omega]^\omega \exists P = (P_k)_{k < n} \in \mathcal{S}_n \forall k < n |X \cap P_k| = \omega.$$

Of course, this definition makes sense for $n = \omega$ as well, but \mathfrak{s}_ω stands for an already defined and studied cardinal invariant. To avoid confusion, let us denote this cardinal by $\mathfrak{s}_{\text{mix}}$.

Then $\mathfrak{s}_n = \mathfrak{s} = \mathfrak{s}_2$ for every $2 \leq n < \omega$. For the non-trivial direction, assume that we have a family \mathcal{S} of splitting partitions of size \mathfrak{s} and consider all possible “ $(n - 1)$ -long iterated nestings” of these partitions. For example, if $n = 3$ then to every pair $(P = (P_0, P_1), Q = (Q_0, Q_1))$ of partitions from \mathcal{S} we associate a partition of ω into three infinite sets as follows: Let $e_0 : \omega \rightarrow Q_0$

be the increasing bijection and take the partition $(e_0[P_0], e_0[P_1], Q_1)$. We obtain $\mathfrak{s}^{n-1} = \mathfrak{s}$ many partitions of ω into n infinite sets, the family \mathcal{S}_n of these partitions satisfies $(*)$, and hence $\mathfrak{s}_n \leq \mathfrak{s}$.

QUESTION 8.8. *Does adding n -splitting partitions ($2 \leq n < \omega$) imply adding $(n + 1)$ -splitting partitions?*

QUESTION 8.9. *Is it true that $\mathfrak{s}_{\text{mix}} = \mathfrak{s}$? Does adding splitting reals (or n -splitting partitions for every n) imply adding mixing reals? What can we say about the Silver forcing? (It is straightforward to see that it adds n -splitting partitions for every n .)*

Acknowledgements. The first author was supported by the Austrian Science Fund (FWF) grant no. P25671, and the Hungarian National Foundation for Scientific Research grants nos. 83726 and 77476. The second author was supported by the Austrian Science Fund (FWF) grant no. P25748. The third author is partially supported by the Hungarian Scientific Foundation grants nos. 104178 and 113047.

References

- [A10] U. Abraham, *Proper forcing*, in: Handbook of Set Theory, Vol. 1, M. Foreman and A. Kanamori (eds.), Springer, 2010, 333–394.
- [BaP10] B. Balcar and T. Pazák, *Quotients of Boolean algebras and regular subalgebras*, Arch. Math. Logic 49 (2010), 329–342.
- [BaSV81] B. Balcar, P. Simon and P. Vojtáš, *Refinement properties and extensions of filters in Boolean algebras*, Trans. Amer. Math. Soc. 267 (1981), 265–283.
- [BaV80] B. Balcar and P. Vojtáš, *Almost disjoint refinement of families of subsets of \mathbb{N}* , Proc. Amer. Math. Soc. 79 (1980), 465–470.
- [BrJ95] T. Bartoszyński and H. Judah, *Set Theory: On the Structure of the Real Line*, AK Peters, 1995.
- [BHM75] J. E. Baumgartner, A. Hajnal and A. Máté, *Weak saturation properties of ideals*, in: Infinite and Finite Sets, Vol. I, Colloq. Math. Soc. János Bolyai 10, A. Hajnal et al. (eds.), North-Holland, 1975, 137–158.
- [Bl10] A. Blass, *Combinatorial cardinal characteristics of the continuum*, in: Handbook of Set Theory, Vol. 1, M. Foreman and A. Kanamori (eds.), Springer, 2010, 395–489.
- [C85] J. Calbrix, *Classes de Baire et espaces d'applications continues*, C. R. Acad. Sci. Paris Sér. I Math. 301 (1985), 759–762.
- [C88] J. Calbrix, *Filtres boréliens sur l'ensemble des entiers et espaces des applications continues*, Rev. Roumaine Math. Pures Appl. 33 (1988), 655–661.
- [E94] F. van Engelen, *On Borel ideals*, Ann. Pure Appl. Logic 70 (1994), 177–203.
- [F00] I. Farah, *Analytic quotients: theory of liftings for quotients over analytic ideals on the integers*, Mem. Amer. Math. Soc. 148 (2000), no. 702.
- [Fl09] J. Flašková, *Ideals and sequentially compact spaces*, Topology Proc. 33 (2009), 107–121.
- [Fl10] J. Flašková, *The relation of rapid ultrafilters and Q -points to van der Waerden ideal*, Acta Univ. Carolin. Math. Phys. 51 (2010), Suppl., 19–27.

- [Hb12] L. J. Halbeisen, *Combinatorial Set Theory. With a Gentle Introduction to Forcing*, Springer, London, 2011.
- [Hr11] M. Hrušák, *Combinatorics of filters and ideals*, in: *Set Theory and Its Applications*, Contemp. Math. 533, Amer. Math. Soc., 2011, 29–69.
- [J03] T. Jech, *Set Theory. The Third Millennium Edition, Revised and Expanded*, Springer Monogr. Math., Springer, Berlin, 2003.
- [K95] A. S. Kechris, *Classical Descriptive Set Theory*, Springer, 1995.
- [Ma91] K. Mazur, *F_σ -ideals and $\omega_1\omega_1^*$ -gaps in the Boolean algebras $P(\omega)/I$* , Fund. Math. 138 (1991), 103–111.
- [Me09] D. Meza-Alcántara, *Ideals and filters on countable sets*, Ph.D. thesis, Universidad Nacional Autónoma de México, 2009.
- [So99] S. Solecki, *Analytic ideals and their applications*, Ann. Pure Appl. Logic 99 (1999), 51–72.
- [Sm77] R. C. Solomon, *Families of sets and functions*, Czechoslovak Math. J. 27 (102) (1977), 556–559.
- [Sk08] L. Soukup, *Nagata’s conjecture and countably compact hulls in generic extensions*, Topology Appl. 155 (2008), 347–353.
- [Sz75] E. Szemerédi, *On sets of integers containing no k elements in arithmetic progression*, Acta Arith. 27 (1975), 199–245.
- [VW98] B. Velickovic and W. H. Woodin, *Complexity of reals in inner models of set theory*, Ann. Pure Appl. Logic 92 (1998), 283–295.
- [Z90] S. Zafrany, *On analytic filters and prefilters*, J. Symbolic Logic 55 (1990), 315–322.

Barnabás Farkas
 Kurt Gödel Research Center for Mathematical Logic
 Vienna, Austria
 E-mail: barnabasfarkas@gmail.com

Yurii Khomskii
 University of Hamburg
 Hamburg, Germany
 E-mail: yurii@deds.nl

Zoltán Vidnyánszky
 Alfréd Rényi Institute of Mathematics
 Budapest, Hungary
 E-mail: vidnyanszky.zoltan@renyi.mta.hu