

Boolean algebras in which every chain and antichain is countable

by

J. E. Baumgartner* (Hanover, N. H.) and P. Komjáth (Budapest)

Abstract. Using the combinatorial axiom \diamond , we deduce the existence of uncountable Boolean algebras in which every chain and antichain is countable. We find examples of such algebras for which the automorphism group has either 1 or 2^{\aleph_0} elements, and we apply a result of Kunen and Tall to conclude that the existence of such algebras is relatively consistent with the negation of the continuum hypothesis.

We also prove that if B is a Boolean algebra in which every antichain is countable then B has a countable dense subalgebra, and if B is atomless then the automorphism group of B has either 1 or 2^{\aleph_0} elements.

0. Introduction. A *chain* in a Boolean algebra is a set which is linearly ordered by the canonical partial ordering associated with the algebra; an *antichain* is a set of pairwise incomparable elements of the algebra. Some authors use the word "antichain" to denote a set of pairwise *disjoint* elements; note that this is not the case here.

Using the combinatorial axiom \diamond , we deduce the existence of uncountable Boolean algebras in which every chain and antichain is countable. We find examples of such algebras for which the automorphism group has either 1 or 2^{\aleph_0} elements, and we apply a result of Kunen and Tall to conclude that the existence of such algebras is relatively consistent with the negation of the continuum hypothesis.

We also prove that if B is a Boolean algebra in which every antichain is countable then B has a countable dense subalgebra, and if B is atomless then the automorphism group of B has either 1 or 2^{\aleph_0} elements.

Our set-theoretic terminology is standard. If $C \subseteq \omega_1$, then C is *closed unbounded* iff $\forall \alpha < \omega_1, \exists \beta \in C, \alpha \leq \beta$ and $\forall \alpha < \omega_1, \sup(C \cap \alpha) \in C$. If $S \subseteq \omega_1$, S is *stationary* iff $S \cap C \neq \emptyset$ for every closed unbounded set C . Fodor's Theorem [4] asserts that if S is stationary, $f: S \rightarrow \omega_1$, and $f(\alpha) < \alpha$ for all $\alpha \in S$, then there is stationary $S' \subseteq S$ such that f is constant on S' .

A *set mapping* on a set X is a function f such that for all $x \in X, f(x) \subseteq X - \{x\}$. A set $Y \subseteq X$ is *free* (with respect to f) iff $\forall y, z \in Y, z \notin f(y)$. Hajnal (see [5] or

* Research partially supported by National Science Foundation grant MCS76-08231.

[3, § 44]) has proved that if X has cardinality κ , $f(x)$ has cardinality $< \lambda$ for all $x \in X$, and $\lambda < \kappa$, then there is a free set of cardinality κ . We will need this result in the next section for $\kappa = \aleph_2$ and $\lambda = \aleph_1$.

The proposition \Diamond asserts that there is a sequence $\langle S_\alpha : \alpha < \omega_1 \rangle$ such that $S_\alpha \subseteq \alpha$ and for any $S \subseteq \omega_1$, $\{\alpha : S \cap \alpha = S_\alpha\}$ is stationary. We refer to $\langle S_\alpha : \alpha < \omega_1 \rangle$ as a \Diamond -sequence. Jensen has shown that \Diamond is true in L , the universe of constructible sets. See [2].

1. Statement of results. An element b of a Boolean algebra B will be called *uncountable* iff $\{a \in B : a \leq b\}$ is uncountable; otherwise b is *countable*.

The main results of the paper are the following:

THEOREM 1. Assume \Diamond . Then there is an atomless field of subsets of ω such that every nonzero element is uncountable and every chain and antichain is countable.

THEOREM 2. Assume \Diamond . Then there is an uncountable atomless field of subsets of ω such that the countable elements form a maximal ideal and every antichain (and hence every chain) is countable.

Theorems 1 and 2 are proved in Sections 2 and 3, respectively.

The next theorem shows that it is no accident that fields of subsets of ω are involved in Theorems 1 and 2.

A subalgebra D of a Boolean algebra B is *dense* in B iff for every nonzero $b \in B$ there is nonzero $d \in D$ such that $d \leq b$.

THEOREM 3. Let B be a Boolean algebra in which every antichain is countable. Then B has a countable dense subalgebra, and hence B is representable as a field of subsets of ω .

Proof. Suppose not. Then it is easy to obtain a sequence $\langle b_\alpha : \alpha < \omega_1 \rangle$ of nonzero elements of B such that for each α , if B_α is the subalgebra of B generated by $\{b_\beta : \beta < \alpha\}$, then there is no nonzero $d \in B_\alpha$ such that $d \leq b_\alpha$. Let

$$Z = \{\alpha < \omega_1 : \exists b \in B_\alpha \ b \wedge b_\alpha, b_\alpha - b \neq 0\}.$$

Case 1. Z is stationary. Let $\langle c_\alpha : \alpha < \omega_1 \rangle$ enumerate $\bigcup \{B_\alpha : \alpha < \omega_1\}$ and for each $\alpha \in Z$ let $f(\alpha)$ be the least ordinal β such that $b_\alpha \wedge c_\beta, b_\alpha - c_\beta \neq 0$. Let $C = \{\alpha : B_\alpha = \{c_\beta : \beta < \alpha\}\}$. It is clear that C is closed unbounded, so $C \cap Z$ is stationary. Moreover, $f(\alpha) < \alpha$ for every $\alpha \in C \cap Z$. By Fodor's Theorem there is stationary $S \subseteq C \cap Z$ on which f is constant.

Let $c = c_{f(\alpha)}$, where $\alpha \in S$ is arbitrary. For each $\alpha \in S$ let $d_\alpha = (c \wedge b_\alpha) \vee (\bar{c} \wedge \bar{b}_\alpha)$ (\bar{c} denotes the complement of c). Then we claim $\{d_\alpha : \alpha \in S\}$ is an uncountable set of pairwise incomparable elements, contradiction. Suppose $\alpha, \beta \in S$ and $\alpha < \beta$. If $d_\alpha \leq d_\beta$, then $c \wedge b_\alpha \leq b_\beta$ and $c \wedge b_\alpha \in B_\beta$, contradicting the choice of b_β . If $d_\beta \leq d_\alpha$ then $\bar{c} \wedge \bar{b}_\beta \leq \bar{c} \wedge \bar{b}_\alpha$ so $\bar{c} \wedge b_\alpha \leq \bar{c} \wedge b_\beta \leq b_\beta$ and we reach a similar contradiction.

Case 2. Z is nonstationary. Then $Y = \omega_1 - Z$ is uncountable, and if $\alpha, \beta \in Y$ and $\alpha < \beta$ then either $b_\alpha \wedge b_\beta = 0$ or $b_\beta \leq b_\alpha$.

Let $W = \{\alpha \in Y : \{\beta \in Y : b_\beta \leq b_\alpha\} \text{ is uncountable}\}$. If W is countable, then we construct $\langle b_{\alpha_\xi} : \xi < \omega_1 \rangle$ inductively so that $\alpha_\xi \in Y - W$ and $b_{\alpha_\xi} \not\leq b_{\alpha_\eta}$ for all $\eta < \xi$.

But then $\{b_{\alpha_\xi} : \xi < \omega_1\}$ is an uncountable pairwise disjoint set. Hence W is uncountable.

If $\{b_\alpha : \alpha \in W\}$ is a chain then let $\langle c_\alpha : \alpha < \omega_1 \rangle$ enumerate $\{b_\alpha : \alpha \in W\}$ in decreasing order (i.e. $\alpha < \beta \Rightarrow c_\beta < c_\alpha$). Then $\{c_\alpha - c_{\alpha+1} : \alpha < \omega_1\}$ is an uncountable pairwise disjoint set, contradiction. Hence $\{b_\alpha : \alpha \in W\}$ is not a chain.

It follows that there are $\alpha, \beta \in W$ with $b_\alpha \wedge b_\beta = 0$. Now let

$$f: \omega_1 \rightarrow \{\gamma \in Y : b_\gamma \leq b_\alpha\}$$

and $g: \omega_1 \rightarrow \{\gamma \in Y : b_\gamma \leq b_\beta\}$ be one-to-one and such that if $\xi < \eta$ then $f(\xi), g(\xi) < f(\eta), g(\eta)$. For each ξ let $d_\xi = b_{f(\xi)} \vee (b_\beta - b_{g(\xi)})$. Then if $\xi \neq \eta$ it is easy to check that d_ξ and d_η are incomparable. This contradiction completes the proof that B has a countable dense subalgebra.

Since any countable Boolean algebra is representable as a field of subsets of ω , the second assertion follows. ■

It should be remarked that this argument generalizes almost *verbatim* to larger cardinals.

It follows from Theorem 3 that every Boolean algebra with no uncountable antichains must have cardinality $\leq 2^{\aleph_0}$. Must every uncountable such algebra have cardinality \aleph_1 ? Without using the continuum hypothesis we have been unable to settle this question, but we can say that the number of countable elements is not too large.

THEOREM 4. Suppose B is a Boolean algebra in which every antichain has cardinality $\leq \aleph_1$. Then B has at most \aleph_1 countable elements.

Proof. Suppose X were a set of \aleph_2 countable elements of B . For each $b \in X$, let $f(b) = \{c \in X : c < b\}$. By Hajnal's set-mapping theorem there is a free set $Y \subseteq X$ of cardinality \aleph_2 . But the elements of Y are clearly pairwise incomparable, contradiction. ■

A Boolean algebra is *rigid* if it has no automorphisms except the identity.

THEOREM 5. Let B be a Boolean algebra in which every antichain is countable.

(a) If every nonzero element of B is uncountable, then B is rigid.

(b) If B is atomless and B contains a nonzero countable element, then B has exactly 2^{\aleph_0} automorphisms.

Proof. (a) If B had a nontrivial automorphism f , then there would be nonzero $b \in B$ such that $f(b) \wedge b = 0$. But then $\{c \vee (f(b) - f(c)) : c < b\}$ would be an uncountable set of pairwise incomparable elements, contradiction.

(b) Any countable atomless Boolean algebra has 2^{\aleph_0} automorphisms so if $b \in B$ is nonzero and countable, there are at least 2^{\aleph_0} automorphisms of B fixing b . On the other hand, by Theorem 3, B has a countable dense subalgebra D . Every automorphism of B is completely determined by its values on D , and there are only 2^{\aleph_0} possible functions mapping D into B . ■

COROLLARY 6. Assuming \Diamond , there are uncountable Boolean algebras with no

uncountable chains or antichains which are rigid, and there are such algebras with the property that every uncountable atomless subalgebra has exactly 2^{\aleph_0} automorphisms.

Finally, we observe that by a result of Kunen and Tall [7, Theorem 11], if B is a Boolean algebra with no uncountable chain or antichain then B remains such an algebra in any extension of the universe via a property (K) notion of forcing. Since the continuum hypothesis may be violated by a property (K) notion of forcing (by adding many Cohen reals, for example), the conclusions of Theorems 1 and 2 and Corollary 6 are relatively consistent with the negation of the continuum hypothesis.

Several problems remain.

PROBLEM 1. Can our uses of \diamond be replaced by CH? Using CH, E. S. Berney [1] has constructed an uncountable Boolean algebra with no uncountable antichains, but his algebra has uncountable chains.

PROBLEM 2. Is it consistent that every uncountable Boolean algebra has an uncountable antichain? It is conceivable that Martin's Axiom implies that there is no uncountable collection of subsets of ω in which all chains and antichains (with respect to inclusion) are countable. Kunen [6] (and the first author, independently) used CH to construct a sequence $\langle A_\alpha : \alpha < \omega_1 \rangle$ of subsets of ω such that if $\alpha < \beta$ then $A_\beta - A_\alpha$ is finite and $\{A_\alpha : \alpha < \omega_1\}$ has no uncountable chains or antichains, and Kunen proved that under Martin's Axiom no such sequence exists, but that is the best result to date.

PROBLEM 3. Is it provable in ZFC that every uncountable Boolean algebra with no uncountable antichains has cardinality \aleph_1 ?

PROBLEM 4. Can Theorems 1 and 2 be generalized to larger cardinals?

2. Proof of Theorem 1. The Boolean algebra of Theorem 1 will be obtained as the union of a sequence $\langle B_\alpha : \alpha < \omega_1 \rangle$ of countable atomless fields of subsets of ω . For each α , $B_{\alpha+1}$ will be generated by B_α together with a single subset x_α of ω , and if α is a limit ordinal then B_α will be the union of the preceding B_β 's. The proposition \diamond will be used to ensure that every potential uncountable chain or antichain is considered at some point.

The only difficulty lies in showing that if a countable set M of maximal chains and antichains in B_α is specified, then x_α can be chosen so that every element of M remains maximal in $B_{\alpha+1}$. This is done in Lemma 2.6. The set x_α is constructed essentially by a forcing argument, as the reader familiar with forcing will see, but no knowledge of forcing is necessary to follow the proof.

Given a partial ordering (P, \leq) , a set $D \subseteq P$ is called *dense* in P if $\forall p \in P \exists q \in D q \leq p$.

For the purpose of the following lemmas, B is always a countable atomless field of subsets of ω . We shall be interested in the partial ordering

$$P = \{(a, b) : a, b \in B, a \subseteq b, b - a \neq \emptyset\},$$

where $(a_1, b_1) \leq (a_2, b_2)$ iff $a_2 \subseteq a_1$ and $b_1 \subseteq b_2$.

LEMMA 2.1. Let m be a maximal antichain in B , and let $D_1(m) = \{(a, b) \in P : \forall x \subseteq \omega \text{ if } a \subseteq x \subseteq b \text{ then } x \text{ is comparable with some element of } m\}$. Then $D_1(m)$ is dense in P .

Proof. Let $(a, b) \in P$. Since B is atomless there are disjoint non-empty $a_1, a_2 \in B$ such that $a_1 \cup a_2 = b - a$. Let $c \in m$ be such that $a \cup a_1$ is comparable with c . If $c \subseteq a \cup a_1$, then $(a \cup a_1, b) \in D_1(m)$. If $a \cup a_1 \subseteq c$, then $(a, a \cup a_1) \in D_1(m)$. ■

LEMMA 2.2. Let m be a maximal chain in B , and let $D_2(m) = \{(a, b) \in P : \forall x \subseteq \omega \text{ if } a \subseteq x \subseteq b \text{ then } x \text{ is incomparable with some element of } m\}$. Then $D_2(m)$ is dense in P .

Proof. Let $(a, b) \in P$. Since B is atomless, there are disjoint non-empty $a_1, a_2, a_3 \in B$ such that $a_1 \cup a_2 \cup a_3 = b - a$. If $a \cup a_1 \in m$ then $(a \cup a_2, b - a_1) \in D_2(m)$. If $a \cup a_1 \notin m$ then there is $c \in m$ such that c and $a \cup a_1$ are incomparable. If $c \cap (a_2 \cup a_3) = \emptyset$ then $(a \cup a_1, b) \in D_2(m)$. If $c \cap a_2 \neq \emptyset$ then $(a \cup a_1, b - a_2) \in D_2(m)$, and if $c \cap a_3 \neq \emptyset$ then $(a \cup a_1, b - a_3) \in D_2(m)$. ■

LEMMA 2.3. Let m be a maximal antichain in B , and let $e, f \in B$. Let $D_1(m, e, f) = \{(a, b) \in P : \forall x \subseteq \omega \text{ if } a \subseteq x \subseteq b \text{ then } (e \cap x) \cup (f - x) \text{ is comparable with some element of } m\}$. Then $D_1(m, e, f)$ is dense in P .

Proof. Let $(a, b) \in P$. It is easy to see that there is $(a', b') \leq (a, b)$ such that one of the following holds.

- (1) $b' - a' \subseteq \omega - (e \cup f)$,
- (2) $b' - a' \subseteq e \cap f$,
- (3) $b' - a' \subseteq e - f$,
- (4) $b' - a' \subseteq f - e$.

If (1) or (2) holds, then clearly $\forall x \subseteq \omega$ if $a' \subseteq x \subseteq b'$ then $(e \cap x) \cup (f - x) \in B$, so $(a', b') \in D_1(m, e, f)$.

If (3) or (4) holds, then let $c = (e \cap a') \cup (f - b')$ and let $d = (e \cap b') \cup (f - a')$. Then $d - c \neq \emptyset$ so $(c, d) \in P$. By Lemma 2.1 there is $(c', d') \in D_1(m)$ such that $(c', d') \leq (c, d)$. If (3) holds, then $(a' \cup (c' - c), b' - (d - d')) \in D_1(m, e, f)$, while if (4) holds, then $(a' \cup (d - d'), b' - (c' - c)) \in D_1(m, e, f)$. ■

LEMMA 2.4. Let m be a maximal chain in B , and let $e, f \in B$. Let $D_2(m, e, f) = \{(a, b) \in P : \forall x \subseteq \omega \text{ if } a \subseteq x \subseteq b \text{ then } (e \cap x) \cup (f - x) \text{ either lies in } m \text{ or else is incomparable with some element of } m\}$. Then $D_2(m, e, f)$ is dense in P .

Proof. Like Lemma 2.3, but using Lemma 2.2 instead of Lemma 2.1. ■

LEMMA 2.5. Let $c \in B$, and let $D_3(c) = \{(a, b) \in P : \forall x \subseteq \omega \text{ if } a \subseteq x \subseteq b \text{ then } x \neq c\}$.

Proof. By Lemma 2.2, letting m be a maximal chain containing c . ■

LEMMA 2.6. Let M be a countable collection of maximal chains and antichains in B . Then for any $(a, b) \in P$ there is $x \notin B$ such that $a \subseteq x \subseteq b$ and if B' is the field of sets generated by $B \cup \{x\}$, then every element of M remains maximal in B' .

Proof. Let D_0, D_1, \dots enumerate all sets of the form $D_1(m, e, f)$, $D_2(m, e, f)$, and $D_3(c)$ where $m \in M$ and $c, e, f \in B$. Form a sequence $\langle (a_n, b_n) : n \in \omega \rangle$ as follows: Let $a_0 = a$, $b_0 = b$. Given (a_n, b_n) , let $(a_{n+1}, b_{n+1}) \in D_n$ be such that $(a_{n+1}, b_{n+1}) \leq (a_n, b_n)$. Let $x = \bigcup \{a_n : n \in \omega\}$. Note that every element of B' has the form $(e \cap x) \cup (f - x)$ for some $e, f \in B$. But now by Lemmas 2.3–2.5 it is clear that

$x \notin B$ and every element of M remains maximal in B' . It is easy to see that B' is atomless. ■

Proof of Theorem 1. Let $\langle S_\alpha: \alpha < \omega_1 \rangle$ be a \diamond -sequence. Since \diamond implies the continuum hypothesis, there is an enumeration $\langle a_\alpha: \alpha < \omega_1 \rangle$ of the power set of ω in which each element occurs uncountably many times.

By induction, we construct a sequence $\langle B_\alpha: \alpha < \omega_1 \rangle$ of countable atomless fields of subsets of ω , and a sequence $\langle M_\alpha: \alpha < \omega_1 \rangle$ where M_α is a finite or countably infinite set of maximal chains and antichains in B_α .

Let B_0 be arbitrary, and let $M_0 = \emptyset$. If α is a limit ordinal, let $B_\alpha = \bigcup \{B_\beta: \beta < \alpha\}$ and $M_\alpha = \bigcup \{M_\beta: \beta < \alpha\}$. Finally, suppose $\alpha = \beta + 1$. If $\{a_\xi: \xi \in S_\beta\}$ is a maximal chain or antichain in B_β then let $M_{\beta+1} = M_\beta \cup \{\{a_\xi: \xi \in S_\beta\}\}$; otherwise let $M_{\beta+1} = M_\beta$. Choose $x_\beta \notin B_\beta$ by Lemma 2.6 so that if $B_{\beta+1}$ is generated by $B_\beta \cup \{x_\beta\}$ then every chain or antichain in $M_{\beta+1}$ remains maximal in $B_{\beta+1}$. Moreover, if $a_\beta \in B_\beta$ then we may assume $x_\beta \leq a_\beta$.

Let $B = \bigcup \{B_\alpha: \alpha < \omega_1\}$. It is clear that every nonzero element of B is uncountable. Suppose $m \subseteq B$ is a maximal chain. Let $S = \{\alpha: a_\alpha \in m\}$. Then it is easy to see that $\{\alpha: \{a_\beta: \beta \in S \cap \alpha\}$ is a maximal chain in $B_\alpha\}$ is closed and unbounded. Since $\{\alpha: S_\alpha = S \cap \alpha\}$ is stationary there is α such that $S_\alpha = S \cap \alpha$ and $\{a_\beta: \beta \in S_\alpha\}$ is a maximal chain in B_α . But then $\{a_\beta: \beta \in S_\alpha\} \in M_\gamma$ for every $\gamma \geq \alpha$ so $\{a_\beta: \beta \in S_\alpha\}$ is maximal in B . Hence $m = \{a_\beta: \beta \in S_\alpha\}$, and m is countable. It can be shown similarly that all antichains are countable. This completes the proof. ■

3. Proof of Theorem 2. We will obtain our Boolean algebra as the union of a sequence $\langle B_\alpha: \alpha < \omega_1 \rangle$ as before, but in addition we will construct a sequence $\langle I_\alpha: \alpha < \omega_1 \rangle$, where I_α is a maximal ideal in B_α and $I_\alpha \subseteq I_\beta$ whenever $\alpha \leq \beta$. It will turn out that $\bigcup \{I_\alpha: \alpha < \omega_1\}$ is the maximal ideal of countable elements in the Boolean algebra.

In Lemmas 3.1–3.7, B will always be a countable atomless field of subsets of ω , and I will be a maximal ideal in B . The partial ordering P to be used this time is $P = \{(a, b): a \in I, b \in B - I, \text{ and } a \subseteq b\}$, ordered as before, i.e., $(a_1, b_1) \leq (a_2, b_2)$ iff $a_1 \supseteq a_2$ and $b_1 \subseteq b_2$.

LEMMA 3.1. Let $m \subseteq I$, and let $D_1(m) = \{(a, b) \in P: \text{either } \forall c \in m \ c \not\subseteq b \text{ or } \exists c \in m \ c \subseteq a\}$. Then $D_1(m)$ is dense in P .

Proof. Let $(a, b) \in P$. If $(a, b) \notin D_1(m)$, then $\exists c \in m \ c \subseteq b$. But then $(a \cup c, b)$ is in $D_1(m)$. ■

LEMMA 3.2. Let $c \in B$ and let $D_2(c) = \{(a, b) \in P: \forall x \subseteq \omega \text{ if } a \subseteq x \subseteq b \text{ then } x \neq c\}$. Then $D_2(c)$ is dense in P .

LEMMA 3.3. Let $c \in I$ and let $D_3(c) = \{(a, b) \in P: c \subseteq a \cup (\omega - b)\}$. Then $D_3(c)$ is dense in P .

LEMMA 3.4. Suppose D is dense in P . Let $S(D) = \{(a, b) \in P: (\omega - b, \omega - a) \in D\}$. Then $S(D)$ is dense in P .

LEMMA 3.5. Suppose D is dense in P and $e, f \in I$. Let

$$T(D, e, f) = \{(a, b) \in P: ((a - e) \cup f, (b - e) \cup f) \in D\}.$$

Then $T(D, e, f)$ is dense in P .

Proof. Let $(a, b) \in P$. By Lemma 3.3 there is $(a', b') \leq (a, b)$ such that $e \cup f \subseteq a' \cup (\omega - b')$. Let $e' = a' \cap e$, $f' = a' \cap f$. Find $(x, y) \in D$ such that $(x, y) \leq ((a' - e) \cup f, (b' - e) \cup f)$. Now let $a'' = (x - (e \cup f)) \cup e' \cup f'$, and $b'' = (y - (e \cup f)) \cup e' \cup f'$. Then it is easy to see $(a'', b'') \in T(D, e, f)$. ■

Now suppose M is a countable (or finite) collection of subsets of B , and let \mathcal{D} be the smallest collection of dense sets such that

- (a) every set of the form $D_1(m)$, $D_2(c)$, $D_3(e)$, for $m \in M$, $c \in B$, $e \in I$, lies in \mathcal{D} , and
- (b) if $D \in \mathcal{D}$ and $e, f \in I$, then $S(D)$, $T(D, e, f) \in \mathcal{D}$.

Let us call $x \subseteq \omega$ (M, I) -generic over B iff $\forall D \in \mathcal{D} \exists (a, b) \in D \ a \subseteq x \subseteq b$. Since \mathcal{D} is countable it is clear that there exists $x \subseteq \omega$ which is (M, I) -generic over B .

LEMMA 3.6. Let x be (M, I) -generic over B , and let B' be the field of sets generated by $B \cup \{x\}$. Then every element of $B' - B$ is (M, I) -generic over B .

Proof. It is evident from condition (b) above and Lemmas 3.4 and 3.5 that if x is (M, I) -generic over B , then so are $(x - e) \cup f$ and $\omega - x$, if $e, f \in I$. To complete the proof, it will suffice to show that any element of $B' - B$ has the form $(x - e) \cup f$ or $((\omega - x) - e) \cup f$ for some $e, f \in I$.

First note that if x is (M, I) -generic over B , then for any $a \in I$, both $a \cap x$ and $a - x$ are in I . This follows immediately from Lemma 3.3.

We observed in the last section that every element of $B' - B$ has the form $(e \cap x) \cup (f - x)$ for some $e, f \in B$. But by genericity of x and $\omega - x$, the remark in the preceding paragraph shows that we cannot have both $e, f \in I$, or both $e, f \notin I$, since otherwise $(e \cap x) \cup (f - x) \in B$.

If $f \in I$ and $e \notin I$ then $(e \cap x) \cup (f - x) = (x - (\omega - e)) \cup (f - x)$, where both $\omega - e$ and $f - x$ are in I . If $e \in I$ and $f \notin I$ then $(e \cap x) \cup (f - x) = ((\omega - x) - (\omega - f)) \cup (e \cap x)$ where $\omega - f$, $e \cap x \in I$, and we are done. ■

LEMMA 3.7. Under the hypotheses of Lemma 3.6, $I \cup \{x\}$ generates a maximal ideal I' in B' and $\forall a \in I \ \forall b \in B'$ if $b \subseteq a$ then $b \in B$.

Proof. We check the last assertion first. If $b \in B' - B$, then b is (M, I) -generic over B , so as observed in the preceding proof, if $a \in I$ then $a \cap b \in I$, so we cannot have $b \subseteq a$.

For the rest, suppose $x \cup a = \omega$ for some $a \in I$. Then $x = \omega - (a - x)$, and since $a - x \in I$, we would have $x \in B$, contradiction. Hence I' is a proper ideal. It remains only to check maximality. Let $y \in B' - B$. If $y = (x - e) \cup f$ for some $e, f \in I$ then $y \in I$. If not, then $y = ((\omega - x) - e) \cup f$ for $e, f \in I$. But then $\omega - y = (x - f) \cup (e - f) \in I$. ■

Proof of Theorem 2. Let $\langle S_\alpha: \alpha \in \omega_1 \rangle$ be a \diamond -sequence, and let $\langle A_\alpha: \alpha < \omega_1 \rangle$ enumerate the power set of ω . For each $\beta < \omega_1$, let $m_\beta = \{a_\alpha: \alpha \in S_\beta\}$.

We obtain sequences $\langle B_\alpha: \alpha < \omega_1 \rangle$, $\langle I_\alpha: \alpha < \omega_1 \rangle$, and $\langle M_\alpha: \alpha < \omega_1 \rangle$ by induction as follows: Let B_0 be an arbitrary countable atomless field of subsets of ω , and let I_0 be a maximal ideal in B_0 . Given B_α and I_α , a maximal ideal in B_α , let $M_\alpha = \{m_\beta: \beta \leq \alpha \text{ and } m_\beta \subseteq B_\alpha\}$. Let x_α be (M_α, I_α) -generic over B_α and let $B_{\alpha+1}$ be generated by $B_\alpha \cup \{x_\alpha\}$, and let $I_{\alpha+1}$ be generated by $I_\alpha \cup \{x_\alpha\}$. If α is a limit ordinal, let $B_\alpha = \bigcup \{B_\beta: \beta < \alpha\}$ and $I_\alpha = \bigcup \{I_\beta: \beta < \alpha\}$. Let $B = \bigcup \{B_\alpha: \alpha < \omega_1\}$ and $I = \bigcup \{I_\alpha: \alpha < \omega_1\}$.

It is easy to see that each B_α is atomless so B is an atomless field of subsets of ω . It follows from Lemma 3.7 that I is a maximal ideal of countable elements of B . It remains only to show that B has no uncountable antichains.

Suppose $m \subseteq I$ were an uncountable antichain. Let $S = \{\alpha: a_\alpha \in m\}$. Let $Z = \{\alpha: \{a_\beta: \beta \in S \cap \alpha\} = m \cap B_\alpha \text{ and } (\forall b \in B_\alpha - I_\alpha) \text{ if } \exists c \in m \text{ } c \subseteq b \text{ then } (\exists \beta \in S \cap \alpha) a_\beta \subseteq b\}$. It is easy to see that Z is closed and unbounded. Hence there is $\alpha \in Z$ such that $S_\alpha = S \cap \alpha$.

We assert that $m = m_\alpha$. This will show that there are no uncountable antichains included in I , and hence that there are no uncountable antichains in B (since if $m \subseteq B - I$ were an antichain, so would be $\{\omega - a: a \in m\} \subseteq I$).

Clearly $m_\alpha \subseteq m$. For each $c \in B - B_0$, let $q(c)$ be the least ordinal β such that $c \in B_{\beta+1} - B_\beta$. Suppose there exists $c \in m - m_\alpha$. Choose such a c with $q(c)$ minimal. Let $q(c) = \beta \geq \alpha$. Then c is (M_β, I_β) -generic over B_β so since $m_\alpha \in M_\beta$ there exists $(a, b) \in D_1(m_\alpha)$ (computed in B_β) such that $a \subseteq c \subseteq b$. Since c is not comparable with any element of m_α it must be the case that $\forall c' \in m_\alpha \text{ } c' \not\subseteq b$. We may assume that $q(b)$ is minimal for $b \in B_\beta$ such that $a \subseteq c \subseteq b$ and $\forall c' \in m_\alpha \text{ } c' \not\subseteq b$.

Now $q(b) > \alpha$ since otherwise $\alpha \in Z$ implies that $\forall c' \in m \text{ } c' \not\subseteq b$, contradiction. Say $q(b) = \gamma \geq \alpha$. Then b is (M_γ, I_γ) -generic over B_γ , and there must be $(a', b') \in D_1(m_\alpha)$ (computed in B_γ) such that $a' \subseteq b \subseteq b'$. But then clearly we must have $\forall c' \in m_\alpha \text{ } c' \not\subseteq b'$ and $q(b') < \gamma$, contradicting minimality of $q(b)$. Hence B has no uncountable antichains.

It follows that B has no uncountable chains, for if m were an uncountable chain we could, as above, assume that $m \subseteq I$. But since all the elements of I are countable, m must have a subset well-ordered in type ω_1 , and this contradicts the fact that B has no uncountable antichains. ■

Added in proof December 3, 1980. Most of the problems have been solved. Shelah found an uncountable algebra with no uncountable chains or antichains using only CH; the first author showed it consistent that every uncountable algebra has an uncountable antichain; Shelah and Van Wesep independently answered Problem 3 negatively; and Shelah has generalized the \diamond -style constructions to larger cardinals.

References

- [1] E. S. Berney, preprint.
- [2] K. Devlin, *Aspects of Constructibility*, Lecture Notes in Mathematics, vol. 354.

- [3] P. Erdős, A. Hajnal, A. Máté and R. Rado, *Combinatorial Set Theory: Partition Relations for Cardinals*, manuscript.
- [4] G. Fodor, *Eine Bemerkung zur Theorie der regressiven Funktionen*, Acta Sci. Math. 17 (1956), pp. 139–142.
- [5] A. Hajnal, *Proof of a conjecture of S. Ruziewicz*, Fund. Math. 50 (1961), pp. 123–128.
- [6] K. Kunen, mimeographed notes.
- [7] — and F. Tall, *Between Martin's Axiom and Souslin's Hypothesis*, preprint.

DARTMOUTH COLLEGE
Hanover, New Hampshire
EÖTVÖS LORAND UNIVERSITY
Budapest

Accepté par la Rédaction le 26. 10. 1978