ON SMALL DIVISORS

A. D. BRUNO

*Institute of Applied Mathematics, Academy of Sciences of the U.S.S.R.
 Moscow, U.S.S.R.*

Let us consider a one-dimensional complex point mapping

$$(1) \quad x^* = ax + \varphi(x), \quad x \in \mathbb{C}$$

near the fixed point $x = 0$, i.e. let $\varphi(0) = \varphi'(0) = 0$. We shall assume that φ is an analytic function at zero, i.e. it can be expanded into a power series which converges in a domain $|x| < \varepsilon$.

Let us consider the case where

$$a^n \neq 1, \quad n = 1, 2, 3, \dots$$

for all natural numbers n . Then there exists a formal invertible change of the coordinate

$$(2) \quad x = y + \xi(x), \quad \xi(0) = \xi'(0) = 0,$$

which reduces the mapping (1) to a normal form

$$(3) \quad y^* = ay.$$

Here $\xi(y)$ is a formal power series which can converge in some neighbourhood $|y| < \varepsilon_1$ of zero or can diverge in every such neighbourhood. We will discuss conditions for convergence.

If $|a| \neq 1$ then the series ξ always converges. Thus we need only consider the case of

$$|a| = 1.$$

Cremer [1] proved that if the number a satisfies

$$(4) \quad \lim_{n \rightarrow \infty} \sqrt[n]{|a^n - 1|} = 0$$

then there exists an analytic function $\varphi(x)$ in (1), for which the series ξ in (2) diverges. Siegel [2] proved the convergence of ξ under the following condi-

tion: there exist positive constants ε, ν such that for all integral p and natural q

$$(5) \quad |p - \lambda q| > \varepsilon q^{-\nu}$$

where $a = \exp(2\pi i \lambda)$. Write $\beta_m = \min |p - \lambda q|$ for all integral p and natural $q < 2^m$. In papers [3; 4, § 5; 5] the convergence of the series ξ was established under the assumption that

$$(6) \quad \sum_{m=1}^{\infty} -\frac{\ln \beta_m}{2^m} < \infty$$

In terms of the numbers β_m , Siegel's condition (5) takes the form $\beta_m > \varepsilon 2^{-m\nu}$ for $m = 1, 2, 3, \dots$. Hence

$$\sum_{m=1}^{\infty} -\frac{\ln \beta_m}{2^m} < \sum_{m=1}^{\infty} \frac{\nu m \ln 2 - \ln \varepsilon}{2^m} < \nu \ln 2 \sum_{m=1}^{\infty} \frac{m}{2^m} - \ln \varepsilon < \infty$$

So the restriction (6) is weaker than (5). Now let us consider condition (4) in terms of β_m . For small $a^n - 1$, we have

$$a^n - 1 \approx 2\pi i (n\lambda - p)$$

for some integer p , and

$$\sqrt[n]{|a^n - 1|} \geq \sqrt[n]{2\pi} \sqrt[n]{\beta_m} \quad \text{for } n < 2^m.$$

But $\sqrt[n]{2\pi} \rightarrow 1$ if $n \rightarrow \infty$. Hence, condition (4) means that

$$(7) \quad \lim_{m \rightarrow \infty} -\frac{\ln \beta_m}{2^m} = \infty,$$

i.e. the members of the series (6) are not bounded from above.

So we have the following results.

1. If the series (6) converges then the series ξ converges and the point mapping (1) is analytically equivalent to the linear mapping (3).
2. If the members of the series (6) are not bounded then the series ξ can diverge and the mapping (1) has a more complicated structure than a linear map.

Note that there is a gap between conditions (6) and (7), when all members of the series (6) are bounded and the series (6) diverges. Up to-day (1986) we do not know whether the series ξ converge or can diverge if the number a belongs to the gap.

Now I want to explain the meaning of the term "small divisors" and the sense of conditions (6) and (7) (or (4)). The series ξ must satisfy the Schröder

equation

$$(8) \quad \xi(ay) = a\xi(y) + \varphi(y + \xi(y)).$$

Denote

$$\varphi = x \sum_{n=1}^{\infty} f_n x^n, \quad \xi = y \sum_{n=1}^{\infty} h_n y^n.$$

We shall solve equation (8) by comparison of coefficients. Then y^{n+1} occurs with the coefficient

$$(9) \quad (a^n - 1)h_n = \frac{1}{a} [f_n + P_n(f_j, h_k)]$$

Here P_n is a polynomial in $f_1, h_1, \dots, f_{n-1}, h_{n-1}$.

Let us omit the summand P_n in (9), we get

$$(10) \quad h_n = \frac{1}{a(a^n - 1)} f_n$$

The Cauchy condition for convergence of the series ξ is

$$(11) \quad \lim 1/\sqrt[n]{|h_n|} > 0$$

Since φ is an analytic function

$$(12) \quad \lim 1/\sqrt[n]{|f_n|} > 0$$

If the coefficients h_n and f_n are connected by (10) and condition (4) is fulfilled, then there exists a series φ with the property (12) such that property (11) is not fulfilled. For example, this is the case when all $f_n = 1$. Here the coefficient h_n is obtained from f_n on dividing it by one "small divisor" $a^n - 1$. Such a relation (10) is typical for linear problems and leads to conditions (4), (7).

Now let us take into account the summand P_n in (9). Then

$$h_n = \frac{1}{(a^n - 1)a} \left[f_n + \sum c_{k_1 k_2 \dots k_l} \frac{f_{k_1}}{(a^{k_1} - 1)} \frac{f_{k_2}}{(a^{k_2} - 1)} \dots \frac{f_{k_l}}{(a^{k_l} - 1)} \right]$$

where $k_1 + k_2 + \dots + k_l = n$. Here "small divisors" $a^{k_i} - 1$ appear in some products, this is typical of nonlinear problems. We can estimate such products and establish the convergence of the series ξ if we use condition (6).

Conditions (6) and (7) can be expressed in terms of the continued fraction of the number λ . Let p_k/q_k be the k th convergent of the continued fraction of λ . Then the conditions (6) and (7) are equivalent respectively to

$$(6') \quad \sum_{k=1}^{\infty} \frac{\ln q_{k+1}}{q_k} < \infty$$

and

$$(7') \quad \overline{\lim} \frac{\ln q_{k+1}}{q_k} < \infty, \quad ,$$

If condition (7) is fulfilled and the series ξ diverges then the mappings (1) and (3) are not analytically equivalent but they can be related a smooth or topological equivalence. However for a special number λ , which is too well approximated by rationals, the mapping (1) can have very complicated structure and can be nonequivalent to the linear mapping (3) in any sense. Some examples of similar situations are given in [6–9]. Here we do not have so clear conditions as for the case of analytical equivalence.

Above we have considered one of the simplest problem with “small divisors”. General problems and corresponding conditions on small divisors are discussed in my article [10]. Some proofs and a comparison of different conditions on small divisors can be found in [4, § 5].

Concerning the mapping (1) I want to add that the number a always is its topological invariant. If condition (6) is satisfied this follows from the analytical reducibility to the normal form (3). For other cases it was proved in [11]. If $a^q = 1$ and $a^n \neq 1$ for $n < q$, then the normal form of the mapping (1) is

$$y^* = ay \exp [by^{qm}(1 + cy^{qm})],$$

where b is a complex number, m is a natural number and $c = 0$ or 1. Hence the mapping (1) has three additional formal invariants b , m and c . But it has infinitely many analytical invariants (see [12–14]).

References

- [1] H. Cremer, *Über die Häufigkeit der Nichtzentren*, Math. Ann. 115 (1938), 573–580.
- [2] C. L. Siegel, *Iteration of analytic functions*, Ann. of Math. 43 (1942), 607–612. Also: *Vorlesungen über Himmelsmechanik*, Springer-Verlag, 1956, § 24 = Moscow, Inost. Literatura, 1959 (in Russian).
- [3] A. D. Bruno, *Convergence of transformations of differential equations to normal form*, Dokl. Akad. Nauk SSSR 165 (1965), 987–989 (in Russian) = Soviet Math. Dokl. 6 (1965), 1536–1538 (English translation).
- [4] —, *Analytical form of differential equations*, Trudy Moskov. Mat. Obšč. 25 (1971), 119–262 (in Russian) = Trans. Moscow Math. Soc. 25 (1971) 131–288 (in English).
- [5] H. Rüssmann, *Über die Iteration analytischer Funktionen*, J. Math. Mech. 17 (1967) 523–532.
- [6] A. N. Kolmogorov, *On dynamical systems with integral invariant on a torus*, Dokl. Akad. Nauk SSSR 93 (1953), 763–766 (in Russian).
- [7] D. V. Anosov and A. B. Katok, *New examples in the smooth ergodic theory. Ergodic diffeomorphisms*, Trudy Moskov. Mat. Obšč. 23 (1970), 3–36 (in Russian) = Trans. Moscow Math. Soc. 23 (1970), 3–37 (English translation).

- [8] D. V. Anosov, *On an additive functional equation connected with an ergodic turn of a circumference*, *Izvestija Akad. Nauk SSSR Ser. Mat.* 37 (1973), 1259–1274 (in Russian) = *Math. of the USSR – Izvestija* 7 (1973) (English translation).
  - [9] A. S. Pjartly, *The birth of complex invariant manifolds near a singular point of a vector field depending of a parameter*, *Funkcional. Anal. i Prilozen.* 6 (1972) no. 4, 95–96 (in Russian) = *Functional Anal. Appl.* 6 (1972) (English translation).
  - [10] A. D. Bruno, *Small divisors*, in *Mathematical Encyclopaedia*, vol. 3, Moscow, Soviet Encyclopaedia, 1982, 505–507 (in Russian). (Mathematical Encyclopaedia will be published in English by Reidel.)
  - [11] V. A. Naishul, *Topological invariants of analytical and area-preserving mappings and their applications to analytical differential equations in C^2 and CP^2* , *Trudy Moscov Mat. Obšč.* 44 (1982), 234–245 (in Russian) = *Trans. Moscow Math. Soc.* 44 (1982), 240–251 (English translation).
  - [12] S. M. Voronin, *The analytical classification of germs of conformal mappings $(C, 0) \rightarrow (C, 0)$ with the identical linear part*, *Funkcional. Anal. i Prilozen.* 15 (1981), no. 1, 1–17 (in Russian) = *Functional Anal. Appl.* 15 (1981), 1–15 (English translation).
  - [13] J. Ecalle, *Les fonctions resurgentes*, tt. I, II. *Publ. Math. d'Orsay*, 8105, 8106 (1981), 1–531.
  - [14] J. Martinet et J.-P. Ramis, *Problemes du modules pour des equations differentielles non lineaires du premier orde*, *Publ. Math. I.H.E.S.* 55 (1982) 63–164.
-