

Dynamics of inner functions revisited

by

JON AARONSON and MAHENDRA NADKARNI

Abstract. We study the circle restrictions of inner functions on the unit disc showing that the local invertibility of a restriction is independent of its singularity set and proving a local characterization of analytic conditional expectations.

We establish central limit properties for some stochastic processes driven by probability preserving restrictions via spectral analysis of their perturbed transfer operators.

1. Introduction.

1.1. Inner functions and their restrictions. An *inner function* on the *unit disc* $\mathbb{D} := \{z \in \mathbb{C} : |z| < 1\}$ is an analytic endomorphism $\phi : \mathbb{D} \hookrightarrow \mathbb{D}$ such that for Lebesgue almost every $\xi \in \partial\mathbb{D}$,

$$\phi(r\xi) \xrightarrow{r \rightarrow 1^-} \phi(\xi) \in \partial\mathbb{D}.$$

The *restriction* of ϕ to $\partial\mathbb{D}$ is defined λ -a.s. (where λ is Lebesgue measure on $\partial\mathbb{D}$) and is a *nonsingular transformation* of $(\partial\mathbb{D}, \lambda)$ in the sense that λ and $\lambda \circ \phi^{-1}$ have the same null sets.

This follows from Nordgren's theorem (in §2.3), which also shows the connection between the ergodic theory of an inner function restriction on $(\partial\mathbb{D}, \lambda)$ and the dynamics of the action of the inner function on $\mathbb{D}$.

For this, and more discussion of the ergodic theory of restrictions, see e.g. [Aar97, Ch. 6] and references therein.

1.2. Overview of the paper. This paper deals with the structure and properties of inner functions, the spectral theory of their transfer operators and the central limit theory of stochastic processes driven by their restrictions.

2020 *Mathematics Subject Classification*: Primary 30J05; Secondary 37C30, 60F05, 28A50.
Key words and phrases: nonsingular transformation, inner function, restriction, Clark measure, multiplicity, angular derivative, forward nonsingular, locally invertible, transfer operator, quasi-compact, characteristic function operator, perturbation, central limit theorem.

Received 30 July 2024; revised 26 October 2025.

Published online 14 July 2026.

The rest of this subsection is devoted to a description of the main results.

Local invertibility vs. singularity set. In §2.2 we consider *local invertibility* of general nonsingular maps showing i.a. that a nonsingular transformation of a standard, nonatomic probability space is locally invertible iff it is *forward nonsingular* (Theorem 2.2).

This enables an elementary proof of a multiplicity result of Aleksandrov (Proposition 2.7): an inner function has a locally invertible restriction iff it admits angular derivatives a.e. on $\partial\mathbb{D}$.

The *singularity set* of an inner function ⁽¹⁾ (see §2.6) was studied in [Sei34]. Evidently, if an inner function has a Lebesgue-null singularity set, then its restriction, being analytic a.e. on $\mathbb{T}$, is locally invertible. However (by Proposition 2.8), any closed set of $\partial\mathbb{D}$ appears as the singularity set of an inner function whose restriction is locally invertible.

Spectrum of the transfer operator and central limits. In §3 & §4, we restrict attention to non-Möbius inner functions $\phi : \mathbb{D} \hookrightarrow \mathbb{D}$ preserving an absolutely continuous probability, showing in §3 that their transfer operators have spectral gaps ⁽²⁾ on weighted Hilbert spaces (see §3.3).

In §4 we consider central limit properties of stochastic processes $(\psi \circ \tau^n : n \geq 1)$ ($\psi : \mathbb{T} \rightarrow \mathbb{R}^d$) driven by such restrictions $\tau = \tau(\phi)$. In particular, if $\phi : \mathbb{D} \hookrightarrow \mathbb{D}$ has a singularity on $\partial\mathbb{D}$ and $\psi : \mathbb{T} \rightarrow \mathbb{R}$ is real analytic and not constant, then $(\psi \circ \tau^n : n \geq 1)$ satisfies the conditional central limit theorem (as in (32)). These results rely on smooth perturbations of the quasicompact transfer operators under consideration.

The central limit theorem for measure preserving inner function restrictions (as in (22)) is established in [IU23] for e.g. nonconstant Hölder continuous functions. See also [NSiG22] for different forms of central limit theorem for measure preserving inner function restrictions.

Analytic conditional expectations. Aleksandrov [Ale86] proved that conditional expectation with respect to a sub- σ -algebra $\mathcal{C} \subset \mathcal{B}(\mathbb{T})$ is *analytic* in the sense that the conditional expectation projection commutes with the *Riesz projection* (orthogonal projection $L^2 \rightarrow H_0^2 := \{f \in H^2 : \hat{f}(0) = 0\}$) if and only if $\mathcal{C} = \tau^{-1}\mathcal{B}(\mathbb{T})$, where $\tau = \tau(\phi)$ with ϕ inner, $\phi(0) = 0$.

In §5 we prove a local version of this (Theorem 5.2).

2. Structure and basic properties of inner functions.

2.1. Nonsingular maps and transformations. A *nonsingular map* $\pi : (X, m) \rightarrow (Y, \mu)$ between the nonatomic, Polish probability spaces (X, m)

⁽¹⁾ The (closed) set of points of $\partial\mathbb{D}$ where it is not analytic.

⁽²⁾ That is, are Doeblin–Fortet operators (see Section 3.1).

and (Y, μ) is a measurable map $\pi : X \rightarrow Y$ satisfying $m \circ \pi^{-1} \sim \mu$. It is *probability preserving* if $m \circ T^{-1} = \mu$.

A *nonsingular* [probability preserving] *transformation* (X, m, T) is a nonsingular [probability preserving] self-map $T : (X, m) \rightarrow (X, m)$.

The *transfer operator* (also called “adjoint” or “predual”) of the nonsingular map $\pi : (X, m) \rightarrow (Y, \mu)$ is $\hat{\pi} : L^1(m) \rightarrow L^1(\mu)$ defined by

$$\int_A \hat{\pi} f d\mu = \int_{\pi^{-1}A} f dm, \quad A \in \mathcal{B}(X).$$

The following is a standard application of the disintegration theorem ([DM78, Ch. III]; see also [Aar97, Ch. 1]):

PROPOSITION 2.1 (Preimage measures). *Let $\pi : (X, m) \rightarrow (Y, \mu)$ be a nonsingular map. Then there exist $Y_0 \in \mathcal{B}(X)$ with $\mu(Y_0) = 1$ and $\nu = \nu^{(\pi)} : Y_0 \rightarrow \mathcal{M}(X)$ such that*

- (1) $\hat{\pi} 1_A(y) = \nu_y(A)$ for $y \in Y_0$, $A \in \mathcal{B}(X)$, and $\nu_y(X) = \frac{dm \circ \pi^{-1}}{d\mu}(x)$;
- (2) $\nu_y(X \setminus \pi^{-1}\{y\}) = 0$ and $\nu_x \perp m$ for μ -a.e. $y \in Y_0$.

The function $x \mapsto \nu_x$ is known as the *transition kernel* of $\hat{\pi}$ and the ν_x are known as *preimage measures* or *fiber measures*.

2.2. Local invertibility and forward nonsingularity. We will say that a nonsingular map $\pi : (X, m) \rightarrow (Y, \mu)$ is

- *almost countable-to-one* if there exists $Y_0 \in \mathcal{B}(Y)$ with $\mu(Y \setminus Y_0) = 0$ such that $\pi^{-1}\{y\} \cap Y_0$ is at most countable for all $y \in Y_0$;
- *locally invertible* if there exists an at most countable partition $\alpha \subset \mathcal{B}(Y)$ such that $\pi : a \rightarrow \pi a$ is invertible and nonsingular for all $a \in \alpha$;
- *forward nonsingular* if there exists $X_0 \in \mathcal{B}(X)$ and $m(X \setminus X_0) = 0$ such that

$$[A \in \mathcal{B}(X_0), m(A) = 0] \implies \mu(\pi A) = 0.$$

As shown in [Rok61] (see also [Nad81] and [Aar97, Ch. 1]), for the nonsingular map $\pi : (X, m) \rightarrow (Y, \mu)$ the “almost countable-to-one” and “locally invertible” conditions are both equivalent to the pure atomicity of the transition kernel (i.e. almost every preimage measure is purely atomic).

THEOREM 2.2. *A nonsingular map $\pi : (X, m) \rightarrow (Y, \mu)$ is forward nonsingular iff it is locally invertible.*

Proof. It is standard that local invertibility implies forward nonsingularity. We will show that forward nonsingularity implies pure atomicity of the transition kernel.

Let $y \mapsto \nu_y$ ($Y_0 \rightarrow \mathcal{M}(X)$) be the transition kernel of $\hat{\pi}$.

We claim first that the function $x \mapsto \nu_{\pi x}(\{x\})$ is measurable.

To see this let d be a Polish metric on X and let α_n be a sequence of partitions of X such that $\alpha_{n+1} \succ \alpha_n$, $\sup_{a \in \alpha_n} \text{diam } a \xrightarrow{n \rightarrow \infty} 0$. Thus $\alpha_n(y) \downarrow \{y\}$ for all $y \in X$ (where $y \in \alpha_n(y) \in \alpha_n$) and the function $y \mapsto \nu_{\pi y}(\{y\}) = \lim_{n \rightarrow \infty} \nu_{\pi y}(\alpha_n(y))$ is measurable.

It follows that $W := \{x \in X : \nu_{\pi x}(\{x\}) = 0\} \in \mathcal{B}(X)$.

If the transition kernel is not purely atomic, then $m(W) > 0$.

Next, πW is analytic, whence universally measurable, and since $\pi^{-1}\pi W \supseteq W$, we have $m(\pi^{-1}\pi W) \geq m(W) > 0$, whence (by nonsingularity of π) $\mu(\pi W) > 0$.

For $V \in \mathcal{B}(W)$,

$$\begin{aligned} m(V) &= m(V \cap W) = m(V \cap \pi^{-1}\pi W) \quad (\because \pi^{-1}\pi W \supset W) \\ &= \int_{\pi W} \widehat{\pi} 1_V d\mu = \int_{\pi W} \nu_x(V) d\mu(x). \end{aligned}$$

By the analytic section theorem [Lus30] (see also [Jan41], [vN49], [Sri98, Thm. 5.5.2], [Coh80, §8.5]), and Luzin's continuity theorem, there exists $A \subset \pi W$ compact with $\mu(A) > 0$ and $\zeta : A \rightarrow B := \zeta A \subset W$ continuous such that $\pi \circ \zeta = \text{Id}$. It follows that

$$B \cap \pi^{-1}\{x\} = \begin{cases} \emptyset, & x \notin A, \\ \{\zeta(x)\} & x \in A, \end{cases}$$

whence

$$\nu_x(B) = 1_A(x) \nu_x(\{\zeta(x)\}) = 0 \quad (\because \pi(x) \in W \text{ \& } x = \pi(\zeta(x))).$$

Thus

$$m(B) = \int_A \nu_x(\{\zeta(x)\}) d\mu(x) = 0,$$

whereas $\pi B = \pi \zeta A = A$, $\mu(\pi B) = \mu(A) > 0$ and forward nonsingularity fails. ■

2.3. Structure of inner function restrictions. It will be convenient to identify $\partial\mathbb{D}$ with $\mathbb{T} := \mathbb{R}/\mathbb{Z} \cong [0, 1)$ via

$$x \in [0, 1) \leftrightarrow \chi(t) := e^{2\pi i x} \in \partial\mathbb{D}.$$

Indeed, $\chi : (\mathbb{T}, m) \rightarrow (\partial\mathbb{D}, \lambda)$ is an isomorphism of measure spaces where m is Lebesgue measure on $[0, 1)$.

Let $\phi : \mathbb{D} \hookrightarrow \mathbb{D}$ be inner with restriction $\phi : \partial\mathbb{D} \hookrightarrow \partial\mathbb{D}$. Define $\tau = \tau(\phi) : \mathbb{T} \hookrightarrow \mathbb{T}$ m -a.e. by $\tau(t) := \chi^{-1}(\varphi(\chi(t)))$.

The following shows that $(\mathbb{T}, m, \tau)$ is a nonsingular transformation, whence so is the isomorphic $(\partial\mathbb{D}, \lambda, \varphi)$.

NORDGREN'S THEOREM ([Nor68]). For $\phi : \mathbb{D} \hookrightarrow \text{inner}$,

$$(3) \quad \pi_z \circ \tau^{-1} = \pi_{\phi(z)}, \quad \text{or equivalently} \quad \widehat{\tau} p_z = p_{\phi(z)},$$

where $\pi_z [p_z]$ is the Poisson measure [density] at z defined by

$$(4) \quad d\pi_z(x) := p_z(x)dx, \quad p_z(x) := \operatorname{Re} \frac{\chi(x) + z}{\chi(x) - z}.$$

REMARK. Equation (3) is also known as *Boole's formula* because a version for rational inner functions of the upper half-plane appears in [Boo57, p. 787] (see also [Gla77, §8]). For a converse to Nordgren's theorem, see [Let77].

2.4. Invariant probabilities for inner function restrictions

DENJOY–WOLFF THEOREM ([Den26, Wol26]). Suppose that $\phi : \mathbb{D} \hookrightarrow$ is analytic, nonconstant and non-Möbius. Then there is a (unique) point $\mathfrak{d} = \mathfrak{d}(\phi) \in \overline{\mathbb{D}}$ such that

$$\frac{1 - |\phi(z)|^2}{|1 - \overline{\mathfrak{d}}\phi(z)|^2} \geq \frac{1 - |z|^2}{|1 - \overline{\mathfrak{d}}z|^2} \quad \forall z \in \mathbb{D},$$

and

$$\phi^n(z) \rightarrow \mathfrak{d} \quad \text{as } n \rightarrow \infty \quad \forall z \in \mathbb{D}.$$

The point $\mathfrak{d}(\phi)$ is called the *Denjoy–Wolff point* of ϕ .

COROLLARY 2.3 (see e.g. [Aar78, Neu78, DM91]). The restriction τ of an inner function ϕ has an absolutely continuous, invariant probability iff the Denjoy–Wolff point of ϕ belongs to $\mathbb{D}$, and in this case $(\mathbb{T}, \pi_{\mathfrak{d}(\phi)}, T)$ either is conjugate to a circle rotation or is an exact probability preserving transformation.

2.4.1. Clark measures. As in [Sak07], the *Clark measure* of an analytic endomorphism $\phi : \mathbb{D} \hookrightarrow$ at $\xi \in \partial\mathbb{D}$ is the representing measure $\mu_\xi = \mu_\xi^{(\phi)} \in \mathcal{M}(\mathbb{T})$ of the positive harmonic function $z \mapsto \operatorname{Re} \frac{\xi + \phi(z)}{\xi - \phi(z)}$ satisfying

$$(5) \quad \operatorname{Re} \frac{\xi + \phi(z)}{\xi - \phi(z)} = \int_{\mathbb{T}} p_z d\mu_\xi.$$

It follows that $\xi \mapsto \mu_\xi$ is weak* continuous $\partial\mathbb{D} \rightarrow \mathcal{M}(\mathbb{T})$.

Now let $\phi : \mathbb{D} \hookrightarrow$ be inner. Then $\operatorname{Re} \frac{\xi + \phi(r\chi)}{\xi - \phi(r\chi)} \xrightarrow{r \rightarrow 1^-} 0$ a.s., whereas by Fatou's theorem,

$$\int_{\mathbb{T}} p_{r\chi} d\mu_\xi \xrightarrow{r \rightarrow 1^-} \frac{d\mu_\xi}{dm} \text{ a.s.} \quad \text{and} \quad \mu_\xi \perp m \quad \forall \xi \in \partial\mathbb{D}.$$

Moreover, since $p_z(t) = \operatorname{Re} \frac{\chi(t) + z}{\chi(t) - z}$, it follows from (3) that $\nu_t^{(\tau)} = \mu_{\chi(t)}$ where $\nu^{(\tau)} : \mathbb{T} \rightarrow \mathcal{M}(\mathbb{T})$ are the preimage measures of the restriction $\tau = \tau(\phi)$. See also [Sak07, §2].

2.5. Factorization of inner functions

2.5.1. Blaschke products. Suppose that $Z \subset \mathbb{D}$ is countable and that $\mathbf{m} : Z \rightarrow \mathbb{N}$ is such that $\sum_{a \in Z} \mathbf{m}(a)(1 - |a|) < \infty$.

The *Blaschke product* with zero set Z and *multiplicity function* $\mathbf{m}$ is $B = B_{Z, \mathbf{m}} : \mathbb{D} \rightarrow \mathbb{C}$ defined by

$$B(z) := \prod_{\alpha \in Z} b_{\alpha}(z)^{\mathbf{m}(\alpha)} \quad \text{with} \quad b_{\alpha}(z) := c_{\alpha} \frac{z - \alpha}{1 - \bar{\alpha}z}, \quad c_{\alpha} = \frac{-\bar{\alpha}}{\alpha} \quad (\alpha \neq 0), \quad c_0 = 1.$$

The product converges locally uniformly on $\mathbb{D}$ because

$$|1 - b_{\alpha}(z)^{\mathbf{m}(\alpha)}| \leq \frac{1 + |z|}{1 - |z|} \cdot \mathbf{m}(\alpha)(1 - |\alpha|).$$

It can be shown that B is inner, $\{a \in \mathbb{D} : B(a) = 0\} = Z$ and for $a \in Z$, $B(z)/b_a(z)^{\mathbf{m}(a)} =: H(z)$ is bounded and analytic with $H(a) \neq 0$.

2.5.2. Singular inner functions. An inner function $S : \mathbb{D} \hookrightarrow \mathbb{C}$ is *singular* (i.e. without zeros) iff

$$\log S(z) = - \int_{\mathbb{T}} \frac{\chi(t) + z}{\chi(t) - z} d\sigma(t),$$

where $\sigma \in \mathfrak{M}(\mathbb{T})$ with $\sigma \perp m$.

In this situation, we write $S = S_{\sigma}$ (&/or $\sigma = \sigma_S$).

FACTORIZATION THEOREM ([Smi29]; see also [Rud74, Thm. 17.15]). *Let $\phi : \mathbb{D} \hookrightarrow \mathbb{C}$ be an inner function, let $\mathcal{Z}(\phi) := \{z \in \mathbb{D} : \phi(z) = 0\}$ and, for $a \in \mathcal{Z}(\phi)$, let*

$$\mathbf{m}(a) := \max \{k \geq 0 : \phi/b_a^k \text{ is bounded on } \mathbb{D}\}.$$

Then $\sum_{a \in \mathcal{Z}(\phi)} \mathbf{m}(a)(1 - |a|) < \infty$ and $\phi = \lambda B_{\mathcal{Z}(\phi), \mathbf{m}} \cdot S$, where $\lambda \in \mathbb{S}^1$ and S is a singular inner function.

2.6. Regular points and singularities. A *regular point* of the inner function $\phi : \mathbb{D} \hookrightarrow \mathbb{C}$ is a point $z \in \partial\mathbb{D}$ such that there exists $U \subset \mathbb{C}$ open with $z \in U$ and an analytic function $F : U \rightarrow \mathbb{C}$ such that $F|_{U \cap \mathbb{D}} \equiv \phi|_{U \cap \mathbb{D}}$.

A nonregular point in $\partial\mathbb{D}$ is called a *singularity*.

The *regularity set* of ϕ is $\mathbf{r}_{\phi} := \{\text{regular points of } \phi\}$ and the *singularity set* of ϕ is $\mathbf{s}_{\phi} := \partial\mathbb{D} \setminus \mathbf{r}_{\phi}$.

- If ν_x is any Clark measure for the inner function ϕ (as in §2.4.1) then $\chi^{-1}\mathbf{s}_{\phi} = (\mathbf{spt} \nu_x)'$.
- If $S = S_{\sigma}$ is a singular inner function then $\chi^{-1}\mathbf{s}_S = \mathbf{spt} \sigma$.

2.6.1. Derivative of a restriction at a regular point. Let $\phi : \mathbb{D} \hookrightarrow \mathbb{C}$ with restriction $\tau = \tau(\phi)$. Then τ is differentiable at each $\theta \in \chi^{-1}\mathbf{r}_{\phi}$ with

$$(6) \quad \tau'(\theta) = \partial\phi(\chi(\theta)) = |\phi'(\chi(\theta))| \quad \text{with} \quad \partial\phi(z) := \frac{z\phi'(z)}{\phi(z)} = z(\log \phi)'(z).$$

If $\phi = \lambda B_{\mathcal{Z}(\phi), \mathbf{m}} S_\sigma$, then

$$\mathbf{s}_\phi = \mathcal{Z}(\phi)' \cup \mathbf{spt} \sigma$$

and, as in [Mar89],

$$\begin{aligned} (7) \quad \tau'(\theta) &= \sum_{\alpha \in \mathcal{Z}(\phi)} \mathbf{m}(\alpha) p_\alpha(\theta) + \frac{1}{2} \int_{\mathbb{T}} \frac{d\sigma(t)}{\sin^2(\pi(\theta - t))} \\ &\geq \sum_{\alpha \in \mathcal{Z}(\phi)} \frac{1 - |\alpha|}{1 + |\alpha|} + \frac{\sigma(\mathbb{T})}{2} =: \eta \geq 0 \quad \forall \theta \in \mathbb{T} \text{ such that } \chi(\theta) \in \mathbf{r}_\phi, \end{aligned}$$

where $p_\alpha(\theta) := \operatorname{Re} \frac{\alpha + \chi(t)}{\alpha - \chi(t)}$ and $\sum_{\alpha \in \emptyset} := 0$. Since either $\mathcal{Z}(\phi) \neq \emptyset$ or $\sigma \neq 0$ (or both), we have $\eta > 0$.

2.6.2. Arc maps. An *arc map* is a triple $(\mathbb{T}, T, \alpha)$, where α is a finite or countable partition mod m of $\mathbb{T}$ into *open arcs* (open, connected subsets $A \subsetneq \mathbb{T}$) ⁽³⁾ and $T : \mathbb{T} \rightarrow \mathbb{T}$ is a map such that

- for each $A \in \alpha$, $T : A \rightarrow T(A)$ is a bi-absolutely continuous homeomorphism,
- $\sigma(\bigcup_{n \geq 0} T^{-n} \alpha) = \mathcal{B}(\mathbb{T})$.

It is called *piecewise C^k* ($k \geq 1$) [*analytic*] if each $T : A \rightarrow TA$ is a C^k -diffeomorphism [bi-analytic].

PROPOSITION 2.4 (Arc map restrictions). *Let $\phi : \mathbb{D} \hookrightarrow \mathbb{D}$ be inner with $m(\mathbf{s}_\phi) = 0$ and suppose the Denjoy–Wolff point of ϕ belongs to $\mathbb{D}$. Then:*

- (i) $\tau : \mathbb{T} \hookrightarrow \mathbb{T}$ defines a piecewise analytic, arc map $(\mathbb{T}, \tau, \alpha)$ such that

$$\begin{aligned} (8) \quad \tau'(x) &\geq \eta > 0 \quad \forall x \in \chi^{-1} \mathbf{r}_\phi, \\ \exists d \geq 1 : |(\tau^d)'(x)| &\geq \beta > 1 \quad \forall x \in \bigcap_{j=0}^{d-1} \tau^{-j} \chi^{-1} \mathbf{r}_\phi. \end{aligned}$$

- (ii) If in addition $\#\mathcal{Z}(\phi), \#\mathbf{s}_\phi < \infty$, then the partition α may be chosen to be surjective:

$$(9) \quad \tau(A) = \mathbb{T} \bmod m \quad \forall A \in \alpha.$$

Moreover,

$$(10) \quad \sup_{x \in \chi^{-1} \mathbf{r}_\phi} \Delta \tau(x) < \infty \quad \text{with} \quad \Delta \tau(x) := \frac{|\tau''(x)|}{\tau'(x)^2}.$$

We will call piecewise onto arc maps satisfying (8)–(10) *Adler maps*.

Adler interval maps (Adler arc maps with surjective partitions into intervals) are considered in [Adl73].

⁽³⁾ Of the form $\chi^{-1} A = (a, b)$ with $0 \leq a < b \leq 1$ or $[0, a) \cup (b, 1]$ with $0 < a \leq b < 1$.

Any Adler arc map is conjugate by rotation to an Adler interval map.

Proof of Proposition 2.5

Proof of (8). There is a Möbius transformation $\Psi : \mathbb{D} \hookrightarrow$ such that $\mathfrak{d}(g) = 0$, where $g = \Psi^{-1} \circ \phi \circ \Psi$. If the restrictions of g & Ψ are U & ψ respectively, then $\tau^n = \psi \circ U^n \circ \psi^{-1}$ for all $n \geq 1$.

Since $g(0) = 0$, by (7) we have

$$U'(\theta) = \partial g(\chi(\theta)) = 1 + \partial h(\chi(\theta)) \geq 1 + \delta_h =: \rho > 1.$$

For $n \geq 1$,

$$\begin{aligned} \tau^{n'} &= \psi' \circ U^n \circ \psi^{-1} \cdot U^{n'} \circ \psi^{-1} \cdot \psi^{-1'} \\ &\geq \rho^n \min_{\mathbb{T}} \psi' \cdot \min_{\mathbb{T}} \psi^{-1'} \\ &\geq B > 1 \quad \text{for large enough } n \geq 1. \end{aligned}$$

Proof that $\sigma(\bigcup_{n \geq 0} T^{-n}\alpha) = \mathcal{B}(\mathbb{T})$. By (8) we see that $\alpha_n := \bigvee_{k=0}^{n-1} \tau^{-1}\alpha$ is also a mod 0 partition of $\mathbb{T}$ into open arcs satisfying

$$\max \{m(a) : a \in \alpha_n\} \leq (1/\eta)^d \cdot (1/\beta)^{n/d},$$

whence $\sigma(\bigcup_{n \geq 0} \tau^{-n}\alpha) \stackrel{m}{=} \mathcal{B}(\mathbb{T})$.

Proof of (9) in case $\mathfrak{s}_\phi \neq \emptyset$. We construct a mod 0 partition α of $\mathbb{T}$ into open arcs satisfying (9).

Since $\#\mathfrak{s}_\phi < \infty$, formula (7) now has the form

$$(11) \quad \mathfrak{T}'(\theta) = \sum_{a \in \mathcal{Z}(\phi)} m(a)p_a(\theta) + \frac{1}{2} \sum_{t \in \mathfrak{s}_\phi} \frac{\sigma(\{t\})}{\sin^2(\pi(\theta - t))},$$

Suppose that $J \subset \mathbb{T}$ is an open arc and $f : J \rightarrow \mathbb{T}$ is continuously differentiable on J with $\min_J f' > 0$. Then f has a *lifting*: there is an interval $\tilde{J} \subset \mathbb{R}$ such that $m\tilde{J} = J$, where $m : \mathbb{R} \rightarrow \mathbb{T}$, $m(x) = x \bmod 1$; and there exists $F : \tilde{J} \rightarrow \mathbb{R}$ continuously differentiable such that

$$m(F(x)) = f(m(x)) \quad \text{for } x \in \tilde{J}.$$

In particular, $F'(x) = f'(m(x))$.

Let $J \subset \chi^{-1}\mathfrak{r}_\phi$ be a maximal open arc (i.e. $\partial J \subset \chi^{-1}\mathfrak{s}_\phi$) and let $\mathfrak{T} : \tilde{J} \rightarrow \mathbb{R}$ be the lifting of $\tau : J \rightarrow \mathbb{T}$. Write $\tilde{J} = (a_-, a_+)$. Then by (11),

$$\mathfrak{T}'(\theta) \xrightarrow{\theta \rightarrow \{a_-, a_+\}, \theta \in (a_-, a_+)} \infty,$$

whence

$$(12) \quad \mathfrak{T}(\theta) \xrightarrow{\theta \rightarrow a_\pm, \theta \in (a_-, a_+)} \pm \infty$$

and there is a countable mod 0 partition $\mathcal{P}_{\tilde{J}}$ of $\tilde{J}$ into open arcs such that for each $A \in \mathcal{P}_{\tilde{J}}$, $\mathfrak{T}A$ is an interval of length 1.

It follows that $\alpha_J := m\mathfrak{p}_{\tilde{J}}$ is a mod 0 partition of J into open arcs such that $\tau A = \mathbb{T} \bmod 0$ for all $A \in \alpha_J$.

Since $\#\mathfrak{s}_\phi < \infty$ we know that $\chi^{-1}\mathfrak{r}_\phi$ is a finite union of maximal open arcs as above and so there is a mod 0 partition α of $\chi^{-1}\mathfrak{r}_\phi$ into open arcs such that $\tau A = \mathbb{T} \bmod 0$ for all $A \in \alpha$.

Proof of (10). Since $\#\mathcal{Z}(\phi)$ and $\mathfrak{s}_\phi$ are both finite, by (7), $\tau(\theta) = \mathfrak{T}(\theta) \bmod 1$ with

$$(13) \quad \mathfrak{T}(\theta) = \mathfrak{b}(\theta) - \frac{1}{2\pi} \sum_{t \in \mathfrak{s}_\phi} \sigma(\{t\}) \cot(\pi(\theta - t))$$

where $\mathfrak{b} \equiv 0$ when $\mathcal{Z}(\phi) = \emptyset$; and when $1 \leq \#\mathcal{Z}(\phi) < \infty$,

$$\mathfrak{b}(\theta) = \int_0^\theta \left(\sum_{a \in \mathcal{Z}(\phi)} \mathfrak{m}(a) p_a(t) \right) dt \bmod 1$$

defines an analytic endomorphism of $\mathbb{T}$.

Since either $\mathfrak{b} = 0 = \Delta(\mathfrak{b})$ or $\mathfrak{b} : \mathbb{T} \hookrightarrow \mathbb{T}$ is analytic expanding, in which case $\Delta(\mathfrak{b}) : \mathbb{T} \rightarrow \mathbb{R}$ is analytic, we have $\|\Delta(\mathfrak{b})\|_\infty < \infty$.

In case $\mathfrak{s}_\phi \neq \emptyset$, by (11) we have

$$(14) \quad \tau'(\theta) = \mathfrak{b}'(\theta) + \frac{1}{2} \sum_{t \in \mathfrak{s}_\phi} \frac{\sigma(\{t\})}{\sin^2(\pi(\theta - t))} =: \mathfrak{b}'(\theta) + s'(\theta)$$

and

$$|\tau''(\theta)| \leq |\mathfrak{b}''(\theta)| + |s''(\theta)| \leq |\mathfrak{b}''(\theta)| + \pi \sum_{t \in \mathfrak{s}_\phi} \sigma(\{t\}) \left| \frac{\cos(\pi(\theta - t))}{\sin^3(\pi(\theta - t))} \right|,$$

whence

$$\begin{aligned} \Delta\tau(\theta) &\leq \|\Delta\mathfrak{b}\|_\infty + \pi \frac{\sum_{t \in \mathfrak{s}_\phi} \sigma(\{t\}) \left| \frac{\cos(\pi(\theta - t))}{\sin^3(\pi(\theta - t))} \right|}{s'(\theta)^2} \\ &\leq \|\Delta\mathfrak{b}\|_\infty + \pi \sum_{t \in \mathfrak{s}_\phi} \frac{\frac{\sigma(\{t\}) |\cos(\pi(\theta - t))|}{|\sin^3(\pi(\theta - t))|}}{\left(\frac{\sigma(\{t\})}{\sin^2(\pi(\theta - t))} \right)^2} \\ &= \|\Delta\mathfrak{b}\|_\infty + \frac{\pi}{2} \sum_{t \in \mathfrak{s}_\phi} \frac{|\sin(2\pi(\theta - t))|}{\sigma(\{t\})} \leq \|\Delta\mathfrak{b}\|_\infty + \frac{\pi}{2} \sum_{t \in \mathfrak{s}_\phi} \frac{1}{\sigma(\{t\})} \\ &=: M < \infty. \blacksquare \end{aligned}$$

EXAMPLE 2.5. We exhibit an inner function $\phi : \mathbb{D} \hookrightarrow \mathbb{D}$ with $\phi(0) = 0$ and singularity set $\mathfrak{s}_\phi = \{\chi(a), \chi(b)\}$ with $0 \leq a < b \leq 1$ such that $\tau = \tau(\phi) : [a, b] \rightarrow [\tau(a+), \tau(b-)] \subset (0, 1)$ is a homeomorphism. Thus the restriction is not an Adler arc map. Indeed, such an inner function must be a Blaschke product since otherwise either $\chi(a)$ is a singularity of the singular factor of

ϕ and by (12), $\tau(a-) = 0$, or $\chi(b)$ is and $\tau(b-) = 1$. This would contradict $[\tau(a+), \tau(b-)] \subsetneq (0, 1)$.

To construct ϕ , first define an inner function of $\mathbb{R}^{2+}$ (the upper half-plane), $B : \mathbb{R}^{2+} \hookrightarrow$, by

$$B(z) := \sum_{n=1}^{\infty} \frac{1}{2^{n+1}} \left(\frac{1 + s_n z}{s_n - z} + \frac{1 + t_n z}{t_n - z} \right)$$

with $s_n := -\frac{1}{n}$ and $t_n = 1 + \frac{1}{n}$; then B is an inner function of $\mathbb{R}^{2+}$ and $B(i) = i$, $B : (0, 1) \rightarrow \mathbb{R}$ is continuous, increasing with $B(0+), B(1-) \in \mathbb{R}$. Let $\psi : \mathbb{D} \rightarrow \mathbb{R}^{2+}$, $\psi(z) = i\frac{1+z}{1-z}$. Then $\phi := \psi^{-1} \circ B \circ \psi$ is as advertised with $a = 1/2$ and $b = 3/4$.

2.7. Radial limit set. The *radial limit set* of ϕ is

$$\Lambda_\phi := \{\beta \in \partial\mathbb{D} : \phi(r\beta) \xrightarrow{r \rightarrow 1-} \phi(\beta) \in \partial\mathbb{D}\}.$$

For example, if $\phi(z) := \exp\left[-\frac{1+z}{1-z}\right]$, then

$$\phi(r\xi) \xrightarrow{r \rightarrow 1-} \begin{cases} e^{-i \cot(\theta/2)}, & \xi = \chi(\theta) \neq 1, \\ 0, & \xi = 1, \end{cases}$$

and $\Lambda_\phi = \partial\mathbb{D} \setminus \{1\}$.

2.7.1. Angular derivatives. The inner function $\phi : \mathbb{D} \hookrightarrow$ has an *angular derivative* $\beta \in \mathbb{C}$ at $\xi \in \Lambda_\phi$ if there exists $\beta \in \mathbb{C}$ such that

$$\frac{\phi(z) - \phi(\xi)}{z - \xi} \xrightarrow{z \rightarrow \xi} \beta, \quad \text{that is,} \quad \frac{\phi(z) - \phi(\xi)}{z - \xi} \xrightarrow{z \rightarrow \xi, |\xi - z| \leq K(1 - |z|)} \beta \quad \forall K > 0.$$

Denote the angular derivative at ξ by $\beta =: \phi'_\angle(\xi)$.

PROPOSITION 2.6 ([Sak07, §3]). *Let $\phi : \mathbb{D} \hookrightarrow$ be inner with $\tau = \tau(\phi)$. The following are equivalent for $\xi = \chi(x) \in \partial\mathbb{D}$:*

- ϕ has an angular derivative at ξ ;
- $\phi'(z) \xrightarrow{z \rightarrow \xi} \beta \in \mathbb{C}$ and in this case $\beta = \phi'_\angle(\xi)$;
- $\int_{\mathbb{T}} \frac{d\nu_w(t)}{|\xi - \chi(t)|^2} < \infty$ for some (hence all) $w \in \mathbb{T}$, $w \neq x$.

PROPOSITION 2.7 ([Ale87], also [Sak07, Theorem 9.6]). *Let $\phi : \mathbb{D} \hookrightarrow$ be inner and let $E \in \mathcal{B}(\mathbb{T})$ with $m(E) > 0$. Then $\tau = \tau(\phi)$ is locally invertible on E iff ϕ has an angular derivative at $\chi(x)$ for m -a.e. $x \in E$.*

Proof. $\Rightarrow$ Local invertibility on E entails forward nonsingularity on E , whence, by [Hei77], existence of angular derivatives a.s. on E .

$\Leftarrow$ Suppose that ϕ has an angular derivative at $\chi(x)$ for m -a.e. $x \in E$. By [Cra91, Lemma 1.5], τ is *almost uniformly differentiable* on E in the sense that there are $E_k \in \mathcal{B}(\mathbb{T})$ with $E_k \uparrow E \bmod m$ such that for all k and $\varepsilon > 0$

there exists $\delta = \delta(k, \varepsilon) > 0$ such that

$$(15) \quad |\tau(x) - \tau(y) - (x - y)g(x)| \leq \varepsilon|x - y| \quad \forall x, y \in E_k, |x - y| < \delta,$$

where $g(x) = |\phi'(\chi(x))|$ with $\phi'(\chi(x))$ the angular derivative of ϕ at $\chi(x)$.

To see that τ is forward nonsingular on E , we note that by possibly shrinking the E_k (as in (15)), we may assume in addition that there exists $M_k > 0$ ($k \geq 1$) such that $g \leq M_k$ on E_k .

Suppose that $k \geq 1$ and $S \in \mathcal{B}(E_k)$ with $m(S) = 0$, and fix $\varepsilon > 0$. There are intervals $\{I_n : n \geq 1\}$ such that

$$S \subset \bigcup_{n \geq 1} I_n, \quad |I_n| \leq \delta(k, 1), \quad \sum_{n \geq 1} |I_n| < \frac{\varepsilon}{M_k + 1}.$$

By (15),

$$|\tau(x) - \tau(y)| \leq (g(x) + 1)|x - y| \leq (M_k + 1)|x - y|$$

and $m(\tau(E_k \cap I_n)) \leq (M_k + 1)m(I_n)$.

It follows that $\tau(S) \subset \bigcup_{n \geq 1} \tau(E_k \cap I_n)$, whence

$$m(\tau(S)) \leq \sum_{n \geq 1} m(\tau(E_k \cap I_n)) \leq (M_k + 1) \sum_{n \geq 1} m(I_n) < \varepsilon$$

and thus $m(\tau(S)) = 0$.

For $S \in \mathcal{B}(E)$ with $m(S) = 0$,

$$m(\tau(S)) \xleftarrow[k \rightarrow \infty]{} m(\tau(E_k \cap S)) = 0$$

and τ is forward nonsingular on E , whence locally invertible on E by Theorem 2.2. ■

An example of a probability preserving restriction which is “a.e. continuum-to-one” (a.e. Clark measure is nonatomic) was constructed in [Don65] (see also [Sak07, Ex. 9.7]).

In particular, for inner functions ϕ , $\tau(\phi)$ is locally invertible on $\mathbb{T}$ iff ϕ has an angular derivative at a.e. point on $\partial\mathbb{D}$.

The next result shows that this property is independent of the singularity set.

PROPOSITION 2.8. *Let $E \subseteq \mathbb{T}$ be a closed set. Then there exists an inner function $\phi : \mathbb{D} \hookrightarrow \mathbb{D}$ with locally invertible restriction $\tau = \tau(\phi)$ such that $\phi(0) = 0$ and $\mathfrak{s}_\phi = \chi(E)$.*

Proof. Let $\Gamma \subset \mathbb{T}$ be countable with $\Gamma' = E$. We will construct ϕ via a Clark measure.

First fix $\varepsilon : \Gamma \rightarrow \mathbb{R}_+$ such that $\sum_{\gamma \in \Gamma} \varepsilon(\gamma) < \infty$ and then fix $\pi \in \mathcal{P}(\Gamma)$ with

$$\sum_{\gamma \in \Gamma} \frac{\pi_\gamma}{\varepsilon(\gamma)^2} < \infty.$$

We claim that

$$(16) \quad \int_{\mathbb{T}} \frac{d\pi(t)}{|t-x|^2} < \infty \quad \text{for } m\text{-a.e. } x \in \mathbb{T},$$

where $\pi := \sum_{\gamma \in \Gamma} \pi_{\gamma} \delta_{\gamma} \in \mathcal{P}(\mathbb{T})$.

Indeed, since

$$\sum_{\gamma \in \Gamma} m(B(\gamma, \varepsilon(\gamma))) = 2 \sum_{\gamma \in \Gamma} \varepsilon(\gamma) < \infty$$

with $B(x, \varepsilon) := (x - \varepsilon, x + \varepsilon)$, by the Borel–Cantelli lemma there exists $K \in \mathcal{B}(\mathbb{T})$ with $K \cap \Gamma = \emptyset$ and $m(K) = 1$ such that for all $x \in K$, there exists $\Gamma_0(x) \in \Gamma$ finite such that

$$(17) \quad |x - \gamma| \geq \varepsilon(\gamma) \quad \forall \gamma \notin \Gamma_0(x).$$

Let $x \in K$. Then

$$\int_{\mathbb{T}} \frac{d\pi(t)}{|t-x|^2} = \sum_{\gamma \in \Gamma} \frac{\pi_{\gamma}}{|x-\gamma|^2} = \left(\sum_{\gamma \in \Gamma_0(x)} + \sum_{\gamma \notin \Gamma_0(x)} \right) \frac{\pi_{\gamma}}{|x-\gamma|^2}.$$

Since $K \subset \mathbb{T} \setminus \Gamma$, we have

$$\sum_{\gamma \in \Gamma_0(x)} \frac{\pi_{\gamma}}{|x-\gamma|^2} < \infty \quad \forall x \in K$$

and

$$\sum_{\gamma \notin \Gamma_0(x)} \frac{\pi_{\gamma}}{|x-\gamma|^2} \leq \sum_{\gamma \notin \Gamma_0(x)} \frac{\pi_{\gamma}}{\varepsilon(\gamma)^2} < \infty,$$

proving (16).

Next, we define $F : \mathbb{D} \rightarrow \mathbb{C}$ by

$$F(z) := \int_{\mathbb{T}} \frac{\chi(t) + z}{\chi(t) - z} d\pi(t).$$

Note that

$$\operatorname{Re} \left(\frac{\chi(t) + z}{\chi(t) - z} \right) = \frac{1 - |z|^2}{|\chi(t) - z|^2} > 0 \quad \forall z \in \mathbb{D},$$

so $F : \mathbb{D} \rightarrow \mathbb{R}_+ \times \mathbb{R}$. Moreover, since $\pi(\mathbb{T}) = 1$, we have $F(0) = 1$.

To construct the inner function, define $\phi := \frac{F-1}{F+1} : \mathbb{D} \rightarrow \mathbb{C}$. Since $F(0) = 1$, we have $\phi(0) = 0$ and since $\operatorname{Re} F > 0$ on $\mathbb{D}$, it follows that $\phi : \mathbb{D} \hookrightarrow$.

Since $\pi \perp m$, we see that for m -a.e. $x \in \mathbb{T}$,

$$\lim_{r \rightarrow 1^-} F(r\chi(x)) =: F(\chi(t)) \in i\mathbb{R},$$

whence for such $x \in \mathbb{T}$ we have

$$\phi(r\chi(x)) \xrightarrow{r \rightarrow 1^-} \frac{F(\chi(x)) - 1}{F(\chi(x)) + 1} \in \partial\mathbb{D}$$

and $\phi : \mathbb{D} \hookrightarrow$ is inner.

Moreover, $\nu_0 = \pi$, whence by (16) and Proposition 2.6, ϕ has an angular derivative at $\chi(t)$ for a.e. $t \in \mathbb{T}$. By Proposition 2.7, $\tau(\phi)$ is locally invertible.

To finish, we note that $\mathfrak{s}_\phi = \mathbf{spt} \pi' = I' = \chi(E)$. ■

The Baire category situation is different:

PROPOSITION 2.9. *Suppose that the inner function $\phi : \mathbb{D} \hookrightarrow \mathbb{D}$ has an angular derivative at a residual set of points. Then $\mathfrak{s}_\phi$ is nowhere dense.*

Proof. Fix $w \in \mathbb{T}$ and define $F : \mathbb{T} \rightarrow (0, \infty]$ by

$$F(x) := \int_{\mathbb{T}} \frac{d\nu_w(t)}{|\chi(x) - \chi(t)|^2}$$

and, for $r \in (0, 1)$, define $F_r : \mathbb{T} \rightarrow \mathbb{R}_+$ by

$$F_r(x) := \int_{\mathbb{T}} \frac{d\nu_w(t)}{|r\chi(x) - \chi(t)|^2}.$$

Then for each $0 < r < 1$, $F_r : \mathbb{T} \rightarrow \mathbb{R}_+$ is continuous and $F_r(x) \xrightarrow{r \rightarrow 1^-} F(x) \in (0, \infty]$ for all $x \in \mathbb{T}$, by dominated convergence in case $F(x) < \infty$ and by Fatou's lemma if $F(x) = \infty$.

By Baire's simple limit theorem, the set

$$C_\infty := \{x \in \mathbb{T} : F : \mathbb{T} \rightarrow (0, \infty] \text{ is continuous at } x\}$$

is residual in $\mathbb{T}$. By assumption, $\Lambda := [F < \infty]$ is residual, whence so is $C := C_\infty \cap \Lambda$. Thus for every $x \in C$, there exists $0 < a_x < b_x$ and an open interval $J_x \ni x$ such that

$$a_x < F < b_x \quad \text{on } J_x.$$

By Seidel's theorem [Sei34] (see also [Zyg02, Thm. 7.48]),

$$\mathfrak{s}_\phi \cap \chi(J_x) = \emptyset,$$

whence $\mathfrak{r}_\phi$ is open and dense and $\mathfrak{s}_\phi$ is nowhere dense. ■

3. Quasicompactness of transfer operators. Let $\mathcal{L}$ be a Banach space. An operator $P \in \text{hom}(\mathcal{L}, \mathcal{L})$ is called *quasicompact* if there exists $A = A(P) \in \text{hom}(\mathcal{L}, \mathcal{L})$ of the form

$$A = \sum_{k=1}^N \lambda_k E_k$$

with $N \geq 1$, $E_1, \dots, E_N \in \text{hom}(\mathcal{L}, \mathcal{L})$ finite-dimensional projections, and $\lambda_1, \dots, \lambda_N \in \mathbb{S}^1 := \{z \in \mathbb{C} : |z| = 1\}$ such that the spectral radius satisfies

$$\rho(P - A) := \lim_{n \rightarrow \infty} \|(P - A)^n\|_{\text{hom}(\mathcal{L}, \mathcal{L})}^{1/n} < 1.$$

Let (X, m, T) be a nonsingular transformation with transfer operator $\hat{T} : L^1(m) \hookrightarrow$. We look for Banach spaces $\mathcal{L} \subset L^1(m)$ on which $\hat{T} : \mathcal{L} \hookrightarrow$ acts quasicompactly.

If (X, m, T) is a weakly mixing, probability preserving transformation with transfer operator $\hat{T}$ acting quasicompactly on $\mathcal{L} \subset L^1(m)$, then $A(\hat{T})f = \mathbb{E}(f)$ and for all $\theta \in (\rho(\hat{T} - \mathbb{E}), 1)$ there exists $M > 0$ such that

$$(18) \quad \|\hat{T}^n f - \mathbb{E}(f)\|_{\mathcal{L}} \leq M\theta^n \|f\|_{\mathcal{L}} \quad \forall f \in \mathcal{L}.$$

The property (18) is called *exponential decay of correlations* because it entails

$$\left| \int_X u \cdot v \circ T^n dm - \mathbb{E}(u)\mathbb{E}(v) \right| \leq M\theta^n \|u\|_{\mathcal{L}} \|v\|_{L^1(m)}.$$

3.1. Doeblin–Fortet operators on an adapted pair. Let $\mathcal{L} \subset \mathcal{C} \subset L^1(m)$ be Banach spaces such that $(\mathcal{C}, \mathcal{L})$ is an *adapted pair* in the sense that

- $\|\cdot\|_{L^1(m)} \leq \|\cdot\|_{\mathcal{C}} \leq \|\cdot\|_{\mathcal{L}}$, $(\overline{\mathcal{L}})_{L^1(m)} = L^1(m)$, and $\mathcal{L}$ -closed, bounded sets are $\mathcal{C}$ -compact.

For example both $(L^1(m), \text{Lip}(\mathbb{T}))$ and $(L^1(m), \text{BV}(\mathbb{T}))$ are adapted pairs. Here $\text{Lip}(\mathbb{T})$ denotes the Lipschitz functions on $\mathbb{T}$ (equivalently, the absolutely continuous functions with essentially bounded derivative), with norm

$$\|f\|_{\text{Lip}} := \|f\|_1 + \|f'\|_{\infty};$$

Lip -closed, bounded sets are L^1 -compact by the Arzelà–Ascoli theorem. Further, $\text{BV}(\mathbb{T})$ denotes the functions of bounded variation on $\mathbb{T}$ with norm

$$\|f\|_{\text{BV}} := \|f\|_1 + \bigvee f, \quad \text{where}$$

$$\bigvee f := \sup \left\{ \sum_{k=0}^{n-1} |f(t_{k+1}) - f(t_k)| : 0 < t_1 < \dots < t_n = 1 \right\};$$

BV -closed, bounded sets are L^1 -precompact by Helly’s theorem.

As in [Nor72, Ch. 3], we say that an operator $P \in \text{hom}(\mathcal{L}, \mathcal{L}) \cap \text{hom}(\mathcal{C}, \mathcal{C})$ is *Doeblin–Fortet* (D-F) on $(\mathcal{C}, \mathcal{L})$ if

$$(19) \quad \|P^n f\|_{\mathcal{C}} \leq H \|f\|_{\mathcal{C}} \quad \forall n \in \mathbb{N}, f \in L^1(m),$$

$$(20) \quad \exists \kappa \geq 1 : \|P^\kappa f\|_{\mathcal{L}} \leq \theta \|f\|_{\mathcal{L}} + R \|f\|_{\mathcal{C}} \quad \forall f \in \mathcal{L}.$$

where $R, H \in \mathbb{R}_+$ and $\theta \in (0, 1)$.

EXAMPLE 3.1 (Adler arc maps). Let $(\mathbb{T}, m = \text{Leb}, T)$ be an Adler map. It is a well known folklore result that the transfer operator $\hat{T}$ is D-F on the adapted pair $(L^1(m), \text{Lip}(\mathbb{T}))$.

It is also D-F on $(L^1(m), \text{BV}(\mathbb{T}))$ because by [Zwe98, Cor. 1] an Adler map satisfies the assumptions of [Ryc83, Prop. 1], which proves the D-F inequality on $(L^1(m), \text{BV})$.

The following lemma is a well-known consequence of the Yosida–Kakutani mean ergodic theorem [YK41, Theorem 1]. See also [ITM50], [Nor72, Chapter 3], [HH01, PP90], [LY73, Theorem 1].

LEMMA 3.2 (rigidity of Doeblin–Fortet operators). *Suppose that P is a Doeblin–Fortet operator on the adapted pair $(\mathcal{C}, \mathcal{L})$. If $f \in \mathcal{C}$ and $Pf = f$, then $f \in \mathcal{L}$.*

It is shown in [ITM50] (see also [Nor72, Ch. 3], [HH01, PP90]) that a D-F operator $P \in \text{hom}(\mathcal{L}, \mathcal{L})$ has spectral radius $\rho(P) \leq 1$ and that, if $\rho(P) = 1$, then P is quasicompact.

3.1.1. Quasicompactness and the central limit theorem. If $\psi \in \mathcal{L}$ and $\mathbb{E}(\psi) = 0$, then by Leonov’s theorem [Leo61],

$$(21) \quad \exists \lim_{n \rightarrow \infty} \frac{1}{n} \mathbb{E}(\psi_n^2) =: \sigma_\psi^2 \geq 0, \quad \text{where} \quad \psi_n := \sum_{k=0}^{n-1} \psi \circ T^k,$$

with equality iff $\psi = g - g \circ T$ for some $g \in \psi$.

If in addition $\sigma_\psi > 0$, then (see [Gor04]) the *stationary process* (X, m, T, ψ) satisfies the central limit theorem:

$$(22) \quad m\left(\left[\frac{\psi_n}{\sigma_\psi \sqrt{n}} \leq t\right]\right) \xrightarrow{n \rightarrow \infty} \frac{1}{\sqrt{2\pi}} \int_{-\infty}^t e^{-s^2/2} ds.$$

3.2. Hardy spaces. The *harmonization* of $f \in L^p(m)$ ($1 \leq p \leq \infty$) is $\tilde{f} : \mathbb{D} \rightarrow \mathbb{C}$ defined by

$$\tilde{f}(z) := \int_{\mathbb{T}} p_z f \, dm,$$

where p_z is as in (4). The function $\tilde{f}$ is harmonic in $\mathbb{D}$ and satisfies

$$\sup_{r \in (0,1)} \|\tilde{f}(r\chi)\|_p = \|f\|_p.$$

It is classical that the Hardy spaces consist of harmonizations:

$$h^p(\mathbb{D}) := \left\{ F : \mathbb{D} \rightarrow \mathbb{C} \text{ harmonic} : \sup_{r \in (0,1)} \|F(r\chi)\|_p < \infty \right\} = \{\tilde{f} : f \in L^p(m)\},$$

$$H^p(\mathbb{D}) := \{f \in h^p : \tilde{f} \text{ analytic on } \mathbb{D}\} \cong \{f \in L^p(\mathbb{T}) : \hat{f}(n) = 0 \, \forall n < 0\}.$$

Let

$$\Lambda_{\tilde{f}} := \left[\exists \lim_{r \rightarrow 1^-} \tilde{f}(r\chi) =: \tilde{f}(\chi) \in \mathbb{C} \right].$$

Then by Fatou’s theorem, $m(\Lambda_{\tilde{f}}) = 1$ and $\tilde{f}(\chi) = f$ a.e.

3.2.1. Action of the transfer operator. Let $\phi : \mathbb{D} \hookrightarrow \mathbb{D}$ be inner with restriction $\tau = \tau(\phi)$. Then (see [Ale87], and also [Sak07, Thm. 3.1]) $\hat{\tau}H^p(\mathbb{D}) \subset H^p(\mathbb{D})$, and if $\phi(0) = 0$, then for $d \geq 1$, $\hat{\tau}(\chi^d)$ is a polynomial in χ of degree at most d .

LEMMA 3.3 (Basic Proposition). *Let $\phi : \mathbb{D} \hookrightarrow \mathbb{D}$ be inner with $\phi^{(k)}(0) = 0$ for all $0 \leq k < \kappa$. Then for $N \geq 1$ we have $\hat{\tau}^N(\chi^d) = 0$ for all $1 \leq d < \kappa^N$, and for $d \geq \kappa^N$,*

$$\hat{\tau}^N(\chi^d) = \sum_{\ell=1}^{\lfloor d/\kappa^N \rfloor} a^{(N)}(d, \ell) \chi^\ell \quad \text{where} \quad a^{(N)}(k, \ell) = \overline{(\phi^{[N]})^\ell}(k).$$

Proof. Let $d \geq 1$ and write $\hat{\tau}^N(\chi^d) = \sum_{\ell \in \mathbb{Z}} a^{(N)}(d, \ell) \chi^\ell$. Then

$$a^{(N)}(d, \ell) = \langle \hat{\tau}^N(\chi^d), \chi^\ell \rangle = \langle \chi^d, \chi^\ell \circ \tau^N \rangle = \langle \chi^d, \phi^N(\chi)^\ell \rangle = \overline{(\phi^N)^\ell}(d) \quad (4).$$

Thus

$$(23) \quad a^{(N)}(d, \ell) = 0 \quad \text{unless} \quad \ell \geq 1 \text{ \& } d \geq \kappa^N \ell. \quad \blacksquare$$

3.3. Weighted Hilbert spaces. A (Hilbert space) *weight* is a sequence $\underline{w} \in \mathbb{R}_+^{\mathbb{N}_0}$ satisfying $1 = w(0) < w(1) < \dots < w(n) \uparrow \infty$. The associated *weighted Hilbert space* is

$$\mathcal{H}_{\underline{w}} := \left\{ f \in L^2(m) : \|f\|_{\underline{w}}^2 := \sum_{n \in \mathbb{Z}} w(|n|) |\hat{f}(n)|^2 < \infty \right\},$$

equipped with the inner product

$$\langle u, v \rangle_{\underline{w}} := \sum_{n \in \mathbb{Z}} w(|n|) \hat{u}(n) \overline{\hat{v}(n)}.$$

Classical examples

(i) For $w(n) = n^2$, $\mathcal{H}_{\underline{w}}$ is isomorphic to the Sobolev space

$$W^{1,2}(\mathbb{T}) := \{f \in C(\mathbb{T}) : f \text{ a.c. \& } f' \in L^2(m)\}.$$

(ii) For $b > 1$, let $w_{1,b}(n) := b^n$. Then

$$\mathcal{H}_b = \mathcal{H}_{\underline{w}_{1,b}} \cong h^2(B_{\mathbb{C}}(0, b)^o).$$

We call a weight $\underline{w}$ *summable* if $\sum_{n \geq 1} \frac{1}{w(n)} < \infty$. Both examples above are summable.

PROPOSITION 3.4. *If $\underline{w}$ is a summable weight, then $(L^2(\mathbb{T}), \mathcal{H}_{\underline{w}})$ is an adapted pair.*

(4) Note that here $\phi^{[N]} := \underbrace{\phi \circ \dots \circ \phi}_{N \text{ times}}$, whereas $(\phi^{[N]})^\ell := \underbrace{\phi^{[N]} \dots \phi^{[N]}}_{\ell \text{ times}}$.

Proof. We show that $B(R) := \{f \in \mathcal{H}_{\underline{w}} : \|f\|_{\underline{w}} \leq R\}$ is strongly compact in $L^2(m)$. To see this, let $f_j \in B(R)$ ($j \geq 1$). Then for $n \in \mathbb{Z}$ and $j \geq 1$,

$$|\widehat{f_j}(n)| \leq \frac{\|f_j\|_{\underline{w}}}{\sqrt{w(|n|)}} \leq \frac{R}{\sqrt{w(|n|)}}$$

and there exist $j_\ell \rightarrow \infty$ and $a \in \ell^2(\mathbb{Z})$ such that

$$\widehat{f_{j_\ell}}(n) \xrightarrow{\ell \rightarrow \infty} a(n).$$

We claim that $\sum_{n \in \mathbb{Z}} w(|n|)|a(n)|^2 \leq R^2$. Indeed,

$$\begin{aligned} R^2 &\geq \sum_{n \in \mathbb{Z}} w(|n|)|\widehat{f_{j_\ell}}(n)|^2 \leq \sum_{|n| \geq N} w(|n|)|\widehat{f_{j_\ell}}(n)|^2 \quad \forall N \geq 1, \\ &\xrightarrow{\ell \rightarrow \infty} \sum_{|n| \leq N} w(|n|)|a(n)|^2 \xrightarrow{N \rightarrow \infty} \sum_{n \in \mathbb{Z}} w(|n|)|a(n)|^2. \end{aligned}$$

Let $A := \sum_{n \in \mathbb{Z}} a(n)\chi^n \in \mathcal{H}_{\underline{w}}$ with $\|A\|_{\underline{w}} \leq R$.

To see that $f_\ell \xrightarrow[\ell \rightarrow \infty]{L^2(m)} A$, by the Riesz–Fischer theorem we get

$$\begin{aligned} \|f_{j_\ell} - A\|_{L^2(m)}^2 &= \sum_{n \in \mathbb{Z}} |\widehat{f_{j_\ell}}(n) - a(n)|^2 \xrightarrow{\ell \rightarrow \infty} 0 \\ &(\because 0 \xleftarrow{\ell \rightarrow \infty} |\widehat{f_{j_\ell}}(n) - a(n)|^2 \ll 1/w(|n|)). \quad \blacksquare \end{aligned}$$

The rest of this section is devoted to showing that the transfer operators of probability preserving, non-Möbius inner functions act quasicompactly on certain weighted Hilbert spaces.

Ivrii and Urbański [IU23] obtained spectral gaps for the action of $\widehat{\tau}$ on $W^{1,2}(\mathbb{T})$ ($\tau = \tau(\phi)$ with ϕ inner, $\phi(0) = 0$) and we obtain them on $\mathcal{H}_b$ (Proposition 3.5 below). In both cases, the minimal essential radius (as in §3.3.2) is the “Koenigs eigenvalue” $|\phi'(0)|$ (Proposition 3.7 below). However, we obtain superexponential decay of correlations on (e.g.) $\mathcal{H}_b$ when $\phi'(0) = 0$ (Proposition 3.6 below).

3.3.1. Admissible weighted Hilbert spaces. Call a summable weight $\underline{w} \in \mathbb{R}_+^{\mathbb{N}}$ and its associated weighted Hilbert space $\mathcal{H}_{\underline{w}}$ *admissible* if

$$(24) \quad \exists C = C_{\underline{w}} > 0 : W\left(\left\lfloor \frac{n}{K} \right\rfloor\right) \leq C \frac{w(n)}{w(K)} \quad \forall n \geq K \geq 1,$$

where $W(n) := \sum_{k=1}^n w(k)$.

We call any $C_{\underline{w}}$ satisfying (24) an *admissibility constant* for $\underline{w}$.

For example, for $b > 1$, $w_{1,b}(n) = b^n$ defines an admissible weight with e.g. $C_{w_{1,b}} = \frac{b}{b-1}$. Also, for $b, s > 1$, $\underline{w}_{s,b}$ defined by $w_{s,b}(n) := b^{n^s}$ defines an admissible weight.

On the other hand, for $t > 0$, $\underline{v}_t$ defined by $v_t(n) := n^t$ is not admissible (although summable for $t > 1$).

Recall from [Kat04, Def. I.2.10] that a Banach space $B \subset L^1(\mathbb{T}, m)$ is *homogeneous* if

$$(25) \quad \begin{aligned} f \in B, s \in \mathbb{T} &\implies f_s \in B, \|f_s\|_B = \|f\|, \\ \|f - f_s\|_B &\xrightarrow{s \rightarrow 0} 0 \quad \text{with } f_s(x) := f(x - s). \end{aligned}$$

Consequently (by [Kat04, Thm. I.2.11]), if B is homogeneous, then for $f \in B$,

$$(26) \quad f * p_r \in B, \quad \|f * p_r\|_B = \|f\|_B \quad \forall 0 < r < 1 \quad \& \quad f * p_r \xrightarrow[r \rightarrow 1]{B} f.$$

Any summably weighted Hilbert space $\mathcal{H}_{\underline{w}}$ is homogeneous.

PROPOSITION 3.5 (Exponential decay of correlations). *Let $\phi : \mathbb{D} \hookrightarrow \mathbb{D}$ be inner, non-Möbius with $\phi(0) = 0$ and $\tau = \tau(\phi)$ and let $\underline{w} \in \mathbb{R}_+^{\mathbb{N}}$ be admissible.*

If there exists $0 < R = R_{\underline{w}} < 1$ such that $\sum_{n \geq 1} \frac{1}{R^{2n}w(n)} < \infty$, then for every $|\phi'(0)| < \rho < 1$ there exists $M > 0$ such that

$$(27) \quad \|\widehat{\tau}^N u - \mathbb{E}(u)\|_{\underline{w}} \leq M \rho^N \|u\|_{\underline{w}} \quad \forall u \in \mathcal{H}_{\underline{w}}, N \geq 0,$$

where $\mathbb{E}(u) := \int_{\mathbb{T}} u \, dm = \widehat{u}(0)$.

Proof. Fix $\rho \in (|\phi'(0)|, 1)$. We first show that

$$(28) \quad \forall 0 < r < 1, \exists M > 0 : |\phi^{[N]}(z)| \leq M \rho^N \quad \forall z \in \mathbb{D}, |z| \leq r,$$

where $\phi^{[N]} = \underbrace{\phi \circ \cdots \circ \phi}_{N \text{ times}}$.

Proof of (28). By assumption, $\phi(z) = zg(z)$, where $g : \mathbb{D} \hookrightarrow \mathbb{D}$ is inner. Thus $\phi'(0) = g(0)$ and there exists $\mathfrak{r} = \mathfrak{r}_\rho$ such that

$$|g(z)| \leq \rho \quad \& \quad |\phi(z)| \leq \rho|z| \quad \forall |z| \leq \mathfrak{r}.$$

Next, there exists N_ρ such that $|\phi^{[N_\rho]}(z)| \leq \mathfrak{r}$ for $|z| \leq r$, whence for $|z| \leq r$,

$$|\phi^{[n]}(z)| \leq \rho^{n-N_\rho} \mathfrak{r} \quad \forall n > N_\rho,$$

and (28) follows.

Proof of (27) for $u \in \mathcal{H}_{\underline{w}} \cap H_0^2$. For $u \in \mathcal{H}_{\underline{w}} \cap H_0^2$ set $v_n := \widehat{u}(n)/R^n$ where $R = R_{\underline{w}}$. Then

$$\sum_{n \geq 1} |v_n| = \sum_{n \geq 1} \sqrt{w(n)} |\widehat{u}(n)| \cdot \frac{1}{R^n \sqrt{w(n)}} \leq \|u\|_{\underline{w}}^2 \sum_{n \geq 1} \frac{1}{R^{2n}w(n)} < \infty,$$

whence $v := \sum_{n \geq 1} v_n \chi^n \in C(\mathbb{T})$ and $v * p_R = u$.

Now

$$u = v * p_R = \int_{\mathbb{T}} v(t) p_{R\chi(t)} \, dm,$$

whence

$$u_N := \widehat{\tau}^N u = \int_{\mathbb{T}} v(t) \widehat{\tau}^N(p_{R\chi(t)}) dm = \int_{\mathbb{T}} v(t) p_{\phi^{[N]}(R\chi(t))} dm \quad \text{by (3)}$$

and

$$\begin{aligned} \widehat{u}_N(\ell) &= \int_{\mathbb{T}} \int_{\mathbb{T}} v(t) p_{\phi^{[N]}(R\chi(t))} \chi^{-\ell} dm dm \\ &= 1_{\mathbb{N}}(\ell) \int_{\mathbb{T}} v(t) \widehat{p}_{\phi^{[N]}(R\chi(t))}(\ell) dm \\ &= 1_{\mathbb{N}}(\ell) \int_{\mathbb{T}} v(t) \overline{(\phi^{[N]}(R\chi(t)))}^\ell dm. \end{aligned}$$

Thus

$$|\widehat{u}_N(\ell)|^2 \leq 1_{\mathbb{N}}(\ell) \int_{\mathbb{T}} |v|^2 dm \int_{\mathbb{T}} |\phi^{[N]}(R\chi(t))|^{2\ell} dt \leq \|u\|_{\underline{w}}^2 (M\rho^N)^{2\ell}.$$

where $M > 0$ is as in (28).

Let N_0 be such that $\mathfrak{b}_n := M^2 \rho^{2n} < \frac{1}{2}$ for $n \geq N_0$. For $N \geq N_0$,

$$\|\widehat{\tau}^N u\|_b^2 = \sum_{\ell \geq 1} b^\ell |\widehat{u}_N(\ell)|^2 \leq \|u\|_b^2 \sum_{\ell \geq 1} \mathfrak{b}_N^\ell = \frac{\mathfrak{b}_N}{1 - \mathfrak{b}_N} \|u\|_b^2 \leq 2M^2 b \rho^{2N} \|u\|_b^2,$$

proving (27).

To continue, let $w \in \mathfrak{H}_w$. Then

$$w = F + \overline{G} + \mathbb{E}(w) \quad \text{for some } F, G \in \mathfrak{H}_w \cap H_0^2$$

and

$$\widehat{\tau}^N w = \widehat{\tau}^N F + \overline{\widehat{\tau}^N G} + \mathbb{E}(w).$$

By (27) for F and G ,

$$\begin{aligned} \|\widehat{\tau}^N w - \mathbb{E}(w)\|_b^2 &= \|\widehat{\tau}^N F\|_b^2 + \|\widehat{\tau}^N G\|_b^2 \leq M\rho^{2N} (\|F\|_b^2 + \|G\|_b^2) \\ &\leq M\rho^{2N} \|w\|_{\underline{w}}^2. \blacksquare \end{aligned}$$

PROPOSITION 3.6 (Superexponential decay of correlations). *Suppose that $\phi(z) = z^\kappa \Phi(z)$ with $\kappa > 1$ and $\Phi : \mathbb{D} \hookrightarrow \text{inner}$ and let $\underline{w} \in \mathbb{R}_+^{\mathbb{N}}$ be admissible. Then*

$$(29) \quad \|\widehat{\tau}^N u - \mathbb{E}(u)\|_{\underline{w}} \leq \frac{\sqrt{C_{\underline{w}}}}{\sqrt{w(\kappa^N)}} \|u\|_{\underline{w}} \quad \forall u \in \mathfrak{H}_w, N \geq 0.$$

Proof. By Lemma 3.3, for $N, k \geq 1$,

$$\widehat{\tau}^N(\chi^k) = \sum_{\ell=1}^{\lfloor d/\kappa^N \rfloor} a^{(N)}(k, \ell) \chi^\ell,$$

where $a^{(N)}(k, \ell) = \overline{(\phi^N)^\ell(k)}$ and $\sum_{\ell=1}^0 = 0$. Thus also $\sum_{k \geq \kappa^N \ell} |a^{(N)}(k, \ell)|^2 = 1$.

Proof of (29) for $u \in \mathcal{H}_{\underline{w}} \cap H_0^2$. We have

$$\begin{aligned} \hat{\tau}^N u &= \sum_{k \geq 1} u_k \hat{\tau}^N(\chi^k) \quad (\text{where } u_k = \hat{u}(k)) \\ &= \sum_{\ell \geq 1, k \geq \kappa^N \ell} u_k a^{(N)}(k, \ell) \chi^\ell \\ &= \sum_{\ell \geq 1} \left(\sum_{k \geq \kappa^N \ell} u_k a^{(N)}(k, \ell) \right) \chi^\ell. \end{aligned}$$

Thus, using Cauchy–Schwarz and (23), we get

$$|(\hat{\tau}^N u)_\ell|^2 = \left| \sum_{k \geq \kappa^N \ell} u_k a^{(N)}(k, \ell) \right|^2 \leq \sum_{k \geq \kappa^N \ell} |u_k|^2$$

and

$$\begin{aligned} \|\hat{\tau}^N u\|_{\underline{w}}^2 &= \sum_{\ell \geq 1} w(\ell) |(\hat{\tau}^N u)_\ell|^2 \leq \sum_{\ell \geq 1} w(\ell) \sum_{k \geq \kappa^N \ell} |u_k|^2 \\ &\leq \sum_{k \geq \kappa^N} |u_k|^2 \sum_{1 \leq \ell \leq k/\kappa^N} w(\ell) = \sum_{k \geq \kappa^N} |u_k|^2 W(\lfloor k/\kappa^N \rfloor) \\ &= \sum_{k \geq \kappa^N} |u_k|^2 w(k) \cdot \frac{W(\lfloor k/\kappa^N \rfloor)}{w(k)} \leq \frac{C_{\underline{w}}}{w(\kappa^N)} \|u\|_{\underline{w}}^2 \end{aligned}$$

by (24), as desired.

To continue, let $u \in \mathcal{H}_{\underline{w}}$. Then

$$u = F + \overline{G} + \mathbb{E}(u) \quad \text{for some } F, G \in \mathcal{H}_{\underline{w}} \cap H_0^2$$

and

$$\hat{\tau}^N u = \hat{\tau}^N F + \overline{\hat{\tau}^N G} + \mathbb{E}(u).$$

By the previous step,

$$\begin{aligned} \|\hat{\tau}^N u - \mathbb{E}(u)\|_{\underline{w}}^2 &= \|T^N F\|_{\underline{w}}^2 + \|\hat{\tau}^N G\|_{\underline{w}}^2 \\ &\leq \frac{C_{\underline{w}}}{w(\kappa^N)} (\|F\|_{\underline{w}}^2 + \|G\|_{\underline{w}}^2) = \frac{C_{\underline{w}}}{w(\kappa^N)} \|u\|_{\underline{w}}^2. \quad \blacksquare \end{aligned}$$

3.3.2. Essential spectral radius. For $(\mathbb{T}, m, T)$ a weakly mixing, probabilistic preserving transformation, suppose that $\hat{T}$ is a Doeblin–Fortet operator on the adapted pair $(L^1(\mathbb{T}, m), \mathcal{L})$.

The *essential spectral radius* of $\hat{T} : \mathcal{L} \hookrightarrow \mathcal{L}$ is

$$\rho_{\text{ess}}(\hat{T}, \mathcal{L}) := \rho(T, \mathcal{L}_0) = \lim_{n \rightarrow \infty} \|\hat{T}^n\|_{\text{hom}(\mathcal{L}_0, \mathcal{L}_0)}^{1/n},$$

where $\mathcal{L}_0 := \{f \in \mathcal{L} : \mathbb{E}(f) = 0\}$. Equivalently, $\rho(\widehat{T}, \mathcal{L})$ is the greatest lower bound of the collection of $\theta \in (0, 1)$ satisfying (18).

PROPOSITION 3.7 (Minimal essential spectral radius ⁽⁵⁾). *Let $\phi : \mathbb{D} \hookrightarrow$ be non-Möbius, inner with $\phi(0) = 0$, and suppose that $\widehat{\tau}$ is a Doeblin–Fortet operator on the adapted pair $(L^1(m), \mathcal{L})$, where there exists $R > 0$ such that $p_z \in \mathcal{L}$ for all $z \in \mathbb{D}$ with $|z| < R$. Then $\rho_{\text{ess}}(\widehat{\tau}, \mathcal{L}) \geq |\phi'(0)|$.*

Proof. By Koenigs’ theorem ([Koe84], see also [Sha93, §6.1]), there exists $z \in \mathbb{D}$ with $|z| < R$ such that $|\phi^n(z)|^{1/n} \xrightarrow{n \rightarrow \infty} |\phi'(0)|$, and by (3), if $\widehat{\tau}$ satisfies (18) with constant θ on $\mathcal{L} \in \mathfrak{h}$, then

$$M\theta^n \geq \|\widehat{\tau}^n(p_z) - 1\|_{\mathcal{L}} \geq \|\widehat{\tau}^n(p_z) - 1\|_1 \geq \varepsilon |\phi^n(z)| = |\phi'(0)|^{n+o(n)}. \blacksquare$$

4. Perturbations and central limits. Let (X, m, T) be a weakly mixing, probability preserving transformation and let $\psi : X \rightarrow \mathbb{C}$ be measurable.

For $z \in \mathbb{C}$ such that $e^{z\psi} \in L^\infty(m)$, define the *perturbed* (or *twisted*) *transfer operator* $\Pi_{z,\psi} : L^1(m) \hookrightarrow$ by

$$\Pi_{z,\psi} f := \widehat{T}(e^{z\psi} f).$$

The operators $P_t = P_{t,\psi} := \widehat{T}(e^{it\psi} f)$ are also known as *characteristic function operators*, because $\mathbb{E}(P_t \mathbb{1}) = \mathbb{E}(e^{it\psi})$.

Let $\mathcal{L}$ be a Banach space of functions on $\mathbb{T}$. A function $f : \mathbb{T} \rightarrow \mathbb{R}$ is an $\mathcal{L}$ -multiplier if $f \cdot u \in \mathcal{L}$ for all $u \in \mathcal{L}$. Let $M(\mathcal{L}) := \{\mathcal{L}\text{-multipliers}\}$. By the Resonance Theorem,

$$\|f\|_{M(\mathcal{L})} := \sup \{\|fu\|_{\mathcal{L}} : u \in \mathcal{L}, \|u\|_{\mathcal{L}} = 1\} < \infty \quad \forall f \in M(\mathcal{L}).$$

Evidently, if $\mathbb{1} \in \mathcal{L}$, then $M(\mathcal{L}) \subseteq \mathcal{L}$. Indeed, for $\mathcal{L} = \text{Lip}$ or BV , $M(\mathcal{L}) = \mathcal{L}$.

In general, if $f \in M(\mathcal{L})$, then $f^N \in M(\mathcal{L})$ for all $N \geq 1$, with $\|f^N\|_{M(\mathcal{L})} \leq \|f\|_{M(\mathcal{L})}^N$, and $e^{zf} \in M(\mathcal{L})$ for all $z \in \mathbb{C}$, with $\|e^{zf}\|_{M(\mathcal{L})} \leq e^{|z| \|f\|_{M(\mathcal{L})}}$.

THEOREM 4.1 (Nagaev’s Theorem [Nag57, RE83] ⁽⁶⁾). *Suppose that $\widehat{T}$ is a Doeblin–Fortet operator on the adapted pair $(L^1(m), \mathcal{L})$ and that $\psi \in M(\mathcal{L}) \cap \mathcal{L}$ satisfies $\mathbb{E}(\psi) = 0$ and*

$$(30) \quad \begin{aligned} &\exists \varepsilon > 0 : P_{t,\psi} \in \text{hom}(\mathcal{L}, \mathcal{L}) \quad \forall |t| < \varepsilon; \\ &t \mapsto P_{t,\psi} \text{ is } C^2 \text{ from } (-\varepsilon, \varepsilon) \text{ to } \text{hom}(\mathcal{L}, \mathcal{L}). \end{aligned}$$

Then:

- (i) *There exists $0 < \mathcal{E} < \varepsilon$ such that P_t is a Doeblin–Fortet operator on $(L^2(m), \mathcal{L})$ for $|t| < \mathcal{E}$.*

⁽⁵⁾ Cf. [BCJ23].

⁽⁶⁾ See also [PP90, HH01], [AD01, Lemma 4.2].

(ii) *There are constants $K > 0$ and $\theta \in (0, 1)$, and C^2 functions $\lambda : B(0, \mathcal{E}) \rightarrow B_{\mathbb{C}}(0, 1)$ and $N : B(0, \mathcal{E}) \rightarrow \text{hom}(\mathcal{L}, \mathcal{L})$, such that*

$$(31) \quad \|P_t^n h - \lambda(t)^n N(t) h\|_{\mathcal{L}} \leq K \lambda(t)^n \|h\|_{\mathcal{L}} \quad \forall |t| < \mathcal{E}, n \geq 1, h \in \mathcal{L},$$

where for $|t| < \mathcal{E}$, $N(t)$ is a projection onto a one-dimensional subspace.

(iii) *If $\sigma_\psi := \lim_{n \rightarrow \infty} \frac{1}{n} \mathbb{E}(\psi_n^2) > 0$ as in (21), then $\lambda(t) = 1 - \sigma_\psi^2 t^2 + o(t^2)$ as $t \rightarrow 0$ and the conditional central limit theorem holds:*

$$(32) \quad \widehat{T}^n 1_{[\frac{\psi_n}{\sigma\sqrt{n}} \leq t]} \xrightarrow{n \rightarrow \infty} \frac{1}{\sqrt{2\pi}} \int_{-\infty}^t e^{-s^2/2} ds.$$

REMARKS. 1. If, in addition, P_t is a Doeblin–Fortet operator for all $t \in \mathbb{R}$ and ψ is T -aperiodic in the sense that $e^{it\psi} = \lambda g \circ \tau / g$ with $t \in \mathbb{R}$, $\lambda \in \mathbb{C}$, $|\lambda| = 1$ & $g : \mathbb{T} \rightarrow \mathbb{C}$ entails $t = 0$, $\lambda = 1$ & g constant, then the conditional local limit theorem holds: for $I \subset \mathbb{R}$ an interval, and $k_n \in \mathbb{Z}$ with $\frac{k_n}{\sigma\sqrt{n}} \rightarrow \kappa \in \mathbb{R}$,

$$(33) \quad \sigma\sqrt{n} \widehat{T}^n (1_{[\psi_n \in n\mathbb{E}(\psi) + k_n + I]}) \xrightarrow{n \rightarrow \infty} \frac{|I|}{\sqrt{2\pi}} e^{-\kappa^2/2}$$

and the skew product $(X \times \mathbb{R}, m \times \text{Leb}, T_\psi)$ is ergodic, where $T_\psi(x, y) = Tx, y + \psi(x)$. See [Gui89, GH88, AD01].

2. If (X, m, T) is an AFU map as in [Zwe98] (e.g. an Adler arc map), then for $\psi \in \text{BV}$, the characteristic function operator $P_{t,\psi}$ is a Doeblin–Fortet operator on $(L^1(m), \text{BV})$ for all $t \in \mathbb{R}$. See [ADSZ04, §5].

To obtain (32) for the stationary process (X, m, T, ψ) via Nagaev’s theorem, we verify (30) and then $\sigma_\psi > 0$.

THEOREM 4.2 (Analyticity of perturbation). *Suppose that $\phi : \mathbb{D} \hookrightarrow$ is non-Möbius inner with $\phi(0) = 0$.*

Let $b > 1$ and let $\underline{w}$ be a summable weight $\sum_{n \geq 1} b^n / w(n) < \infty$.

If $\psi \in \mathcal{H}_{\underline{w}}$ then for all $z \in \mathbb{C}$,

$$\Pi_z := \Pi_{z,\psi} \in \text{hom}(\mathcal{K}_b, \mathcal{K}_b).$$

Moreover, $\mathbb{C} \ni z \mapsto \Pi_z \in \text{hom}(\mathcal{K}_b, \mathcal{K}_b)$ is holomorphic with

$$\frac{d^n \Pi_z}{dz^n}(f) = \Pi_z(\psi^n f) =: \Pi_z^{(n)}(f).$$

In particular, (30) holds.

LEMMA 4.3 (Multiplier lemma).

(i) *Let $b > 1$ and let $\underline{w}$ be a summable weight with $\sum_{n \geq 1} b^n / w(n) < \infty$. Then $\mathcal{H}_{\underline{w}} \subseteq M(\mathcal{K}_b)$ with*

$$(34) \quad \|f\|_{M(\mathcal{K}_b)} \leq R_{b,\underline{w}} \|f\|_{\underline{w}} \quad \forall f \in \mathcal{H}_{\underline{w}},$$

where $R_{b,\underline{w}} := \sqrt{1 + 2 \sum_{n \geq 1} b^n / w(n)}$.

- (ii) If $\underline{w}$ is a weight with $\overline{\lim}_{n \rightarrow \infty} w(n+1)/w(n) = \infty$, then there are no non-constant multipliers of $\mathcal{H}_{\underline{w}}$.

Note that if $w(n) = B^n$ with $B > b$ then $R_{b,\underline{w}} = \sqrt{\frac{B+b}{B-b}}$.

Proof of (i). We have

$$\begin{aligned}
 \|fg\|_{\mathcal{H}_b}^2 &= \sum_{n \in \mathbb{Z}} b^{|n|} |\widehat{fg}(n)|^2 = \sum_{n \in \mathbb{Z}} b^{|n|} \left| \sum_{k \in \mathbb{Z}} \widehat{f}(k) \widehat{g}(n-k) \right|^2 \\
 &= \sum_{n \in \mathbb{Z}} b^{|n|} \left| \sum_{k \in \mathbb{Z}} \sqrt{w(|k|)} |\widehat{f}(k)| \frac{|\widehat{g}(n-k)|}{\sqrt{w(|k|)}} \right|^2 \\
 &\leq \sum_{n \in \mathbb{Z}} b^{|n|} \sum_{k \in \mathbb{Z}} w(|k|) |\widehat{f}(k)|^2 \sum_{\ell \in \mathbb{Z}} \frac{|\widehat{g}(n-\ell)|^2}{w(|\ell|)} \\
 &= \|f\|_{\underline{w}}^2 \sum_{n, \ell \in \mathbb{Z}} b^{|n|} \frac{|\widehat{g}(n-\ell)|^2}{w(|\ell|)} = \|f\|_{\underline{w}}^2 \sum_{n, \ell \in \mathbb{Z}} \frac{b^{|\ell|}}{w(|\ell|)} b^{|n|-|\ell|} |\widehat{g}(n-\ell)|^2 \\
 &\leq \|f\|_{\underline{w}}^2 \sum_{n, \ell \in \mathbb{Z}} \frac{b^{|\ell|}}{w(|\ell|)} b^{|n-\ell|} |\widehat{g}(n-\ell)|^2 = \|f\|_{\underline{w}}^2 \|g\|_{\mathcal{H}_b}^2 \sum_{\ell \in \mathbb{Z}} \frac{b^{|\ell|}}{w(|\ell|)} \\
 &= R_{b,\underline{w}}^2 \|f\|_{\underline{w}}^2 \|g\|_{\mathcal{H}_b}^2.
 \end{aligned}$$

Proof of (ii). Suppose otherwise. Then there exist $f \in M(\mathcal{H}_{\underline{w}})$ and $\ell \geq 1$ with $\widehat{f}(\ell) \neq 0$. Suppose that $\nu_k \uparrow \infty$ with $w(\nu_k + 1) \geq kw(\nu_k)$ for $k \geq 1$. It follows that

$$\infty > \|f\|_{M(\mathcal{H}_{\underline{w}})}^2 \geq \frac{\|f\chi^{\nu_k}\|_{\mathcal{H}_{\underline{w}}}^2}{w(\nu_k)} \geq |\widehat{f}(\ell)|^2 \frac{w(\ell + \nu_k)}{w(\nu_k)} \geq k |\widehat{f}(\ell)|^2 \xrightarrow{k \rightarrow \infty} \infty. \blacksquare$$

Proof of Theorem 4.2. It suffices to show that for every $\omega \in \mathbb{C}$, there exists $\varepsilon = \varepsilon_\omega > 0$ such that

$$(35) \quad \sum_{n=0}^N \frac{(z - \omega)^n}{n!} \Pi_\omega^{(n)} \xrightarrow[N \rightarrow \infty]{\text{hom}(\mathcal{H}_b, \mathcal{H}_b)} \Pi_z \quad \forall z \in B(\omega, \varepsilon).$$

To this end, fix $\omega \in \mathbb{C}$ and $1 < b < b_1 < b_2$ with $\sum_{n \geq 1} b_2^n / w(n) < \infty$. Let $f \in \mathcal{H}_b$. Then for $k \geq 1$,

$$\begin{aligned}
 \|\Pi_\omega^{(k)}(f)\|_{\mathcal{H}_b} &= \|\widehat{\tau}(\psi^k e^{\omega\psi} f)\|_{\mathcal{H}_b} \leq M\rho \|\psi^k e^{\omega\psi} f\|_{\mathcal{H}_b} \quad \text{by (27)} \\
 &\leq M\rho R_{b, \underline{w}_1, b_1} \|f\|_{\mathcal{H}_b} \|\psi^k e^{\omega\psi}\|_{b_1} \quad \text{by (34)} \\
 &= M\rho \sqrt{\frac{b_1 + b}{b_1 - b}} \|f\|_{\mathcal{H}_b} \|\psi^k e^{\omega\psi}\|_{b_1}.
 \end{aligned}$$

To continue, by repeated application of (34),

$$\begin{aligned} \|\psi^k e^{\omega\psi}\|_{b_1} &\leq \left(\sqrt{\frac{b_2 + b_1}{b_2 - b_1}} \right)^{k+1} \|\psi\|_{b_2}^k \|e^{\omega\psi}\|_{b_2} \\ &\leq \left(\sqrt{\frac{b_2 + b_1}{b_2 - b_1}} \right)^{k+1} \|\psi\|_{b_2}^k \exp[|\omega| R_{b_2, \underline{w}} \|\psi\|_{\underline{w}}^2]. \end{aligned}$$

Thus

$$\|II_\omega^{(k)}\|_{\text{hom}(\mathcal{K}_b, \mathcal{K}_b)} \ll \left(\sqrt{\frac{b_2 + b_1}{b_2 - b_1}} \right)^k$$

and (35) holds with $\varepsilon_\omega = \sqrt{\frac{b_2 - b_1}{b_2 + b_1}}$. ■

4.1. Periodicity. Let $\phi : \mathbb{D} \hookrightarrow \mathbb{D}$ be non-Möbius inner with $\phi(0) = 0$ and let $\psi : \mathbb{T} \rightarrow \mathbb{R}$ be measurable.

We call $t \in \mathbb{R}$ a τ -period of ψ if there are $t \in \mathbb{R}$, $\lambda \in \mathbb{C}$ with $|\lambda| = 1$ and $g : \mathbb{T} \rightarrow \mathbb{C}$ measurable such that $e^{it\psi} = \lambda g \circ \tau / g$.

We denote the collection of τ -periods of ψ by $\mathcal{Q}(\psi)$, and we call ψ τ -aperiodic if $\mathcal{Q}(\psi) = \{0\}$ and τ -periodic otherwise.

It is standard to show that for $t \in \mathbb{R}$, $\lambda \in \mathbb{C}$ with $|\lambda| = 1$ and $f \in L^1(m)$,

$$(36) \quad e^{it\psi} f = \lambda f \circ \tau \iff P_t(f) := \widehat{\tau}(e^{it\psi} f) = \lambda f,$$

and also, if the limit $\lim_{n \rightarrow \infty} \frac{1}{n} \mathbb{E}(\psi_n^2) =: \sigma_\psi^2 \geq 0$ exists, then $\sigma_\psi > 0$ if $\mathcal{Q}(\psi)$ is discrete.

THEOREM 4.4. *Let $\phi : \mathbb{D} \hookrightarrow \mathbb{D}$ be non-Möbius inner with with Denjoy–Wolff point in $\mathbb{D}$ and nonempty singularity set.*

- (i) *If $\psi : \mathbb{T} \rightarrow \mathbb{R}$ is nonconstant and real analytic, then $\mathcal{Q}(\psi)$ is discrete and the stationary process $(\mathbb{T}, \pi_{\mathfrak{d}_\phi}, \tau, \psi)$ satisfies (32).*
- (ii) *If both $\mathcal{Z}(\phi)$ and $\mathfrak{s}_\phi$ are finite, and $\psi : \mathbb{T} \rightarrow \mathbb{R}$ is nonconstant and of bounded variation, then ψ is τ -aperiodic and the stationary process $(\mathbb{T}, \pi_{\mathfrak{d}_\phi}, \tau, \psi)$ satisfies (33).*

The assumption $\mathfrak{s}_\phi \neq \emptyset$ is essential. If ϕ is a finite Blaschke product with $\mathfrak{d}(\phi) \in \mathbb{D}$ and $g : \mathbb{T} \rightarrow \mathbb{R}$ nonconstant and real analytic, then so is $\psi := g - g \circ \tau : \mathbb{T} \rightarrow \mathbb{R}$, whence $P_t(\chi(tg)) = \chi(tg)$ for all $t \in \mathbb{R}$ and $\mathcal{Q}(\psi) = \mathbb{R}$.

Proof of (i). Let $\mathcal{E} > 0$ be as in (i). Fix $1 < b < B$ with $\psi \in \mathcal{K}_B$. If $\mathcal{Q}(\psi)$ is not discrete, then there are $t \in (0, \mathcal{E}) \cap \mathcal{Q}(\psi)$, $f \in L^1(m)$ and $\lambda \in \mathbb{C}$ with $|\lambda| = 1$ such that $P_t(f) = \lambda f$. For $t \in (0, \mathcal{E}) \cap \mathcal{Q}(\psi)$, P_t is a Doeblin–Fortet operator on $(L^2(m), \mathcal{K}_b)$. Write $Q := \bar{\lambda} P_t$. Then $Q : \mathcal{K}_b \hookrightarrow \mathcal{K}_b$ is a Doeblin–Fortet operator, $Qf = f$ and by Lemma 3.1, $f \in \mathcal{K}_b$.

By (36) we have

$$(37) \quad f(\tau x) = \bar{\lambda} f(x) e^{it\psi(x)} \quad \text{for } m\text{-a.e. } x \in \mathbb{T}.$$

Using harmonization, we see that (37) holds for all $x \in \Lambda_\phi$.

Now suppose that v is a point of discontinuity of τ . Then $\chi(v) \in \mathfrak{s}_\phi$.

By Seidel's theorem [Sei34] (see also [Zyg02, Thm. 7.48]), for every $w \in \mathbb{T}$ there exists $z_n = z_n(w) \in \Lambda_\phi$ such that $z_n \rightarrow v$ and $\tau(z_n) = w$. Thus

$$f(w) = f(\tau(z_n)) = \bar{\lambda} f(z_n) \psi(z_n) \xrightarrow{n \rightarrow \infty} \bar{\lambda} f(v) \psi(v),$$

so f is constant, $e^{it\psi} \equiv \lambda$, whence ψ (being continuous) is constant. ■

Proof of (ii). Under the assumptions of (ii), $(\mathbb{T}, \pi_{\mathfrak{d}(\phi)}, \tau)$ is an Adler arc map, whence an AFU map as in [Zwe98] and $P_{t,\psi}$ is a Doeblin–Fortet operator on $(L^1(m), \mathbf{BV}(\mathbb{T}))$ for all $t \in \mathbb{R}$. Using [ADSZ04, §5], it suffices to prove that ψ is τ -aperiodic.

To this end, suppose that $f : \mathbb{T} \rightarrow \mathbb{C}$ is measurable and satisfies (37). By Lemma 3.1, $f \in \mathbf{BV}$. We must show that f is constant. Suppose otherwise; then there exist $\xi, \zeta \in \mathbb{T}$ such that $|f(\xi) - f(\zeta)| =: \eta > 0$.

Let $v \in \chi^{-1}\mathfrak{s}_\phi$. Then there exists $z_n \rightarrow v$ monotonically with $\tau(z_{2n}) = \xi$ and $\tau(z_{2n+1}) = \zeta$. The function $G(x) := \bar{\lambda} f(x) e^{it\psi(x)}$ is in $\mathbf{BV}$, thus so is $f \circ \tau = G$. Therefore

$$\infty > \bigvee f \circ \tau \geq \sum_{n \geq 1} |f(\tau(z_n)) - f(\tau(z_{n+1}))| = \sum_{n \geq 1} \eta = \infty. \quad \blacksquare$$

5. A local version of Aleksandrov's theorem. Fix $A \in \mathcal{B}(\mathbb{T})$ with $m(A) > 0$ and a sub- σ -algebra $\mathcal{C} \subseteq \mathcal{B}(A)$.

For $1 \leq p \leq \infty$ write

$$H_0^p(A, \mathcal{C}) := \{f \in H_0^p : f|_A \text{ is } \mathcal{C}\text{-measurable}\}.$$

By [Rud74, Thm. 17.18] if $f, g \in H_0^2(m)$ and $f|_A \equiv g|_A$, then $f \equiv g$.

Let $P : H_0^2 \rightarrow H_0^2(A, \mathcal{C})$ be the orthogonal projection in the sense that

$$(38) \quad (\text{Id} - P)H_0^2 \subset H_0^2(A, \mathcal{C})^\perp.$$

Call $(A, \mathcal{C})$ an *analytic pair* if

$$\mathbb{E}_{m_A}^{\mathcal{C}}(f) = (Pf)|_A \quad \text{for } f \in H_0^2,$$

where $m_A(B) := m(A \cap B)$ and $\mathbb{E}_{m_A}^{\mathcal{C}}(f)$ is conditional expectation on the measure space (A, m_A) with respect to the sub- σ -algebra $\mathcal{C} \subseteq \mathcal{B}(A)$.

Let $A \in \mathcal{B}(\mathbb{T})$ with $m(A) > 0$. It is easy to see that $(A, \mathcal{C})$ is an analytic pair if either $\mathcal{C} \stackrel{m_A}{=} \mathcal{B}(A)$ (in which case $H_0^2(A, \mathcal{C}) = H_0^2$), or $\mathcal{C} \stackrel{m_A}{=} \{\emptyset, A\}$ (in which case $H_0^2(A, \mathcal{C}) = \{0\}$).

Next, we give an example which turns out to be general.

EXAMPLE 5.1. If $A \in \tau^{-1}\mathcal{B}(\mathbb{T})$ and $\mathcal{C} = \tau^{-1}\mathcal{B}(\mathbb{T}) \cap A$, where $\tau = \tau(\phi)$ with $\phi : \mathcal{D} \hookrightarrow \text{inner}$ with $\phi(0) = 0$, then $(A, \mathcal{C})$ is an analytic pair.

Proof. We have

$$\begin{aligned} H_0^2(A, \tau^{-1}\mathcal{B}) &:= \{f \in H_0^2 : f|_A \text{ is } \tau^{-1}\mathcal{B}\text{-measurable}\} \\ &= \{f \in H_0^2 : \exists g \in H_0^2, f|_A = g \circ \tau|_A\} \\ &= H_0^2 \circ \tau \quad \text{by [Rud74, Thm. 17.18]}. \end{aligned}$$

To continue, note that because $m \circ \tau^{-1} = m$,

$$\mathbb{E}_m^{\tau^{-1}\mathcal{B}(\mathbb{T})}(f) = \hat{\tau}(f) \circ \tau.$$

Since $\hat{\tau} : H_0^2 \hookrightarrow$,

$$P := \mathbb{E}_m^{\tau^{-1}\mathcal{B}(\mathbb{T})} = \hat{\tau}(f) \circ \tau : H_0^2 \rightarrow H_0^2 \circ \tau$$

is an orthogonal projection.

Let $A = \tau^{-1}B$. Then

$$\mathbb{E}_{m_A}^{\mathcal{C}}(f) = \mathbb{E}_m^{\tau^{-1}\mathcal{B}}(1_B \circ \tau f)|_A = ((1_B \hat{\tau} f) \circ \tau)|_A = (Pf)|_A. \blacksquare$$

THEOREM 5.2. *If $(A, \mathcal{C})$ is an analytic pair, then either $\mathcal{C} = \{\emptyset, A\}$, or there exists $\phi : \mathcal{D} \hookrightarrow \text{inner}$ with $\phi(0) = 0$ such that $A \in \tau^{-1}\mathcal{B}$ and $\mathcal{C} = A \cap \tau^{-1}\mathcal{B}$, where $\tau = \tau(\phi)$.*

The cases with $C = \mathbb{T}$ are established in [Ale86] ⁽⁷⁾.

Proof. We divide the proof into several claims.

CLAIM 1. $P(gf) = gP(f)$ for all $f \in H_0^\infty$ and $g \in H_0^2(A, \mathcal{C})$.

Proof. We have

$$P(gf)|_A = \mathbb{E}_{m_A}^{\mathcal{C}}(fg) = g|_A \mathbb{E}_{m_A}^{\mathcal{C}}(f) = (gP(f))|_A,$$

and by [Rud74, Thm. 17.18], $P(gf) = gP(f)$ a.s. on $\mathbb{T}$.

CLAIM 2. *For every $f \in L^2(A, \mathcal{C}, m_A)$ there are $g, h \in H_0^2(A, \mathcal{C})$ and $\gamma \in \mathbb{C}$ such that*

$$f = (g + \bar{h} + \gamma)|_A.$$

Proof. There are $G, H \in H_0^2$ and $\gamma \in \mathbb{C}$ such that

$$f = G + \overline{H} + \gamma.$$

Next, $g = PG$, $h = PH \in H_0^2(A, \mathcal{C})$ and a.s. on A ,

$$(g + \bar{h} + \gamma) = (PG + \overline{PH} + \gamma) = \mathbb{E}_{m_A}^{\mathcal{C}}(f) = f.$$

CLAIM 3. *Let $A \in \mathcal{B}(\mathbb{T})$ and let $\mathcal{C}, \mathcal{C}' \subseteq \mathcal{B}(A)$ be such that both $(A, \mathcal{C})$ and $(A, \mathcal{C}')$ are analytic pairs. Then*

$$\mathcal{C} \leq \mathcal{C}' \iff H_0^2(A, \mathcal{C}) \subset H_0^2(A, \mathcal{C}').$$

Proof. $\Rightarrow$ follows from the definition and $\Leftarrow$ follows from Claim 2.

⁽⁷⁾ See also [Sak07, Thm. 5.6].

CLAIM 4. $L := \overline{\text{span}}\{fg : f \in H_0^2(A, \mathcal{C}), g \in H_0^\infty(A, \mathcal{C})\} \subsetneq H_0^2(A, \mathcal{C})$.

Proof. Let

$$d := \min \{k \geq 0 : \chi^{-k}g \in H^2 \ \forall g \in H_0^2(A, \mathcal{C})\}.$$

Then $d \in \mathbb{N}$.

By Claim 2, there exists nonconstant $G \in H_0^\infty(A, \mathcal{C})$ with $\chi^{-1}G \in H^2$. Thus $\chi^{-d-1}g \in H^2$ for all $g \in L$, whence $H_0^2(A, \mathcal{C}) \setminus L \neq \emptyset$.

CLAIM 5. Every $\phi \in H_0^2(A, \mathcal{C}) \cap L^\perp$ with $\|\phi\|_2 = 1$ is inner with $\phi(0) = 0$.

Proof. For $n \geq 1$, $P(\phi\chi^n) \stackrel{\text{Claim 1}}{=} \phi P(\chi^n) \in L$, whence $\phi \perp \phi P(\chi^n)$ ($\because \phi \in L^\perp$). Thus for $n \geq 1$,

$$(\widehat{|\phi|^2})(n) = \langle \phi, \phi\chi^n \rangle = \langle \phi, P(\phi\chi^n) \rangle = \langle \phi, \phi P(\chi^n) \rangle = 0,$$

whence also $(\widehat{|\phi|^2})(-n) = \overline{(\widehat{|\phi|^2})(n)} = 0$ and

$$|\phi|^2 = (\widehat{|\phi|^2})(0) = \|\phi\|_2^2 = 1. \blacksquare$$

Fix $\phi \in H_0^2(A, \mathcal{C}) \cap L^\perp$ (inner with $\phi(0) = 0$).

CLAIM 6. $H_0^2(A, \mathcal{C}) = H_0^2 \circ \tau = H_0^2(\mathbb{T}, \tau^{-1}\mathcal{B})$, where $\tau = \tau(\phi)$.

Proof. Since $\phi \in H_0^\infty(A, \mathcal{C})$, we have (i) $A \cap \tau^{-1}\mathcal{B}(\mathbb{T}) \subseteq \mathcal{C}$ and (ii) $\phi^n \in H_0^\infty(A, \mathcal{C})$ for all $n \geq 1$. It follows that every $F \in H_0^2 \circ \tau$ is in $H_0^2(A, \mathcal{C})$ having the form $F \circ \chi = \sum_{n \geq 1} a_n \phi^n$ with $(a_k : k \geq 1) \in \ell^2$. Thus $H_0^2(A, \mathcal{C}) \supseteq H_0^2 \circ \tau$.

To show equality we will prove that

$$M := H_0^2(A, \mathcal{C}) \cap (H_0^2 \circ \tau)^\perp = \{0\}.$$

To this end, we show first that

$$(39) \quad \overline{\phi}M \subset M.$$

Indeed, let $g \in M$; then $g \perp \phi^j$ for all $j \geq 1$ and

$$(40) \quad \langle \overline{\phi}g, \phi^j \rangle = 0 \quad \forall j \geq 0.$$

For $k \geq 1$,

$$\langle \overline{\phi}g, \chi^{-k} \rangle = \langle g\chi^k, \phi \rangle \stackrel{(38)}{=} \langle P(g\chi^k), \phi \rangle = \langle gP(\chi^k), \phi \rangle = 0.$$

Thus $\overline{\phi}g \in H_0^2(A, \mathcal{C})$.

Each $H \in H_0^2(A, \tau^{-1}\mathcal{B})$ is of the form $H = h \circ \tau$ with $h \in H_0^2$. By (40),

$$\overline{\phi}g \perp \sum_{k \geq 0} \hat{h}(k)\phi^k = h \circ \tau = H$$

and $\overline{\phi}g \in M$, proving (39).

To see that $M = \{0\}$, suppose otherwise: let $g \in M$, $g \neq 0$; then by (39) (used repeatedly) $\overline{\phi}^j g \in M$ for all $j \geq 1$, which is impossible unless $g \equiv 0$. $\blacksquare$

CLAIM 7. $A \in \tau^{-1}\mathcal{B}(\mathbb{T})$.

Proof. By Claim 6, $H_0^2(A, \mathcal{C}) = H_0^2 \circ \tau$ and $Pf = \widehat{\tau}(f) \circ \tau$ and

$$\mathbb{E}_{m_A}^{\mathcal{C}}(f) = P(1_A f)|_A$$

for all $f \in H_0^2$ and hence $f \in L^2$.

In particular, $(\widehat{\tau}(1_A) \circ \tau)|_A = 1$ and we claim that $\widehat{\tau}(1_A) \circ \tau = 1_A$.

To see this, note that

$$\widehat{\tau}(1_A) \circ \tau = 1_A + J,$$

where $J := \widehat{\tau}(1_A) \circ \tau \cdot 1_{A^c} \geq 0$.

Now,

$$m(A) = \mathbb{E}_m(\widehat{\tau}(1_A) \circ \tau) = m(A) + \mathbb{E}_m(J),$$

whence $\mathbb{E}_m(J) = 0$, $J = 0$ a.s., $\widehat{\tau}(1_A) \circ \tau = 1_A$ and $A \in \tau^{-1}\mathcal{B}(\mathbb{T})$. ■

References

- [Aar78] J. Aaronson, *Ergodic theory for inner functions of the upper half plane*, Ann. Inst. H. Poincaré Sect. B (N.S.) 14 (1978), 233–253.
- [Aar97] J. Aaronson, *An Introduction to Infinite Ergodic Theory*, Math. Surveys Monogr. 50, Amer. Math. Soc., Providence, RI, 1997.
- [AD01] J. Aaronson and M. Denker, *Local limit theorems for partial sums of stationary sequences generated by Gibbs–Markov maps*, Stoch. Dynam. 1 (2001), 193–237.
- [ADSZ04] J. Aaronson, M. Denker, O. Sarig, and R. Zweimüller, *Aperiodicity of cocycles and conditional local limit theorems*, Stoch. Dynam. 4 (2004), 31–62.
- [Adl73] R. Adler, *F-expansions revisited*, in: Recent Advances in Topological Dynamics (New Haven, CN, 1972; in honor of Gustav Arnold Hedlund), Lecture Notes in Math. 318, Springer, Berlin, 1973, 1–5.
- [Ale86] A. B. Aleksandrov, *Measurable partitions of the circle induced by inner functions*, Zap. Nauchn. Sem. Leningrad. Otdel. Mat. Inst. Steklov. 149 (1986), 103–106, 188 (in Russian).
- [Ale87] A. B. Aleksandrov, *Multiplicity of boundary values of inner functions*, Izv. Akad. Nauk Armyan. SSR Ser. Mat. 22 (1987), 490–503, 515 (in Russian).
- [Boo57] G. Boole, *On the comparison of transcendents, with certain applications to the theory of definite integrals*, Philos. Trans. Roy. Soc. London 147 (1857), 745–803.
- [BCJ23] O. Butterley, G. Canestrari, and S. Jain, *Discontinuities cause essential spectrum*, Comm. Math. Phys. 398 (2023), 627–653.
- [Coh80] D. L. Cohn, *Measure Theory*, Birkhäuser, Boston, MA, 1980.
- [Cra91] M. Craizer, *Entropy of inner functions*, Israel J. Math. 74 (1991), 129–168.
- [DM78] C. Dellacherie and P.-A. Meyer, *Probabilities and Potential*, North-Holland Math. Stud. 29, North-Holland, Amsterdam, 1978.
- [Den26] A. Denjoy, *Sur l'itération des fonctions analytiques*, C. R. Acad. Sci. Paris 182 (1926), 255–257.
- [DM91] C. I. Doering and R. Mañé, *The Dynamics of Inner Functions*, Ensaios Mat. 3, Sociedade Brasileira de Matemática, Rio de Janeiro, 1991.
- [Don65] W. F. Donoghue, Jr., *On the perturbation of spectra*, Comm. Pure Appl. Math. 18 (1965), 559–579.

- [Gla77] J. W. L. Glaisher, *Note on certain theorems in definite integration*, Messenger (2) 7 (1877), 63–74.
- [Gor04] M. I. Gordin, *A remark on the martingale approximation method for proving the central limit theorem for stationary random sequences*, Zap. Nauchn. Sem. POMI 311 (2004), 124–132, 299–300 (in Russian).
- [Gui89] Y. Guivarc’h, *Propriétés ergodiques, en mesure infinie, de certains systèmes dynamiques fibrés*, Ergodic Theory Dynam. Systems 9 (1989), 433–453.
- [GH88] Y. Guivarc’h et J. Hardy, *Théorèmes limites pour une classe de chaînes de Markov et applications aux difféomorphismes d’Anosov*, Ann. Inst. H. Poincaré Probab. Statist. 24 (1988), 73–98.
- [Hei77] M. Heins, *On the finite angular derivatives of an analytic function mapping the open unit disk into itself*, J. London Math. Soc. (2) 15 (1977), 239–254.
- [HH01] H. Hennion and L. Hervé, *Limit Theorems for Markov Chains and Stochastic Properties of Dynamical Systems by Quasi-Compactness*, Lecture Notes in Math. 1766, Springer, Berlin, 2001.
- [ITM50] C. T. Ionescu Tulcea et G. Marinescu, *Théorie ergodique pour des classes d’opérations non complètement continues*, Ann. of Math. (2) 52 (1950), 140–147.
- [IU23] O. Ivrii and M. Urbański, *Inner functions, composition operators, symbolic dynamics and thermodynamic formalism*, arXiv:2308.16063v2 (2025).
- [Jan41] W. Jankoff, *Sur l’uniformisation des ensembles (A)*, C. R. (Dokl.) Acad. Sci. URSS N. Ser. 30 (1941), 597–598.
- [Kat04] Y. Katznelson, *An Introduction to Harmonic Analysis*, 3rd ed., Cambridge Math. Library, Cambridge Univ. Press, Cambridge, 2004.
- [Koe84] G. Koenigs, *Investigations of the integrals of certain functional equations*, Ann. Sci. École Norm. Sup. (3) 1 (1884), 1–41.
- [LY73] A. Lasota and J. A. Yorke, *On the existence of invariant measures for piecewise monotonic transformations*, Trans. Amer. Math. Soc. 186 (1973), 481–488 (1974).
- [Leo61] V. P. Leonov, *On the dispersion of time means of a stationary stochastic process*, Teor. Veroyatnost. i Primenen. 6 (1961), 93–101 (in Russian).
- [Let77] G. Letac, *Which functions preserve Cauchy laws?* Proc. Amer. Math. Soc. 67 (1977), 277–286.
- [Lus30] N. Lusin, *Sur le problème de M. Jacques Hadamard d’uniformisation des ensembles*, Mathematica (Cluj) 4 (1930), 54–66.
- [Mar89] N. F. G. Martin, *On ergodic properties of restrictions of inner functions*, Ergodic Theory Dynam. Systems 9 (1989), 137–151.
- [Nad81] M. G. Nadkarni, *Some measure-theoretic remarks on inner functions and related results*, Sankhyā Ser. A 43 (1981), 251–259.
- [Nag57] S. V. Nagaev, *Some limit theorems for stationary Markov chains*, Teor. Veroyatnost. i Primenen. 2 (1957), 389–416 (in Russian).
- [Neu78] J. H. Neuwirth, *Ergodicity of some mappings of the circle and the line*, Israel J. Math. 31 (1978), 359–367.
- [NSiG22] A. Nicolau and O. Soler i Gibert, *A central limit theorem for inner functions*, Adv. Math. 401 (2022), art. 108318, 39 pp.
- [Nor68] E. A. Nordgren, *Composition operators*, Canad. J. Math. 20 (1968), 442–449.
- [Nor72] M. F. Norman, *Markov Processes and Learning Models*, Math. Sci. Engrg. 84, Academic Press, New York, 1972.
- [PP90] W. Parry and M. Pollicott, *Zeta functions and the periodic orbit structure of hyperbolic dynamics*, Astérisque 187–188 (1990), 268 pp.

- [Rok61] V. A. Rokhlin, *Exact endomorphisms of a Lebesgue space*, Izv. Akad. Nauk SSSR Ser. Mat. 25 (1961), 499–530 (in Russian); English transl.: Amer. Math. Soc. Transl. (2) 39 (1964), 1–36.
- [RE83] J. Rousseau-Egele, *Un théorème de la limite locale pour une classe de transformations dilatantes et monotones par morceaux*, Ann. Probab. 11 (1983), 772–788.
- [Rud74] W. Rudin, *Real and Complex Analysis*, 2nd ed., McGraw-Hill, New York, 1974.
- [Ryc83] M. Rychlik, *Bounded variation and invariant measures*, Studia Math. 76 (1983), 69–80.
- [Sak07] E. Saksman, *An elementary introduction to Clark measures*, in: Topics in Complex Analysis and Operator Theory, Univ. Málaga, Málaga, 2007, 85–136.
- [Sei34] W. Seidel, *On the distribution of values of bounded analytic functions*, Trans. Amer. Math. Soc. 36 (1934), 201–226.
- [Sha93] J. H. Shapiro, *Composition Operators and Classical Function Theory*, Universitext, Springer, New York, 1993.
- [Smi29] V. Smirnov, *Sur les valeurs limites des fonctions, régulières à l'intérieur d'un cercle*, J. Soc. Phys.-Math. Leningrad 2 (1929), 22–37; English transl.: Topics in Interpolation Theory, Birkhäuser, 1997, 481–494.
- [Sri98] S. M. Srivastava, *A Course on Borel Sets*, Grad. Texts in Math. 180, Springer, New York, 1998.
- [vN49] J. von Neumann, *On rings of operators. Reduction theory*, Ann. of Math. (2) 50 (1949), 401–485.
- [Wol26] J. Wolff, *Sur l'itération des fonctions bornées*, C. R. Acad. Sci. Paris 182 (1926), 200–201.
- [YK41] K. Yosida and S. Kakutani, *Operator-theoretical treatment of Markoff's process and mean ergodic theorem*, Ann. of Math. (2) 42 (1941), 188–228.
- [Zwe98] R. Zweimüller, *Ergodic structure and invariant densities of non-Markovian interval maps with indifferent fixed points*, Nonlinearity 11 (1998), 1263–1276.
- [Zyg02] A. Zygmund, *Trigonometric Series. Vols. I, II*, 3rd ed., Cambridge Math. Library, Cambridge Univ. Press, Cambridge, 2002.

Jon Aaronson
 School of Mathematical Sciences
 Tel Aviv University
 69978 Tel Aviv, Israel
 E-mail: aaro@tau.ac.il

Mahendra Nadkarni
 Department of Mathematics
 University of Mumbai
 Kalina, Mumbai 400098, India
 E-mail: mgnadkarni@gmail.com