

Proof of a conjecture of Hirschhorn and Sellers on overpartitions

by

WILLIAM Y. C. CHEN (Tianjin) and ERNEST X. W. XIA (Zhenjiang)

1. Introduction. The objective of this paper is to give a proof of a conjecture of Hirschhorn and Sellers on the number of overpartitions. We shall use the technique of dissections of theta functions.

Let us begin with some notation and terminology on q -series and partitions. We adopt the common notation

$$(1.1) \quad (a; q)_{\infty} = \prod_{n=0}^{\infty} (1 - aq^n),$$

where $|q| < 1$, and we write

$$(1.2) \quad (a_1, \dots, a_n; q)_{\infty} = (a_1; q)_{\infty} \cdots (a_n; q)_{\infty}.$$

Recall that the *Ramanujan theta function* $f(a, b)$ is defined by

$$(1.3) \quad f(a, b) = \sum_{n=-\infty}^{\infty} a^{n(n+1)/2} b^{n(n-1)/2},$$

where $|ab| < 1$. The Jacobi triple product identity can be restated as

$$(1.4) \quad f(a, b) = (-a, -b, ab; ab)_{\infty}.$$

In particular, let $f(-q) = f(-q, -q^2)$. Then we have

$$(1.5) \quad f(-q) = \sum_{n=-\infty}^{\infty} (-1)^n q^{n(3n-1)/2} = (q; q)_{\infty}.$$

For any positive integer n , we use f_n to denote $f(-q^n)$, that is,

$$(1.6) \quad f_n = (q^n; q^n)_{\infty} = \prod_{k=1}^{\infty} (1 - q^{nk}).$$

2010 *Mathematics Subject Classification*: Primary 11P83; Secondary 05A17.

Key words and phrases: overpartition, congruence, theta function, dissection formula, (p, k) -parametrization.

The function f_n is related to the generating function of overpartitions. A *partition* of a positive integer n is a nonincreasing sequence of positive integers whose sum is n . An *overpartition* of n is a partition in which the first occurrence of a number may be overlined (see Corteel and Lovejoy [CL]). For $n \geq 1$, we let $\bar{p}(n)$ denote the number of overpartitions of n , and set $\bar{p}(0) = 1$. Corteel and Lovejoy [CL] showed that the generating function for $\bar{p}(n)$ is given by

$$(1.7) \quad \sum_{n=0}^{\infty} \bar{p}(n)q^n = \frac{f_2}{f_1^2}.$$

Hirschhorn and Sellers [HS-1] and Fortin, Jacob and Mathieu [FJM] obtained the following Ramanujan-type generating function formulas for $\bar{p}(2n+1)$, $\bar{p}(4n+3)$ and $\bar{p}(8n+7)$:

$$(1.8) \quad \sum_{n=0}^{\infty} \bar{p}(2n+1)q^n = 2 \frac{f_2^2 f_8^2}{f_1^4 f_4},$$

$$(1.9) \quad \sum_{n=0}^{\infty} \bar{p}(4n+3)q^n = 8 \frac{f_2 f_4^6}{f_1^8},$$

$$(1.10) \quad \sum_{n=0}^{\infty} \bar{p}(8n+7)q^n = 64 \frac{f_2^{22}}{f_1^{23}}.$$

The above identities lead to congruences modulo 2, 8 and 64 for the overpartition function. Mahlborg [M] proved that $\bar{p}(n)$ is divisible by 64 for almost all n by using relations between $\bar{p}(n)$ and the number of representations of n as a sum of squares. Using the theory of modular forms, Treneer [T] showed that the coefficients of a wide class of weakly holomorphic modular forms have infinitely many congruence relations for powers of every prime p other than 2 and 3. In particular, he proved that $\bar{p}(5m^3n) \equiv 0 \pmod{5}$ for any n that is coprime to m , where m is a prime satisfying $m \equiv -1 \pmod{5}$.

The following conjecture was posed by Hirschhorn and Sellers [HS-1].

CONJECTURE 1.1. For $n \geq 0$,

$$(1.11) \quad \bar{p}(40n+35) \equiv 0 \pmod{40}.$$

To prove the conjecture, we derive a generating function for $\bar{p}(40n+35)$ modulo 5 by using 2-dissection formulas for quotients of theta functions given by Ramanujan [B], and Hirschhorn and Sellers [HS-2]. Then we use the (p, k) -parametrization of theta functions due to Alaca, Alaca and Williams [AAW, AW, W] to show that $\bar{p}(40n+35) \equiv 0 \pmod{5}$ for $n \geq 0$. Combining this congruence and $\bar{p}(4n+3) \equiv 0 \pmod{8}$ for $n \geq 0$, we confirm the conjecture.

2. The generating function. In this section, we derive a generating function of $\bar{p}(40n+35)$ modulo 5. We first recall several 2-dissection formulas for quotients of theta functions due to Ramanujan [B] and Hirschhorn and Sellers [HS-2].

The following relations are consequences of dissection formulas of Ramanujan collected in Entry 25 in Berndt's book [B, p. 40]. Recall that $f_n = (q^n; q^n)_\infty$, as given by (1.6).

LEMMA 2.1. *We have*

$$(2.1) \quad f_1^2 = \frac{f_2 f_8^5}{f_4^2 f_{16}^2} - 2q \frac{f_2 f_{16}^2}{f_8},$$

$$(2.2) \quad \frac{1}{f_1^2} = \frac{f_8^5}{f_2^5 f_{16}^2} + 2q \frac{f_4^2 f_{16}^2}{f_2^5 f_8},$$

$$(2.3) \quad f_1^4 = \frac{f_4^{10}}{f_2^2 f_8^4} - 4q \frac{f_2^2 f_8^4}{f_4^2},$$

$$(2.4) \quad \frac{1}{f_1^4} = \frac{f_4^{14}}{f_2^{14} f_8^4} + 4q \frac{f_4^2 f_8^4}{f_2^{10}}.$$

Hirschhorn and Sellers [HS-2] established the following 2-dissection formula.

LEMMA 2.2. *We have*

$$(2.5) \quad \frac{f_5}{f_1} = \frac{f_8 f_{20}^2}{f_2^2 f_{40}} + q \frac{f_4^3 f_{10} f_{40}}{f_2^3 f_8 f_{20}}.$$

By Lemmas 2.1 and 2.2, we are led to a generating function of $\bar{p}(40n+35)$ modulo 5.

THEOREM 2.3. *We have*

$$(2.6) \quad \sum_{n=0}^{\infty} \bar{p}(40n+35)q^n \\ \equiv 2 \frac{f_2^{122}}{f_1^{63} f_4^{40}} + 3 \frac{f_1 f_2^{26}}{f_8^8} + 4q \frac{f_2^{98}}{f_1^{55} f_4^{24}} + 3q f_1^9 f_2^2 f_4^8 + 4q^2 \frac{f_2^{74}}{f_1^{47} f_4^8} \\ + 4q^3 \frac{f_2^{50} f_4^8}{f_1^{39}} + 4q^4 \frac{f_2^{26} f_4^{24}}{f_1^{31}} + 2q^5 \frac{f_2^2 f_4^{40}}{f_1^{23}} \pmod{5}.$$

Proof. Recall that the theta functions $\varphi(q)$ and $\psi(q)$ are defined by

$$(2.7) \quad \varphi(q) = f(q, q) = \sum_{n=-\infty}^{\infty} q^{n^2},$$

$$(2.8) \quad \psi(q) = f(q, q^3) = \sum_{n=0}^{\infty} q^{n(n+1)/2}.$$

By the Jacobi triple product identity, we find

$$(2.9) \quad \varphi(q) = \frac{f_2^5}{f_1^2 f_4^2},$$

$$(2.10) \quad \psi(q) = \frac{f_2^2}{f_1}.$$

Replacing q by $-q$ in (2.9) and (2.10), and using the fact that

$$(2.11) \quad f(q) = (-q; -q)_{\infty} = \frac{f_2^3}{f_1 f_4},$$

we deduce that

$$(2.12) \quad \varphi(-q) = \frac{f_1^2}{f_2},$$

$$(2.13) \quad \psi(-q) = \frac{f_1 f_4}{f_2}.$$

In view of (1.3), (1.4) and (2.7), we see that

$$(2.14) \quad \begin{aligned} \varphi(q) &= \sum_{n=-\infty}^{\infty} q^{n^2} = \sum_{n=-\infty}^{\infty} q^{25n^2} + 2q \sum_{n=-\infty}^{\infty} q^{25n^2+10n} + 2q^4 \sum_{n=-\infty}^{\infty} q^{25n^2+20n} \\ &= \varphi(q^{25}) + 2qD(q^5) + 2q^4E(q^5), \end{aligned}$$

where

$$(2.15) \quad D(q) = \sum_{n=-\infty}^{\infty} q^{5n^2+2n} = (-q^3, -q^7, q^{10}; q^{10})_{\infty},$$

$$(2.16) \quad E(q) = \sum_{n=-\infty}^{\infty} q^{5n^2+4n} = (-q, -q^9, q^{10}; q^{10})_{\infty}.$$

It is easily checked that

$$(2.17) \quad D(q)E(q) = \frac{f_2^2 f_5 f_{20}}{f_1 f_4}.$$

By the binomial theorem, we see that for any positive integer k

$$(2.18) \quad (1 - q^k)^5 \equiv (1 - q^{5k}) \pmod{5},$$

which implies that

$$(2.19) \quad \varphi^5(q) \equiv \varphi(q^5) \pmod{5}.$$

Based on (1.7), (2.9) and (2.19), we have

$$(2.20) \quad \begin{aligned} & \sum_{n=0}^{\infty} \bar{p}(n)(-q)^n \\ &= \frac{1}{\varphi(q)} = \frac{\varphi^4(q)}{\varphi^5(q)} \\ &\equiv \frac{\varphi^4(q)}{\varphi(q^5)} \equiv \frac{(\varphi(q^{25}) + 2qD(q^5) + 2q^4E(q^5))^4}{\varphi(q^5)} \\ &\equiv \frac{1}{\varphi(q^5)} (\varphi^4(q^{25}) + 3q\varphi^3(q^{25})D(q^5) + 4q^2\varphi^2(q^{25})D^2(q^5) \\ &\quad + 2q^3\varphi(q^{25})D^3(q^5) + 3q^4\varphi^3(q^{25})E(q^5) + q^4D^4(q^5) \\ &\quad + 3q^5\varphi^2(q^{25})D(q^5)E(q^5) + q^6\varphi(q^{25})D^2(q^5)E(q^5) \\ &\quad + 4q^7D^3(q^5)E(q^5) + 4q^8\varphi^2(q^{25})E^2(q^5) + q^9\varphi(q^{25})D(q^5)E^2(q^5) \\ &\quad + q^{10}D^2(q^5)E^2(q^5) + 2q^{12}\varphi(q^{25})E^3(q^5) \\ &\quad + 4q^{13}D(q^5)E^3(q^5) + q^{16}E^4(q^5)^4) \pmod{5}, \end{aligned}$$

which yields

$$(2.21) \quad \begin{aligned} & \sum_{n=0}^{\infty} \bar{p}(5n)(-q)^n \\ &\equiv \frac{\varphi^4(q^5) + 3q\varphi^2(q^5)D(q)E(q) + q^2D^2(q)E^2(q)}{\varphi(q)} \pmod{5}. \end{aligned}$$

Plugging (2.9) and (2.17) into (2.21), we get

$$(2.22) \quad \sum_{n=0}^{\infty} \bar{p}(5n)(-q)^n \equiv \frac{f_1^2 f_4^2 f_{10}^{20}}{f_2^5 f_5^8 f_{20}^8} + 3q \frac{f_1 f_4 f_{10}^{10}}{f_2^3 f_5^3 f_{20}^3} + q^2 \frac{f_5^2 f_{20}^2}{f_2} \pmod{5}.$$

Replacing q by $-q$ in (2.22) and invoking (2.11), we arrive at

$$(2.23) \quad \sum_{n=0}^{\infty} \bar{p}(5n)q^n \equiv \frac{f_2 f_5^8}{f_1^2 f_{10}^4} - 3q \frac{f_5^3 f_{10}}{f_1} + q^2 \frac{f_{10}^6}{f_2 f_5^2} \pmod{5}.$$

According to the 2-dissection formulas (2.1), (2.2), (2.3), (2.5) and congru-

ence (2.23), we obtain

$$\begin{aligned}
 (2.24) \quad & \sum_{n=0}^{\infty} \bar{p}(5n)q^n \\
 & \equiv \frac{f_2}{f_{10}^4} \left(\frac{f_8^5}{f_2^5 f_{16}^2} + 2q \frac{f_4 f_{16}^2}{f_2^5 f_8} \right) \left(\frac{f_{20}^{10}}{f_{10}^2 f_{40}^4} - 4q^5 \frac{f_{10}^2 f_{40}^4}{f_{20}^2} \right)^2 \\
 & \quad - 3q f_{10} \left(\frac{f_{10} f_{40}^5}{f_{20}^2 f_{80}^2} - 2q^5 \frac{f_{10} f_{80}^2}{f_{40}} \right) \left(\frac{f_8 f_{20}^2}{f_2^2 f_{40}} + q \frac{f_4^3 f_{10} f_{40}}{f_2^3 f_8 f_{20}} \right) \\
 & \quad + q^2 \frac{f_{10}^6}{f_2} \left(\frac{f_{40}^5}{f_{10}^5 f_{80}^2} + 2q^5 \frac{f_{20}^2 f_{80}^2}{f_{10}^5 f_{40}} \right) \\
 & \equiv \frac{f_8^5 f_{20}^{20}}{f_2^4 f_{10}^8 f_{16}^2 f_{40}^8} + 2q \frac{f_4^2 f_{16}^2 f_{20}^{20}}{f_2^4 f_8 f_{10}^8 f_{40}^8} - 3q \frac{f_8 f_{10}^2 f_{40}^4}{f_2^2 f_{80}^2} - 3q^2 \frac{f_4^3 f_{10}^3 f_{40}^6}{f_2^3 f_8 f_{20}^3 f_{80}^2} \\
 & \quad + q^2 \frac{f_{10} f_{40}^5}{f_2 f_{80}^2} - 3q^5 \frac{f_8^5 f_{20}^8}{f_2^4 f_{10}^4 f_{16}^2} + q^6 \frac{f_8 f_{10}^2 f_{20}^2 f_{80}^2}{f_2^2 f_{40}^2} - q^6 \frac{f_4^2 f_{16}^2 f_{20}^8}{f_2^4 f_8 f_{40}^4} \\
 & \quad + q^7 \frac{f_4^3 f_{10}^3 f_{80}^2}{f_2^3 f_8 f_{20}} + 2q^7 \frac{f_{10} f_{20}^2 f_{80}^2}{f_2 f_{40}} + q^{10} \frac{f_8^5 f_{40}^8}{f_2^4 f_{16}^2 f_{20}^4} \\
 & \quad + 2q^{11} \frac{f_4^2 f_{16}^2 f_{40}^8}{f_2^4 f_8 f_{40}^4} \pmod{5}.
 \end{aligned}$$

Extracting the terms of odd powers of q on both sides of (2.24), we have

$$\begin{aligned}
 (2.25) \quad & \sum_{n=0}^{\infty} \bar{p}(10n+5)q^{2n+1} \\
 & \equiv 2q \frac{f_4^2 f_{16}^2 f_{20}^{20}}{f_2^4 f_8 f_{10}^8 f_{40}^8} - 3q \frac{f_8 f_{10}^2 f_{40}^4}{f_2^2 f_{80}^2} - 3q^5 \frac{f_8^5 f_{20}^8}{f_2^4 f_{10}^4 f_{16}^2} + q^7 \frac{f_4^3 f_{10}^3 f_{80}^2}{f_2^3 f_8 f_{20}} \\
 & \quad + 2q^7 \frac{f_{10} f_{20}^2 f_{80}^2}{f_2 f_{40}} + 2q^{11} \frac{f_4^2 f_{16}^2 f_{40}^8}{f_2^4 f_8 f_{40}^4} \pmod{5}.
 \end{aligned}$$

Dividing both sides of (2.25) by q and replacing q^2 by q , we get

$$\begin{aligned}
 (2.26) \quad & \sum_{n=0}^{\infty} \bar{p}(10n+5)q^n \\
 & \equiv 2 \frac{f_2^2 f_8^2 f_{10}^{20}}{f_1^4 f_4 f_5^8 f_{20}^8} - 3 \frac{f_4 f_5^2 f_{40}^4}{f_1^2 f_{40}^2} - 3q^2 \frac{f_4^5 f_{10}^8}{f_1^4 f_5^8 f_8^2} \\
 & \quad + q^3 \frac{f_2^3 f_5^3 f_{40}^2}{f_1^3 f_4 f_{10}} + 2q^3 \frac{f_5 f_{10}^2 f_{40}^2}{f_1 f_{20}} + 2q^5 \frac{f_2^2 f_8^2 f_{20}^8}{f_1^4 f_4 f_{10}^4} \pmod{5}.
 \end{aligned}$$

Employing the 2-dissection formulas (2.4), (2.5) and congruence (2.26), we

deduce that

$$\begin{aligned}
(2.27) \quad & \sum_{n=0}^{\infty} \bar{p}(10n+5)q^n \\
& \equiv 2 \frac{f_2^2 f_8^2 f_{10}^{20}}{f_4 f_8^8 f_{20}} \left(\frac{f_4^{14}}{f_2^{14} f_8^4} + 4q \frac{f_4^2 f_8^4}{f_2^{10}} \right) \left(\frac{f_{20}^{14}}{f_{10}^{14} f_4^4} + 4q^5 \frac{f_{20}^2 f_4^4}{f_{10}^{10}} \right)^2 \\
& \quad - 3 \frac{f_4 f_4^4 f_{20}}{f_{40}^2} \left(\frac{f_8 f_{20}^2}{f_2^2 f_{40}} + q \frac{f_4^3 f_{10} f_{40}}{f_2^3 f_8 f_{20}} \right)^2 \\
& \quad - 3q^2 \frac{f_4^5 f_{10}^8}{f_8^2} \left(\frac{f_4^{14}}{f_2^{14} f_8^4} + 4q \frac{f_4^2 f_8^4}{f_2^{10}} \right) \left(\frac{f_{20}^{14}}{f_{10}^{14} f_4^4} + 4q^5 \frac{f_{20}^2 f_4^4}{f_{10}^{10}} \right) \\
& \quad + q^3 \frac{f_2^3 f_4^2 f_{40}}{f_4 f_{10}} \left(\frac{f_8 f_{20}^2}{f_2^2 f_{40}} + q \frac{f_4^3 f_{10} f_{40}}{f_2^3 f_8 f_{20}} \right)^3 \\
& \quad + 2q^3 \frac{f_{10}^2 f_{40}^2}{f_{20}} \left(\frac{f_8 f_{20}^2}{f_2^2 f_{40}} + q \frac{f_4^3 f_{10} f_{40}}{f_2^3 f_8 f_{20}} \right) \\
& \quad + 2q^5 \frac{f_2^2 f_8^2 f_{20}^8}{f_4 f_{10}^4} \left(\frac{f_4^{14}}{f_2^{14} f_8^4} + 4q \frac{f_4^2 f_8^4}{f_2^{10}} \right) \\
& \equiv -3 \frac{f_4 f_8^2 f_{20}^8}{f_2^4 f_4^4} + 2 \frac{f_4^{13} f_{20}^{20}}{f_2^{12} f_8^2 f_{10}^8 f_{40}^8} + 3q \frac{f_4 f_8^6 f_{20}^{20}}{f_8^8 f_{10}^4 f_8^8} - q \frac{f_4^4 f_{10} f_{20}^5}{f_2^5 f_{40}^2} \\
& \quad - 3q^2 \frac{f_4^{19} f_{20}^{14}}{f_2^{14} f_8^6 f_{10}^6 f_{40}^4} - 3q^2 \frac{f_4^7 f_{10}^2 f_{20}^2}{f_2^6 f_8^2} - 2q^3 \frac{f_4^7 f_8^2 f_{20}^{14}}{f_2^{10} f_{10}^6 f_{40}^4} \\
& \quad + 2q^3 \frac{f_8 f_{10}^2 f_{20} f_{40}}{f_2^2} + q^3 \frac{f_8^3 f_{20}^6}{f_2^3 f_4 f_{10} f_{40}} + 2q^4 \frac{f_4^3 f_{10}^3 f_{40}^3}{f_2^3 f_8 f_{20}^2} \\
& \quad + 3q^4 \frac{f_4^2 f_8 f_{20}^3 f_{40}}{f_2^4} + 3q^5 \frac{f_4^5 f_{10} f_{40}^3}{f_2^5 f_8} + 3q^5 \frac{f_4^{13} f_{20}^8}{f_2^{12} f_8^2 f_{10}^4} \\
& \quad + 2q^6 \frac{f_4 f_8^6 f_{20}^8}{f_2^8 f_{10}^4} + q^6 \frac{f_4^8 f_{10}^2 f_{40}^5}{f_2^6 f_8^3 f_{20}^3} - 2q^7 \frac{f_4^{19} f_{20}^2 f_{40}^4}{f_2^{14} f_8^6 f_{10}^2} \\
& \quad - 3q^8 \frac{f_4^7 f_8^2 f_{20}^2 f_{40}^4}{f_2^{10} f_{10}^2} + 2q^{10} \frac{f_4^{13} f_{40}^8}{f_2^{12} f_8^2 f_{20}^4} + 3q^{11} \frac{f_4 f_8^6 f_{40}^8}{f_2^8 f_{40}^2} \pmod{5}.
\end{aligned}$$

Extracting the terms of odd powers of q on both sides of (2.27), then dividing by q and replacing q^2 by q , we find that

$$\begin{aligned}
(2.28) \quad & \sum_{n=0}^{\infty} \bar{p}(20n+15)q^n \\
& \equiv 3 \frac{f_2 f_4^6 f_{10}^{20}}{f_1^8 f_5^8 f_{20}^8} - \frac{f_2^4 f_5 f_{10}^5}{f_1^5 f_{20}^2} - 2q \frac{f_2^7 f_4^2 f_{10}^{14}}{f_1^{10} f_5^6 f_{20}^4} + 2q \frac{f_4 f_5^2 f_{10} f_{20}}{f_1^2}
\end{aligned}$$

$$\begin{aligned}
& + q \frac{f_4^3 f_{10}^6}{f_1^3 f_2 f_5 f_{20}} + 3q^2 \frac{f_2^5 f_5 f_{20}^3}{f_1^5 f_4} + 3q^2 \frac{f_2^{13} f_{10}^8}{f_1^{12} f_4^2 f_5^4} \\
& - 2q^3 \frac{f_2^{19} f_{10}^2 f_{20}^4}{f_1^{14} f_4^6 f_5^2} + 3q^5 \frac{f_2 f_4^6 f_{20}^8}{f_1^8 f_{10}^4} \pmod{5}.
\end{aligned}$$

By (2.18), we see that

$$(2.29) \quad f_5 \equiv f_1^5 \pmod{5}.$$

Substituting (2.29) into (2.28) gives

$$\begin{aligned}
(2.30) \quad & \sum_{n=0}^{\infty} \bar{p}(20n+15)q^n \\
& \equiv 3 \frac{f_2^{101}}{f_1^{48} f_4^{34}} - \frac{f_2^{29}}{f_4^{10}} - 2q \frac{f_2^{77}}{f_1^{40} f_4^{18}} + 2q f_1^8 f_2^5 f_4^6 + q \frac{f_2^{29}}{f_1^8 f_4^2} \\
& + 3q^2 f_2^5 f_4^{14} + 3q^2 \frac{f_2^{53}}{f_1^{32} f_4^2} - 2q^3 \frac{f_2^{29} f_4^{14}}{f_1^2} \\
& + 3q^5 \frac{f_4^{46}}{f_1^8 f_2^{19}} \pmod{5}.
\end{aligned}$$

Combining the 2-dissection formulas (2.3), (2.4) and congruence (2.30), we see that

$$\begin{aligned}
(2.31) \quad & \sum_{n=0}^{\infty} \bar{p}(20n+15)q^n \\
& \equiv 3 \frac{f_2^{101}}{f_4^{34}} \left(\frac{f_4^{14}}{f_2^{14} f_8^4} + 4q \frac{f_4^2 f_8^4}{f_2^{10}} \right)^{12} + 2q f_2^5 f_4^6 \left(\frac{f_4^{10}}{f_2^2 f_8^4} - 4q \frac{f_2^2 f_8^4}{f_4^2} \right)^2 \\
& - \frac{f_2^{29}}{f_4^{10}} - 2q \frac{f_2^{77}}{f_4^{18}} \left(\frac{f_4^{14}}{f_2^{14} f_8^4} + 4q \frac{f_4^2 f_8^4}{f_2^{10}} \right)^{10} \\
& + q \frac{f_2^{29}}{f_4^2} \left(\frac{f_4^{14}}{f_2^{14} f_8^4} + 4q \frac{f_4^2 f_8^4}{f_2^{10}} \right)^2 + 3q^2 f_2^5 f_4^{14} \\
& + 3q^2 \frac{f_2^{53}}{f_4^2} \left(\frac{f_4^{14}}{f_2^{14} f_8^4} + 4q \frac{f_4^2 f_8^4}{f_2^{10}} \right)^8 \\
& - 2q^3 f_2^{29} f_4^{14} \left(\frac{f_4^{14}}{f_2^{14} f_8^4} + 4q \frac{f_4^2 f_8^4}{f_2^{10}} \right)^6 \\
& + 3q^5 \frac{f_4^{46}}{f_2^{19}} \left(\frac{f_4^{14}}{f_2^{14} f_8^4} + 4q \frac{f_4^2 f_8^4}{f_2^{10}} \right)^2
\end{aligned}$$

$$\begin{aligned}
 &\equiv 3 \frac{f_4^{134}}{f_2^{67} f_8^{48}} - \frac{f_2^{29}}{f_4^{10}} + 2q \frac{f_4^{122}}{f_2^{63} f_8^{40}} + 3q \frac{f_2 f_4^{26}}{f_8^8} + q^2 \frac{f_4^{110}}{f_2^{59} f_8^{32}} \\
 &\quad + 4q^3 \frac{f_4^{98}}{f_2^{55} f_8^{24}} + 3q^3 f_2^9 f_4^2 f_8^8 + q^4 \frac{f_4^{86}}{f_2^{51} f_8^{16}} + 4q^5 \frac{f_4^{74}}{f_2^{47} f_8^8} \\
 &\quad + 4q^7 \frac{f_4^{50} f_8^8}{f_2^{39}} + q^8 \frac{f_4^{38} f_8^{16}}{f_2^{35}} + 4q^9 \frac{f_4^{26} f_8^{24}}{f_2^{31}} + q^{10} \frac{f_4^{14} f_8^{32}}{f_2^{27}} \\
 &\quad + 2q^{11} \frac{f_4^2 f_8^{40}}{f_2^{23}} + 3q^{12} \frac{f_8^{48}}{f_2^{19} f_4^{10}} \pmod{5}.
 \end{aligned}$$

Extracting the terms of odd powers of q on both sides of (2.31), then dividing by q and replacing q^2 by q , we obtain (2.6). ■

3. Proof of Conjecture 1.1. In this section, we use the (p, k) -parametrization of theta functions given by Alaca, Alaca and Williams [AAW, AW, W] to represent the generating function of $\bar{p}(40n + 35)$ modulo 5 as a linear combination of functions in p and k , where p and k are defined in terms of the theta function $\varphi(q)$ as follows:

$$(3.1) \quad p := p(q) = \frac{\varphi^2(q) - \varphi^2(q^3)}{2\varphi^2(q^3)}, \quad k := k(q) = \frac{\varphi^3(q^3)}{\varphi(q)}$$

(see Alaca, Alaca and Williams [AAW]). Williams [W] proved that

$$(3.2) \quad p = 2 \frac{f_2^3 f_3^3 f_{12}^6}{f_1 f_4^2 f_9^6}.$$

We have the following congruence.

THEOREM 3.1. For $n \geq 0$,

$$(3.3) \quad \bar{p}(40n + 35) \equiv 0 \pmod{5}.$$

Proof. The following representations of $q^{1/24} f_1$, $q^{1/12} f_2$ and $q^{1/6} f_4$ in terms of p and k are due to Alaca and Williams [AW]:

$$(3.4) \quad q^{1/24} f_1 = 2^{-1/6} p^{1/24} (1 - p)^{1/2} (1 + p)^{1/6} (1 + 2p)^{1/8} (2 + p)^{1/8} k^{1/2},$$

$$(3.5) \quad q^{1/12} f_2 = 2^{-1/3} p^{1/12} (1 - p)^{1/4} (1 + p)^{1/12} (1 + 2p)^{1/4} (2 + p)^{1/4} k^{1/2},$$

$$(3.6) \quad q^{1/6} f_4 = 2^{-2/3} p^{1/6} (1 - p)^{1/8} (1 + p)^{1/24} (1 + 2p)^{1/8} (2 + p)^{1/2} k^{1/2}.$$

Substituting (3.4), (3.5) and (3.6) into (2.6), we find that

$$\begin{aligned}
 (3.7) \quad 2^{19} \sum_{n=0}^{\infty} \bar{p}(40n + 35) q^n \\
 \equiv \frac{\sqrt{2} p^{7/8} (1 + 2p)^{21/8} (2 + p)^{21/8}}{16q^{7/8} (1 - p)^6 (1 + p)^2 \sqrt{k}} F(p, k) \pmod{5},
 \end{aligned}$$

where $F(p, k)$ is given by

$$(3.8) \quad F(p, k) = 5k^{10}(524288 + 6029312p + 88735744p^2 + 840761344p^3 + 5072977920p^4 + 22470361088p^5 + 75791417344p^6 + 196034666496p^7 + 392385622016p^8 + 610286094336p^9 + 731633712128p^{10} + 663209854464p^{11} + 441020946176p^{12} + 204189055872p^{13} + 59086163776p^{14} + 8129694944p^{15} + 138932400p^{16} + 2477318p^{19} - 16585772p^{18} + 33708184p^{17} + 19661p^{20}).$$

By (3.4) and (3.5), we have

$$(3.9) \quad \frac{f_2^{22}}{f_1^{23}} = \frac{\sqrt{2} p^{7/8} (1+2p)^{21/8} (2+p)^{21/8}}{16q^{7/8} (1-p)^6 (1+p)^2 \sqrt{k}}.$$

Hence (3.7) can be rewritten as

$$(3.10) \quad 2^{19} \sum_{n=0}^{\infty} \bar{p}(40n+35)q^n \equiv \frac{f_2^{22}}{f_1^{23}} F(p, k) \pmod{5},$$

where $F(p, k)$ is defined by (3.8). Clearly, f_2^{22}/f_1^{23} is a formal power series in q with integer coefficients. By (3.1) and (3.2), we see that p and k are also formal power series in q with integer coefficients. It can be seen that the coefficients of $F(p, k)$ are divisible by 5. So we reach the assertion. ■

To complete the proof of Conjecture 1.1, we recall that Hirschhorn and Sellers [HS-1], and Fortin, Jacob and Mathieu [FJM] independently, derived the congruence

$$(3.11) \quad \bar{p}(4n+3) \equiv 0 \pmod{8}$$

for $n \geq 0$. This yields

$$(3.12) \quad \bar{p}(40n+35) \equiv 0 \pmod{8}$$

for $n \geq 0$. Combining (3.12) and $\bar{p}(40n+35) \equiv 0 \pmod{5}$ for $n \geq 0$, we come to the conclusion that $\bar{p}(40n+35) \equiv 0 \pmod{40}$ for $n \geq 0$. ■

Acknowledgments. This work was supported by the 973 Project and the National Science Foundation of China.

References

- [AAW] A. Alaca, Ş. Alaca and K. S. Williams, *On the two-dimensional theta functions of the Borweins*, Acta Arith. 124 (2006), 177–195.
- [AW] Ş. Alaca and K. S. Williams, *The number of representations of a positive integer by certain octonary quadratic forms*, Funct. Approx. Comment. Math. 43 (2010), 45–54.

- [B] B. C. Berndt, *Ramanujan's Notebooks. Part III*, Springer, New York, 1991.
- [CL] S. Corteel and J. Lovejoy, *Overpartitions*, Trans. Amer. Math. Soc. 356 (2004), 1623–1635.
- [FJM] J.-F. Fortin, P. Jacob and P. Mathieu, *Jagged partitions*, Ramanujan J. 10 (2005), 215–235.
- [HS-1] M. D. Hirschhorn and J. A. Sellers, *Arithmetic relations for overpartitions*, J. Combin. Math. Combin. Comput. 53 (2005), 65–73.
- [HS-2] M. D. Hirschhorn and J. A. Sellers, *Elementary proofs of parity results for 5-regular partitions*, Bull. Austral. Math. Soc. 81 (2010), 58–63.
- [M] K. Mahlburg, *The overpartition function modulo small powers of 2*, Discrete Math. 286 (2004), 263–267.
- [T] S. Treneer, *Congruences for the coefficients of weakly holomorphic modular forms*, Proc. London Math. Soc. (3) 93 (2006), 304–324.
- [W] K. S. Williams, *Fourier series of a class of eta quotients*, Int. J. Number Theory 8 (2012), 993–1004.

William Y. C. Chen
Center for Applied Mathematics
Tianjin University
Tianjin 300072, P.R. China
E-mail: chenyc@tju.edu.cn

Ernest X. W. Xia
Department of Mathematics
Jiangsu University
Zhenjiang, Jiangsu 212013, P.R. China
E-mail: ernestxwxa@163.com

*Received on 16.7.2013
and in revised form on 1.11.2013*

(7524)

