

## Inverse zero-sum problems II

by

WOLFGANG A. SCHMID (Graz)

**1. Introduction.** In the 1960s H. Davenport popularized the following problem, motivated by an application in algebraic number theory. Let  $G$  be an additive finite abelian group. Determine the smallest integer  $\ell$  such that each sequence over  $G$  of length at least  $\ell$  has a non-empty subsequence the sum of whose terms equals  $0 \in G$ . This integer is now called Davenport's constant of  $G$ , denoted  $D(G)$ . We refer to the recent survey article [10], the lecture notes [14], the monographs [13], in particular Chapters 5 to 7, and [20], in particular Chapter 9, for detailed information on and applications of Davenport's constant, e.g., in investigations of the arithmetic of maximal orders of algebraic number fields.

Parallel to the problem of determining Davenport's constant, a direct problem, the associated inverse problem, i.e., the problem of determining the structure of the longest sequences that do not have a subsequence with sum zero, was intensively investigated as well. On the one hand, solutions to the inverse problem are relevant in the above mentioned applications as well, and on the other hand, inverse results for one type of group can be applied in investigations of the direct problem for other, more complicated, types of groups (see, e.g., [1]).

In this paper, we investigate the inverse problem associated to Davenport's constant for general finite abelian groups of rank two, complementing the investigations of the first paper in this series [12] that focused on groups of the form  $C_m^2$ , i.e., the direct sum of two cyclic groups of order  $m$ . To put this in context, we recall that the value of Davenport's constant for groups of rank two is well-known (cf. Theorem 4.1 and the references there); moreover, for cyclic groups, answers to both the direct and the inverse problem are well-known (cf. Theorems 4.1 and 4.2 and, e.g., [18, 22] for refinements), whereas for groups of rank at least three, both the direct and the inverse problem is in general wide open (see, e.g., [3, 19] for results in special cases).

---

2010 *Mathematics Subject Classification*: Primary 11P70; Secondary 11B30.

*Key words and phrases*: Davenport's constant, zero-sum sequence, inverse problem.

For groups of the form  $C_m^2$  there is a well-known and well-supported conjecture regarding the answer to the inverse problem (see Definition 3.1 for details). For groups of the form  $C_2 \oplus C_{2n}$  and  $C_3 \oplus C_{3n}$  the inverse problem was solved in [8, Section 3] and [5], respectively, and in [7, Section 8] and [15] partial results in the general case were obtained. Here we solve, *assuming* the above mentioned conjecture for groups of the form  $C_m^2$  is true, the inverse problem for general groups of rank two (see Theorem 3.2).

In our proof, we use direct and inverse results for cyclic groups and groups of the form  $C_m^2$ , which we recall in Subsection 4.1, that we combine by using the Inductive Method (cf. [13, Section 5.7]).

**2. Notation and terminology.** We recall some standard notation and terminology (we follow [10] and [13]).

We denote by  $\mathbb{Z}$  the set of integers, and by  $\mathbb{N}$  and  $\mathbb{N}_0$  the positive and non-negative integers, respectively. For  $a, b \in \mathbb{Z}$ , we denote by  $[a, b] = \{z \in \mathbb{Z} : a \leq z \leq b\}$  the interval of integers. For  $k \in \mathbb{Z}$  and  $m \in \mathbb{N}$ , we let  $[k]_m$  be the integer in  $[0, m - 1]$  that is congruent to  $k$  modulo  $m$ .

Let  $G$  denote an additively written finite abelian group. (Throughout, we use additive notation for abelian groups.) For a subset  $G_0 \subset G$ , we denote by  $\langle G_0 \rangle$  the subgroup generated by  $G_0$ . We call elements  $e_1, \dots, e_r \in G \setminus \{0\}$  *independent* if  $\sum_{i=1}^r m_i e_i = 0$  with  $m_i \in \mathbb{Z}$  implies that  $m_i e_i = 0$  for each  $i \in [1, r]$ . We call a subset of  $G$  a *basis* if it generates  $G$  and its elements are independent. For  $n \in \mathbb{N}$ , we denote by  $C_n$  a cyclic group of order  $n$ . For each finite abelian group  $G$ , there exist uniquely determined  $1 < n_1 \mid \dots \mid n_r$  such that  $G \cong C_{n_1} \oplus \dots \oplus C_{n_r}$ ; we refer to  $r$  as the *rank* of  $G$  and to  $\exp(G) = n_r$  as the *exponent* of  $G$  (except for  $|G| = 1$ , where the exponent is 1).

We denote by  $\mathcal{F}(G)$  the (multiplicatively written) free abelian monoid over  $G$ , that is, the monoid of all formal commutative products

$$S = \prod_{g \in G} g^{v_g(S)}$$

with  $v_g(S) \in \mathbb{N}_0$ . We call such an element  $S$  a *sequence over  $G$* . We refer to  $v_g(S)$  as the *multiplicity* of  $g$  in  $S$ . Moreover,  $\sigma(S) = \sum_{g \in G} v_g(S)g \in G$  is called the *sum* of  $S$ ,  $|S| = \sum_{g \in G} v_g(S) \in \mathbb{N}_0$  the *length* of  $S$ , and  $\text{supp}(S) = \{g \in G : v_g(S) > 0\} \subset G$  the *support* of  $S$ .

We denote the unit element of  $\mathcal{F}(G)$  by 1 and call it the *empty sequence*. If  $T \in \mathcal{F}(G)$  and  $T \mid S$  (in  $\mathcal{F}(G)$ ), then we call  $T$  a *subsequence* of  $S$ ; we say that it is a *proper subsequence* if  $1 \neq T \neq S$ . Moreover, we denote by  $T^{-1}S$  its *co-divisor*, i.e., the unique sequence  $R$  with  $RT = S$ .

If  $\sigma(S) = 0$ , then we call  $S$  a *zero-sum sequence* (zss, for short), and if  $\sigma(T) \neq 0$  for each  $1 \neq T \mid S$ , then we say that  $S$  is *zero-sum free*. We call a zss a *minimal zss* (mzss, for short) if it is non-empty and has no proper subsequence with sum zero.

Using the notation recalled above, the definition of Davenport’s constant can be given as follows. For a finite abelian group  $G$ , let  $\ell \in \mathbb{N}$  be minimal with the property that each  $S \in \mathcal{F}(G)$  with  $|S| \geq \ell$  has a subsequence  $1 \neq T \mid S$  such that  $\sigma(T) = 0$ .

It is a simple and well-known fact that  $D(G)$  is the maximal length of a mzss over  $G$  and that each zero-sum free sequence of length  $D(G) - 1$  over  $G$ , i.e., a sequence appearing in the inverse problem associated to  $D(G)$ , is a subsequence of a mzss of length  $D(G)$ . Since it has technical advantages, we thus in fact investigate the structure of mzss of maximal length (ml-mzss, for short) instead of zero-sum free sequences of length  $D(G) - 1$ .

Each map  $f : G \rightarrow G'$  between finite abelian groups extends uniquely to a monoid homomorphism  $\mathcal{F}(G) \rightarrow \mathcal{F}(G')$ , which we denote by  $f$  as well. If  $f$  is a group homomorphism, then  $\sigma(f(S)) = f(\sigma(S))$  for each  $S \in \mathcal{F}(G)$ .

**3. Formulation of result.** In this section we recall the conjecture mentioned in the introduction and formulate our result.

**DEFINITION 3.1.** Let  $m \in \mathbb{N}$ . The group  $C_m^2$  is said to have *Property B* if each ml-mzss equals  $g^{\exp(G)-1}T$  for some  $g \in C_m^2$  and  $T \in \mathcal{F}(C_m^2)$ .

Property **B** was introduced by W. Gao and A. Geroldinger [7, 9]. It is conjectured that for each  $m \in \mathbb{N}$  the group  $C_m^2$  has Property **B** (see the just mentioned papers and, e.g., [10, Conjecture 4.5]). We recall some result on this conjecture.

By a very recent result (see [11], and [9] for an earlier partial result) it is known that to establish Property **B** for  $C_m^2$  for each  $m \in \mathbb{N}$ , it suffices to establish it for  $C_p^2$  for each prime  $p$ . Moreover, Property **B** is known to hold for  $C_m^2$  for  $m \leq 28$  (see [2] and [9] for  $m \leq 7$ ). For further recent results towards establishing Property **B** see [16, 12, 2].

As indicated in the introduction, we characterize ml-mzss for finite abelian groups of rank two, under the assumption that a certain subgroup of the group has Property **B**.

**THEOREM 3.2.** *Let  $G$  be a finite abelian group of rank two, say,  $G \cong C_m \oplus C_{mn}$  with  $m, n \in \mathbb{N}$  and  $m \geq 2$ . The following sequences are minimal zero-sum sequences of maximal length.*

- (i)  $S = e_j^{\text{ord } e_j - 1} \prod_{i=1}^{\text{ord } e_k} (-x_i e_j + e_k)$  where  $\{e_1, e_2\}$  is a basis of  $G$  with  $\text{ord } e_2 = mn$ ,  $\{j, k\} = \{1, 2\}$ , and  $x_i \in \mathbb{N}_0$  with  $\sum_{i=1}^{\text{ord } e_k} x_i \equiv -1 \pmod{\text{ord } e_j}$ .
- (ii)  $S = g_1^{sm-1} \prod_{i=1}^{(n+1-s)m} (-x_i g_1 + g_2)$  where  $s \in [1, n]$ ,  $\{g_1, g_2\}$  is a generating set of  $G$  with  $\text{ord } g_2 = mn$  and, in case  $s \neq 1$ ,  $mg_1 = mg_2$ , and  $x_i \in \mathbb{N}_0$  with  $\sum_{i=1}^{(n+1-s)m} x_i = m - 1$ .

If  $C_m^2$  has Property **B**, then all minimal zero-sum sequences of maximal length over  $G$  are of this form.

The case  $G \cong C_m^2$ , i.e.  $n = 1$ , of this result is well-known and included for completeness only (see, e.g., [13, Theorem 5.8.7]); in particular, note that (ii) is redundant for  $n = 1$ .

This result can be combined with the above mentioned results on Property **B** to yield unconditional results for special types of groups. We do not formulate these explicitly and only point out that, since  $C_2^2$  and  $C_3^2$  have Property **B** (cf. above), the results on  $C_2 \oplus C_{2n}$  and  $C_3 \oplus C_{3n}$  mentioned in the introduction can be obtained in this way.

**4. Proof of the result.** In this section we give the proof of Theorem 3.2. First, we recall some results that we use in the proof.

**4.1. Known results.** The value of  $D(G)$  for  $G$  of rank two, i.e., the answer to the direct problem, is well-known (see [17, 21]).

**THEOREM 4.1.** *Let  $m, n \in \mathbb{N}$ . Then  $D(C_m \oplus C_{mn}) = m + mn - 1$ .*

Next, we recall some results on sequences over cyclic groups: the solution to the inverse problem associated to Davenport’s constant for cyclic groups, a simple special case of [4], and the theorem of Erdős–Ginzburg–Ziv [6].

**THEOREM 4.2.** *Let  $n \in \mathbb{N}$  and  $S \in \mathcal{F}(C_n)$ .*

- (i)  *$S$  is a  $ml$ - $mzss$  if and only if  $S = e^n$  for some  $e \in C_n$  with  $\langle e \rangle = C_n$ .*
- (ii) *If  $|S| \geq 2n - 1$ , then there exists some  $T \mid S$  with  $|T| = n$  and  $\sigma(T) = 0$ .*

The following result is a main tool in the proof of Theorem 3.2. It was obtained in [9, Proposition 4.1, Theorem 7.1]; note that now the additional assumption in the original version (regarding the existence of zss of length  $m$  and  $2m$ ) can be dropped, since by [10, Theorem 6.5] it is known to be fulfilled for each  $m \in \mathbb{N}$  (also note that the second type of sequence requires  $t \geq 3$ ).

**THEOREM 4.3.** *Let  $m, t \in \mathbb{N}$  with  $m \geq 2$  and  $t \geq 2$ . Suppose that  $C_m^2$  has Property **B**. Let  $S \in \mathcal{F}(C_m^2)$  be a zss of length  $tm - 1$  that cannot be written as the product of  $t$  non-empty zss. Then for some basis  $\{f_1, f_2\}$  of  $C_m^2$ ,*

$$S = f_1^{sm-1} \prod_{i=1}^{(t-s)m} (a_i f_1 + f_2)$$

*with  $s \in [1, t - 1]$  and  $a_i \in [0, m - 1]$  where  $\sum_{i=1}^{(t-s)m} a_i \equiv 1 \pmod{m}$ , or*

$$S = f_1^{s_1 m} f_2^{s_2 m-1} (b f_1 + f_2)^{s_3 m-1} (b f_1 + 2 f_2)$$

with  $s_i \in \mathbb{N}$  such that  $s_1 + s_2 + s_3 = t$  and  $b \in [1, m - 1]$  such that  $\gcd\{b, m\} = 1$ .

**4.2. Proof of Theorem 3.2.** We start by establishing that all the sequences are indeed ml-mzss.

Since the length of each sequence is  $mn + n - 1$ , and by Theorem 4.1, it suffices to show that they are mzss. It is readily seen that  $\sigma(S) = 0$ , thus it remains to show minimality. Let  $1 \neq T | S$  be a zss. We assert that  $T = S$ . If  $S$  is as given in (i), then it suffices to note that  $e_j^{\text{ord } e_j - 1}$  is zero-sum free, thus  $(-x_i e_j + e_k) | T$  for some  $i \in [1, \text{ord } e_k]$ , and this implies  $\prod_{i=1}^{\text{ord } e_k} (-x_i e_j + e_k) | T$ , which implies  $S = T$ . Suppose  $S$  is as given in (ii). We first note that  $ag_1 \in \langle g_2 \rangle$  if and only if  $m | a$ . Let  $v \in \mathbb{N}_0$  and  $I \subset [1, (n+1-s)m]$  be such that  $T = g_1^v \prod_{i \in I} (-x_i g_1 + g_2)$ . Since  $\sigma(T) = 0$  and by the above observation, it follows that  $m | (v - \sum_{i \in I} x_i)$ , say  $mb = v - \sum_{i \in I} x_i$ , where  $b \in [0, s - 1]$ . Furthermore, we get  $mbg_1 + |I|g_2 = 0$ . If  $s = 1$ , then  $b = 0$ , implying that  $|I| = mn$  and  $v = m - 1$ , that is,  $S = T$ . If  $s > 1$ , we have  $mg_1 = mg_2$ , thus  $mn ||I| + mb$  and indeed  $mn = |I| + mb$ . Yet,  $mn = |I| + mb$  implies  $|I| = [1, (n + 1 - s)m]$  and  $b = s - 1$ , that is,  $S = T$ . Thus, the sequences are mzss.

Now, we show that if  $C_m^2$  has Property **B**, then each ml-mzss is of this form.

As already mentioned, the case  $n = 1$  is well-known (cf. Theorem 4.3). We thus assume  $n \geq 2$ , that is,  $G \cong C_m \oplus C_{mn}$  with  $m \geq 2$  and  $n \geq 2$ . Furthermore, let  $H = \{mg : g \in G\} \cong C_n$  and let  $\varphi : G \rightarrow G/H$  be the canonical map; we have  $G/H \cong C_m^2$ . We apply the Inductive Method, as in [7, Section 8], with the exact sequence

$$0 \rightarrow H \hookrightarrow G \xrightarrow{\varphi} G/H \rightarrow 0.$$

Let  $S \in \mathcal{F}(G)$  be a ml-mzss. First, we assert that  $\varphi(S)$  cannot be written as the product of  $n + 1$  non-empty zss, in order to apply Theorem 4.3. Suppose this is possible, say  $\varphi(S) = \prod_{i=1}^{n+1} \varphi(S_i)$  with non-empty zss  $\varphi(S_i)$ . Then  $\prod_{i=1}^{n+1} \sigma(S_i) \in \mathcal{F}(H)$  has a proper subsequence that is a zss, yielding a proper subsequence of  $S$  that is a zss.

Thus, by Theorem 4.3 there exists a basis  $\{f_1, f_2\}$  of  $C_m^2$  such that

$$(4.1) \quad \varphi(S) = f_1^{sm-1} \prod_{i=1}^{(n+1-s)m} (a_i f_1 + f_2)$$

with  $s \in [1, n]$ ,  $a_i \in [0, m - 1]$ , and  $\sum_{i=1}^{(n+1-s)m} a_i \equiv 1 \pmod{m}$ , or

$$(4.2) \quad \varphi(S) = f_1^{s_1 m} f_2^{s_2 m - 1} (b f_1 + f_2)^{s_3 m - 1} (b f_1 + 2 f_2)$$

with  $s_i \in \mathbb{N}$  such that  $s_1 + s_2 + s_3 = n + 1$  and  $b \in [1, m - 1]$  such that

$\gcd(b, m) = 1$ . We distinguish two cases, depending on which of the two structures  $\varphi(S)$  has.

CASE 1:  $\varphi(S)$  is of the form given in (4.1). Moreover, we assume the basis  $\{f_1, f_2\}$  is chosen in such a way that  $s$  is maximal. Furthermore, let  $\psi : G/H \rightarrow \langle f_1 \rangle$  denote the projection with respect to  $G/H = \langle f_1 \rangle \oplus \langle f_2 \rangle$ . Let  $S = FT$  be such that  $\varphi(F) = f_1^{sm-1}$  and  $T = \prod_{i=1}^{(n+1-s)m} h_i$  be such that  $\varphi(h_i) = a_i f_1 + f_2$ .

We call a factorization  $T = S_0 S_1 \dots S_{n-s}$  *admissible* if  $\sigma(\varphi(S_i)) = 0$  and  $|S_i| = m$  for  $i \in [1, n - s]$  (then  $\sigma(\varphi(S_0)) = f_1$  and  $|S_0| = m$ ). Since for a sequence  $T' | T$  of length  $m$  the conditions  $\sigma(\varphi(T')) = 0$  and  $\sigma(\psi(\varphi(T'))) = 0$  are equivalent, the existence of admissible factorizations follows using Theorem 4.2.

Let  $T = S_0 S_1 \dots S_{n-s}$  be an admissible factorization such that  $|\text{supp}(S_0)|$  is maximal (among all admissible factorizations of  $T$ ). Moreover, let  $F = F_0 F_1 \dots F_{s-1}$  with  $|F_0| = m - 1$  and  $|F_i| = m$  for  $i \in [1, s - 1]$ . Then  $\sigma(\varphi(F_i)) = 0$  for  $i \in [1, s - 1]$ ,  $\sigma(\varphi(S_i)) = 0$  for  $i \in [1, n - s]$ , and  $\sigma(\varphi(S_0 F_0)) = 0$ . Thus,  $\sigma(S_0 F_0) \prod_{i=1}^{s-1} \sigma(F_i) \prod_{i=1}^{n-s} \sigma(S_i)$  is a sequence over  $H$ , and it is a mzss. Since its length is  $n$ , it follows by Theorem 4.2 that there exists some generating element  $e \in H$  such that this sequence is equal to  $e^n$ .

We show that  $|\text{supp}(F)| = 1$ . We assume to the contrary that there exist distinct  $g, g' \in \text{supp}(F)$ .

First, suppose  $s \geq 2$ . We may assume  $g | F_i$  and  $g' | F_j$  for distinct  $i, j \in [0, s - 1]$ . Now we consider  $F'_i = g^{-1} g' F_i$  and  $F'_j = g'^{-1} g F_j$  and  $F'_k = F_k$  for  $k \notin \{i, j\}$ . As above, we find that  $\sigma(S_0 F'_0) \prod_{i=1}^{s-1} \sigma(F'_i) \prod_{i=1}^{n-s} \sigma(S_i)$  is a ml-mzss over  $H$  and thus equal to  $\bar{e}^n$  for some generating element  $\bar{e} \in H$  and indeed, since at most two elements in the sequence are changed and for  $n = 2$  there is only one generating element of  $H$ , we have  $e = \bar{e}$ . Thus,  $\sigma(F_i) = \sigma(F'_i) = \sigma(F_i) + g' - g$ , a contradiction.

Second, suppose  $s = 1$ . It follows that  $m \geq 3$ , since for  $m = 2$  we have  $|F| = 1$ . We consider  $S_0 S_j$  for some  $j \in [1, n - 1]$ . Let  $S_0 S_j = T' T''$  with  $|T'| = |T''| = m$ . Since  $\sigma(\varphi(T')) + \sigma(\varphi(T'')) = f_1$  and  $\sigma(\varphi(T')), \sigma(\varphi(T'')) \in \langle f_1 \rangle$  it follows that there exists some  $a \in [0, m - 1]$  such that  $\sigma(\varphi(T')) = (a + 1)f_1$  and  $\sigma(\varphi(T'')) = -af_1$ . Let  $F_0 = F' F''$  with  $|F'| = m - (a + 1)$  and  $|F''| = a$ . We note that  $\sigma(T' F') \sigma(T'' F'') \prod_{i=1, i \neq j}^{n-1} \sigma(S_i)$ , is a ml-mzss over  $H$  and again it follows that it is equal to  $e^n$  (with the same element  $e$  as above). If both  $F'$  and  $F''$  are non-empty, we may assume  $g | F'$  and  $g' | F''$ , to obtain a contradiction as above. Thus, it remains to investigate whether there exists a factorization  $S_0 S_j = T' T''$  with  $|T'| = |T''| = m$  such that  $\{\sigma(\varphi(T')), \sigma(\varphi(T''))\} \neq \{0, f_1\}$ . We observe that such a factorization exists except if  $\varphi(S_0 S_j) = (bf_1 + f_2)^{2m-1} (cf_1 + f_2)$  (note that  $\varphi(S_0 S_j) = (bf_1 + f_2)^{2m}$  is impossible, since  $\sigma(\varphi(S_0 S_j)) \neq 0$ ).

Thus, if such a factorization does not exist, for each  $j \in [1, n - 1]$ , then  $\varphi(T) = (bf_1 + f_2)^{mn-1}(cf_1 + f_2)$ . Since  $\sigma(\varphi(T)) = f_1$ , we get  $cf_1 = (b + 1)f_1$ . Thus, with respect to the basis consisting of  $\bar{f}_1 = bf_1 + f_2$  and  $\bar{f}_2 = f_1$ , we have  $\varphi(S) = \bar{f}_1^{mn-1}\bar{f}_2^{m-1}(\bar{f}_1 + \bar{f}_2)$ , contradicting the assumption that the basis  $\{f_1, f_2\}$  maximizes  $s$ .

Therefore, we have  $|\text{supp}(F)| = 1$  and thus

$$S = g_1^{sm-1}T$$

for some  $g_1 \in G$ .

First, we consider the case  $s = n$ . We have  $\text{ord } g_1 = mn$  and thus  $G = \langle g_1 \rangle \oplus H_2$  where  $H_2 \subset G$  is a cyclic group of order  $m$ . Let  $\pi : G \rightarrow H_2$  denote the projection with respect to  $G = \langle g_1 \rangle \oplus H_2$ . We observe that  $\pi(\prod_{i=1}^m h_i) \in \mathcal{F}(H_2)$  is a mzss and consequently it is equal to  $g_2^m$  for some generating element  $g_2$  of  $H_2$ . We note that  $\{g_1, g_2\}$  is a basis of  $G$ . Thus,  $S$  is of the form given in (i).

Thus, we may assume  $s < n$ . Next, we show that if  $\varphi(h_j) = \varphi(h_k)$  for  $j, k \in [1, (n - s + 1)m]$ , then  $h_j = h_k$ . As  $|h_j^{-1}T| = (n - s + 1 - 2)m + 2m - 1$ , using again the projection  $\psi$  introduced above and Theorem 4.1, it follows that there exists an admissible factorization  $T = S'_0 S'_1 \dots S'_{n-s}$  with  $h_j | S'_0$ . Let  $\ell \in [1, n - s]$  such that  $h_j h_k | S'_0 S'_\ell$ . Let  $S'_0 S'_\ell = T'_j T'_k$  such that  $h_j | T'_j$ ,  $h_k | T'_k$  and  $|T'_j| = |T'_k| = m$ . As above, it follows that  $\sigma(\varphi(T'_j)) = (a' + 1)f_1$  and  $\sigma(\varphi(T'_k)) = -a'f_1$  for some  $a' \in [0, m - 1]$ . We note that  $\sigma(T'_j g_1^{m-a'-1})\sigma(T'_k g_1^{a'})\sigma(g_1^m)^{s-1} \prod_{i=1, i \neq \ell}^{n-s} \sigma(S'_i)$  is a ml-mzss over  $H$  and thus equals  $e^m$  for a generating element  $e' \in H$ . As above, it follows that  $\sigma(h_j^{-1}h_k T'_j g_1^{m-a'-1}) = e'$ . Thus,  $h_j = h_k$ .

Consequently, we have

$$S = g_1^{sm-1} \prod_{x \in [0, m-1]} k_x^{v_x}$$

with  $\varphi(k_x) = x f_1 + f_2$  for  $x \in [0, m - 1]$  and suitable  $v_x \in \mathbb{N}_0$ .

In the following we show that  $S$  is of the form given in (ii) or  $\text{ord } g_1 = m$ . At the end we show that if  $\text{ord } g_1 = m$ , then  $S$  is of the form given in (i). We start with the following assertion.

ASSERTION. *Let  $T = \bar{S}_0 \bar{S}_1 \dots \bar{S}_{n-s}$  be an admissible factorization. Let  $k_x | \bar{S}_0$  and let  $k_y | \bar{S}_i$  for some  $i \in [1, n - s]$ . If  $x < y$ , then  $k_y - k_x = (y - x)g_1$  and if  $x > y$ , then  $k_y - k_x = (y - x)g_1 + mg_1$ .*

*Proof of Assertion.* We note that  $\sigma(\varphi(\bar{S}_0 k_x^{-1} k_y)) = (-x + y + 1)f_1$  and  $\sigma(\varphi(\bar{S}_i k_y^{-1} k_x)) = (-y + x)f_1$ . Thus, we have  $\sigma(\varphi(\bar{S}_0 k_x^{-1} k_y g_1^{[x-y-1]_m})) = 0 = \sigma(\varphi(\bar{S}_i k_y^{-1} k_x g_1^{[y-x]_m}))$ . We observe that  $[x - y - 1]_m + [y - x]_m = m - 1$ . Thus, as above,  $\sigma(\bar{S}_i) = \sigma(\bar{S}_i k_y^{-1} k_x g_1^{[y-x]_m})$ . Hence,  $k_y = k_x + [y - x]_m g_1$ .

Consequently, if  $x < y$ , then  $k_y - k_x = (y - x)g_1$ , and if  $x > y$ , then  $k_y - k_x = (y - x)g_1 + mg_1$ , proving the Assertion.

First, we show that  $\text{supp}(S_0^{-1}T) \subset \text{supp}(S_0)$  or  $\text{ord } g_1 = m$ . We assume that there exists some  $i \in [1, n - s]$  and some  $k_t | S_i$  such that  $k_t \nmid S_0$  and show that this implies  $\text{ord } g_1 = m$ .

The sequence  $k_t^{-1}S_iS_0$  has length  $2m - 1$ . Thus, as above, there exists a subsequence  $S_i'' | k_t^{-1}S_iS_0$  such that  $\sigma(\varphi(S_i'')) = 0$  and  $|S_i''| = m$ . Let  $S_0'' = S_i''^{-1}S_iS_0$  and  $S_j'' = S_j$  for  $j \notin \{0, i\}$ . We see that  $S_0''S_1'' \dots S_{n-s}''$  is an admissible factorization of  $T$ . Since  $k_t | S_0''$  and  $k_t \nmid S_0$  and  $|\text{supp}(S_0)|$  is maximal (by assumption), there exists some  $k_u | S_0$  such that  $k_u \nmid S_0''$  and thus  $k_u | S_i''$ . Clearly  $k_u \neq k_t$  and thus  $t \neq u$ .

We apply the Assertion twice. First, to  $k_u | S_0$  and  $k_t | S_i$ . If  $u < t$ , then  $k_t - k_u = (t - u)g_1$  and if  $u > t$ , then  $k_t - k_u = (t - u)g_1 + mg_1$ . Second, to  $k_t | S_0''$  and  $k_u | S_i''$ . If  $t < u$ , then  $k_u - k_t = (u - t)g_1$  and if  $t > u$ , then  $k_u - k_t = (u - t)g_1 + mg_1$ .

Thus, if  $u < t$ , then  $k_t - k_u = (t - u)g_1$  and  $k_u - k_t = (u - t)g_1 + mg_1$ . Adding these two equations, we get  $mg_1 = 0$  and  $\text{ord } g_1 = m$ . And, if  $u > t$ , then  $k_t - k_u = (t - u)g_1 + mg_1$  and  $k_u - k_t = (u - t)g_1$ , again yielding  $\text{ord } g_1 = m$ .

Second, we show that  $|\text{supp}(S_0^{-1}T)| = 1$  or  $\text{ord } g_1 = m$ . We assume that  $|\text{supp}(S_0^{-1}T)| \geq 2$ , say it contains elements  $k_u, k_t$  with  $t > u$ . Let  $i, j \in [1, n - s]$ , not necessarily distinct, be such that  $k_u | S_i$  and  $k_t | S_j$ . By the above argument we may assume that  $\text{supp}(S_0^{-1}T) \subset \text{supp}(S_0)$ . We apply the Assertion with  $k_t | S_0$  and  $k_u | S_i$  to obtain  $k_u - k_t = (u - t)g_1 + mg_1$ . Then we apply the Assertion with  $k_u | S_0$  and  $k_t | S_j$  to obtain  $k_t - k_u = (t - u)g_1$ . Thus, we obtain  $mg_1 = 0$ .

Consequently, we have  $T = k_u^{(n-s)m}S_0$  and  $k_u | S_0$  for some  $u \in [0, m - 1]$  or  $\text{ord } g_1 = m$ .

We assume that  $T = k_u^{(n-s)m}S_0$  and  $k_u | S_0$  for some  $u \in [0, m - 1]$ . Since  $n - s \geq 1$  and  $\sigma(k_u^m) = e$ , it follows that  $\text{ord } k_u = mn$ . Let  $f'_2 = \varphi(k_u) = uf_1 + f_2$ . It follows that  $\{f_1, f'_2\}$  is a basis of  $G/H$ .

If, for  $x \in [0, m - 1]$ , an element  $h \in \text{supp}(T) = \text{supp}(S_0)$  exists with  $\varphi(h) = -xf_1 + f'_2$  (as shown above there is at most one such element), then we denote it by  $k'_{-x}$ . In particular,  $k_u = k'_0$ .

For each  $k'_{-x} \in \text{supp}(S_0)$ , as above,  $\sigma(k_0'^m) = \sigma(k_0'^{m-1}k'_{-x}g_1^x)$ . Thus, we have  $k'_0 = k'_{-x} + xg_1$ . Let  $x_i \in [0, m - 1]$  be such that  $S_0 = \prod_{i=1}^m k'_{-x_i}$ . We know that  $\sum_{i=1}^m (-x_i f_1) = f_1$ , i.e.,  $\sum_{i=1}^m x_i \equiv m - 1 \pmod{m}$ .

We show that  $\sum_{i=1}^m x_i = m - 1$  or  $\text{ord } g_1 = m$ . Assume the former does not hold, and let  $\ell$  be maximal such that  $\sum_{i=1}^\ell x_i = c < m$ . We observe that  $\sigma(k_0'^{m-\ell}(\prod_{i=1}^\ell f'_{-x_i}g_1^c)) = \sigma(k_0'^{m-\ell-1}(\prod_{i=1}^{\ell+1} f'_{-x_i})g_1^{[c+x_{\ell+1}]m})$ . By



the choice of  $\ell$  it follows that  $[c + x_{\ell+1}]_m = c + x_{\ell+1} - m$ . Thus,  $k'_0 + cg_1 = k_{-x_{\ell+1}} + (c + x_{\ell+1} - m)g_1$  and  $k'_0 = k_{-x_{\ell+1}} + (x_{\ell+1} - m)g_1$ , which implies  $mg_1 = 0$ .

So,  $\text{ord } g_1 = m$ , or  $S$  is of the form, where  $k'_0 = g_2$ ,

$$S = g_1^{sm-1} g_2^{(n-s)m} \prod_{i=1}^m (-x_i g_1 + g_2)$$

with  $x_i \in [0, m - 1]$  and  $\sum_{i=1}^m x_i = m - 1$ . Clearly,  $\{g_1, g_2\}$  is a generating set of  $G$ . Moreover, we know that if  $s \geq 2$ , then  $\sigma(g_1^m) = e = \sigma(g_2^m)$ . Thus,  $S$  is of the form given in (ii).

Finally, suppose  $\text{ord } g_1 = m$ . We have  $s = 1$ . Let  $\omega : G \rightarrow G/\langle g_1 \rangle$  denote the canonical map. The sequence  $\omega(\prod_{i=1}^{mn} h_i)$  is a mzss. Thus, by Theorems 4.1 and 4.2,  $G/\langle g_1 \rangle$  is a cyclic group of order  $mn$  and  $\omega(\prod_{i=1}^{mn} h_i) = \omega(g_2)^{mn}$  for some  $g_2 \in G$ , and  $\text{ord } g_2 = mn$ . Thus,  $\{g_1, g_2\}$  is a basis of  $G$  and  $S$  has the form given in (i).

CASE 2:  $\varphi(S)$  is of the form given in (4.2). If  $m = 2$ , then  $bf_1 + 2f_2 = f_1$  and the sequence  $\varphi(S)$  is also of the form given in (4.1). Thus, we assume  $m \geq 3$ . Moreover, we note that with respect to the basis  $f'_1 = f_1$  and  $f'_2 = bf_1 + f_2$ , we have  $f_2 = (m - b)f'_1 + e'_2$  and  $bf_1 + 2f_2 = (m - b)f'_1 + 2f'_2$ . Thus, we may assume that  $b < m/2$ . Let  $S = FT$  with  $\varphi(T) = f_1^m (bf_1 + f_2)^{m-1} f_2^{m-1} (bf_1 + 2f_2)$ . We note that  $F = \prod_{i=1}^{n-2} F_i$  with  $\varphi(F_i) \in \{f_1^m, f_2^m, (bf_1 + f_2)^m\}$  for each  $i \in [1, n - 2]$ . Suppose  $T = T_1 T_2$  such that  $\sigma(\varphi(T_i)) = 0$  and  $T_i \neq 1$  for  $i \in [1, 2]$ . Then  $\sigma(T_1)\sigma(T_2) \prod_{i=1}^{n-2} \sigma(F_i)$  is a ml-mzss over  $H$  and thus equal to  $e^n$  for some generating element  $e$  of  $H$ . It follows that for each factorization  $T = T_1 T_2$  with  $\sigma(\varphi(T_i)) = 0$  and  $T_i \neq 1$  for  $i \in [1, 2]$ , we have  $\sigma(T_1) = \sigma(T_2) = e$ .

Let  $T'_1 | T$  be such that  $\varphi(T'_1) = f_1^b (bf_1 + f_2)^{m-1} f_2$  and let  $T'_2 = T'^{-1}_1 T$ . Suppose that, for some  $i \in [1, 2]$ , there exist distinct elements  $g_i, g'_i \in \text{supp}(S)$  such that  $\varphi(g_i) = \varphi(g'_i) = f_i$ . We may assume that  $g_i | T'_1$  and  $g'_i | T'_2$ . It follows that  $\sigma(g_i^{-1} g'_i T'_1) = e = \sigma(T'_1)$ , a contradiction. Thus,  $\varphi^{-1}(f_i) \cap \text{supp}(S) = \{g_i\}$  for  $i \in [1, 2]$ .

Now, let  $T''_1 | T$  be such that  $\varphi(T''_1) = f_1^{2b} (bf_1 + f_2)^{m-2} f_2^2$  and  $T''_2 = T''^{-1}_1 T$ . We can argue in the same way that  $\varphi^{-1}(bf_1 + f_2) \cap \text{supp}(S) = \{k_b\}$ . Finally, let  $k | S$  be such that  $\varphi(k) = bf_1 + 2f_2$ . It follows that

$$S = g_1^{s_1 m} g_2^{s_2 m-1} k_b^{s_3 m-1} k.$$

We note that  $\text{ord } g_1 = mn$  and that  $\{g_1, g_2\}$  is a generating set of  $G$ .

Since, as above,  $\sigma(k_b^{m-2} k g_1^b) = e = \sigma(k_b^{m-1} k g_2^{m-1})$ , it follows that  $bg_1 = k_b + (m - 1)g_2$ . Moreover,  $\sigma(g_1^{2b} k_b^{m-2} g_2^2) = e = \sigma(g_1^b k_b^{m-1} g_2)$  implies that  $bg_1 + g_2 = k_b$ . Thus,  $mg_2 = 0$  and  $\{g_2, g_1\}$  is a basis of  $G$ . Moreover, we have  $s_2 = 1$ . Additionally, we get  $k + (m - 1)g_2 = bg_1 + g_2$ , i.e.,  $k = bg_1 + 2g_2$ .

We observe that the projection to  $\langle g_1 \rangle$  (with respect to  $G = \langle g_1 \rangle \oplus \langle g_2 \rangle$ ) of the sequence  $g_2^{-(m-1)}S$ , i.e., the sequence  $g_1^{s_1 m}(bg_1)^{s_3 m}$ , is a mzss. Since  $s_1 + s_3 = n$ , this implies  $b = 1$ . Thus,  $S$  is of the form given in (i).

**Acknowledgments.** The author would like to thank the referee for suggestions and corrections, and A. Geroldinger and D. Gryniewicz for valuable discussions related to this paper. This work was supported by the FWF (P18779-N13).

### References

- [1] G. Bhowmik, I. Halupczok, and J.-C. Schlage-Puchta, *Inductive methods and zero-sum free sequences*, *Integers* 9 (2009), 515–536.
- [2] —, —, —, *The structure of maximal zero-sum free sequences*, *Acta Arith.* 143 (2010), 21–50.
- [3] G. Bhowmik and J.-C. Schlage-Puchta, *Davenport’s constant for groups of the form  $\mathbb{Z}_3 \oplus \mathbb{Z}_3 \oplus \mathbb{Z}_{3d}$* , in: *Additive Combinatorics*, Amer. Math. Soc., Providence, RI, 2007, 307–326.
- [4] J. D. Bovey, P. Erdős, and I. Niven, *Conditions for a zero sum modulo  $n$* , *Canad. Math. Bull.* 18 (1975), 27–29.
- [5] F. Chen and S. Savchev, *Minimal zero-sum sequences of maximum length in the group  $C_3 \oplus C_{3k}$* , *Integers* 7 (2007), A42, 6 pp.
- [6] P. Erdős, A. Ginzburg, and A. Ziv, *Theorem in the additive number theory*, *Bull. Res. Council Israel* 10F (1961), 41–43.
- [7] W. Gao and A. Geroldinger, *On long minimal zero sequences in finite abelian groups*, *Period. Math. Hungar.* 38 (1999), 179–211.
- [8] —, —, *On the order of elements in long minimal zero-sum sequences*, *ibid.* 44 (2002), 63–73.
- [9] —, —, *On zero-sum sequences in  $\mathbb{Z}/n\mathbb{Z} \oplus \mathbb{Z}/n\mathbb{Z}$* , *Integers* 3 (2003), A8, 45 pp.
- [10] —, —, *Zero-sum problems in finite abelian groups: a survey*, *Expo. Math.* 24 (2006), 337–369.
- [11] W. Gao, A. Geroldinger, and D. Gryniewicz, *Inverse zero-sum problems III*, *Acta Arith.* 141 (2010), 103–152.
- [12] W. Gao, A. Geroldinger, and W. A. Schmid, *Inverse zero-sum problems*, *ibid.* 128 (2007), 245–279.
- [13] A. Geroldinger and F. Halter-Koch, *Non-unique Factorizations. Algebraic, Combinatorial and Analytic Theory*, Chapman & Hall/CRC, Boca Raton, FL, 2006.
- [14] A. Geroldinger and I. Z. Ruzsa, *Combinatorial Number Theory and Additive Group Theory*, Birkhäuser, Basel, 2009.
- [15] B. Girard, *Inverse zero-sum problems and algebraic invariants*, *Acta Arith.* 135 (2008), 231–246.
- [16] G. Lettl and W. A. Schmid, *Minimal zero-sum sequences in  $C_n \oplus C_n$* , *Eur. J. Combin.* 28 (2007), 742–753.
- [17] J. E. Olson, *A combinatorial problem on finite Abelian groups II*, *J. Number Theory* 1 (1969), 195–199.
- [18] S. Savchev and F. Chen, *Long zero-free sequences in finite cyclic groups*, *Discrete Math.* 307 (2007), 2671–2679.

- [19] W. A. Schmid, *The inverse problem associated to the Davenport constant for  $C_2 \oplus C_2 \oplus C_{2n}$ , and applications to the arithmetical characterization of class groups*, <http://www.uni-graz.at/~wschmid/personal/research.html>.
- [20] T. Tao and V. Vu, *Additive Combinatorics*, Cambridge Univ. Press, Cambridge, 2006.
- [21] P. van Emde Boas, *A combinatorial problem on finite Abelian groups II*, Technical report, Math. Centrum, Amsterdam, Afd. Zuivere Wisk. ZW 1969-007 (1969), 60 pp.
- [22] P. Yuan, *On the index of minimal zero-sum sequences over finite cyclic groups*, J. Combin. Theory Ser. A 114 (2007), 1545–1551.

Wolfgang A. Schmid  
Institut für Mathematik und Wissenschaftliches Rechnen  
Karl-Franzens-Universität Graz  
Heinrichstraße 36  
8010 Graz, Austria  
E-mail: wolfgang.schmid@uni-graz.at

*Received on 3.2.2008  
and in revised form on 14.1.2010*

(5631)