

On the fiber product preserving gauge bundle functors on vector bundles

by WŁODZIMIERZ M. MIKULSKI (Kraków)

Abstract. We present a complete description of all fiber product preserving gauge bundle functors F on the category \mathcal{VB}_m of vector bundles with m -dimensional bases and vector bundle maps with local diffeomorphisms as base maps. Some corollaries of this result are presented.

Introduction. Modern differential geometry has clarified that product preserving bundle functors on the category $\mathcal{M}f$ of manifolds and maps play very important roles. To such bundle functors one can lift some geometric structures as vector fields, forms, connections, etc. To define such lifts the only important property is the product preservation. Such functors have been classified by means of Weil algebras [5].

Research quite similar to that on manifolds has been done on fibered manifolds. A wide class of bundle functors on the category \mathcal{FM}_m of fibered manifolds with m -dimensional bases and fiber preserving maps with local diffeomorphisms as base maps is the class of fiber product preserving functors. Such functors have been classified in [6], and studied in [1]–[4], [8].

In turn research similar to that on fibered manifolds has been done on vector bundles. A wide class of (gauge) bundle functors on the category \mathcal{VB}_m of vector bundles with m -dimensional bases and vector bundle maps with local diffeomorphisms as base maps is the class of fiber product preserving functors. For example the r -jet prolongation functor plays an important role in the theory of higher order connections, Lagrangians, differential equations, etc. Below, we present many examples of such functors. Some of them are well known. It seems natural and useful to classify all such functors. The purpose of the present paper is to describe all fiber product preserving gauge bundle functors on \mathcal{VB}_m .

2000 *Mathematics Subject Classification*: 58A05, 58A20.

Key words and phrases: (fiber product preserving) gauge bundle functors, natural transformations, jets, Weil algebras.

We first recall the following definitions (see e.g. [5]).

Let $F : \mathcal{VB}_m \rightarrow \mathcal{FM}$ be a covariant functor into the category \mathcal{FM} of fibered manifolds and their fibered maps. Let $B_{\mathcal{VB}_m} : \mathcal{VB}_m \rightarrow \mathcal{Mf}$ and $B_{\mathcal{FM}} : \mathcal{FM} \rightarrow \mathcal{Mf}$ be the respective base functors.

A gauge bundle functor on \mathcal{VB}_m is a functor F as above satisfying:

(i) (*Base preservation*) $B_{\mathcal{FM}} \circ F = B_{\mathcal{VB}_m}$. Hence the induced projections form a functor transformation $\pi : F \rightarrow B_{\mathcal{VB}_m}$.

(ii) (*Localization*) For every inclusion of an open vector subbundle $i_{E|U} : E|U \rightarrow E$, $F(E|U)$ is the restriction $\pi^{-1}(U)$ of $\pi : FE \rightarrow B_{\mathcal{VB}_m}(E)$ to U and $F i_{E|U}$ is the inclusion $\pi^{-1}(U) \rightarrow FE$.

(iii) (*Regularity*) F transforms smoothly parametrized systems of \mathcal{VB}_m -morphisms into smoothly parametrized systems of \mathcal{FM} -morphisms.

A gauge bundle functor $F : \mathcal{VB}_m \rightarrow \mathcal{FM}$ is of finite order r if from $j_x^r f = j_x^r g$ it follows that $F_x f = F_x g$ for any \mathcal{VB}_m -objects $E_1 \rightarrow M_1$, $E_2 \rightarrow M_2$, any \mathcal{VB}_m -maps $f, g : E_1 \rightarrow E_2$ and any $x \in M_1$.

Given two gauge bundle functors F_1, F_2 on \mathcal{VB}_m , by a natural transformation $\mu : F_1 \rightarrow F_2$ we mean a system of base preserving fibered maps $\mu : F_1 E \rightarrow F_2 E$ for every vector bundle E satisfying $F_2 f \circ \mu = \mu \circ F_1 f$ for every \mathcal{VB}_m -morphism $f : E \rightarrow G$.

A gauge bundle functor F on \mathcal{VB}_m is fiber product preserving if for any fiber product projections

$$E_1 \xleftarrow{\text{pr}_1} E_1 \times_M E_2 \xrightarrow{\text{pr}_2} E_2$$

in the category \mathcal{VB}_m ,

$$F E_1 \xleftarrow{F \text{pr}_1} F(E_1 \times_M E_2) \xrightarrow{F \text{pr}_2} F E_2$$

are fiber product projections in the category \mathcal{FM} . In other words, we have $F(E_1 \times_M E_2) = F(E_1) \times_M F(E_2)$ modulo the corestriction of $(F \text{pr}_1, F \text{pr}_2)$.

The most important example of a fiber product preserving gauge bundle functor is the *r-jet prolongation functor* $J^r : \mathcal{VB}_m \rightarrow \mathcal{FM}$, where for a \mathcal{VB}_m -object $p : E \rightarrow M$ we have $J^r E = \{j_x^r \sigma \mid \sigma \text{ is a local section of } E, x \in M\}$ and for a \mathcal{VB}_m -map $f : E_1 \rightarrow E_2$ covering $\underline{f} : M_1 \rightarrow M_2$ we have $J^r f : J^r E_1 \rightarrow J^r E_2$, where $J^r f(j_x^r \sigma) = j_{\underline{f}(x)}^r (f \circ \sigma \circ \underline{f}^{-1})$ for $j_x^r \sigma \in J^r E_1$.

Another example is the so-called *vertical r-jet prolongation functor* $J_v^r : \mathcal{VB}_m \rightarrow \mathcal{FM}$, where for a \mathcal{VB}_m -object $p : E \rightarrow M$ we have $J_v^r E = \{j_x^r \gamma \mid \gamma \text{ is a local map } M \rightarrow E_x, x \in M\}$ and for a \mathcal{VB}_m -map $f : E_1 \rightarrow E_2$ covering $\underline{f} : M_1 \rightarrow M_2$ we have $J_v^r f : J_v^r E_1 \rightarrow J_v^r E_2$, where $J_v^r f(j_x^r \gamma) = j_{\underline{f}(x)}^r (f \circ \gamma \circ \underline{f}^{-1})$ for $j_x^r \gamma \in J_v^r E_1$.

Still another example is the *vertical Weil functor* $V^A : \mathcal{VB}_m \rightarrow \mathcal{FM}$ corresponding to a Weil algebra A , where for a \mathcal{VB}_m -object $p : E \rightarrow M$

we have $V^A E = \bigcup_{x \in M} T^A(E_x)$ and for a \mathcal{VB}_m -map $f : E_1 \rightarrow E_2$ we have $V^A f = \bigcup_{x \in M_1} T^A(f_x) : V^A E_1 \rightarrow V^A E_2$. The functor $V^A E$ is equivalent to $E \otimes A$.

One more example is the following vector bundle modification $T_{fl}^{(r)} : \mathcal{VB}_m \rightarrow \mathcal{FM}$ of the vector r -tangent bundle $T^{(r)}$ over manifolds. For a vector bundle $p : E \rightarrow M$ from \mathcal{VB}_m we have a vector bundle $T_{fl}^{r*} E = J_{fl}^r(E, \mathbb{R})_0 = \{j_x^r \gamma \mid \gamma : E \rightarrow \mathbb{R} \text{ is fiber linear, } \gamma_x = 0, x \in M\}$ over M , where γ_x is the restriction of γ to the fiber E_x of E over $x \in M$. Let $T_{fl}^{(r)} E = (T_{fl}^{r*} E)^*$ be the dual vector bundle. Every \mathcal{VB}_m -map $f : E \rightarrow \bar{E}$ covering $\underline{f} : M \rightarrow \bar{M}$ induces a \mathcal{VB}_m -map $T_{fl}^{(r)} f : T_{fl}^{(r)} E \rightarrow T_{fl}^{(r)} \bar{E}$ covering \underline{f} , where $\langle T_{fl}^{(r)} f(\omega), j_{\underline{f}(x)}^r \gamma \rangle = \langle \omega, j_x^r(\gamma \circ f) \rangle$ for $\omega \in (T_{fl}^{(r)})_x E$, and $\gamma : \bar{E} \rightarrow \mathbb{R}$ is fiber linear with $\gamma_{\underline{f}(x)} = 0, x \in M$.

The fiber product $F_1 \times_{B_{\mathcal{VB}_m}} F_2 : \mathcal{VB}_m \rightarrow \mathcal{FM}$ of fiber product preserving gauge bundle functors $F_1, F_2 : \mathcal{VB}_m \rightarrow \mathcal{FM}$ is again a fiber product preserving gauge bundle functor. We recall that $(F_1 \times_{B_{\mathcal{VB}_m}} F_2)(E) = F_1 E \times_M F_2 E$ for any \mathcal{VB}_m -object $E \rightarrow M$, and $(F_1 \times_{B_{\mathcal{VB}_m}} F_2)(f)(v_1, v_2) = (F_1 f(v_1), F_2 f(v_2))$ for any \mathcal{VB}_m -map $f : E \rightarrow G$ and any $(v_1, v_2) \in F_1 E \times_M F_2 E$.

The composition of fiber product preserving gauge bundle functors on \mathcal{VB}_m is again a fiber product preserving gauge bundle functor on \mathcal{VB}_m . (In Lemma 1 it will be proved that every fiber product preserving gauge bundle functor has values in \mathcal{VB}_m . So, the composition is possible.)

If $F : \mathcal{VB}_m \rightarrow \mathcal{FM}$ is a fiber product preserving gauge bundle functor of order r we define a new fiber product preserving gauge bundle functor $(F^*)^* : \mathcal{VB}_m \rightarrow \mathcal{FM}$ by $E \mapsto (FE^*)^*$ and $f \mapsto (Ff^*)^*$, where $(\)^*$ denotes the dualization of \mathcal{VB}_m -objects and \mathcal{VB}_m -maps.

The first main result in this paper is that all fiber product preserving gauge bundle functors F on \mathcal{VB}_m of finite order r are in bijection with so-called admissible triples, i.e. triples (V, H, t) where V is a finite-dimensional vector space over \mathbb{R} , $H : G_m^r \rightarrow GL(V)$ is a smooth group homomorphism from $G_m^r = \text{inv } J_0^r(\mathbb{R}^m, \mathbb{R}^m)_0$ into $GL(V)$ and $t : \mathcal{D}_m^r \rightarrow \text{gl}(V)$ is a G_m^r -equivariant unital associative algebra homomorphism from $\mathcal{D}_m^r = J_0^r(\mathbb{R}^m, \mathbb{R})$ into $\text{gl}(V)$.

The second main result is that natural transformations between two fiber product preserving gauge bundle functors on \mathcal{VB}_m of order r are in bijection with morphisms between the corresponding admissible triples.

The third main result is that any fiber product preserving gauge bundle functor on \mathcal{VB}_m is of finite order.

As corollaries of the above results we describe explicitly all natural endomorphisms $J^r \rightarrow J^r, J_v^r \rightarrow J_v^r$ and $V^A \rightarrow V^A$ for any Weil algebra A .

All manifolds are assumed to be finite-dimensional. All manifolds and maps are assumed to be smooth, i.e. of class \mathcal{C}^∞ .

1. Fiber product preserving gauge bundle functors on \mathcal{VB}_m corresponding to admissible triples

DEFINITION 1. An *admissible triple of order r and dimension m* is a triple (V, H, t) , where V is a finite-dimensional vector space over \mathbb{R} , $H : G_m^r \rightarrow \text{GL}(V)$ is a smooth group homomorphism from the Lie group $G_m^r = \text{inv } J_0^r(\mathbb{R}^m, \mathbb{R}^m)_0$ of invertible r -jets at $0 \in \mathbb{R}^m$ of diffeomorphisms $\mathbb{R}^m \rightarrow \mathbb{R}^m$ preserving 0 into the group $\text{GL}(V)$ of linear isomorphisms of V , and $t : \mathcal{D}_m^r \rightarrow \text{gl}(V)$ is a G_m^r -equivariant unital algebra homomorphism from the unital algebra $\mathcal{D}_m^r = J_0^r(\mathbb{R}^m, \mathbb{R})$ of r -jets at $0 \in \mathbb{R}^m$ of maps $\mathbb{R}^m \rightarrow \mathbb{R}$ into the unital associative algebra $\text{gl}(V)$ of linear endomorphisms of V .

We recall that G_m^r acts on \mathcal{D}_m^r by $j_0^r \varphi \cdot j_0^r \gamma = j_0^r(\gamma \circ \varphi^{-1})$ for $j_0^r \varphi \in G_m^r$, $j_0^r \gamma \in \mathcal{D}_m^r$. This action will be denoted by H_m^r . We also recall that G_m^r acts on $\text{gl}(V)$ by $\xi \cdot A = H(\xi) \circ A \circ H(\xi^{-1})$ for $\xi \in G_m^r$, $A \in \text{gl}(V)$. These actions are by unital algebra isomorphisms.

Let (V, H, t) be an admissible triple of order r and dimension m . We are going to construct a fiber product preserving gauge bundle functor $T^{(V,H,t)} : \mathcal{VB}_m \rightarrow \mathcal{FM}$.

EXAMPLE 1. For a vector bundle $p : E \rightarrow M$ from \mathcal{VB}_m we put

$$T^{(V,H,t)} E = \bigcup_{x \in M} \text{Hom}_{t_x}(J^r \mathcal{C}_x^{\infty, fl}(E), \tilde{V}_x M).$$

Here $\tilde{V} : \mathcal{M}f_m \rightarrow \mathcal{VB}$ is the natural vector bundle corresponding to the G_m^r space V , i.e. $\tilde{V}M = P^r M[V, H]$ (the associated bundle) for any m -manifold M and $\tilde{V}\varphi = P^r \varphi[\text{id}_V] : \tilde{V}M_1 \rightarrow \tilde{V}M_2$ for any embedding $\varphi : M_1 \rightarrow M_2$ between m -manifolds, and $\text{Hom}_{t_x}(J^r \mathcal{C}_x^{\infty, fl}(E), \tilde{V}_x M)$ is the space of module homomorphisms over $t_x : J_x^r(M, \mathbb{R}) \rightarrow \text{gl}(\tilde{V}_x M)$ from the (free) $J_x^r(M, \mathbb{R})$ -module $J^r \mathcal{C}_x^{\infty, fl}(E)$ of r -jets at $x \in M$ of germs at x of fiber linear maps $E \rightarrow \mathbb{R}$ into the $\text{gl}(\tilde{V}_x M)$ -module $\tilde{V}_x M$, where $t_x : J_x^r(M, \mathbb{R}) \rightarrow \text{gl}(\tilde{V}_x M)$ is the unital algebra homomorphism induced by t such that $t_x(j_x^r \gamma) = \tilde{V}_0 \varphi \circ t(j_0^r(\gamma \circ \varphi)) \circ (\tilde{V}_0 \varphi)^{-1}$ for any $\gamma : M \rightarrow \mathbb{R}$ and any embedding $\varphi : \mathbb{R}^m \rightarrow M$ with $\varphi(0) = x$ (t_x is well defined because t is G_m^r -equivariant).

Given a vector bundle trivialization $(x^1 \circ p, \dots, x^m \circ p, y^1, \dots, y^n) : E|U \rightarrow \mathbb{R}^m \times \mathbb{R}^n$ we have an induced fiber bundle trivialization $(\tilde{x}^1, \dots, \tilde{x}^m, \tilde{y}^1, \dots, \tilde{y}^n) : T^{(V,H,t)} E|U \rightarrow \mathbb{R}^m \times V^n$ such that $\tilde{x}^i(\Phi) = x^i(x_0) \in \mathbb{R}$ and $\tilde{y}^j(\Phi) = \Phi(j_{x_0}^r(y^j)) \in \tilde{V}_{x_0} M \cong V$ for any $\Phi \in \text{Hom}_{t_{x_0}}(J^r \mathcal{C}_{x_0}^{\infty, fl}(E), \tilde{V}_{x_0} M)$, $i=1, \dots, m, j=1, \dots, n$, where $V \cong \tilde{V}_{x_0} M$ is given by $v \leftrightarrow \tilde{V}((x^1, \dots, x^m)^{-1}$

$\circ \tau_{(x^i(x_0))}(v)$ for $v \in V$, and $\tau_y : \mathbb{R}^m \rightarrow \mathbb{R}^m$ is the translation by $y \in \mathbb{R}^m$. Then $T^{(V,H,t)}E$ with the obvious projection is a fiber bundle over M .

Every \mathcal{VB}_m -map $f : E_1 \rightarrow E_2$ covering $\underline{f} : M_1 \rightarrow M_2$ induces a fibered map $T^{(V,H,t)}E_1 \rightarrow T^{(V,H,t)}E_2$ covering \underline{f} such that

$$T^{(V,H,t)}f(\Phi)(j_{\underline{f}(x)}^r \xi) = \tilde{V} \underline{f} \circ \Phi(j_x^r(\xi \circ f))$$

for any $\Phi \in \text{Hom}_{t_x}(J^r C_x^{\infty,fl}(E_1), \tilde{V}_x M)$, $x \in M_1$, and any fiber linear map $\xi : E_2 \rightarrow \mathbb{R}$.

If in some vector bundle trivializations $f(x, y) = (\underline{f}(x), \sum_j f^j(x)y_j)$ for $x \in \mathbb{R}^m$, $y = (y_1, \dots, y_n) \in \mathbb{R}^n$, $f^j = (f_l^j) : \mathbb{R}^m \rightarrow \mathbb{R}^k$, $j = 1, \dots, n$, $l = 1, \dots, k$, then in the induced trivializations

$$T^{(V,H,t)}f(x, v) = \left(\underline{f}(x), \left(\sum_j t(j_0^r(f_l^j \circ \underline{f}^{-1} \circ \tau_{\underline{f}(x)})) (H(j_0^r(\tau_{-\underline{f}(x)} \circ \underline{f} \circ \tau_x))(v_j)) \right)_{l=1}^k \right)$$

for $x \in \mathbb{R}^m$ and $v = (v_1, \dots, v_n) \in V^n$.

The correspondence $T^{(V,H,t)} : \mathcal{VB}_m \rightarrow \mathcal{FM}$ is a fiber product preserving gauge bundle functor of order r and it takes values in the category \mathcal{VB}_m .

DEFINITION 2. We call $T^{(V,H,t)} : \mathcal{VB}_m \rightarrow \mathcal{FM}$ the fiber product preserving gauge bundle functor corresponding to the admissible triple (V, H, t) .

2. Admissible triples corresponding to fiber product preserving gauge bundle functors on \mathcal{VB}_m

LEMMA 1. Let $F : \mathcal{VB}_m \rightarrow \mathcal{FM}$ be a f.p.p.g.b. functor.

(a) Given a vector bundle $p : E \rightarrow M$ we have a canonical vector bundle structure on FE .

(b) Given a \mathcal{VB}_m -map $f : E_1 \rightarrow E_2$ over $\underline{f} : M_1 \rightarrow M_2$ the induced map $Ff : FE_1 \rightarrow FE_2$ is a vector bundle map over \underline{f} .

Proof. The fiber sum map $+^E : E \times_M E \rightarrow E$, the fiber scalar multiplication maps $\lambda_t^E : E \rightarrow E$ for $t \in \mathbb{R}$ and the zero map $0^E : E \rightarrow E$ are \mathcal{VB}_m -maps and we can apply the functor F . We obtain $+^{FE} := F(+^E) : FE \times_M FE \cong F(E \times_M E) \rightarrow FE$, $\lambda_t^{FE} := F(\lambda_t^E) : FE \rightarrow FE$ and $0^{FE} := F(0^E) : FE \rightarrow FE$. It is easily seen that $(FE, +^{FE}, \lambda_t^{FE}, 0^{FE})$ is a vector bundle structure on FE . ■

Let $F : \mathcal{VB}_m \rightarrow \mathcal{FM}$ be a f.p.p.g.b. functor of order r . We are going to construct an admissible triple (V^F, H^F, t^F) .

EXAMPLE 2. We put

$$V^F := F_0(\mathbb{R}^m \times \mathbb{R}),$$

the fiber at $0 \in \mathbb{R}^m$ of the vector bundle $F(\mathbb{R}^m \times \mathbb{R})$, where $\mathbb{R}^m \times \mathbb{R}$ is (of course) the trivial vector bundle over \mathbb{R}^m with fiber \mathbb{R} . Then V^F is a finite-dimensional vector space over \mathbb{R} .

We define $H^F : G_m^r \rightarrow \text{GL}(V^F)$ by

$$H^F(\xi)(v) := F_0(\varphi \times \text{id}_{\mathbb{R}})(v), \quad v \in V^F, \xi = j_0^r \varphi \in G_m^r.$$

$H^F(\xi)(v)$ is well defined because F is of order r . By the definition of V^F , $H^F(\xi) \in \text{GL}(V^F)$. By the functoriality of F , H^F is a group homomorphism. By the regularity of F , H^F is smooth.

We define $t^F : \mathcal{D}_m^r \rightarrow \text{gl}(V^F)$ by

$$t^F(\eta)(v) = F_0(\tilde{\gamma})(v), \quad v \in V^F, \eta = j_0^r \gamma \in \mathcal{D}_m^r,$$

where $\tilde{\gamma} : \mathbb{R}^m \times \mathbb{R} \rightarrow \mathbb{R}^m \times \mathbb{R}$ is a \mathcal{VB}_m -map such that $\tilde{\gamma}(x, y) = (x, \gamma(x)y)$ for $x \in \mathbb{R}^m, y \in \mathbb{R}$. Then $t^F(\eta)(v)$ is well defined because F is of order r . By the definition of V^F , $t^F(\eta) \in \text{gl}(V^F)$. By the functoriality of F and the definitions of the actions one can verify in a standard (but long) way that t^F is a G_m^r -equivariant unital algebra homomorphism.

Then (V^F, H^F, t^F) is an admissible triple of order r and dimension m .

DEFINITION 3. We call (V^F, H^F, t^F) the *admissible triple corresponding to F* .

3. Admissible triples corresponding to some fiber product preserving gauge bundle functors on \mathcal{VB}_m . In this section we present admissible triples corresponding to fiber product preserving gauge bundle functors on \mathcal{VB}_m presented in the introduction. The results of this section will not be used to prove the main result.

FACT 1. *The admissible triple corresponding to the r -jet prolongation gauge bundle functor $J^r : \mathcal{VB}_m \rightarrow \mathcal{FM}$ is $(\mathcal{D}_m^r, H_m^r, t_m^r)$, where $H_m^r : G_m^r \rightarrow \text{Aut}(\mathcal{D}_m^r)$ is defined after Definition 1 and $t_m^r : \mathcal{D}_m^r \rightarrow \text{gl}(\mathcal{D}_m^r)$ is given by $t_m^r(\eta)(\varrho) = \eta\varrho$ for $\eta, \varrho \in \mathcal{D}_m^r$.*

FACT 2. *The admissible triple corresponding to the vertical r -jet prolongation gauge bundle functor $J_v^r : \mathcal{VB}_m \rightarrow \mathcal{FM}$ is $(\mathcal{D}_m^r, H_m^r, t_m^r \circ \varepsilon_m^r)$, where $H_m^r : G_m^r \rightarrow \text{Aut}(\mathcal{D}_m^r)$ is defined after Definition 1, $t_m^r : \mathcal{D}_m^r \rightarrow \text{gl}(\mathcal{D}_m^r)$ is defined above and $\varepsilon_m^r : \mathcal{D}_m^r \rightarrow \mathbb{R} \subset \mathcal{D}_m^r$ is the algebra homomorphism.*

FACT 3. *The admissible triple corresponding to the vertical Weil gauge bundle functor $V^A : \mathcal{VB}_m \rightarrow \mathcal{FM}$ corresponding to a Weil algebra A is $(A, \text{id}_A, \varepsilon^A)$, where $\text{id}_A : G_m^r \rightarrow \{\text{id}_A\} \subset \text{GL}(A)$ is the trivial group homomorphism and $\varepsilon^A : \mathcal{D}_m^r \rightarrow \text{gl}(A)$ is given by $\varepsilon^A(\eta)(a) = \gamma(0)a$ for $\eta = j_0^r \gamma \in \mathcal{D}_m^r, a \in A$.*

The gauge bundle functors of Facts 1–3 are “almost restrictions” of fiber product preserving bundle functors on \mathcal{FM}_m . In general, let $F : \mathcal{FM}_m \rightarrow$

\mathcal{FM} be a fiber product preserving bundle functor and (A, H, t) be its corresponding triple in the sense of [6]. Then by “almost restriction” we have a fiber product preserving gauge bundle functor $\tilde{F} : \mathcal{VB}_m \rightarrow \mathcal{FM}$ given by $\tilde{F}E = FE$ for any \mathcal{VB}_m -object E and $\tilde{F}f = Ff$ for any \mathcal{VB}_m -map f .

FACT 4. *The admissible triple corresponding to \tilde{F} is $(\tilde{A}, \tilde{H}, \tilde{t})$, where \tilde{A} is the vector space A , $\tilde{H} : G_m^r \rightarrow \text{GL}(\tilde{A})$ is $H : G_m^r \rightarrow \text{Aut}(A) \subset \text{GL}(\tilde{A})$ and $\tilde{t} : \mathcal{D}_m^r \rightarrow \text{gl}(\tilde{A})$ is given by $\tilde{t}(\eta)(a) = t(\eta)a$ for $a \in \tilde{A}$ and $\eta \in \mathcal{D}_m^r$.*

The gauge bundle functor V^A of Fact 3 is a particular case of the following general construction. Let $\tilde{V} : \mathcal{M}f_m \rightarrow \mathcal{VB}$ be the natural vector bundle corresponding to a group homomorphism $H : G_m^r \rightarrow \text{GL}(V)$, i.e. $\tilde{V}M = P^r M[V, H]$ (the associated bundle) for any m -manifold M and $\tilde{V}\varphi = P^r \varphi[\text{id}_V] : \tilde{V}M_1 \rightarrow \tilde{V}M_2$ for any embedding $\varphi : M_1 \rightarrow M_2$ between m -manifolds. Let $\tilde{V} : \mathcal{VB}_m \rightarrow \mathcal{VB}$ be the fiber product preserving gauge bundle functor such that $\tilde{V}E = E \otimes_M \tilde{V}M$ for any \mathcal{VB}_m -object $p : E \rightarrow M$ and $\tilde{V}f = f \otimes \tilde{V}f : \tilde{V}E_1 \rightarrow \tilde{V}E_2$ for any \mathcal{VB}_m -map $f : E_1 \rightarrow E_2$ covering $f : M_1 \rightarrow M_2$. (For $V = A$ and trivial $H : G_m^r \rightarrow \{\text{id}_V\} \subset \text{GL}(V)$ we obtain V^A .)

FACT 5. *The admissible triple corresponding to the above $\tilde{V} : \mathcal{VB}_m \rightarrow \mathcal{FM}$ is (V, H, ε^V) , where $\varepsilon^V : \mathcal{D}_m^r \rightarrow \text{gl}(V)$ is given by $\varepsilon^V(\eta)(v) = \gamma(0)v$ for $\eta = j_0^r \gamma \in \mathcal{D}_m^r$, $v \in V$.*

FACT 6. *The admissible triple (V, H, t) of order r corresponding to $T_{fl}^{(r)} : \mathcal{VB}_m \rightarrow \mathcal{FM}$ (see the Introduction) is given by $V = T_0^{(r)}\mathbb{R}^m = (J_0^r(\mathbb{R}^m, \mathbb{R})_0)^*$, $H : G_m^r \rightarrow \text{GL}(T_0^{(r)}\mathbb{R}^m)$, $\langle H(\xi)(\omega), j_0^r \gamma \rangle = \langle \omega, j_0^r(\gamma \circ \varphi) \rangle$, $t : \mathcal{D}_m^r \rightarrow \text{gl}(T_0^{(r)}\mathbb{R}^m)$, $\langle t(\varrho)(\omega), j_0^r \gamma \rangle = \langle \omega, j_0^r(\eta\gamma) \rangle$ for $\omega \in (J_0^r(\mathbb{R}^m, \mathbb{R})_0)^*$, $\xi = j_0^r \varphi \in G_m^r$, $j_0^r \gamma \in J_0^r(\mathbb{R}^m, \mathbb{R})_0$, $\varrho = j_0^r \eta \in \mathcal{D}_m^r$.*

Let $F : \mathcal{VB}_m \rightarrow \mathcal{FM}$ be a fiber product preserving gauge bundle functor of order r . Define a new fiber product preserving gauge bundle functor $(F^*)^* : \mathcal{VB}_m \rightarrow \mathcal{FM}$ by $E \mapsto (FE^*)^*$ and $f \mapsto (Ff^*)^*$, where $(\)^*$ denotes the dualization of \mathcal{VB}_m -objects and \mathcal{VB}_m -maps.

FACT 7. *Let (V, H, t) be an admissible triple of order r corresponding to a fiber product preserving gauge bundle functor $F : \mathcal{VB}_m \rightarrow \mathcal{FM}$ of order r . The admissible triple corresponding to $(F^*)^* : \mathcal{VB}_m \rightarrow \mathcal{FM}$ is (V^*, H^*, t^*) , where $H^*(\xi) = (H(\xi^{-1}))^*$ for $\xi \in G_m^r$ and $t^*(\eta) = (t(\eta))^*$ for $\eta \in \mathcal{D}_m^r$.*

Let $F_1, F_2 : \mathcal{VB}_m \rightarrow \mathcal{FM}$ be fiber product preserving gauge bundle functors of order r . Define $F_1 \times_{B_{\mathcal{VB}_m}} F_2 : \mathcal{VB}_m \rightarrow \mathcal{FM}$ by $(F_1 \times_{B_{\mathcal{VB}_m}} F_2)(E) = F_1E \times_M F_2E$ for any \mathcal{VB}_m -object $E \rightarrow M$ and $(F_1 \times_{B_{\mathcal{VB}_m}} F_2)(f)(v_1, v_2) =$

$(F_1f(v_1), F_2f(v_2))$ for any \mathcal{VB}_m -map $f : E \rightarrow G$ and any $(v_1, v_2) \in F_1E \times_M F_2E$. Then $F_1 \times_{B\mathcal{VB}_m} F_2 : \mathcal{VB}_m \rightarrow \mathcal{FM}$ is again a fiber product preserving gauge bundle functor of order r . Let $(V^{F_1}, H^{F_1}, t^{F_1})$, $(V^{F_2}, H^{F_2}, t^{F_2})$ and $(V^{F_1 \times_{B\mathcal{VB}_m} F_2}, H^{F_1 \times_{B\mathcal{VB}_m} F_2}, t^{F_1 \times_{B\mathcal{VB}_m} F_2})$ be the admissible triples corresponding to F_1 , F_2 and $F_1 \times_{B\mathcal{VB}_m} F_2$ respectively. We have a new admissible triple $(V^{F_1} \oplus V^{F_2}, H^{F_1} \oplus H^{F_2}, t^{F_1} \oplus t^{F_2})$ of order r and dimension m such that $(H^{F_1} \oplus H^{F_2})(\xi) = H^{F_1}(\xi) \oplus H^{F_2}(\xi) : V^{F_1} \oplus V^{F_2} \rightarrow V^{F_1} \oplus V^{F_2}$ for any $\xi \in G_m^r$ and $(t^{F_1} \oplus t^{F_2})(\eta) = t^{F_1}(\eta) \oplus t^{F_2}(\eta) : V^{F_1} \oplus V^{F_2} \rightarrow V^{F_1} \oplus V^{F_2}$ for any $\eta \in \mathcal{D}_m^r$.

FACT 8. *We have $(V^{F_1 \times_{B\mathcal{VB}_m} F_2}, H^{F_1 \times_{B\mathcal{VB}_m} F_2}, t^{F_1 \times_{B\mathcal{VB}_m} F_2}) = (V^{F_1} \oplus V^{F_2}, H^{F_1} \oplus H^{F_2}, t^{F_1} \oplus t^{F_2})$.*

Let $F_1, F_2 : \mathcal{VB}_m \rightarrow \mathcal{FM}$ be fiber product preserving gauge bundle functors of orders r_1 and r_2 (respectively). Then F_1 and F_2 are of order $r = r_1 + r_2$. Since F_2 has values in \mathcal{VB}_m (see Lemma 1), we have the composition $F_1 \circ F_2 : \mathcal{VB}_m \rightarrow \mathcal{FM}$ which is a fiber product preserving gauge bundle functor of order r . Let $(V^{F_1}, H^{F_1}, t^{F_1})$, $(V^{F_2}, H^{F_2}, t^{F_2})$ and $(V^{F_1 \circ F_2}, H^{F_1 \circ F_2}, t^{F_1 \circ F_2})$ be the admissible triples of order r and dimension m corresponding to F_1 , F_2 and $F_1 \circ F_2$ respectively. By tensoring over \mathbb{R} we have the admissible triple $(V^{F_1} \otimes V^{F_2}, H^{F_1} \otimes H^{F_2}, t^{F_1} \otimes t^{F_2})$ of order r and dimension m , where (of course) $(H^{F_1} \otimes H^{F_2})(\xi) = H^{F_1}(\xi) \otimes H^{F_2}(\xi) : V^{F_1} \otimes V^{F_2} \rightarrow V^{F_1} \otimes V^{F_2}$ for any $\xi \in G_m^r$ and $(t^{F_1} \otimes t^{F_2})(\eta) = t^{F_1}(\eta) \otimes t^{F_2}(\eta) : V^{F_1} \otimes V^{F_2} \rightarrow V^{F_1} \otimes V^{F_2}$ for any $\eta \in \mathcal{D}_m^r$.

OPEN PROBLEM. Express $(V^{F_1 \circ F_2}, H^{F_1 \circ F_2}, t^{F_1 \circ F_2})$ by $(V^{F_1}, H^{F_1}, t^{F_1})$ and $(V^{F_2}, H^{F_2}, t^{F_2})$. Is $(V^{F_1 \circ F_2}, H^{F_1 \circ F_2}, t^{F_1 \circ F_2})$ canonically isomorphic to $(V^{F_1} \otimes V^{F_2}, H^{F_1} \otimes H^{F_2}, t^{F_1} \otimes t^{F_2})$? In my opinion, it is not. Otherwise, there is a natural exchanging automorphism of $J^1 \circ J^1 : \mathcal{VB}_m \rightarrow \mathcal{FM}$. But this is rather impossible because an exchanging automorphism of $J^1 \circ J^1 : \mathcal{FM}_m \rightarrow \mathcal{FM}$ (over \mathcal{FM}_m) does not exist (see [5], [7]).

4. Classification of fiber product preserving gauge bundle functors on \mathcal{VB}_m of order r in terms of admissible triples of order r and dimension m . The following classification proposition shows that any fiber product preserving gauge bundle functor on \mathcal{VB}_m of order r is equivalent to some fiber product preserving gauge bundle functor as in Example 1.

PROPOSITION 1. *Let $F : \mathcal{VB}_m \rightarrow \mathcal{FM}$ be a fiber product preserving gauge bundle functor of order r . Let (V^F, H^F, t^F) be the admissible triple (of order r and dimension m) corresponding to F . Then we have a natural equivalence $\Theta^F : F \cong T^{(V^F, H^F, t^F)}$.*

Proof. Let $p : E \rightarrow M$ be a \mathcal{VB}_m -object. We construct canonically a diffeomorphism $\Theta^F : FE \rightarrow T^{(V^F, H^F, t^F)}E$ as follows. Given a point $y \in$

$F_x E$, $x \in M$, we define $\Theta^F(y) : J^r \mathcal{C}_x^{\infty, fl}(E) \rightarrow \tilde{V}_x^F M$ by

$$\Theta^F(y)(\xi) = F_x(f)(y) \in F_x(M \times \mathbb{R}) \cong \tilde{V}_x^F M, \quad \xi = j_x^r f \in J^r \mathcal{C}_x^{\infty, fl}(E),$$

where a fiber linear map $f : E \rightarrow \mathbb{R}$ is (in an obvious way) considered as the \mathcal{VB}_m -map $f : E \rightarrow M \times \mathbb{R}$ covering the identity of M and where the identification $F_x(M \times \mathbb{R}) \cong \tilde{V}_x^F M$ is given by $F_x(M \times \mathbb{R}) \ni F(\varphi \times \text{id}_{\mathbb{R}})(v) \cong \langle j_0^r \varphi, v \rangle \in \tilde{V}_x^F M$ for $v \in V^F = F_0(\mathbb{R}^m \times \mathbb{R})$, where $\varphi : \mathbb{R}^m \rightarrow M$ is an embedding with $\varphi(0) = x$. Then $\Theta^F(y)$ is well defined because F is of order r . Recalling the definition of (V^F, H^F, t^F) (see Example 2) and using the functoriality of F one can verify in a standard (but long) way that $\Theta^F(y)$ is a module homomorphism over $t_x^F : J_x^r(M, \mathbb{R}) \rightarrow \text{gl}(\tilde{V}_x^F M)$, i.e. $\Theta^F(y) \in T_x^{(V^F, H^F, t^F)} E$.

It remains to show that $\Theta^F : FE \rightarrow T^{(V^F, H^F, t^F)} E$ is a diffeomorphism.

Because $\Theta^F : F \rightarrow T^{(V^F, H^F, t^F)}$ is natural with respect to \mathcal{VB}_m -maps, and F and $T^{(V^F, H^F, t^F)}$ preserve fiber products, and E is locally a (multi) fiber product of $\mathbb{R}^m \times \mathbb{R}$, we may assume that $E = \mathbb{R}^m \times \mathbb{R}$, the trivial vector bundle over \mathbb{R}^m with fiber \mathbb{R} . But for $E = \mathbb{R}^m \times \mathbb{R}$ the transformation Θ^F is the composition $F(\mathbb{R}^m \times \mathbb{R}) \cong \mathbb{R}^m \times V^F \cong T^{(V^F, H^F, t^F)}(\mathbb{R}^m \times \mathbb{R})$, where the first identification is given by $F_x(\mathbb{R}^m \times \mathbb{R}) \ni v = (x, F(\tau_{-x} \times \text{id}_{\mathbb{R}})(v)) \in \{x\} \times V^F$, $x \in \mathbb{R}^m$, and the second trivialization is induced (see Example 1) by the obvious trivialization of $\mathbb{R}^m \times \mathbb{R}$. ■

5. Classification of admissible triples of order r and dimension m in terms of fiber product preserving gauge bundle functors on \mathcal{VB}_m of order r . The following classification proposition shows that any admissible triple of order r and dimension m is isomorphic to some admissible triple as in Example 2.

PROPOSITION 2. *Let (V, H, t) be an admissible triple of order r and dimension m . Let $F = T^{(V, H, t)}$. Then we have an isomorphism $\mathcal{O}^{(V, H, t)} : (V, H, t) \cong (V^F, H^F, t^F)$ of admissible triples.*

We recall that a morphism $(V_1, H_1, t_1) \rightarrow (V_2, H_2, t_2)$ of admissible triples is a linear map $\mathcal{O} : V_1 \rightarrow V_2$ such that $H_2(\xi) \circ \mathcal{O} = \mathcal{O} \circ H_1(\xi)$ for any $\xi \in G_m^r$ and $t_2(\eta) \circ \mathcal{O} = \mathcal{O} \circ t_1(\eta)$ for any $\eta \in \mathcal{D}_m^r$.

Proof. The composition $\mathcal{O}^{(V, H, t)} : V \rightarrow V^F = \text{Hom}_{t_0}(J^r \mathcal{C}_0^{\infty, fl}(\mathbb{R}^m \times \mathbb{R}), \tilde{V}_0 \mathbb{R}^m) \cong \{0\} \times V$ of $\mathcal{O}^{(V, H, t)}$ with the isomorphism induced (see Example 1) by the usual trivialization of $\mathbb{R}^m \times \mathbb{R}$ is the (almost) identity map. One can show in a standard (but long) way that $\mathcal{O}^{(V, H, t)}$ is a morphism $(V, H, t) \rightarrow (V^F, H^F, t^F)$ of admissible triples. ■

6. Natural transformations of fiber product preserving gauge bundle functors on \mathcal{VB}_m of order r and induced morphisms between admissible triples. Let $F_1, F_2 : \mathcal{VB}_m \rightarrow \mathcal{FM}$ be fiber product preserving gauge bundle functors of order r . Let $(V^{F_1}, H^{F_1}, t^{F_1})$ and $(V^{F_2}, H^{F_2}, t^{F_2})$ be the corresponding admissible triples of order r and dimension m . Let $\mu : F_1 \rightarrow F_2$ be a natural transformation.

EXAMPLE 3. Define $\nu^\mu : V^{F_1} \rightarrow V^{F_2}$ to be the restriction and corestriction of $\mu : F_1(\mathbb{R}^m \times \mathbb{R}) \rightarrow F_2(\mathbb{R}^m \times \mathbb{R})$ to $V^{F_1} = (F_1)_0(\mathbb{R}^m \times \mathbb{R})$ and $V^{F_2} = (F_2)_0(\mathbb{R}^m \times \mathbb{R})$. Then $\nu^\mu : (V^{F_1}, H^{F_1}, t^{F_1}) \rightarrow (V^{F_2}, H^{F_2}, t^{F_2})$ is a morphism of admissible triples. If μ is an isomorphism, then so is ν^μ .

DEFINITION 4. We call ν^μ the *morphism corresponding to μ* .

7. Morphisms between admissible triples and induced natural transformations between fiber product preserving gauge bundle functors. Let (V_1, H_1, t_1) and (V_2, H_2, t_2) be admissible triples of order r and dimension m . Let $\nu : (V_1, H_1, t_1) \rightarrow (V_2, H_2, t_2)$ be a morphism of admissible triples.

EXAMPLE 4. Given a \mathcal{VB}_m -object $p : E \rightarrow M$ define a base preserving fibered map $\mu^\nu : T^{(V_1, H_1, t_1)} E \rightarrow T^{(V_2, H_2, t_2)} E$ as follows. Let $\Phi \in T_x^{(V_1, H_1, t_1)} E = \text{Hom}_{(t_1)_x}(J^r C_x^{\infty, fl}(E), (\tilde{V}_1)_x M)$, $x \in M$. Put $\mu^\nu(\Phi) = \tilde{\nu}_x \circ \Phi : J^r C_x^{\infty, fl}(E) \rightarrow (\tilde{V}_2)_x M$, where $\tilde{\nu}_x : (\tilde{V}_1)_x M \rightarrow (\tilde{V}_2)_x M$, $\tilde{\nu}_x(\langle j_x^r \varphi, v \rangle) = \langle j_x^r \varphi, \nu(v) \rangle$ for $v \in V_1$ and $\varphi : \mathbb{R}^m \rightarrow M$ is an embedding with $\varphi(0) = x$. We see that $\mu^\nu(\Phi) \in \text{Hom}_{(t_2)_x}(J^r C_x^{\infty, fl}(E), (\tilde{V}_2)_x M) = T_x^{(V_2, H_2, t_2)} E$ and that $\mu^\nu : T^{(V_1, H_1, t_1)} \rightarrow T^{(V_2, H_2, t_2)}$ is a natural transformation. If ν is an isomorphism, then so is μ^ν .

DEFINITION 5. We call μ^ν the *natural transformation corresponding to ν* .

8. Object classification theorem. The first main result in this paper is the following theorem.

THEOREM 1. *The correspondence $F \mapsto (V^F, H^F, t^F)$ induces a bijective correspondence between the equivalence classes of fiber product preserving gauge bundle functors F on \mathcal{VB}_m of order r and the equivalence classes of admissible triples (V, H, t) of order r and dimension m . The inverse correspondence is induced by $(V, H, t) \mapsto T^{(V, H, t)}$.*

Proof. The correspondence $[F] \mapsto [(V^F, H^F, t^F)]$ is well defined, for if $\mu : F_1 \rightarrow F_2$ is an isomorphism, then so is $\nu^\mu : (V^{F_1}, H^{F_1}, t^{F_1}) \rightarrow (V^{F_2}, H^{F_2}, t^{F_2})$.

The correspondence $[(V, H, t)] \mapsto [T^{(V, H, t)}]$ is well defined, for if $\nu : (V_1, H_1, t_1) \rightarrow (V_2, H_2, t_2)$ is an isomorphism, then so is $\mu^\nu : T^{(V_1, H_1, t_1)} \rightarrow T^{(V_2, H_2, t_2)}$.

From Proposition 1 it follows that $[F] = [T^{(V^F, H^F, t^F)}]$. From Proposition 2 it follows that $[(V, H, t)] = [(V^F, H^F, t^F)]$ if $F = T^{(V, H, t)}$. ■

9. Morphism classification theorem. Let F_1 and F_2 be fiber product preserving gauge bundle functors on \mathcal{VB}_m of order r . Let $(V^{F_1}, H^{F_1}, t^{F_1})$ and $(V^{F_2}, H^{F_2}, t^{F_2})$ be the corresponding admissible triples of order r and dimension m .

LEMMA 2. Let $\nu : (V^{F_1}, H^{F_1}, t^{F_1}) \rightarrow (V^{F_2}, H^{F_2}, t^{F_2})$ be a morphism of admissible triples. Let $\mu^{[\nu]} : F_1 \rightarrow F_2$ be a natural transformation given by the composition

$$F_1 \xrightarrow{\Theta^{F_1}} T^{(V^{F_1}, H^{F_1}, t^{F_1})} \xrightarrow{\mu^\nu} T^{(V^{F_2}, H^{F_2}, t^{F_2})} \xrightarrow{(\Theta^{F_2})^{-1}} F_2,$$

where Θ^F is as in Proposition 1 and μ^ν is described in Example 4. Then $\mu = \mu^{[\nu]}$ is the unique natural transformation $F_1 \rightarrow F_2$ such that $\nu^\mu = \nu$, where ν^μ is as in Example 3.

Proof. Suppose $\bar{\mu} : F_1 \rightarrow F_2$ is another natural transformation such that $\nu^{\bar{\mu}} = \nu$. Then $\bar{\mu}$ coincides with μ on the vector bundle $\mathbb{R}^m \times \mathbb{R}$. Hence $\bar{\mu} = \mu$ by the same argument as in the proof of Proposition 1. ■

Now, the following second main result of this paper is clear.

THEOREM 2. Let F_1 and F_2 be two fiber product preserving gauge bundle functors on \mathcal{VB}_m of order r . The correspondence $\mu \mapsto \nu^\mu$ is a bijection between natural transformations $F_1 \rightarrow F_2$ and morphisms $(V^{F_1}, H^{F_1}, t^{F_1}) \rightarrow (V^{F_2}, H^{F_2}, t^{F_2})$ between corresponding admissible triples. The inverse correspondence is $\nu \mapsto \mu^{[\nu]}$.

10. Finite order theorem

THEOREM 3. Any fiber product preserving gauge bundle functor $F : \mathcal{VB}_m \rightarrow \mathcal{FM}$ is of finite order.

Proof. Define $A^F : \mathcal{C}^{\infty, fl}(\mathbb{R}^m \times \mathbb{R}) \rightarrow \mathcal{C}^\infty(F(\mathbb{R}^m \times \mathbb{R}), F(\mathbb{R}^m \times \mathbb{R}))$ by $A^F(f) = Ff$, where a fiber linear map $f : \mathbb{R}^m \times \mathbb{R} \rightarrow \mathbb{R}$ is considered as a base preserving \mathcal{VB}_m -map $\mathbb{R}^m \times \mathbb{R} \rightarrow \mathbb{R}^m \times \mathbb{R}$ in an obvious way. Clearly, A^F is π -local, where $\pi : F(\mathbb{R}^m \times \mathbb{R}) \rightarrow \mathbb{R}^m$ is the projection. The zero map $O \in \mathcal{C}^{\infty, fl}(\mathbb{R}^m \times \mathbb{R})$ is invariant with respect to the translations of \mathbb{R}^m . Moreover, given $f \in \mathcal{C}^{\infty, fl}(\mathbb{R}^m \times \mathbb{R})$, $v \in F_0(\mathbb{R}^m \times \mathbb{R})$, a neighborhood W of $j_0^\infty(0)$ and a neighborhood U of $0 \in F_0(\mathbb{R}^m \times \mathbb{R})$, there is $t \in \mathbb{R}_+$ such that $j_0^r(f \circ \lambda_t) \in W$ and $F\lambda_t(v) \in U$, where λ_t is the fiber homothety by t . Then

by the non-linear Peetre theorem [5] we deduce that the operator A^F is of finite order r_1 . In particular, F_0f depends on $j_0^{r_1}f$ only.

We have the bundle functor $G^F : \mathcal{M}f_m \rightarrow \mathcal{FM}$ such that $G^FM = F(M \times \mathbb{R})$ for any m -manifold M and $G^F\varphi = F(\varphi \times \text{id}_{\mathbb{R}}) : G^FM \rightarrow G^FN$ for any embedding $\varphi : M \rightarrow N$ between m -manifolds. By the Palais–Terng theorem (see [5]), G^F has finite order r_2 .

We prove that F is of order $r = \max(r_1, r_2)$. We consider a \mathcal{VB}_m -map $f : E_1 \rightarrow E_2$ and a point $x \in M_1$. It remains to show that $F_x f$ depends on $j_x^r f$.

Using \mathcal{VB}_m -trivializations we can assume that $E_1 = \mathbb{R}^m \times \mathbb{R}^n$, $E_2 = \mathbb{R}^m \times \mathbb{R}^q$ and $x = 0 \in \mathbb{R}^m$. Since F preserves fiber products, we can assume that $q = 1$. For the same reason, we can assume that $n = 1$. (Apply the fact that the map $I : \times_M^n F(\mathbb{R}^m \times \mathbb{R}) \rightarrow F(\mathbb{R}^m \times \mathbb{R}^n)$ inverse to the fiber product identification $F(\mathbb{R}^m \times \mathbb{R}^n) \rightarrow \times_M^n F(\mathbb{R}^m \times \mathbb{R})$ is given by $I(v_1, \dots, v_n) = \sum_j F(g^j)(v^j)$ for $v_1, \dots, v_n \in F_x(\mathbb{R}^m \times \mathbb{R})$, $x \in \mathbb{R}^m$, $g^j : \mathbb{R}^m \times \mathbb{R} \rightarrow \mathbb{R}^m \times \mathbb{R}^n$, $g^j(x, y) = (x, e_j y)$, $e_j = (0, \dots, 0, 1, 0, \dots, 0)$ (1 in position j), $j = 1, \dots, n$, where the sum is the one in the vector bundle $F(\mathbb{R}^m \times \mathbb{R}^n)$; see Lemma 1.) Then we can write $f = \tilde{f} \circ (\varphi \times \text{id}_{\mathbb{R}})$, where $\tilde{f} : \mathbb{R}^m \times \mathbb{R} \rightarrow \mathbb{R}^m \times \mathbb{R}$ is a base preserving \mathcal{VB}_m -map and $\varphi : \mathbb{R}^m \rightarrow \mathbb{R}^m$ is a 0-preserving embedding. Then $F_0f = F_0\tilde{f} \circ G_0^F\varphi$ depends on $j_0^{r_1}\tilde{f}$ and $j_0^{r_2}\varphi$ as we showed above. ■

11. Applications. We give some applications of the main results. We will use the Facts from Section 3.

LEMMA 3. *Let $a : \mathcal{D}_m^r \rightarrow \mathcal{D}_m^r$ be a G_m^r -equivariant linear map, where \mathcal{D}_m^r is a G_m^r -space via H_m^r as in Fact 1. Then there exist unique $k, l \in \mathbb{R}$ such that $a = k \text{id}_{\mathcal{D}_m^r} + l\varepsilon_m^r$, where $\text{id}_{\mathcal{D}_m^r}$ is the identity map and ε_m^r is the map as in Fact 2.*

Proof. Below $\alpha, \beta \in (\mathbb{N} \cup \{0\})^m$ with $|\alpha| \leq r$ and $|\beta| \leq r$. We can write $a(j_0^r x^\alpha) = \sum_\beta a_\beta^\alpha j_0^r x^\beta$ for some unique real numbers a_β^α . By the equivariance of a with respect to $j_0^r(\tau^1 x^1, \dots, \tau^m x^m) \in G_m^r$ we deduce that $\tau^\alpha a_\beta^\alpha = \tau^\beta a_\beta^\alpha$ for any $\tau = (\tau^1, \dots, \tau^m) \in \mathbb{R}_+^m$. Then $a(j_0^r x^\alpha) = a_\alpha^\alpha j_0^r x^\alpha$. If $\alpha \neq (0)$, then by the equivariance of a with respect to $j_0^r(x^1 + x^\alpha, x^2, \dots, x^m) \in G_m^r$ we deduce that $a_{\varepsilon_1}^{e_1} j_0^r x^1 + a_\alpha^\alpha j_0^r x^\alpha = a(j_0^r(x^1 + x^\alpha)) = a_{\varepsilon_1}^{e_1}(j_0^r x^1 + j_0^r x^\alpha)$, i.e. $a(j_0^r x^\alpha) = a_{\varepsilon_1}^{e_1} j_0^r x^\alpha$. So, it remains to put $k = a_{\varepsilon_1}^{e_1}$ and $l = a_{(0)}^{(0)} - a_{\varepsilon_1}^{e_1}$. ■

LEMMA 4. *Suppose that $a : \mathcal{D}_m^r \rightarrow \mathcal{D}_m^r$ is a G_m^r -equivariant linear map such that $t_m^r(\eta) \circ a = a \circ t_m^r(\eta)$ for any $\eta = j_0^r \gamma \in \mathcal{D}_m^r$, where t_m^r is as in Fact 1. Then there exists a unique $k \in \mathbb{R}$ such that $a = k \text{id}_{\mathcal{D}_m^r}$.*

Proof. By Lemma 3, $a = k \text{id}_{\mathcal{D}_m^r} + l\varepsilon_m^r$. Since a commutes with t_m^r , we easily obtain $l = 0$. ■

LEMMA 5. *The space of endomorphisms of (V, H, ε^V) (see Fact 5) is the space of G_m^r -equivariant linear endomorphisms of V , where V is a G_m^r -space via H .*

Proof. This is a simple observation. ■

COROLLARY 1. *Any natural endomorphism $\mu : J^r \rightarrow J^r$ is $k \operatorname{id}_{J^r}$ for some uniquely determined real number k .*

Proof. By Theorem 2 natural endomorphisms $J^r \rightarrow J^r$ are in bijection with endomorphisms of the admissible triple $(\mathcal{D}_m^r, H_m^r, t_m^r)$ corresponding to J^r (see Fact 1). By Lemma 4 these endomorphisms are $k \operatorname{id}_{\mathcal{D}_m^r}$ for $k \in \mathbb{R}$. ■

COROLLARY 2. *Any natural endomorphism $\mu : J_v^r \rightarrow J_v^r$ is $k \operatorname{id}_{J_v^r} + l \mu_m^r$ for some uniquely determined real numbers k, l , where for any \mathcal{VB}_m -object $p : E \rightarrow M$ we have $\mu_m^r : J_v^r E \rightarrow J_v^r E$ with $\mu(j_x^r \gamma) = j_x^r(\gamma(x))$ for $\gamma : M \rightarrow E_x$, $x \in M$, where $\gamma(x) : M \rightarrow E_x$ is the constant map.*

Proof. The proof is quite similar to that of Corollary 1. We use Theorem 2, Lemma 3 and Fact 2. ■

COROLLARY 3. *Any natural endomorphism $\mu : \tilde{V}E \rightarrow \tilde{V}E$ for any \mathcal{VB}_m -object $p : E \rightarrow M$, where \tilde{V} is as in Fact 5, is $\mu = \operatorname{id}_E \otimes \bar{\mu}_M$ for some natural endomorphism $\bar{\mu} : \tilde{V}M \rightarrow \tilde{V}M$.*

In particular, any natural endomorphism $\mu : V^A E = E \otimes A \rightarrow V^A E$, where V^A is the vertical Weil gauge bundle functor corresponding to a Weil algebra A , is $\mu = \operatorname{id}_E \otimes B$ for some $B \in \mathfrak{gl}(A)$.

Proof. The proof is quite similar to that of Corollary 1. We use Theorem 2, Lemma 5 and Fact 5. That $V^A E = E \otimes A$ follows from the fact that the admissible triples corresponding to the functors are isomorphic. ■

References

- [1] A. Cabras and I. Kolář, *Prolongations of projectable tangent valued forms*, Arch. Math. (Brno) 38 (2002), 243–257.
- [2] M. Doupovec and I. Kolář, *Iteration of fiber product preserving bundle functors*, Monatsh. Math. 134 (2001), 39–50.
- [3] I. Kolář, *Bundle functors of the jet type*, in: Differential Geometry and Applications (Brno, 1998), Masaryk Univ., Brno, 1999, 231–237.
- [4] —, *On the geometry of fiber product preserving bundle functors*, in: Differential Geometry and its Applications (Opava, 2001), Math. Publ. 3, Silesian Univ. Opava, Opava, 2001, 85–92.
- [5] I. Kolář, P. Michor and J. Slovák, *Natural Operations in Differential Geometry*, Springer, 1993.
- [6] I. Kolář and W. M. Mikulski, *On the fiber product preserving bundle functors*, Differential Geom. Appl. 11 (1999), 105–115.

- [7] I. Kolář and M. Modugno, *Natural maps on the iterated jet prolongation of a fibered manifold*, Ann. Mat. Pura Appl. (4) 158 (1991), 151–165.
- [8] W. M. Mikulski, *The natural operators lifting projectable vector fields to some fiber product preserving bundles*, Ann. Polon. Math. 81 (2003), 261–271.

Institute of Mathematics
Jagiellonian University
Reymonta 4
30-059 Kraków, Poland
E-mail: mikulski@im.uj.edu.pl

Reçu par la Rédaction le 11.2.2003
Révisé le 2.4.2003

(1428)