

Topologically Invertible Elements and Topological Spectrum

by

Mati ABEL and Wiesław ŻELAZKO

Presented by Stanisław KWAPIEŃ

Summary. Properties of topologically invertible elements and the topological spectrum of elements in unital semitopological algebras are studied. It is shown that the inversion $x \mapsto x^{-1}$ is continuous in every invertive Fréchet algebra, and singly generated unital semitopological algebras have continuous characters if and only if the topological spectrum of the generator is non-empty. Several open problems are presented.

1. Introduction

1.1. A *semitopological algebra* is a real or complex linear topological space with an associative separately continuous multiplication. An element x in a semitopological algebra A with unit element e_A is said to be *left* (respectively, *right*) *topologically invertible* if $e_A \in \overline{Ax}$ (respectively, $e_A \in \overline{xA}$)⁽¹⁾, or equivalently, if there is a net $(z_\alpha)_{\alpha \in A}$ in A such that $\lim_\alpha z_\alpha x = e_A$ (respectively, $\lim_\alpha x z_\alpha = e_A$). An element x in A is *topologically invertible* if it is left and right topologically invertible. The set of all left (respectively, right) invertible elements in A will be denoted by $G_l(A)$ (respectively, $G_r(A)$), and the set of all topologically left (respectively, right) invertible elements in A by $G_l^t(A)$ (respectively, $G_r^t(A)$). Hence, the set $G(A)$ of all invertible elements equals $G_l(A) \cap G_r(A)$, and the set $G^t(A)$ of all topologically invertible elements equals $G_l^t(A) \cap G_r^t(A)$. It is easy to see that $G(A) \subset G^t(A)$. In

2000 *Mathematics Subject Classification*: Primary 46H05; Secondary 46H20.

Key words and phrases: topologically invertible elements, topological spectrum, spectral mapping property, discontinuity of inversion, functional topological spectrum.

Research supported in part by Estonian Science Foundation grant 6205.

⁽¹⁾ Here and later on \overline{U} denotes the closure of U in A .

particular, when $G(A) = G^t(A)$, A is called an *invertive algebra* ⁽²⁾ (see [2, p. 14]); and a topologically invertible element is said to be *proper* (see [20, p. 323]) if it is non-invertible. Properties of topologically invertible elements have been studied in several papers, for example, in [2], [6]–[9], [11], [13]–[16] and [18]–[20].

1.2. Let A be a semitopological algebra, and $m(A)$ the set of all closed two-sided ideals in A which are maximal as left or right ideals in A . A semitopological algebra A over \mathbb{K} is called a *Gelfand–Mazur algebra* if A/M (in the quotient topology) is topologically isomorphic to \mathbb{K} for each $M \in m(A)$, and a *simplicial algebra* if every closed regular left (respectively, right) ideal of A is contained in some closed maximal left (respectively, right) ideal of A . Main classes of Gelfand–Mazur algebras have been described in [1] or [4]. It is known (see [5, Corollary 5] ⁽³⁾) that every commutative unital locally m -pseudoconvex Hausdorff algebra is simplicial.

1.3. Let A be a semitopological algebra over \mathbb{K} with unit element e_A , $\mathfrak{M}(A)$ the set of all continuous \mathbb{K} -valued characters (non-zero multiplicative linear functionals) on A ,

$$\sigma_A(x) = \{\lambda \in \mathbb{K} : x - \lambda e_A \notin G(A)\}$$

the (algebraic) *spectrum* of $x \in A$, and

$$\varrho_A(x) = \sup\{|\lambda| : \lambda \in \sigma_A(x)\}$$

the (algebraic) *spectral radius* of x . The (algebraic) spectrum $\sigma_A(x)$ and the spectral radius $\varrho_A(x)$ of x play an important role in invertive algebras. For non-invertive algebras a similar role is played by the *topological spectrum*

$$\sigma_A^t(x) = \{\lambda \in \mathbb{K} : x - \lambda e_A \notin G^t(A)\}$$

and the *topological spectral radius*

$$\varrho^t(x) = \sup\{|\lambda| : \lambda \in \sigma_A^t(x)\}.$$

Similarly to the spectral radius, we set $\varrho_A^t(x) = 0$ if $\sigma_A^t(x) = \emptyset$, and $\varrho_A^t(x) = \infty$ if $\sigma_A^t(x)$ is unbounded. It is easy to see that $\sigma_A^t(x) \subset \sigma_A(x)$ and $\varrho_A^t(x) \leq \varrho_A(x)$ for each $x \in A$. Moreover, A is invertive if and only if $\sigma_A^t(x) = \sigma_A(x)$ for each $a \notin G(A)$. Indeed, if A is invertive, then $\sigma_A^t(x) = \sigma_A(x)$ for each $a \in A$. Conversely, let $a \in A \setminus G(A)$. Then $0 \in \sigma_A(a)$. If now $\sigma_A(a) = \sigma_A^t(a)$, then $a \notin G^t(A)$. Hence, $G(A) = G^t(A)$, and so A is invertive.

⁽²⁾ It is known (see [2, Corollary 2]) that every complete unital locally m -pseudoconvex algebra is an invertive algebra, but any commutative unital F -algebra with a discontinuous inverse is not (see [19, Proposition 4]).

⁽³⁾ For complete algebras see [3, Proposition 2] or [10, Corollary 7.1.14], and for locally m -convex Hausdorff algebras see [12, pp. 321–322].

Let $x \in A$, $\varphi \in \mathfrak{M}(A)$ and $x - \varphi(x)e_A \in G^t(A)$. Then there are nets $(x_\alpha)_{\alpha \in A}$ and $(y_\mu)_{\mu \in \mathcal{M}}$ in A such that both $((x - \varphi(x)e_A)x_\alpha)_{\alpha \in A}$ and $(y_\mu(x - \varphi(x)e_A))_{\mu \in \mathcal{M}}$ converge to e_A . Since φ is continuous, the nets $(\varphi(x - \varphi(x)e_A)\varphi(x_\alpha))_{\alpha \in A}$ and $(\varphi(x - \varphi(x)e_A)\varphi(y_\mu))_{\mu \in \mathcal{M}}$ then converge to 1, but this is not possible, because $\varphi(x - \varphi(x)e_A) = 0$. Hence, $x - \varphi(x)e_A \notin G^t(A)$ for each $\varphi \in \mathfrak{M}(A)$. Consequently,

$$\{\varphi(x) : \varphi \in \mathfrak{M}(A)\} \subset \sigma_A^t(x)$$

for each $x \in A$. In particular, if $\mathfrak{M}(A)$ is not empty and

$$\sigma_A^t(x) = \{\varphi(x) : \varphi \in \mathfrak{M}(A)\}$$

for each $x \in A$, we will say that x has *functional topological spectrum*.

1.4. In this paper, we will study the properties of topologically invertible elements and of the topological spectrum of elements in unital semitopological algebras. Among other results, it is shown that (a) the set of all topologically invertible elements in any unital Fréchet (that is, complete and metrizable) algebra is a G_δ -set; (b) if the inversion in a unital Fréchet algebra A is discontinuous, then A has left and right dense ideals; (c) if A is a commutative complex unital semitopological algebra, then the topological spectrum has the spectral mapping property; (d) every element of a commutative unital simplicial Gelfand–Mazur algebra has functional topological spectrum; and (e) if every element in a unital semitopological algebra A has functional topological spectrum, then the topological spectral radius is a submultiplicative seminorm on A . Several open problems are presented at the end of paper.

2. Properties of $G^t(A)$ for a Fréchet algebra A . Let A be a unital (real or complex) non-commutative Fréchet algebra with F -norm $\|\cdot\|$, i.e. with a function $x \mapsto \|x\|$ on A such that

- (i) $\|x\| \geq 0$ for each $x \in A$, and $\|x\| = 0$ if and only if $x = \theta_A$ ⁽⁴⁾;
- (ii) $\|x + y\| \leq \|x\| + \|y\|$ for all $x, y \in A$;
- (iii) $(\lambda, x) \mapsto \|\lambda x\|$ is a jointly continuous map from $\mathbb{K} \times A$ to \mathbb{R}_+ .

Let $\|e_A\| = 1$, and

$$g_l(x) = \inf_{u \in A} \|ux - e_A\| \quad \text{and} \quad g_r(x) = \inf_{u \in A} \|xu - e_A\|$$

for each $x \in A$. Then

$$G_l^t(A) = \{x \in A : g_l(x) = 0\} \quad \text{and} \quad G_r^t(A) = \{x \in A : g_r(x) = 0\}.$$

⁽⁴⁾ Here and below, θ_A denotes the null element of A .

LEMMA 1. *Let A be a unital non-commutative Fréchet algebra. Then the sets*

$$S_{l,\lambda} = \{x \in A : g_l(x) < \lambda\} \quad \text{and} \quad S_{r,\lambda} = \{x \in A : g_r(x) < \lambda\}$$

are open in A for any $\lambda > 0$.

Proof. The proof is similar to the proof of Lemma 1 in [19]. ■

COROLLARY 1. *The function g_l is continuous at all points of $G_l^t(A)$, and g_r is continuous at all points of $G_r^t(A)$.*

It is known (see [19, Proposition 3]) that $G^t(A)$ is a G_δ -set for any commutative unital F -algebra A . In the non-commutative case, we have

PROPOSITION 1. *Let A be a non-commutative unital F -algebra. Then $G_l^t(A)$, $G_r^t(A)$ and $G^t(A)$ are G_δ -sets.*

Proof. Since

$$G_l^t(A) = \bigcap_{n \in \mathbb{N}} \{x \in A : g_l(x) < 1/n\}, \quad G_r^t(A) = \bigcap_{n \in \mathbb{N}} \{x \in A : g_r(x) < 1/n\},$$

$G_l^t(A)$ and $G_r^t(A)$ are G_δ -sets by Lemma 1. Therefore so is $G^t(A)$ as the intersection of two G_δ -sets. ■

COROLLARY 2. *Let A be a unital F -algebra. If the inversion $x \mapsto x^{-1}$ is discontinuous in A , then $G^t(A) \setminus G(A) \neq \emptyset$.*

Proof. It is known (see, for example, [17, Theorem 1.6]) that the inversion is continuous in A if and only if $G(A)$ is a G_δ -set and $G(A) \subset G^t(A)$. If the inversion is discontinuous, then $G^t(A) \setminus G(A) \neq \emptyset$ by Proposition 1. ■

COROLLARY 3. *In every invertive Fréchet algebra the inversion is continuous.*

COROLLARY 4. *If the inversion is discontinuous in a unital F -algebra A , then A has a dense left and a dense right ideal.*

Proof. By Corollary 2 there is an element

$$x \in (G_l^t(A) \cap G_r^t(A)) \setminus G(A).$$

Suppose that $x \in G_l(A)$. Then there exists $a \in A$ such that $ax = e_A$. Since $x \in G_r^t(A)$ as well, there is a sequence $(x_n) \subset A$ such that (xx_n) converges to e_A . This implies that (x_n) converges to a and $xa = e_A$. Thus $x \in G(A)$, which is not the case. Similarly, from $x \in G_r(A)$ it follows that $x \in G(A)$, which is again impossible. Consequently,

$$x \in (G_l^t(A) \setminus G_l(A)) \cap (G_r^t(A) \setminus G_r(A)).$$

This means that the left ideal Ax and the right ideal xA are dense in A . ■

3. Singly generated topological algebras. We say that a semitopological algebra A is *singly generated by an element* $x_0 \in A$ (the generator) if the set of all polynomials in x_0 (with real coefficients in the case of a real algebra) is dense in A . We now show how the topological spectrum can be used to decide when a unital singly generated semitopological algebra A has continuous characters ⁽⁵⁾. In this case A is commutative.

PROPOSITION 2. *Let A be a unital singly generated (real or complex) semitopological algebra and x_0 the generator of A . Then there is a one-to-one correspondence between the elements in $\mathfrak{M}(A)$ and the scalars in $\sigma_A^t(x_0)$.*

Proof. Suppose that $\sigma_A^t(x_0) \neq \emptyset$ and fix $\lambda \in \sigma_A^t(x_0)$. Let A_0 be the subalgebra of A consisting of all polynomials p in x_0 (then A_0 is a dense subalgebra of A) and let

$$(a) \quad f_\lambda(p(x_0)) = p(\lambda).$$

We show that f_λ is a character on A_0 . To this end, we first have to show that f_λ is well defined, i.e. if $p_1(x_0) = p_2(x_0)$, then $p_1(\lambda) = p_2(\lambda)$, or that $p(x) = 0$ implies $p(\lambda) = 0$. If this is not so, i.e. if $p(x_0) = 0$ but $p(\lambda) \neq 0$, then the relation

$$p(x_0) - p(\lambda)e_A = (x_0 - \lambda e_A)q(x_0, \lambda)$$

for some polynomial q in two variables implies that

$$e_A = -[p(\lambda)]^{-1}(x_0 - \lambda e_A)q(x_0, \lambda),$$

i.e. λ is not in the (algebraic) spectrum $\sigma_A(x_0)$. So, λ cannot be in the (smaller) spectrum $\sigma_A^t(x_0)$, which gives a contradiction. Hence, the formula (a) defines a multiplicative linear functional on A_0 .

We now show that f_λ is continuous. Indeed, if not, then $f_\lambda^{-1}(0)$ is dense in A_0 , and so there is a net $(p_\alpha)_{\alpha \in A} \subset f_\lambda^{-1}(0)$ such that $(p_\alpha(x_0))_{\alpha \in A}$ tends to e_A . Since

$$p_\alpha(x_0) = (x_0 - \lambda e_A)q_\alpha(x_0, \lambda) = r_\alpha(x_0, \lambda)(x_0 - \lambda e_A)$$

for some polynomials q_α and r_α in two variables, $x_0 - \lambda e_A$ is topologically invertible, which is a contradiction, since $\lambda \in \sigma_A^t(x_0)$. Thus, f_λ is a continuous character on A_0 and extends by continuity to an element of $\mathfrak{M}(A)$, which we also denote by f_λ . Then

$$(b) \quad f_\lambda(x_0) = \lambda,$$

so that to different scalars λ correspond different functionals in $\mathfrak{M}(A)$.

Conversely, assume that $\mathfrak{M}(A) \neq \emptyset$, let $f \in \mathfrak{M}(A)$, and put $\lambda = f(x_0)$. Then $x = x_0 - \lambda e_A \in f^{-1}(0)$ and x is not topologically invertible, as other-

⁽⁵⁾ This result has been extended by the second author to finitely generated commutative topological algebras (see [21]).

wise there is a net $(u_\alpha)_{\alpha \in \mathcal{A}}$ with $(u_\alpha x)_{\alpha \in \mathcal{A}} \rightarrow e_A$, which is impossible (because then $0 = f(u_\alpha x) \rightarrow f(e_A) = 1$ for each $\alpha \in \mathcal{A}$). Thus $\lambda \in \sigma_A^t(x_0)$ and the conclusion follows (the announced one-to-one correspondence is given by (b)). ■

Thus we have a situation somewhat similar to that of singly generated (unital) Banach algebras, where the spectrum of the generator can be identified with the maximal ideal space (equipped with the Gelfand topology) both as a set and as a compact space. The example below shows that the two sets may not be homeomorphic in general.

EXAMPLE. Let A be the Cartesian product of continuum many copies of the field of scalars, provided with the coordinatewise algebra operations and the product topology. We can identify A with the algebra of all scalar-valued functions on the complex plane \mathbb{C} with the topology of pointwise convergence. This topology is given by the seminorms $\|x\|_S = \max_{\zeta \in S} |x(\zeta)|$ for each $x \in A$, where S runs over the family \mathcal{F} of all finite subsets of \mathbb{C} . Clearly, A is a complete multiplicatively convex algebra, and all its multiplicative linear functionals are continuous and are of the form $f_\zeta(x) = x(\zeta)$, where $\zeta \in \mathbb{C}$ (see, e.g., [14]). The algebra A is singly generated by the function x_0 defined by $x_0(\zeta) = \zeta$. In fact, for any x in A and S in \mathcal{F} , there is a polynomial p with scalar coefficients such that $p(x)(\zeta) = x(\zeta)$ for all ζ in S . Thus $\|p(x) - x\|_S = 0$ and x can be approximated by polynomials of x . We can identify $\mathfrak{M}(A)$ with \mathbb{C} as a set, but, as shown below, not as a topological space, because the Gelfand topology of $\mathfrak{M}(A)$ is discrete. To show this, consider the element x_1 of A given by

$$x_1(\zeta) = \begin{cases} 1 & \text{if } \zeta = 0, \\ 0 & \text{otherwise,} \end{cases}$$

and the character $f_0(x) = x(0)$. Its neighbourhood

$$U(f; x_1, 1/2) = \{f \in \mathfrak{M}(A) : |f(x_1) - f_0(x_1)| < 1/2\}$$

in the Gelfand topology consists of f_0 only, and our claim follows.

As a corollary to Proposition 2 we obtain the following

PROPOSITION 3. *A singly generated unital semitopological (real or complex) algebra has a continuous character if and only if the topological spectrum of its generator is non-empty.*

REMARK. The above result fails to be true if we replace the topological spectrum by the algebraic spectrum. The B_0 -algebra $L^\omega[0, 1]$ of Arens (see [8] or [20, Example 10.5]), equal to $\bigcap_{p=1}^\infty L_p[0, 1]$, is singly generated ⁽⁶⁾ by the function x_0 , where $x_0(t) = t$ for each $t \in \mathbb{C}$ (that is, the subalgebra of

⁽⁶⁾ All algebraic operations in $L^\omega[0, 1]$ are defined pointwise.

all polynomials in x_0 is dense in $L^\omega[0, 1]$). Then $\sigma_{L^\omega[0,1]}(x_0) = [0, 1] \neq \emptyset$. On the other hand (see [20, Subsection 14.2]), $L^\omega[0, 1]$ has no characters and so no continuous characters. It can be easily seen that the translates $x_0 - \lambda e$ with $0 \leq \lambda \leq 1$ are not invertible and not topologically invertible. Therefore, $\sigma_{L^\omega[0,1]}^t(x_0) = \emptyset$ (this also follows from Proposition 3).

4. Topological spectrum

4.1. Properties of the topological spectrum. Let A be a unital semitopological algebra. The topological spectrum of elements of A has several properties similar to the (algebraic) spectrum. It is easy to see that $\sigma_A^t(\theta_A) = \{0\}$, $\sigma_A^t(e_A) = \{1\}$ and $\sigma_A^t(e) = \{0, 1\}$ for any idempotent e of A with $e \neq \theta_A$ and $e \neq e_A$. Moreover, we have

PROPOSITION 4. *Let A be a unital semitopological algebra. For any $x, y \in A$ with $\sigma_A^t(x) \neq \emptyset$ and $\sigma_A^t(xy) \neq \emptyset$ and $\mu \in \mathbb{K}$ we have*

- (a) $\sigma_A^t(\mu x) = \mu \sigma_A^t(x)$;
- (b) $\sigma_A^t(\mu e_A + x) = \mu + \sigma_A^t(x)$;
- (c) if $x \in G(A)$, then $\sigma_A^t(x^{-1}) = \{\lambda^{-1} : \lambda \in \sigma_A^t(x)\}$;
- (d) $\sigma_A^t(xy) \cup \{0\} = \sigma_A^t(yx) \cup \{0\}$.

Proof. Since $\mu x - \lambda e_A = \mu(x - \mu^{-1}\lambda e_A)$ for all $\mu \neq 0$ and $\lambda \in \mathbb{K}$, it follows that $\lambda \in \sigma_A^t(\mu x)$ if and only if $\mu^{-1}\lambda \in \sigma_A^t(x)$. Therefore

$$\sigma_A^t(\mu x) \subset \mu \sigma_A^t(x) = \mu \sigma_A^t(\mu^{-1}(\mu x)) \subset \sigma_A^t(\mu x)$$

for $\mu \neq 0$. Thus, we have (a) for $\mu \neq 0$ (if $\mu = 0$, then (a) also holds). Similarly to (a), the equality (b) follows from the identity

$$(\mu e_A + x) - \lambda e_A = x - (\lambda - \mu)e_A,$$

and (c) follows from

$$x^{-1} - \lambda^{-1}e_A = (-\lambda^{-1}x^{-1})(x - \lambda e_A) = (x - \lambda e_A)(-\lambda^{-1}x^{-1}),$$

because $(-\lambda^{-1}x^{-1})(x - \lambda e_A) = (x - \lambda e_A)(-\lambda^{-1}x^{-1})$ for each $x \in G(A)$.

To prove (d), we first show that $e_A - xy \in G^t(A)$ if and only if $e_A - yx \in G^t(A)$. Let $e_A - xy \in G^t(A)$. Then there exist nets $(u_\alpha)_{\alpha \in A}$ and $(v_\beta)_{\beta \in B}$ in A such that $((e_A - xy)u_\alpha)_{\alpha \in A}$ and $(v_\beta(e_A - xy))_{\beta \in B}$ converge to e_A . Since

$$(yux + e_A)(e_A - yx) = y[u(e_A - xy)]x + e_A - yx$$

and

$$(e_A - yx)(yux + e_A) = y[(e_A - xy)u]x + e_A - yx$$

for all $x, y, u \in A$ and the multiplication in A is separately continuous, the nets $((yv_\beta x + e_A)(e_A - yx))_{\beta \in B}$ and $((e_A - yx)(yu_\alpha + e_A))_{\alpha \in A}$ converge to e_A . Hence, $e_A - yx \in G^t(A)$. By symmetry, $e_A - yx \in G^t(A)$ implies $e_A - xy \in G^t(A)$.

Let now $\lambda \in \sigma_A^t(xy)$. Then $xy - \lambda e_A \notin G^t(A)$. Since

$$xy - \lambda e_A = -\lambda(e_A - (\lambda^{-1}x)y)$$

for $\lambda \neq 0$, we have $e_A - (\lambda^{-1}x)y \notin G^t(A)$. Hence, $\lambda \in \sigma_A^t(yx)$. Thus

$$\sigma_A^t(xy) \cup \{0\} \subset \sigma_A^t(yx) \cup \{0\}.$$

Interchanging x and y , we get the reverse inclusion. ■

LEMMA 2. *Let A and B be unital semitopological algebras and π a continuous homomorphism from A into B such that $\pi(e_A) = e_B$. Then*

$$\sigma_B^t(\pi(x)) \subset \sigma_A^t(x)$$

for all $x \in A$.

Proof. Let $\lambda \in \sigma_B^t(\pi(x))$. Then $\pi(x - \lambda e_A) = \pi(x) - \lambda e_B \notin G^t(B)$. Therefore $x - \lambda e_A \notin G^t(A)$. Thus $\lambda \in \sigma_A^t(x)$. ■

PROPOSITION 5. *Let A and B be unital semitopological algebras and π a continuous open homomorphism from A onto B . If there exists a neighbourhood O of zero in A such that $e_A + O + \ker \pi \subset G^t(A)$, then*

$$\sigma_B^t(\pi(x)) = \sigma_A^t(x)$$

for each $x \in A$.

Proof. By Lemma 2, the inclusion $\sigma_B^t(\pi(x)) \subset \sigma_A^t(x)$ holds. To prove the opposite inclusion, assume that $\lambda \notin \sigma_B^t(\pi(x))$ and O is as in the statement. Then $\pi(x) - \lambda e_B \in G^t(B)$. Therefore there exist nets $(x_\mu)_{\mu \in M}$ and $(y_\nu)_{\nu \in V}$ in A such that the nets $((\pi(x) - \lambda e_B)\pi(x_\mu))_{\mu \in M}$ and $(\pi(y_\nu)(\pi(x) - \lambda e_B))_{\nu \in V}$ converge to e_B in B . Since π is open, $\pi(O)$ is a neighbourhood of zero in B . Hence there exist $\mu_0 \in M$ and $\nu_0 \in V$ such that

$$(x - \lambda e_A)x_\mu - e_A \in \pi^{-1}(\pi(O)) = O + \ker \pi$$

whenever $\mu > \mu_0$, and

$$y_\nu(x - \lambda e_A) - e_A \in \pi^{-1}(\pi(O)) = O + \ker \pi$$

whenever $\nu > \nu_0$. This means that

$$(x - \lambda e_A)x_\mu \in e_A + O + \ker \pi \subset G^t(A)$$

whenever $\mu > \mu_0$ and

$$y_\nu(x - \lambda e_A) \in e_A + O + \ker \pi \subset G^t(A)$$

whenever $\nu > \nu_0$.

Now we fix $\mu_1 > \mu_0$ and $\nu_1 > \nu_0$. Then there exist nets $(u_\alpha)_{\alpha \in A}$ and $(v_\beta)_{\beta \in B}$ in A such that $((x - \lambda e_A)(x_{\mu_1}u_\alpha))_{\alpha \in A}$ and $((v_{\beta}y_{\nu_1})(x - \lambda e_A))_{\beta \in B}$ converge to e_A in A . Consequently, $x - \lambda e_A \in G^t(A)$, i.e. $\lambda \notin \sigma_A^t(x)$. Thus, $\sigma_A^t(x) \subset \sigma_B^t(\pi(x))$ holds as well. ■

COROLLARY 5. Let A be a unital semitopological algebra, I a two-sided ideal of A and π the canonical homomorphism from A onto A/I . If there exists a neighbourhood O of zero in A such that $e_A + O + I \subset G^t(A)$, then

$$\sigma_{A/I}^t(\pi(x)) = \sigma_A^t(x)$$

for each $x \in A$.

COROLLARY 6. Let A and B be unital semitopological algebras and π a homomorphism and homeomorphism from A into B such that $\pi(e_A) = e_B$. If

(a) $G^t(A)$ is open in A

or

(b) $G^t(\pi(A)) = G^t(B) \cap \pi(A)$,

then

$$\sigma_B^t(\pi(x)) = \sigma_A^t(x)$$

for each $x \in A$.

Proof. The inclusion $\sigma_B^t(\pi(x)) \subset \sigma_A^t(x)$ holds for each $x \in A$, by Lemma 2. If A satisfies condition (a) and π is one-to-one, then the opposite inclusion also holds for each $x \in A$, by Proposition 5. Let now A and π satisfy condition (b). Let $x \in A$ and $\lambda \notin \sigma_B^t(\pi(x))$. Then $\pi(x - \lambda e_A) \in G^t(\pi(A))$ by (b). Therefore there exist nets $(x_\mu)_{\mu \in M}$ and $(y_\nu)_{\nu \in V}$ in A such that both $(\pi[(x - \lambda e_A)x_\mu])_{\mu \in M}$ and $(\pi[y_\nu(x - \lambda e_A)])_{\nu \in V}$ converge to $\pi(e_A)$ in $\pi(A)$. Since π^{-1} is continuous, $((x - \lambda e_A)x_\mu)_{\mu \in M}$ and $(y_\nu(x - \lambda e_A))_{\nu \in V}$ converge to e_A in A . Hence, $x - \lambda e_A \in G^t(A)$, i.e. $\lambda \notin \sigma_A^t(x)$. Consequently, $\sigma_A^t(x) \subset \sigma_B^t(\pi(x))$. ■

PROPOSITION 6. Let A and B be unital semitopological algebras and π a continuous homomorphism from A into B . If B is a Hausdorff space and $\pi(A)$ is dense in B , then

$$G^t(\pi(A)) = G^t(B) \cap \pi(A).$$

Proof. It is easy to see that $G^t(\pi(A)) \subset G^t(B) \cap \pi(A)$. To prove the opposite inclusion, let O be a neighbourhood of zero in $\pi(A)$ (then there is a neighbourhood O' of zero in B with $O = O' \cap \pi(A)$) and let $y \in G^t(B) \cap \pi(A)$. Then there are $x \in A$ such that $y = \pi(x)$ and a neighbourhood O_1 of zero in B such that $yO_1 + O_1, O_1y + O_1 \subset O$. Since $y \in G^t(B)$, there exist nets $(y_\mu)_{\mu \in M}$ and $(z_\nu)_{\nu \in V}$ in B such that $(yy_\mu)_{\mu \in M}$ and $(z_\nu y)_{\nu \in V}$ converge to e_B . Therefore, there are $\mu_0 \in M$ and $\nu_0 \in V$ such that $yy_{\mu_0} - e_B \in O_1$ whenever $\mu > \mu_0$, and $z_{\nu_0}y - e_B \in O_1$ whenever $\nu > \nu_0$. Let $\mu_1 \in M$ and $\nu_1 \in V$ be such that $\mu_1 > \mu_0$ and $\nu_1 > \nu_0$. Then $yy_{\mu_1} - e_B \in O_1$ and $z_{\nu_1}y - e_B \in O_1$. Since $\pi(A)$ is dense in B , there are nets $(u_\alpha)_{\alpha \in A}$ and $(v_\beta)_{\beta \in B}$ such that $(\pi(u_\alpha))_{\alpha \in A}$ converges to y_{μ_1} and $(\pi(v_\beta))_{\beta \in B}$ converges to z_{ν_1} . Hence, there

are $\lambda_0 \in \mathcal{A}$ and $\beta_0 \in \mathcal{B}$ such that $\pi(u_\alpha) - y_{\mu_1} \in O_1$ whenever $\alpha > \alpha_0$, and $\pi(v_\beta) - z_{\nu_1} \in O_1$ whenever $\beta > \beta_0$. Taking this into account ⁽⁷⁾,

$\pi(x)\pi(u_\alpha) - \pi(e_A) = \pi(x)[\pi(u_\alpha) - y_{\mu_1}] + \pi(x)y_{\mu_1} - e_B \in yO_1 + O_1 \subset O'$
whenever $\alpha > \alpha_0$, and

$\pi(v_\beta)\pi(x) - \pi(e_A) = [\pi(v_\beta) - z_{\nu_1}]\pi(x) + z_{\nu_1}\pi(x) - e_B \in O_1y + O_1 \subset O'$
whenever $\beta > \beta_0$. Consequently, $(\pi(x)\pi(u_\alpha))_{\alpha \in A}$ and $(\pi(v_\beta)\pi(x))_{\beta \in B}$ converge to $\pi(e_A)$ in $\pi(A)$. Thus, $y \in G^t(\pi(A))$. ■

COROLLARY 7. *Let A and B be unital semitopological algebras and π a continuous homomorphism from A into B . If B is a Hausdorff space and $\pi(A)$ is dense in B , then*

$$\sigma_{\pi(A)}^t(\pi(a)) = \sigma_B^t(\pi(a))$$

for each $a \in A$.

Proof. Let $a \in A$. If $\lambda \in \sigma_{\pi(A)}^t(\pi(a))$, then $\pi(a) - \lambda\pi(e_A) \notin G^t(\pi(A))$. Hence, $\pi(a) - \lambda\pi(e_A) \notin G^t(B)$, by Proposition 6. Therefore, $\lambda \in \sigma_B^t(\pi(a))$. Thus,

$$\sigma_{\pi(A)}^t(\pi(a)) \subset \sigma_B^t(\pi(a)).$$

The opposite inclusion holds similarly, by Proposition 6. ■

COROLLARY 8. *Let A be a unital semitopological Hausdorff algebra, \tilde{A} the completion of A and τ the homeomorphism from A into \tilde{A} defined by the completion of A . Then*

$$\sigma_{\tilde{A}}^t(\tau(a)) = \sigma_A^t(a)$$

for each $a \in A$ ⁽⁸⁾.

Proof. Since $\tau(A)$ is dense in \tilde{A} , the assertion holds by Corollary 6. ■

4.2. Spectral mapping property for the topological spectrum. To prove the spectral mapping theorem for the topological spectrum, we need the following

LEMMA 3. *Let A be a commutative unital semitopological algebra, $n \in \mathbb{N}$ and $x_1, \dots, x_n \in A$. Then $x_1 \cdots x_n \in G^t(A)$ if and only if $x_i \in G^t(A)$ for each $i \in \mathbb{N}_n$ ⁽⁹⁾.*

⁽⁷⁾ Since B is a Hausdorff space and π is continuous, and $\pi(A)$ is dense in B , we have $\pi(e_A) = e_B$.

⁽⁸⁾ For unital complex metrizable algebras this result is known (see [19, Proposition 16]).

⁽⁹⁾ Here and below, $\mathbb{N}_n = \{1, 2, \dots, n\}$ for each $n \in \mathbb{N}$.

Proof. Let $x = x_1 \cdots x_n \in G^t(A)$. Then there exists a net $(z_\alpha)_{\alpha \in A}$ in A such that $(xz_\alpha)_{\alpha \in A}$ converges to e_A . Therefore

$$(x_i(x_1 \cdots x_{i-1} \cdot x_{i+1} \cdots x_n z_\alpha))_{\alpha \in A}$$

converges to e_A for each $i \in \mathbb{N}_n$. Now for each fixed $\alpha \in \mathcal{A}$ we put $w_\alpha^1 = x_2 \cdots x_n z_\alpha$, $w_\alpha^i = x_1 \cdots x_{i-1} \cdot x_{i+1} \cdots x_n z_\alpha$ for each i with $2 \leq i \leq n - 2$ and $w_\alpha^n = x_1 \cdots x_{n-1} z_\alpha$. Then for any $i \in \mathbb{N}_n$ the net $(x_i w_\alpha^i)_{\alpha \in A}$ converges to e_A . Hence, $x_i \in G^t(A)$ for $1 \leq i \leq n$.

Conversely, let $x_i \in G^t(A)$ for any $1 \leq i \leq n$. Using induction, it is sufficient to consider the case $n = 2$. Since $x_1 \in G^t(A)$, there exists a net $(z_\alpha)_{\alpha \in A}$ in A such that $(x_1 z_\alpha)_{\alpha \in A}$ converges to e_A , so for any (open) neighbourhood U of zero in A there is $\alpha_0 \in \mathcal{A}$ such that $x_1 z_\alpha \in e_A + U$ whenever $\alpha \geq \alpha_0$. Moreover, since $x_2 \in G^t(A)$, there exists a net $(v_\beta)_{\beta \in \mathcal{B}}$ in A such that $(v_\beta x_2)_{\beta \in \mathcal{B}}$ converges to e_A . Now $(v_\beta x_2(x_1 z_\alpha))_{\beta \in \mathcal{B}}$ converges to $x_1 z_\alpha$ for each fixed $\alpha \in \mathcal{A}$. Since $e_A + U$ is a neighbourhood of $x_1 z_\alpha$ for each $\alpha \geq \alpha_0$, there exists $\beta_0 \in \mathcal{B}$ such that $(v_\beta z_\alpha)x_1 x_2 \in e_A + U$ for each $\alpha \geq \alpha_0$ and $\beta \geq \beta_0$. Hence, there is a net $(v_\beta z_\alpha)_{(\beta, \alpha) \in \mathcal{B} \times \mathcal{A}}$ in A such that $((v_\beta z_\alpha)(x_1 x_2))_{(\beta, \alpha) \in \mathcal{B} \times \mathcal{A}}$ converges to e_A . Consequently, $x_1 x_2 \in G^t(A)$. ■

PROPOSITION 7. *Let A be a commutative complex unital semitopological algebra. Then*

$$\sigma_A^t(p(x)) = p(\sigma_A^t(x))$$

for any complex non-constant polynomial p and element $x \in A$.

Proof. Fix a polynomial p of degree $n > 0$ and $x \in A$. If $\lambda \in \sigma_A^t(x)$, then $x - \lambda e_A \notin G^t(A)$. Since $p(x) - p(\lambda)e_A = (x - \lambda e_A)r(x, \lambda)$ (where r is a polynomial in two variables), we have $p(x) - p(\lambda)e_A \notin G^t(A)$ by Lemma 3, i.e. $p(\lambda) \in \sigma_A^t(p(x))$. Thus, $p(\sigma_A^t(x)) \subset \sigma_A^t(p(x))$.

Conversely, let $\mu \in \sigma_A^t(p(x))$. Then $g(t) = p(t) - \mu = \alpha(t - \mu_1) \cdots (t - \mu_n)$, where $\alpha, \mu_1, \dots, \mu_n \in \mathbb{C}$, $\alpha \neq 0$ and μ_1, \dots, μ_n are the zeros of g . It follows that $p(\mu_i) = \mu$ for each $1 \leq i \leq n$. Now we have the factorization $g(x) = p(x) - \mu e_A = \alpha(x - \mu_1 e_A) \cdots (x - \mu_n e_A)$. Since $\mu \in \sigma_A^t(p(x))$, it follows that $g(x) \notin G^t(A)$. Hence, $x - \mu_k e_A \notin G^t(A)$ for some k by Lemma 3, i.e. $\mu_k \in \sigma_A^t(x)$. Therefore, $\mu = p(\mu_k) \in p(\sigma_A^t(x))$. Thus $\sigma_A^t(p(x)) \subset p(\sigma_A^t(x))$. ■

4.3. *Functional topological spectrum.* Next we describe the class of semitopological algebras, all elements of which have functional topological spectrum.

PROPOSITION 8. *Let A be a commutative unital simplicial Gelfand–Mazur algebra. If $\varphi(a) \neq 0$ for each $\varphi \in \mathfrak{M}(A)$ ⁽¹⁰⁾, then $a \in G^t(A)$.*

⁽¹⁰⁾ Since A is a commutative unital simplicial Gelfand–Mazur algebra, $\mathfrak{M}(A)$ is not empty.

Proof. Let $a \in A$ be such that $\varphi(a) \neq 0$ for each $\varphi \in \mathfrak{M}(A)$. Suppose that $a \notin G^t(A)$. Then there is a closed maximal ideal M in A such that $\overline{aA} \subset M$ (because A is simplicial). Since A is also a Gelfand–Mazur algebra, there exists $\varphi \in \mathfrak{M}(A)$ such that $M = \ker \varphi$. Hence, $\varphi(a) = 0$, contrary to assumption. ■

COROLLARY 9. *Every element of a commutative unital simplicial Gelfand–Mazur algebra has functional topological spectrum.*

Proof. Let $a \in A$ and $\lambda \in \sigma_A^t(a)$. Then $a - \lambda e_A \notin G^t(A)$. Hence, by Proposition 8, there is a $\varphi \in \mathfrak{M}(A)$ such that $\varphi(a) = \lambda$. Thus, $\sigma_A^t(a) \subset \{\varphi(x) : \varphi \in \mathfrak{M}(A)\}$. ■

COROLLARY 10. *Every element of a commutative unital locally m -pseudoconvex Hausdorff algebra has functional topological spectrum.*

Proof. Every commutative unital locally m -pseudoconvex Hausdorff algebra is a simplicial algebra (see [5, Corollary 5]) and a Gelfand–Mazur algebra (see, for example, [1, Corollary 2]). Now apply Corollary 9. ■

4.4. Properties of topological spectral radius. Now we list some properties of the topological spectral radius.

PROPOSITION 9. *Let A be a semitopological algebra. The topological spectral radius ϱ_A^t has the following properties:*

- (a) $\varrho_A^t(\mu x) = |\mu| \varrho_A^t(x)$ for all $x \in A$ and $\mu \in \mathbb{K}$;
- (b) $\varrho_A^t(xy) = \varrho_A^t(yx)$ for all $x, y \in A$;
- (c) if A is commutative, then $\varrho_A^t(x^n) = \varrho_A^t(x)^n$ for all $x \in A$ and $n \in \mathbb{N}$.

In particular, if every element in A has functional topological spectrum, then

- (d) $\varrho_A^t(x + y) \leq \varrho_A^t(x) + \varrho_A^t(y)$ for all $x, y \in A$;
- (e) $\varrho_A^t(xy) \leq \varrho_A^t(x) \varrho_A^t(y)$ for all $x, y \in A$.

Proof. Let $x \in A$ be such that $\sigma_A^t(x) \neq \emptyset$, and $\mu \in \mathbb{K} \setminus \{0\}$ (the case when $\mu = 0$ is trivial). Since $\sigma_A^t(\mu x) = \mu \sigma_A^t(x)$ for each $\mu \in \mathbb{K}$ by Proposition 4(a), we have $|\lambda| \leq |\mu| \varrho_A^t(x)$ for all $\lambda \in \sigma_A^t(\mu x)$ and $\mu \in \mathbb{K}$. Therefore, the relations

$$\varrho_A^t(\mu x) \leq |\mu| \varrho_A^t(x) = |\mu| \varrho_A^t(\mu^{-1}(\mu x)) \leq \varrho_A^t(\mu x)$$

yield (a). Statements (b) and (c) follow from Proposition 4(d) and Proposition 7 respectively. If A has functional topological spectrum, then we can represent every $\lambda \in \sigma_A^t(x + y)$ in the form $\lambda = \varphi_0(x) + \varphi_0(y)$ for some $\varphi_0 \in \mathfrak{M}(A)$. Since

$$|\lambda| \leq |\varphi_0(x)| + |\varphi_0(y)| \leq \varrho_A^t(x) + \varrho_A^t(y),$$

statement (d) follows. Statement (e) is proved similarly. ■

COROLLARY 11. *Let A be a unital semitopological algebra. If every element of A has functional topological spectrum, then the topological spectral radius is a submultiplicative seminorm on A .*

COROLLARY 12. *Let A be a commutative unital locally m -pseudoconvex Hausdorff algebra. Then the topological spectral radius is a submultiplicative seminorm on A .*

5. Open problems. Now we state several open problems, connected with topological invertibility of elements.

5.1. Let A be a semitopological algebra and $x \in G^t(A)$. Then there exist nets $(x_\alpha)_{\alpha \in A}$ and $(y_\mu)_{\mu \in M}$ such that $\lim_\alpha x_\alpha x = e_A$ and $\lim_\mu xy_\mu = e_A$. Let $G_0^t(A) = \{x \in G^t(A) : \text{there exists a net } (x_\alpha)_{\alpha \in A} \text{ such that } \lim_\alpha x_\alpha x = e_A \text{ and } \lim_\alpha xx_\alpha = e_A\}$. If A is commutative, then $G_0^t(A) = G^t(A)$.

PROBLEM 1. *Does there exist a non-commutative semitopological algebra A for which $G_0^t(A) = G^t(A)$?*

5.2. By Corollary 2, if A is a unital Fréchet algebra with discontinuous inversion $x \mapsto x^{-1}$, then $G^t(A) \setminus G(A) \neq \emptyset$, but the converse is open so far.

PROBLEM 2. *Let A be a unital Fréchet algebra. Does the discontinuity of the inversion $x \mapsto x^{-1}$ on $G(A)$ follow from $G^t(A) \setminus G(A) \neq \emptyset$?*

5.3. By [18, Theorem 2], there is a semitopological algebra A with $\overline{G^t(A)} = A$.

PROBLEM 3. *For which semitopological algebras A does $\overline{G^t(A)} = A$ hold?*

PROBLEM 4. *For which semitopological algebras A does $\overline{G(A)} = G^t(A)$ hold?*

5.4. We have the following result.

PROPOSITION 10. *A semitopological algebra A is a Q -algebra ⁽¹¹⁾ if and only if $G^t(A)$ is open and $\sigma_A^t(a) = \sigma_A(a)$ for each $a \in A \setminus G(A)$.*

Proof. If A is a Q -algebra, then $G^t(A) = G(A)$. Therefore, $G^t(A)$ is open in A and $\sigma_A^t(a) = \sigma_A(a)$ for each $a \in A \setminus G(A)$.

Conversely, let A be a semitopological algebra such that $G^t(A)$ is open and $\sigma_A^t(a) = \sigma_A(a)$ for each $a \in A \setminus G(A)$. If now $a \in A \setminus G(A)$, then $0 \in \sigma_A(a)$. Therefore there exists a sequence (λ_n) in $\sigma_A^t(a)$ which converges to 0. Since $a - \lambda_n e_A \in A \setminus G^t(A)$ and $A \setminus G^t(A)$ is closed in A , we have $a \in A \setminus G^t(A)$. Thus, $G(A) = G^t(A)$. Consequently, A is a Q -algebra. ■

PROBLEM 5. *Describe the elements of a semitopological algebra A for which $\sigma_A^t(a) = \sigma_A(a)$.*

⁽¹¹⁾ That is, $G(A)$ is open in A .

5.5. If A is an algebra and B a maximal commutative subalgebra of A , then $\sigma_A(a) = \sigma_B(a)$. It is not clear whether a similar result holds for the topological spectrum.

PROBLEM 6. *Let A be a semitopological algebra. Does there exist a maximal commutative subalgebra such that $\sigma_A^t(a) \neq \sigma_B^t(a)$ for some $a \in A$?*

References

- [1] M. Abel, *Gelfand–Mazur algebras*, in: Topological Vector Spaces, Algebras and Related Areas (Hamilton, ON, 1994), Longman, Harlow, 1994, 116–129.
- [2] —, *Advertive topological algebras*, in: General Topological Algebras (Tartu, 1999), Math. Stud. (Tartu) 1, Est. Math. Soc., Tartu, 2001, 14–24.
- [3] —, *Descriptions of the topological radical in topological algebras*, *ibid.*, 25–31.
- [4] —, *Survey of results on Gelfand–Mazur algebras*, in: Non-Normed Topological Algebras (Rabat, 2000), E. N. S. Takaddoum Publ., Rabat, 2004, 14–25.
- [5] —, *Inductive limits of Gelfand–Mazur algebras*, *Int. J. Pure Appl. Math.* 16 (2004), 363–378.
- [6] M. Akkar, A. Beddaa et M. Oudadess, *Sur une classe d’algèbres topologiques*, *Bull. Belg. Math. Soc. Simon Stevin* 3 (1996), 13–24.
- [7] —, —, —, *Topologically invertible elements in metrizable algebras*, *Indian J. Pure Appl. Math.* 27 (1996), 1–5.
- [8] R. F. Arens, *The space L^ω and convex topological rings*, *Bull. Amer. Math. Soc.* 52 (1946), 931–935.
- [9] H. Arizmendi, A. Carillo and L. Palacios, *On Q_t -algebras*, manuscript.
- [10] V. K. Balachandran, *Topological Algebras*, North-Holland Math. Stud. 185, North-Holland, Amsterdam, 2000.
- [11] S. J. Batt and A. D. Thatte, *On topological invertibility*, *Indian J. Pure Appl. Math.* 15 (1984), 1308–1312.
- [12] B. Beckenstein, L. Narici and Ch. Suffel, *Topological Algebras*, North-Holland Math. Stud. 24, North-Holland, Amsterdam, 1977.
- [13] A. Beddaa, *Algèbres localement convexes advertiblement complètes et continuité automatique de morphismes*, Thèse Sci. Math., Univ. Mohamed V, Rabat, 1997.
- [14] A. Białynicki-Birula and W. Żelazko, *On the multiplicative-linear functionals on the Cartesian product of algebras*, *Bull. Acad. Polon. Sci. Sér. Sci. Math. Astronom. Phys.* 5 (1957), 201–203.
- [15] R. Choukri, A. El Kinani et M. Oudadess, *Inversibilité topologique et problème de l’idéal fermé*, *Bol. Soc. Mat. Mexicana* (3) 9 (2003), 109–117.
- [16] K. Luha, *Topological invertibility in topological algebras*, *Tartu Üli. Toimetised* 940 (1992), 71–74.
- [17] W. Żelazko, *Selected Topics in Topological Algebras*, Aarhus Univ. Lecture Notes 31, Aarhus Univ., Aarhus, 1971.
- [18] —, *Topological simplicity of a certain \mathcal{LF} -algebra*, *Period. Math. Hungar.* 35 (1997), 145–148.
- [19] —, *When a commutative unital F -algebra has a dense principal ideal*, in: Topological Algebras and Their Applications, *Contemp. Math.* 341, Amer. Math. Soc., Providence, RI, 2004, 133–137.
- [20] —, *F -algebras: some results and open problems*, in: Functional Analysis and its Applications, North-Holland Math. Stud. 197, Elsevier, Amsterdam, 2004, 317–326.

- [21] W. Żelazko, *Continuous characters and joint topological spectrum*, submitted.

Mati Abel
Institute of Pure Mathematics
University of Tartu
Liivi 2-614, 50409 Tartu, Estonia
E-mail: mati.abel@ut.ee

Wiesław Żelazko
Institute of Mathematics
Polish Academy of Sciences
P.O. Box 137, Śniadeckich 8
00-956 Warszawa, Poland
E-mail: zelazko@impan.gov.pl

Received June 30, 2006

(7537)