

COMPOSITION AND L^2 -BOUNDEDNESS OF FLAG KERNELS

BY

PAWEŁ GŁOWACKI (Wrocław)

Abstract. We prove the composition and L^2 -boundedness theorems for the Nagel–Ricci–Stein flag kernels related to the natural gradation of homogeneous groups.

In [3], Nagel, Ricci, and Stein introduce the notion of a flag kernel which generalizes that of a singular integral kernel of Calderón–Zygmund as a tool in their investigation of operators naturally associated with the $\bar{\partial}_b$ on some CR submanifolds of $\mathbb{C}^n \times \mathbb{C}^n$. A *flag kernel* K on a Euclidean vector space V endowed with a family of dilations and a corresponding homogeneous norm $x \mapsto |x|$ is a tempered distribution associated with gradations

$$V = \bigoplus_{j=1}^R V_j, \quad V^* = \bigoplus_{j=1}^R V_j^*$$

of the space and its dual. The Fourier transform of K is required to be smooth for $\xi_R \neq 0$ and satisfy

$$(0.1) \quad |D^\alpha \widehat{K}(\xi)| \leq C_\alpha |\xi_1|^{-|\alpha_1|} \cdots |\xi_R|^{-|\alpha_R|},$$

where

$$|\xi|_j = \sum_{k=j}^R |\xi_k|, \quad \xi = \sum_{k=1}^R \xi_k \in V^*,$$

and α_j are submultiindices corresponding to the spaces V_j^* . Actually, the authors define the flag kernels directly in terms of the smoothness and cancellation properties of the kernels, and then prove that the multiplier condition (0.1) is an equivalent possibility of definition.

They prove that if V is the Lie algebra of a homogeneous group identified with the group itself, dilations are automorphisms of the group, the spaces V_j are homogeneous, and $[V_j, V_k] = \{0\}$ for $j \neq k$, then the composition of flag kernels associated with the same gradation is still a flag kernel. Moreover,

2010 *Mathematics Subject Classification*: Primary 22E30; Secondary 35S05.

Key words and phrases: singular integrals, flag kernels, symbolic calculus, homogeneous groups, Fourier transform.

under the same hypotheses any flag kernel K defines a bounded operator

$$Kf(x) = f \star \tilde{K}(x) = \int_V f(xy)K(y) dy$$

on $L^p(V)$ for $1 < p < \infty$.

A natural question arises whether the composition and boundedness properties still hold if the underlying gradation is the natural gradation of a homogeneous group. Note that for homogeneous groups of step greater than 2 the commutator condition is no longer satisfied. We provide an answer in the affirmative under somewhat relaxed assumptions on the kernels and $p = 2$.

The results presented here depend heavily on the symbolic calculus of [1] and can be regarded as an example of usefulness of such a calculus. There occurs a striking resemblance between the estimates defining flag kernels and those of the calculus which has been created and developed quite independently.

The problem of the L^p -boundedness of flag kernels on arbitrary homogeneous groups will be dealt with in another paper [2].

I wish to thank Fulvio Ricci for a fruitful conversation concerning the subject of this paper.

Even though I had no opportunity to discuss the subject matter of this paper with Andrzej Hulanicki, it unavoidably bears signs of his influence which can be traced back throughout the whole of my mathematical work.

1. Background. Let \mathfrak{g} be a nilpotent Lie algebra with a fixed Euclidean structure and \mathfrak{g}^* its dual. Let $\{\delta_t\}_{t>0}$ be a family of group dilations on \mathfrak{g} and let

$$\mathfrak{g}_j = \{x \in \mathfrak{g} : \delta_t x = t^{d_j} x\}, \quad 1 \leq j \leq R,$$

where $1 = d_1 < \dots < d_R$. Then

$$(1.1) \quad \mathfrak{g} = \bigoplus_{j=1}^R \mathfrak{g}_j, \quad \mathfrak{g}^* = \bigoplus_{j=1}^R \mathfrak{g}_j^*,$$

and

$$[\mathfrak{g}_i, \mathfrak{g}_j] \subset \begin{cases} \mathfrak{g}_k & \text{if } d_i + d_j = d_k, \\ \{0\} & \text{if } d_i + d_j \notin \mathcal{D}, \end{cases}$$

where $\mathcal{D} = \{d_j : 1 \leq j \leq R\}$. Let

$$\xi \mapsto |\xi| = \sum_{j=1}^R \|\xi_j\|^{1/d_j} = \sum_{j=1}^R |\xi_j|$$

be a homogeneous norm on \mathfrak{g}^* . We say that \mathfrak{g} is *homogeneous of step R* .

We shall also regard \mathfrak{g} as a Lie group with the Campbell–Hausdorff multiplication

$$x_1 x_2 = x_1 + x_2 + r(x_1, x_2),$$

where

$$\begin{aligned} r(x_1, x_2) &= \frac{1}{2} [x_1, x_2] + \frac{1}{12} ([x_1, [x_1, x_2]] + [x_2, [x_2, x_1]]) \\ &\quad + \frac{1}{24} [x_2, [x_1, [x_2, x_1]]] + \cdots \end{aligned}$$

is the (finite) sum of terms of order at least 2 in the Campbell–Hausdorff series for \mathfrak{g} .

Let

$$|\xi|_j = \sum_{k=j}^R |\xi_k|, \quad 1 \leq j \leq R,$$

and let $|\xi|_{R+1} = 0$. Let

$$\mathbf{q}_\xi(\eta) = \sum_{j=1}^R \frac{\|\eta_j\|}{1 + |\xi|_{j+1}}, \quad \xi, \eta \in \mathfrak{g}^*,$$

be a family of norms (a Hörmander metric) on \mathfrak{g}^* . Let g_j be a family of functions on \mathfrak{g}^* satisfying

$$|\xi|_{j+1} \leq g_j(\xi) \leq |\xi|, \quad 1 \leq j \leq R,$$

and

$$\left(\frac{1 + g_j(\xi)}{1 + g_j(\eta)} \right)^{\pm 1} \leq C(1 + \mathbf{q}_\xi(\xi - \eta))^M$$

for some $C > 0$ and $M > 0$. The metric \mathbf{q} is fixed throughout the paper (cf. [1]).

The class $S^m(\mathfrak{g})$, where $m \in \mathbb{R}$, is defined as the space of all $A \in \mathcal{S}'(\mathfrak{g})$ whose Fourier transforms are smooth and satisfy

$$|D^\alpha \widehat{A}(\xi)| \leq C_\alpha (1 + |\xi|)^m \prod_{j=1}^R (1 + g_j(\xi))^{-|\alpha_j|}, \quad \xi \in \mathfrak{g}^*,$$

where $\alpha = (\alpha_1, \dots, \alpha_R)$ is a multiindex of length equal to the dimension of \mathfrak{g}^* , and α_j are submultiindices corresponding to the subspaces \mathfrak{g}_j^* . Note that the elements of $S^m(\mathfrak{g})$ have no singularity at infinity.

The space $S^m(\mathfrak{g})$ is a Fréchet space if equipped with the seminorms

$$\|A\|_\alpha = \sup_{\xi \in \mathfrak{g}^*} (1 + |\xi|)^{-m} \prod_{k=1}^R (1 + g_k(\xi))^{|\alpha|} |D^\alpha \widehat{A}(\xi)|.$$

The class $S^0(\mathfrak{g})$ is known to be a subalgebra of $\mathcal{B}(L^2(\mathfrak{g}))$. More precisely, we have the following two propositions proved in [1].

PROPOSITION 1.2. *The mapping*

$$S^{m_1}(\mathfrak{g}) \times S^{m_2}(\mathfrak{g}) \ni (A, B) \mapsto A \star B \in S^{m_1+m_2}(\mathfrak{g})$$

is continuous.

PROPOSITION 1.3. *If $A \in S^0(\mathfrak{g})$, then*

$$\text{Op}(A)f(x) = \int_{\mathfrak{g}} f(xy) A(dy), \quad f \in \mathcal{S}(\mathfrak{g}),$$

extends to a bounded operator on $L^2(\mathfrak{g})$, and the mapping

$$S^0(\mathfrak{g}) \ni A \mapsto \text{Op}(A) \in \mathcal{B}(L^2(\mathfrak{g}))$$

is continuous.

2. Main results. We extend the definition of a flag kernel of Nagel–Ricci–Stein to include all $K \in \mathcal{S}'(\mathfrak{g})$ whose Fourier transforms are smooth for $\xi_R \neq 0$ and satisfy

$$(2.1) \quad |D^\alpha \widehat{K}(\xi)| \leq C_\alpha \prod_{j=1}^R g_j(\xi)^{-|\alpha_j|}, \quad \xi_R \neq 0.$$

Note that for $g_j(\xi) = |\xi|_j$ we get the usual flag kernels. Another interesting choice is $g_j(\xi) = |\xi|_{j+1}$. In the latter case the estimates of the derivatives in the direction of ξ_R are irrelevant. Observe that if

$$\langle K_t, f \rangle = \int_{\mathfrak{g}} f(tx)K(x) dx,$$

then the flag kernels K_t satisfy the estimates (2.1) uniformly in $t > 0$.

We shall need two cut-off functions. Let $\phi \in C^\infty(\mathfrak{g}_R^*)$ be equal to 1 for $1 \leq |\xi_R| \leq 2$ and vanish for $|\xi_R| \geq 4$ and $|\xi_R| \leq 1/2$. Let $\psi \in C^\infty(\mathfrak{g}_R^*)$ be equal to 1 for $1/2 \leq |\xi_R| \leq 4$ and vanish for $|\xi_R| \geq 8$ and $|\xi_R| \leq 1/4$. Thus, in particular, $\phi \cdot \psi = \phi$.

THEOREM 2.2. *A composition of flag kernels is also a flag kernel.*

Proof. Let $K = K_1 \star K_2$, where K_j are flag kernels. Then

$$\widehat{K}(\xi) = \widehat{A_1 \star A_2}(\xi), \quad 1 \leq |\xi_R| \leq 2,$$

where

$$\widehat{A_j}(\xi) = \widehat{K_j}(\xi)\phi(\xi_R),$$

and $A_j \in S^0(\mathfrak{g})$. Therefore, by Proposition 1.2,

$$(2.3) \quad |D^\alpha \widehat{K}(\xi)| \leq C_\alpha \prod_{j=1}^R g_j(\xi)^{-|\alpha_j|}, \quad 1 \leq |\xi_R| \leq 2.$$

Since

$$(2.4) \quad \widehat{(K_j)_t}(\xi) = \widehat{K}_j(t\xi), \quad t > 0,$$

satisfy (2.1) uniformly, we get the estimate (2.3) for $1/t \leq |\xi_R| \leq 2/t$, $t > 0$, that is, for all $\xi_R \neq 0$. ■

THEOREM 2.5. *Let K be a flag kernel. The operator $f \mapsto f \star \widetilde{K}$ defined initially on $\mathcal{S}(\mathfrak{g})$ extends uniquely to a bounded operator on $L^2(\mathfrak{g})$.*

Proof. For $f \in \mathcal{S}(\mathfrak{g})$ let

$$\widehat{f}_n(\xi) = \widehat{f}(\xi)\phi(2^{-n}\xi_R), \quad n \in \mathbb{Z}.$$

Then

$$m\|f\|_2 \leq \sum_{n=-\infty}^{\infty} \|f_n\|_2 \leq M\|f\|_2, \quad f \in \mathcal{S}(\mathfrak{g}),$$

for some $m, M > 0$. Let K_n be defined by

$$\widehat{K}_n(\xi) = \widehat{K}(\xi)\psi(2^{-n}\xi_R).$$

Then the flag kernels K_n are uniformly in $S^0(\mathfrak{g})$, and

$$\|Kf\|_2 \leq \frac{1}{m} \sum_{n \in \mathbb{Z}} \|K_n f_n\| \leq \frac{C}{m} \sum_{n \in \mathbb{Z}} \|f_n\|_2 \leq \frac{CM}{m} \|f\|_2$$

for a $C > 0$, which completes the proof. ■

REFERENCES

- [1] P. Głowacki, *The Melin calculus for general homogeneous groups*, Ark. Mat. 45 (2007), 31–48.
- [2] —, *L^p -boundedness of singular integral operators with flag kernels via symbolic calculus*, to appear.
- [3] A. Nagel, F. Ricci and E. M. Stein, *Singular integrals with flag kernels and analysis on quadratic CR manifolds*, J. Funct. Anal. 181 (2001), 29–118.

Paweł Głowacki
 Institute of Mathematics
 University of Wrocław
 Pl. Grunwaldzki 2/4
 50-384 Wrocław, Poland
 E-mail: glowacki@math.uni.wroc.pl

Received 28 March 2009;
 revised 16 August 2009