Upper and lower estimates for Schauder frames and atomic decompositions

by

Kevin Beanland (Lexington, VA), Daniel Freeman (St. Louis, MO) and **Rui Liu** (Tianjin and College Station, TX)

Abstract. We prove that a Schauder frame for any separable Banach space is shrinking if and only if it has an associated space with a shrinking basis, and that a Schauder frame for any separable Banach space is shrinking and boundedly complete if and only if it has a reflexive associated space. To obtain these results, we prove that the upper and lower estimate theorems for finite-dimensional decompositions of Banach spaces can be extended and modified to Schauder frames. We show as well that if a separable infinitedimensional Banach space has a Schauder frame, then it also has a Schauder frame which is not shrinking.

1. Introduction. Frames for Hilbert spaces were introduced by Duffin and Schaeffer in 1952 [DS] to address some questions in non-harmonic Fourier series. However, the current popularity of frames is largely due to their successful application to signal processing which was initiated by Daubechies, Grossmann, and Meyer in 1986 [DGM]. A frame for an infinitedimensional separable Hilbert space H is a sequence of vectors $(x_i)_{i=1}^{\infty} \subset H$ for which there exist constants $0 \le A \le B$ such that for any $x \in H$,

(1)
$$A\|x\|^{2} \leq \sum_{i=1}^{\infty} |\langle x, x_{i} \rangle|^{2} \leq B\|x\|^{2}.$$

If A = B = 1, then $(x_i)_{i=1}^{\infty}$ is called a *Parseval frame*. Given any frame $(x_i)_{i=1}^{\infty}$ for a Hilbert space H, there exists a frame $(f_i)_{i=1}^{\infty}$ for H, called an alternate dual frame, such that for all $x \in H$,

(2)
$$x = \sum_{i=1}^{\infty} \langle x, f_i \rangle x_i.$$

2010 Mathematics Subject Classification: Primary 46B20; Secondary 41A65.

Key words and phrases: Banach spaces, frames, Schauder frames, atomic decompositions. [161]

The equality in (2) allows the reconstruction of any vector x in the Hilbert space from the sequence of coefficients $(\langle x, f_i \rangle)_{i=1}^{\infty}$. The standard method to construct such a frame $(f_i)_{i=1}^{\infty}$ is to take $f_i = S^{-1}x_i$ for all $i \in \mathbb{N}$, where S is the positive, self-adjoint invertible operator on H defined by Sx = $\sum_{i=1}^{\infty} \langle x, x_i \rangle x_i$ for all $x \in H$. The operator S is called the *frame operator* and the frame $(S^{-1}x_i)_{i=1}^{\infty}$ is called the *canonical dual frame* of $(x_i)_{i=1}^{\infty}$.

In their AMS memoir [HL], Han and Larson initiated studying the dilation viewpoint of frames, that is, analyzing frames as orthogonal projections of Riesz bases, where a *Riesz basis* is a semi-normalized unconditional basis for a Hilbert space. To start this approach, they proved the following theorem.

THEOREM 1.1 ([HL]). If $(x_i)_{i=1}^{\infty}$ is a frame for a Hilbert space H, then there exists a larger Hilbert space $Z \supset H$ and a Riesz basis $(z_i)_{i=1}^{\infty}$ for Zsuch that $P_X z_i = x_i$ for all $i \in \mathbb{N}$, where P_X is the orthogonal projection onto X. Furthermore, if $(x_i)_{i=1}^{\infty}$ is Parseval, then $(z_i)_{i=1}^{\infty}$ can be taken to be an orthonormal basis.

Recently, Theorem 1.1 was extended to operator-valued measures and bounded linear maps [HLLL]. Recently as well, a continuous version of Theorem 1.1 was given for vector bundles and Riemannian manifolds [FPWW].

The concept of a frame was extended to Banach spaces in 1988 by Feichtinger and Gröchenig [FG] (and more generally in [G]) through the introduction of atomic decompositions. The main goal of [G] was to obtain for Banach spaces the unique association of a vector with the natural set of frame coefficients. In 2008, Schauder frames for Banach spaces were developed [CDOSZ] with the goal of creating a procedure to represent vectors using quantized coefficients. A Schauder frame essentially takes as its definition an extension of the equation (2) to Banach spaces.

DEFINITION 1.2. Let X be an infinite-dimensional separable Banach space. A sequence $(x_i, f_i)_{i=1}^{\infty} \subset X \times X^*$ is called a *Schauder frame* for X if $x = \sum_{i=1}^{\infty} f_i(x)x_i$ for all $x \in X$.

In particular, if $(x_i)_{i=1}^{\infty}$ and $(f_i)_{i=1}^{\infty}$ are frames for a Hilbert space H, then $(f_i)_{i=1}^{\infty}$ is an alternate dual frame for $(x_i)_{i=1}^{\infty}$ if and only if $(x_i, f_i)_{i=1}^{\infty}$ is a Schauder frame for H. As noted in [CDOSZ], a separable Banach space has a Schauder frame if and only if it has the bounded approximation property. By the uniform boundedness principle, for any Schauder frame $(x_i, f_i)_{i=1}^{\infty}$ of a Banach space X, there exists a constant $C \geq 1$ such that $\sup_{n\geq m} \|\sum_{i=m}^n f_i(x)x_i\| \leq C \|x\|$ for all $x \in X$. The least such value C is called the frame constant of $(x_i, f_i)_{i=1}^{\infty}$. A Schauder frame $(x_i, f_i)_{i=1}^{\infty}$ is called unconditional if the series $x = \sum_{i=1}^{\infty} f_i(x)x_i$ converges unconditionally for all $x \in X$. The following definitions allow the dilation viewpoint of Han and Larson to be extended to Schauder frames.

DEFINITION 1.3. Let $(x_i, f_i)_{i=1}^{\infty}$ be a frame for a Banach space X, and let Z be a Banach space with basis $(z_i)_{i=1}^{\infty}$ and coordinate functionals $(z_i^*)_{i=1}^{\infty}$. We call Z an associated space to $(x_i, f_i)_{i=1}^{\infty}$, and $(z_i)_{i=1}^{\infty}$ an associated basis, if the operators $T: X \to Z$ and $S: Z \to X$ are bounded, where T(x) = $T(\sum f_i(x)x_i) = \sum f_i(x)z_i$ for all $x \in X$ and $S(z) = S(\sum z_i^*(z)z_i) =$ $\sum z_i^*(z)x_i$ for all $z \in Z$.

Essentially, Theorem 1.1 states that a frame for a Hilbert space has an associated basis which is a Riesz basis for a Hilbert space. Furthermore, the proof in [HL] actually involves constructing the operators T and S given in Definition 1.3, and thus the definition of an associated space is a very natural way to extend the notion of dilation of Hilbert space frames to dilation of Schauder frames. In [CHL], it is shown that every Schauder frame has an associated space, which is referred to as the minimal associated space in [L]. Being able to dilate a Parseval frame for a Hilbert space to an orthonormal basis is very useful in understanding and working with frames for Hilbert spaces. The minimal associated basis may be used similarly for studying Schauder frames. However, if a Schauder frame has some useful property, then an associated basis with a corresponding property should be used to study the Schauder frame. To take full advantage of this approach, we need to characterize Schauder frame properties in terms of associated bases. This would allow the large literature on Schauder basis properties to be applied to Schauder frames. For example, in [CHL] it is proven that a Schauder frame is unconditional if and only if it has an unconditional associated space.

If $(x_i, f_i)_{i=1}^{\infty}$ is a Schauder frame, then the reconstruction operator, S, for the minimal associated space contains c_0 in its kernel if and only if a finite number of vectors can be removed from $(x_i)_{i=1}^{\infty}$ to make it a Schauder basis [LZ]. This implies in particular that except in trivial cases, the minimal associated basis will not be boundedly complete and the minimal associated space will not be reflexive. Thus, to work with a reflexive associated space, we will need to consider a new method of constructing associated spaces. A Banach space with a basis is reflexive if and only if the basis is shrinking and boundedly complete. In order to characterize when a Schauder frame has a reflexive associated space, it is then natural to consider a generalization of the properties of being shrinking and boundedly complete from the context of bases to that of Schauder frames.

DEFINITION 1.4. Given a Schauder frame $(x_i, f_i)_{i=1}^{\infty} \subset X \times X^*$, let $T_n : X \to X$ be the operator $T_n(x) = \sum_{i \ge n} f_i(x)x_i$. The frame $(x_i, f_i)_{i=1}^{\infty}$ is called *shrinking* if $||x^* \circ T_n|| \to 0$ for all $x^* \in X^*$. The frame $(x_i, f_i)_{i=1}^{\infty}$ is called *boundedly complete* if $\sum_{i=1}^{\infty} x^{**}(f_i)x_i$ converges in norm to an element of X for all $x^{**} \in X^{**}$.

K. Beanland et al.

As noted in [CL], if $(x_i)_{i=1}^{\infty}$ is a Schauder basis and $(x_i^*)_{i=1}^{\infty}$ are the biorthogonal functionals of $(x_i)_{i=1}^{\infty}$, then the frame $(x_i, x_i^*)_{i=1}^{\infty}$ is shrinking if and only if the basis $(x_i)_{i=1}^{\infty}$ is shrinking, and the frame $(x_i, x_i^*)_{i=1}^{\infty}$ is boundedly complete if and only if the basis $(x_i)_{i=1}^{\infty}$ is boundedly complete. Thus the definition of a frame being shrinking or boundedly complete is consistent with that of a basis. In [L], the frame properties "shrinking" and "boundedly complete".

It is not difficult to see that if a Schauder frame has a shrinking associated basis, then the frame must be shrinking as well, and that if a Schauder frame has a boundedly complete associated basis, then the frame must be boundedly complete. In [L], the minimal and maximal associated spaces are defined, and it is proven that if a frame is shrinking and satisfies some strong local conditions, then the minimal associated basis is shrinking, and if a frame is boundedly complete and satisfies some strong local conditions, then the maximal associated basis is boundedly complete. The advantage of these theorems is that explicit associated bases are constructed with the desired properties of being shrinking or boundedly complete. On the other hand, not every Schauder frame with a shrinking or boundedly complete associated basis satisfies the local conditions given in [L]. Thus, this is not a complete characterization of which frames have a shrinking associated basis or of which frames have a boundedly complete associated basis. In the following theorem, we give a complete characterization.

THEOREM 1.5. Let $(x_i, f_i)_{i=1}^{\infty}$ be a Schauder frame for a Banach space X. Then $(x_i, f_i)_{i=1}^{\infty}$ is shrinking if and only if $(x_i, f_i)_{i=1}^{\infty}$ has a shrinking associated basis. Furthermore, $(x_i, f_i)_{i=1}^{\infty}$ is shrinking and boundedly complete if and only if $(x_i, f_i)_{i=1}^{\infty}$ has a reflexive associated space.

The properties of being shrinking and boundedly complete do not immediately give us much structure to directly create an associated space. To obtain Theorem 1.5, we will prove a stronger result, involving upper and lower estimates. In [OSZ2], it is shown that every Banach space with separable dual satisfies certain upper estimates and that every separable reflexive Banach space satisfies certain upper and lower estimates. As we will be using these estimates, we will define them precisely in Section 3. One of the main results of [OSZ2] is that every separable reflexive Banach space embeds into a reflexive Banach space with a finite-dimensional decomposition (FDD) satisfying the same upper and lower estimates, and one of the main results of [FOSZ] is that every Banach space with separable dual embeds into a Banach space with a shrinking FDD satisfying the same upper estimate. We extend the techniques developed in [OSZ1] to Schauder frames, and prove the following theorems which are extensions of the results from [OSZ2] and [FOSZ], and will be used to prove Theorem 1.5. THEOREM 1.6. Let X be a Banach space with a shrinking Schauder frame $(x_i, f_i)_{i=1}^{\infty}$. Let $(v_i)_{i=1}^{\infty}$ be a normalized, 1-unconditional, block stable, right dominant, and shrinking basic sequence. If X satisfies subsequential $(v_i)_{i=1}^{\infty}$ -upper tree estimates, then there exist $(n_i)_{i=1}^{\infty}, (K_i)_{i=1}^{\infty} \in [\mathbb{N}]^{\omega}$ and an associated space Z with a shrinking basis $(z_i)_{i=1}^{\infty}$ such that $(\operatorname{span}_{j \in [n_i, n_{i+1})} z_i)_{i=1}^{\infty}$ is an FDD that satisfies subsequential $(v_{K_i})_{i=1}^{\infty}$ -upper block estimates.

THEOREM 1.7. Let X be a Banach space with a shrinking and boundedly complete Schauder frame $(x_i, f_i)_{i=1}^{\infty}$. Let $(u_i)_{i=1}^{\infty}$ be a normalized, 1-unconditional, block stable, right dominant, and shrinking basic sequence, and let $(v_i)_{i=1}^{\infty}$ be a normalized, 1-unconditional, block stable, left dominant, and boundedly complete basic sequence such that (u_i) dominates (v_i) . Then X satisfies subsequential $(u_i)_{i=1}^{\infty}$ -upper tree estimates and subsequential $(v_i)_{i=1}^{\infty}$ -lower tree estimates if and only if there exist $(n_i)_{i=1}^{\infty}, (K_i)_{i=1}^{\infty} \in [\mathbb{N}]^{\omega}$ and a reflexive associated space Z with associated basis $(z_i)_{i=1}^{\infty}$ such that the FDD $(\operatorname{span}_{j \in [n_i, n_{i+1}]} z_j)_{i=1}^{\infty}$ satisfies subsequential $(u_{K_i})_{i=1}^{\infty}$ -upper block estimates and subsequential $(v_{K_i})_{i=1}^{\infty}$ -lower block estimates.

Using the theory of trees and branches in Banach spaces developed by Odell and Schlumprecht, we find that Theorem 1.5 follows immediately from Theorems 1.6 and 1.7. The basic idea is that every Banach space with separable dual satisfies subsequential $(v_i)_{i=1}^{\infty}$ -upper tree estimates for some normalized, 1-unconditional, block stable, right dominant, and shrinking basic sequence $(v_i)_{i=1}^{\infty}$. If X is a Banach space with a shrinking Schauder frame, then it is known that X must have separable dual, and hence we may apply Theorem 1.6 to obtain a shrinking associated basis to the shrinking Schauder frame. Similarly, every separable reflexive Banach space satisfies subsequential $(u_i)_{i=1}^{\infty}$ -upper tree estimates and subsequential $(v_i)_{i=1}^{\infty}$ -lower tree estimates for some basic sequences $(u_i)_{i=1}^{\infty}$ and $(v_i)_{i=1}^{\infty}$ satisfying the hypotheses of Theorem 1.7. If X is a Banach space with a shrinking and boundedly complete Schauder frame, then it is known that X must be reflexive, and hence we may apply Theorem 1.7 to obtain a reflexive associated space for the Schauder frame. Because Theorems 1.5–1.7 are our main theorems, at the end of Section 3 we will prove explicitly (assuming some well known properties about Tsirelson spaces) how Theorem 1.5 follows from Theorems 1.6 and 1.7.

One of the main goals in proving the upper and lower estimate theorems in [OSZ1] and [FOSZ] was to obtain embedding results that preserve quantitative bounds on the Szlenk index. In [Z], Zippin proves that every Banach space with separable dual embeds into a Banach space with a shrinking basis, and that every separable reflexive Banach space embeds into a Banach space with a shrinking and boundedly complete basis. On the other hand, there does not exist a single Banach space Z with a shrinking basis such that every reflexive Banach space embeds into Z [Sz]. To prove this result, Szlenk created an ordinal index Sz on the set of separable Banach spaces such that: (1) Sz(X) is countable if and only if X has separable dual, (2) if X embeds into Y then Sz(X) \leq Sz(Y), and (3) for every countable ordinal α there exists a Banach space with separable dual X such that Sz(X) > α . Thus, if Z is a separable Banach space such that every Banach space with separable dual embeds into Z, then Sz(Z) is uncountable and hence Z does not have separable dual. There have been many further results in this direction, for example Bourgain [B] proved that if Z is a separable Banach space such that every separable reflexive Banach space embeds into Z, then also every separable Banach space embeds into Z. Bourgain then asked if there exists a separable reflexive Banach space Z such that every separable uniformly convex Banach space embeds into Z. In [OS2] Odell and Schlumprecht construct such a space answering Bourgain's question.

Inspired by [OS2], Pełczyński asked if for every countable ordinal α there exists a reflexive Banach space Z such that every separable reflexive Banach space X with $\max(S_Z(X), S_Z(X^*)) \leq \alpha$ embeds into Z, which naturally leads to the question if there exists a Banach space Z with separable dual such that every separable Banach space X with $S_Z(X) \leq \alpha$ embeds into Z. There have been two separate methods of answering these problems. The proofs of Argyros and Dodos [AD] and Dodos and Ferenczi [DF] used the descriptive set theory framework of studying sets of Banach spaces which was initiated in Bossard's PhD thesis [Bos]. The proofs in [OSZ2] and [FOSZ] used instead the equivalence of a Banach space X satisfying $Sz(X) \leq \omega^{\alpha\omega}$ with the Banach space having separable dual and satisfying certain upper $T_{\alpha,c}$ -estimates, where $T_{\alpha,c}$ is the Tsirelson space of order α and constant c. This approach was later generalized to more ordinals than just $\omega^{\alpha\omega}$ by Causey [C1], [C2]. We will rely on many of the techniques used in these papers, but there will be some notable distinctions. In all of [OS1], [OS2], [OSZ1], [OSZ2], [C1], and [C2], the very first step in the embedding theorems is to use Zippin's theorem to embed the Banach space into either a reflexive Banach space with a basis or a Banach space with a shrinking basis. In our case, obtaining a reflexive associated space or shrinking associated basis is the conclusion of our Theorem 1.5 rather than the first step in the proof. This is a significant obstacle, and much of the theorems and lemmas that we prove in Section 2 are aimed at overcoming it. Using the equivalence between upper Tsirelson estimates and bounds on the Szlenk index (namely [FOSZ, Theorem 1.3] and [OSZ2, Theorem 21]), we obtain the following corollaries as immediate applications of Theorems 1.6 and 1.7 (the necessary definitions will be given in Section 3).

COROLLARY 1.8. Let $(x_i, f_i)_{i=1}^{\infty}$ be a shrinking Schauder frame for a Banach space X and let α be a countable ordinal. Then the following are equivalent:

- (a) X has Szlenk index at most $\omega^{\alpha\omega}$.
- (b) X satisfies subsequential $T_{\alpha,c}$ -upper tree estimates for some constant 0 < c < 1, where $T_{\alpha,c}$ is the Tsirelson space of order α and constant c.
- (c) $(x_i, f_i)_{i=1}^{\infty}$ has an associated shrinking basis $(z_i)_{i=1}^{\infty}$ such that there exist $(n_i)_{i=1}^{\infty}, (K_i)_{i=1}^{\infty} \in [\mathbb{N}]^{\omega}$ and 0 < c < 1 such that the FDD $(\operatorname{span}_{j\in[n_i,n_{i+1})}z_i)_{i=1}^{\infty}$ satisfies subsequential $(t_{K_i})_{i=1}^{\infty}$ -upper block estimates, where $(t_i)_{i=1}^{\infty}$ is the unit vector basis for $T_{\alpha,c}$.
- (d) $(x_i, f_i)_{i=1}^{\infty}$ has an associated Banach space with a shrinking basis and Szlenk index at most $\omega^{\alpha\omega}$.

COROLLARY 1.9. Let $(x_i, f_i)_{i=1}^{\infty}$ be a shrinking and boundedly complete Schauder frame for a Banach space X and let α be a countable ordinal. Then the following are equivalent:

- (a) X and X^* both have Szlenk index at most $\omega^{\alpha\omega}$.
- (b) X satisfies subsequential $T_{\alpha,c}$ -upper tree estimates and subsequential $T^*_{\alpha,c}$ -lower tree estimates for some constant 0 < c < 1.
- (c) $(x_i, f_i)_{i=1}^{\infty}$ has an associated shrinking and boundedly complete basis $(z_i)_{i=1}^{\infty}$ such that there exist $(n_i)_{i=1}^{\infty}, (K_i)_{i=1}^{\infty} \in [\mathbb{N}]^{\omega}$ and $0 < \infty$ c < 1 such that the FDD $(\operatorname{span}_{j \in [n_i, n_{i+1})} z_i)_{i=1}^{\infty}$ satisfies subsequential $(t_{K_i})_{i=1}^{\infty}$ upper block estimates and subsequential $(t_{K_i}^*)_{i=1}^{\infty}$ -lower block estimates, where $(t_i)_{i=1}^{\infty}$ is the unit vector basis for $T_{\alpha,c}$.
- (d) $(x_i, f_i)_{i=1}^{\infty}$ has an associated reflexive Banach space Z such that Z and Z^{*} both have Szlenk index at most $\omega^{\alpha\omega}$.

Both Schauder frames and atomic decompositions are natural extensions of frame theory into the study of Banach space. These two concepts are directly related, and some papers in the area, such as [CHL], [CL], and [CLS], are stated in terms of atomic decompositions, while others, such as [CDOSZ], [L], and [LZ], are stated in terms of Schauder frames.

DEFINITION 1.10. Let X be a Banach space and Z be a Banach sequence space. We say that a sequence of pairs $(x_i, f_i)_{i=1}^{\infty} \subset X \times X^*$ is an *atomic* decomposition of X with respect to Z if there exist positive constants Aand B such that for all $x \in X$:

- (a) $(f_i(x_i))_{i=1}^{\infty} \in \mathbb{Z},$
- (b) $A||x|| \leq ||(f_i(x_i))_{i=1}^{\infty}||_Z \leq B||x||,$ (c) $x = \sum_{i=1}^{\infty} f_i(x)x_i.$

If the unit vectors in the Banach space Z given in Definition 1.10 form a basis for Z, then an atomic decomposition is simply a Schauder frame with a specified associated space Z. We choose to use the terminology of Schauder frames for this paper instead of atomic decomposition, as, to us, an associated space is an object which is useful for studying the Schauder frame but is external to the space X and frame $(x_i, f_i)_{i=1}^{\infty}$. Our goals are, essentially, to construct "nice" associated spaces, given a particular Schauder frame. However, our theorems can be stated in terms of atomic decompositions. In particular, Theorem 1.5 can be formulated as follows.

THEOREM 1.11. Let X be a Banach space and Z be a Banach sequence space whose unit vectors form a basis for Z. Let $(x_i, f_i)_{i=1}^{\infty}$ be an atomic decomposition of X with respect to Z. Then $(x_i, f_i)_{i=1}^{\infty}$ is shrinking if and only if there exists a Banach sequence space Z' whose unit vectors form a shrinking basis for Z' such that $(x_i, f_i)_{i=1}^{\infty}$ is an atomic decomposition of X with respect to Z'. Furthermore, $(x_i, f_i)_{i=1}^{\infty}$ is shrinking and boundedly complete if and only if there exists a reflexive Banach sequence space Z' whose unit vectors form a basis for Z' such that $(x_i, f_i)_{i=1}^{\infty}$ is an atomic decomposition of X with respect to Z'.

2. Shrinking and boundedly complete Schauder frames. It is well known that a basis (x_i) for a Banach space X is shrinking if and only if the biorthogonal functionals (x_i^*) form a boundedly complete basis for X^* . The following theorem extends this useful characterization to Schauder frames.

THEOREM 2.1 ([CL, Proposition 2.3], [L, Proposition 4.8]). Let X be a Banach space with a Schauder frame $(x_i, f_i)_{i=1}^{\infty} \subset X \times X^*$. The frame $(x_i, f_i)_{i=1}^{\infty}$ is shrinking if and only if $(f_i, x_i)_{i=1}^{\infty}$ is a boundedly complete Schauder frame for X^* .

It is a classic and fundamental result of James that a basis for a Banach space is both shrinking and boundedly complete if and only if the Banach space is reflexive. The following theorem shows that one side of James' characterization holds for frames.

THEOREM 2.2 ([CL, Proposition 2.4], [L, Proposition 4.9]). If $(x_i, f_i)_{i=1}^{\infty}$ is a shrinking and boundedly complete Schauder frame of a Banach space X, then X is reflexive.

It was left as an open question in [CL] whether the converse of Theorem 2.2 holds. The following theorem shows that this is false for any Banach space X, and is evidence of how general Schauder frames can exhibit fairly unintuitive structure.

THEOREM 2.3. Let X be a Banach space which admits a Schauder frame (i.e. has the bounded approximation property). Then X has a Schauder frame which is not shrinking.

Proof. Let $(x_i, f_i)_{i=1}^{\infty}$ be a Schauder frame for X. If $(x_i, f_i)_{i=1}^{\infty}$ is not shrinking, then we are done. Thus we assume that $(x_i, f_i)_{i=1}^{\infty}$ is shrinking. Fix $x \in X$ such that $x \neq 0$, and choose $(y_i^*)_{i=1}^{\infty} \subset S_{X^*}$ such that $y_i^* \to_{w^*} 0$. For all $n \in \mathbb{N}$, we define elements $(x'_{3n-2}, f'_{3n-2}), (x'_{3n-1}, f'_{3n-1}), (x'_{3n}, f'_{3n}) \in X \times X^*$ in the following way:

$$\bar{x}_{3n-2} = x_n, \quad \bar{x}_{3n-1} = x, \quad \bar{x}_{3n} = x, \\ \bar{f}_{3n-2} = f_n, \quad \bar{f}_{3n-1} = -y_n^*, \quad \bar{f}_{3n} = y_n^*$$

As $y_i^* \to_{w^*} 0$, it is not difficult to see that $(\bar{x}_i, \bar{f}_i)_{i=1}^{\infty}$ is a frame for X. However, $(\bar{x}_i, \bar{f}_i)_{i=1}^{\infty}$ is not shrinking. Indeed, let $x^* \in X^*$ be such that $x^*(x) = 1$. As $(x_i, f_i)_{i=1}^{\infty}$ is shrinking, there exists $N_0 \in \mathbb{N}$ such that $|\sum_{i=M}^{\infty} x^*(x_i) f_i(y)| \le \frac{1}{4} ||y||$ for all $y \in X$ and $M \ge N_0$. Let $M \ge N_0$ and choose $y \in B_X$ such that $y_M^*(y) > 3/4$. We now have the following estimate:

$$\|x^* \circ T_{3M}\| \ge x^* \circ T_{3M}(y) = x^*(x)y^*_M(y) + \sum_{i=M+1}^{\infty} f_i(y)x^*(x_i) > \frac{3}{4} - \frac{1}{4} = \frac{1}{2}$$

Thus $||x^* \circ T_N|| \not\rightarrow 0$, and hence $(\bar{x}_i, \bar{f}_i)_{i=1}^{\infty}$ is not shrinking.

As a Schauder frame must be shrinking in order to have a shrinking associated basis, Theorem 2.3 implies that not every Schauder frame for a reflexive Banach space has a reflexive associated space.

DEFINITION 2.4. If $(x_i, f_i)_{i=1}^{\infty}$ is a Schauder frame for a Banach space X and $(z_i)_{i=1}^{\infty}$ is an associated basis, then $(x_i, f_i)_{i=1}^{\infty}$ is called *strongly shrinking* relative to $(z_i)_{i=1}^{\infty}$ if

$$||x^* \circ S_n|| \to 0$$
 for all $x^* \in X^*$,

where $S_n: Z \to X$ is defined by $S_n(z) = \sum_{i=n}^{\infty} z_i^*(z) x_i$.

It is clear that if a Schauder frame is strongly shrinking relative to some associated basis, then the Schauder frame must be shrinking. Also, if a Schauder frame has a shrinking associated basis, then the frame is strongly shrinking relative to the basis. In [CL], examples of shrinking Schauder frames are given which are not strongly shrinking relative to some given associated spaces. However, we will prove later that for any given shrinking Schauder frame, there exists an associated basis such that the frame is strongly shrinking relative to the basis. Before proving this, we state the following theorem which illustrates why the concept of strongly shrinking will be important to us, and allows us to use frames in duality arguments.

THEOREM 2.5 ([CL, Lemma 1.7, Theorem 1.8]). If $(x_i, f_i)_{i=1}^{\infty}$ is a Schauder frame for a Banach space X, and $(z_i)_{i=1}^{\infty}$ is an associated basis for $(x_i, f_i)_{i=1}^{\infty}$, then $(z_i^*)_{i=1}^{\infty}$ is an associated basis for $(f_i, x_i)_{i=1}^{\infty}$ if and only if $(x_i, f_i)_{i=1}^{\infty}$ is strongly shrinking relative to $(z_i)_{i=1}^{\infty}$. Furthermore, given operators $T: X \to Z$ and $S: Z \to X$ defined by $T(x) = \sum f_i(x)z_i$ for all $x \in X$ and $S(z) = \sum z_i^*(z)x_i$ for all $z \in Z$, if $(x_i, f_i)_{i=1}^{\infty}$ is strongly shrinking relative to $(z_i)_{i=1}^{\infty}$, then $S^*: X^* \to [z_i^*]$ and $T^*: [z_i^*] \to X^*$ are given by $S^*(x^*) = \sum x^*(x_i)z_i^*$ for all $x^* \in X^*$ and $T^*(z^*) = \sum z^*(z_i)f_i$ for all $z^* \in [z_i^*]$.

Applying Theorem 2.5 to reflexive Banach spaces gives the following corollary.

COROLLARY 2.6. If $(x_i, f_i)_{i=1}^{\infty}$ is a shrinking frame for a reflexive Banach space X and $(z_i)_{i=1}^{\infty}$ is an associated basis such that $(x_i, f_i)_{i=1}^{\infty}$ is strongly shrinking relative to $(z_i)_{i=1}^{\infty}$, then $(f_i, x_i)_{i=1}^{\infty}$ is strongly shrinking relative to $(z_i^*)_{i=1}^{\infty}$.

Before proceeding further, we need some stability lemmas. Note that if $(z_i)_{i=1}^{\infty}$ is a basis for a Banach space Z, with projection operators $P_{(n,k)} : Z \to Z$ given by $P_{(n,k)}(\sum a_i z_i) = \sum_{i \in (n,k)} a_i z_i$, then $P_{(1,k)} \circ P_{(n,\infty)} = 0$ and $P_{(n,\infty)} \circ P_{(1,k)} = 0$ for all k < n. The analogous property fails when working with frames. However, the following lemmas will essentially allow us to obtain this property within some given $\varepsilon > 0$ if n is chosen sufficiently larger than k.

LEMMA 2.7. Let $(x_i, f_i)_{i=1}^{\infty}$ be a Schauder frame for a Banach space X. Then for all $\varepsilon > 0$ and $k \in \mathbb{N}$, there exists $N \in \mathbb{N}$ such that N > k and

$$\sup_{n \ge m \ge N > k \ge n_0 \ge m_0} \left\| \sum_{i=m}^n f_i \left(\sum_{j=m_0}^{n_0} f_j(x) x_j \right) x_i \right\| \le \varepsilon \|x\| \quad \text{for all } x \in X.$$

Proof. As $(x_i, f_i)_{i=1}^{\infty}$ is a Schauder frame, for each $1 \leq \ell \leq k$ with $f_\ell \neq 0$, there exists $N_\ell > k$ such that $\sup_{n \geq m \geq N_\ell} \|\sum_{i=m}^n f_i(x_\ell) x_i\| < \varepsilon/(k \|f_\ell\|)$. Let $N = \max_{1 \leq \ell \leq k} N_\ell$. We now obtain the following estimate for $n \geq m \geq N > k \geq n_0 \geq m_0$ and $x \in X$:

$$\begin{split} \left\|\sum_{i=m}^{n}\sum_{j=m_{0}}^{n_{0}}f_{j}(x)f_{i}(x_{j})x_{i}\right\| &\leq k \max_{1\leq\ell\leq k}\left\|\sum_{i=m}^{n}f_{\ell}(x)f_{i}(x_{\ell})x_{i}\right\| \quad \text{as } k\geq n_{0} \\ &\leq k \max_{1\leq\ell\leq k}\left\|\sum_{i=m}^{n}f_{i}(x_{\ell})x_{i}\right\|\left\|f_{\ell}\right\|\left\|x\right\|\leq \varepsilon\|x\| \quad \text{as } n\geq m\geq N_{\ell}. \end{split}$$

In terms of operators, Lemma 2.7 can be stated as follows: for all $k \in \mathbb{N}$, if $(x_i, f_i)_{i=1}^{\infty}$ is a Schauder frame then $\lim_{N\to\infty} ||T_N \circ (\mathrm{Id}_X - T_k)|| = 0$, where $T_n : X \to X$ is given by $T_n(x) = \sum_{i=n}^{\infty} f_i(x)x_i$ for all $n \in \mathbb{N}$. We now prove that for all $k \in \mathbb{N}$, if $(x_i, f_i)_{i=1}^{\infty}$ is a shrinking Schauder frame then $\lim_{N\to\infty} ||(\mathrm{Id}_X - T_k) \circ T_N|| = 0$. The frame given in the proof of Theorem 2.3 shows that we cannot drop the condition of shrinking. LEMMA 2.8. Let $(x_i, f_i)_{i=1}^{\infty}$ be a shrinking Schauder frame for a Banach space X. Then for all $\varepsilon > 0$ and $k \in \mathbb{N}$, there exists $N \in \mathbb{N}$ such that N > kand

$$\sup_{n \ge m \ge N > k \ge n_0 \ge m_0} \left\| \sum_{i=m_0}^{n_0} f_i \left(\sum_{j=m}^n f_j(x) x_j \right) x_i \right\| \le \varepsilon \|x\| \quad \text{for all } x \in X.$$

Proof. By Theorem 2.1, $(f_i, x_i)_{i=1}^{\infty}$ is a Schauder frame for X^* . Thus for each $1 \leq \ell \leq k$ with $x_\ell \neq 0$, there exists an integer $N_\ell > k$ such that $\sup_{n \geq m \geq N_\ell} \|\sum_{j=m}^n f_\ell(x_j)f_j\| < \varepsilon/(k\|x_\ell\|)$. Let $N = \max_{1 \leq \ell \leq k} N_\ell$. We now obtain the following estimate for $n \geq m \geq N > k \geq n_0 \geq m_0$ and $x \in X$:

$$\begin{split} \left\| \sum_{i=m_0}^{n_0} f_i \Big(\sum_{j=m}^n f_j(x) x_j \Big) x_i \right\| \\ &\leq k \sup_{1 \leq \ell \leq k} \left\| f_\ell \Big(\sum_{j=m}^n f_j(x) x_j \Big) x_\ell \right\| \quad \text{as } k \geq n_0 \\ &= k \sup_{1 \leq \ell \leq k} \left\| \sum_{j=m}^n f_j(x) f_\ell(x_j) \right\| \|x_\ell\| \\ &\leq k \sup_{1 \leq \ell \leq k} \left\| \sum_{j=m}^n f_\ell(x_j) f_j \right\| \|x\| \|x_\ell\| \leq \varepsilon \|x\| \quad \text{as } n \geq m \geq N_\ell. \end{split}$$

Our method for proving that every shrinking frame has a shrinking associated basis is to first prove that every shrinking frame is strongly shrinking with respect to some associated basis, and then renorm that associated basis to make it shrinking. The following theorem is thus our first major step.

THEOREM 2.9. Let $(x_i, f_i)_{i=1}^{\infty}$ be a shrinking Schauder frame for a Banach space X. Then $(x_i, f_i)_{i=1}^{\infty}$ has an associated basis $(z_i)_{i=1}^{\infty}$ such that $(x_i, f_i)_{i=1}^{\infty}$ is strongly shrinking relative to $(z_i)_{i=1}^{\infty}$.

Proof. We repeatedly apply Lemma 2.8 to obtain a subsequence $(N_k)_{k=1}^{\infty}$ of \mathbb{N} such that for all $k \in \mathbb{N}$,

(3)
$$\sup_{n \ge m \ge N_k} \left\| \sum_{i=1}^k f_i \Big(\sum_{j=m}^n f_j(x) x_j \Big) x_i \right\| \le 2^{-2k} \|x\| \quad \text{for all } x \in X$$

We assume without loss of generality that $x_i \neq 0$ for all $i \in \mathbb{N}$. We denote the unit vector basis of c_{00} by $(z_i)_{i=1}^{\infty}$, and define the following norm $\|\cdot\|_Z$ for all $(a_i) \in c_{00}$:

(4)
$$\left\|\sum a_i z_i\right\|_Z = \max_{n \ge m} \left\|\sum_{i=m}^n a_i x_i\right\| \vee \max_{k \in \mathbb{N}, n \ge m \ge N_k} 2^k \left\|\sum_{i=1}^k f_i\left(\sum_{j=m}^n a_j x_j\right) x_i\right\|.$$

K. Beanland et al.

It follows easily that $(z_i)_{i=1}^{\infty}$ is a bimonotone basic sequence, and thus $(z_i)_{i=1}^{\infty}$ is a bimonotone basis for the completion of c_{00} under $\|\cdot\|_Z$, which we denote by Z. We first prove that $(z_i)_{i=1}^{\infty}$ is an associated basis for $(x_i, f_i)_{i=1}^{\infty}$. Let C be the frame constant of $(x_i, f_i)_{i=1}^{\infty}$, that is, $\max_{n\geq m} \|\sum_{i=m}^n f_i(x)x_i\| \leq C \|x\|$ for all $x \in X$. By (3) and (4), the operator $T: X \to Z$, defined by $T(x) = \sum f_i(x)z_i$ for all $x \in X$, is bounded and $\|T\| \leq C$. We know that $\|\sum_{i=m}^n a_i z_i\|_Z \geq \|\sum_{i=m}^n a_i x_i\|$, and hence the operator $S: Z \to X$ defined by $S(z) = \sum z_i^*(z)x_i$ is bounded and $\|S\| = 1$. Thus $(z_i)_{i=1}^{\infty}$ is an associated basis for $(x_i, f_i)_{i=1}^{\infty}$.

We now prove that $(x_i, f_i)_{i=1}^{\infty}$ is strongly shrinking relative to $(z_i)_{i=1}^{\infty}$. Let $\varepsilon > 0$ and $x^* \in B_{X^*}$. As $(x_i, f_i)_{i=1}^{\infty}$ is shrinking, we may choose $k \in \mathbb{N}$ such that $2^{-k} < \varepsilon/2$ and $\|\sum_{j=k+1}^{\infty} x^*(x_j) f_j\| < \varepsilon/2$. We obtain the following estimate for any $N \ge N_k$ and $z = \sum a_i z_i \in Z$:

$$x^* \left(\sum_{i=N}^{\infty} a_i x_i\right) = \sum_{j=1}^{\infty} x^* (x_j) f_j \left(\sum_{i=N}^{\infty} a_i x_i\right) \quad \text{as } (f_i, x_i)_{i=1}^{\infty} \text{ is a frame for } X^*$$
$$= \sum_{j=1}^k x^* (x_j) f_j \left(\sum_{i=N}^{\infty} a_i x_i\right) + \sum_{j=k+1}^{\infty} x^* (x_j) f_j \left(\sum_{i=N}^{\infty} a_i x_i\right)$$
$$\leq \|x^*\| \left\| \sum_{j=1}^k f_j \left(\sum_{i=N}^{\infty} a_i x_i\right) x_j \right\| + \left\| \sum_{j=k+1}^{\infty} x^* (x_j) f_j \right\| \|z\|_Z$$
$$\leq \|x^*\| 2^{-k} \|z\|_Z + \left\| \sum_{j=k+1}^{\infty} x^* (x_j) f_j \right\| \|z\|_Z \quad \text{by } (4) \text{ as } N \geq N_k$$
$$< (\varepsilon/2) \|z\|_Z + (\varepsilon/2) \|z\|_Z.$$

Thus, for all $x^* \in X^*$ and $\varepsilon > 0$, there exists $M \in \mathbb{N}$ such that $|x^*(\sum_{i=N}^{\infty} z_i^*(z)x_i)| < \varepsilon$ for all $N \ge M$ and $z^* \in B_{Z^*}$. Hence, $(x_i, f_i)_{i=1}^{\infty}$ is strongly shrinking relative to $(z_i)_{i=1}^{\infty}$.

The following lemmas incorporate an associated basis into the tail and initial segment estimates of Lemmas 2.7 and 2.8.

LEMMA 2.10. Let X be a Banach space with a shrinking Schauder frame $(x_i, f_i)_{i=1}^{\infty}$. Let Z be a Banach space with a basis $(z_i)_{i=1}^{\infty}$ such that $(x_i, f_i)_{i=1}^{\infty}$ is strongly shrinking relative to $(z_i)_{i=1}^{\infty}$. Then for all $k \in \mathbb{N}$ and $\varepsilon > 0$, there exists $N \in \mathbb{N}$ such that

$$\sup_{k\geq n\geq m} \left\|\sum_{i=N}^{\infty} \left(\sum_{j=m}^{n} x^*(x_j) f_j(x_i)\right) z_i^*\right\| < \varepsilon \|x^*\| \quad \text{for all } x^* \in X^*.$$

Proof. Let $k \in \mathbb{N}$ and $\varepsilon > 0$. By renorming Z, we may assume without loss of generality that $(z_i)_{i=1}^{\infty}$ is bimonotone. Let $K \ge 1$ be the frame constant of the frame $(f_i, x_i)_{i=1}^{\infty}$ for X^{*}. We choose a finite $\frac{\varepsilon}{2\|S\|}$ -net $(y_{\alpha}^*)_{\alpha \in A}$ in

 $\begin{cases} y^* \in K \cdot B_{Y^*} : y^* \in \operatorname{span}_{1 \leq i \leq k} f_i \}. \text{ By Theorem 2.5, the bounded operator} \\ S^* : X^* \to [z_i^*] \text{ is given by } S^*(x^*) = \sum_{i=1}^{\infty} x^*(x_i) z_i^* \text{ for all } x^* \in X^*. \text{ As} \\ (z_i^*)_{i=1}^{\infty} \text{ is a basis for } [z_i^*]_{i=1}^{\infty}, \text{ for each } \alpha \in A, \text{ there exists } N_{\alpha} \in \mathbb{N} \text{ such that} \\ \|P_{[N_{\alpha},\infty)} \circ S^*(y_{\alpha}^*)\| = \|\sum_{i=N_{\alpha}}^{\infty} y_{\alpha}^*(x_i) z_i^*\| < \varepsilon/2. \text{ We set } N = \max_{\alpha \in A} N_{\alpha}. \\ \text{Given, } x^* \in B_X^* \text{ and } m, n \in \mathbb{N} \text{ such that } k \geq n \geq m, \text{ we choose } \alpha \in A \text{ for which } \|y_{\alpha}^* - \sum_{j=m}^n x^*(x_j) f_j\| < \varepsilon/(2\|S\|). \\ \end{cases}$

$$\begin{split} \left\| \sum_{i=N}^{\infty} \left(\sum_{j=m}^{n} x^*(x_j) f_j(x_i) \right) z_i^* \right\| &= \left\| P_{[N,\infty)} \circ S^* \left(\sum_{j=m}^{n} x^*(x_j) f_j \right) \right\| \\ &\leq \left\| P_{[N,\infty)} \circ S^*(y_\alpha^*) \right\| + \left\| P_{[N,\infty)} \circ S^* \left(y_\alpha^* - \sum_{j=m}^{n} x^*(x_j) f_j \right) \right\| \\ &\leq \left\| P_{[N,\infty)} \circ S^*(y_\alpha^*) \right\| + \left\| P_{[N,\infty)} \right\| \left\| S \right\| \left\| y_\alpha^* - \sum_{j=m}^{n} x^*(x_j) f_j \right\| \\ &< \frac{\varepsilon}{2} + \left\| S \right\| \frac{\varepsilon}{2\|S\|} = \varepsilon. \quad \bullet \end{split}$$

LEMMA 2.11. Let X be a Banach space with a shrinking Schauder frame $(x_i, f_i)_{i=1}^{\infty}$, and let Z be a Banach space with a basis $(z_i)_{i=1}^{\infty}$. Then for all $k \in \mathbb{N}$ and $\varepsilon > 0$, there exists $N \in \mathbb{N}$ such that

$$\sup_{n \ge m \ge N} \left\| \sum_{i=1}^k \left(\sum_{j=m}^n x^*(x_j) f_j(x_i) \right) z_i^* \right\| < \varepsilon \|x^*\| \quad \text{for all } x^* \in X^*.$$

Proof. Let $k \in \mathbb{N}$ and $\varepsilon > 0$. As $(x_j, f_j)_{j=1}^{\infty}$ is a Schauder frame for X, the series $\sum_{j=1}^{\infty} f_j(x_i) x_j$ converges in norm to x_i for all $i \in \mathbb{N}$. Thus there exists $N \in \mathbb{N}$ such that $\sup_{n \ge m \ge N} \|\sum_{j=m}^n f_j(x_i) x_j\| < \varepsilon/(k \|z_i^*\|)$ for all $1 \le i \le k$. For $x^* \in B_{X^*}$ and $n \ge m \ge N$, we have

$$\left\|\sum_{i=1}^{k} \left(\sum_{j=m}^{n} x^*(x_j) f_j(x_i)\right) z_i^*\right\| \le \sum_{i=1}^{k} \left\|\sum_{j=m}^{n} f_j(x_i) x_j\right\| \|z_i^*\|$$
$$< \sum_{i=1}^{k} \frac{\varepsilon}{k \|z_i^*\|} \|z_i^*\| = \varepsilon. \quad \bullet$$

The following lemma and theorem are based on an idea of W. B. Johnson [J], and are analogous to Proposition 3.1 in [FOSZ], and Lemma 4.3 in [OS1]. Their importance comes from allowing us to use arguments that require "skipping coordinates", and in particular, will allow us to apply Proposition 2.14.

LEMMA 2.12. Let X be a Banach space with a boundedly complete Schauder frame $(x_i, f_i)_{i=1}^{\infty} \subset X \times X^*$. Let $\varepsilon_i \searrow 0$ and $(p_i)_{i=1}^{\infty} \in [\mathbb{N}]^{\omega}$. There exists $(k_i)_{i=1}^{\infty} \in [\mathbb{N}]^{\omega}$ such that for all $x^{**} \in X^{**}$ and all $N \in \mathbb{N}$ there exists $M \in \mathbb{N}$ such that $k_N < M < k_{N+1}$ and

$$\sup_{p_{M+1} > n \ge m \ge p_{M-1}} \left\| \sum_{i=m}^{n} x^{**}(f_i) x_i \right\| < \varepsilon_N \|x^{**}\|.$$

Proof. Assume not; then there exist $\varepsilon > 0$ and $K_0 \in \mathbb{N}$ such that for all $K > K_0$ there exists $x_K^{**} \in B_{X^{**}}$ such that for all $K_0 < M < K$ there exists $n_{K,M}, m_{K,M} \in \mathbb{N}$ with $p_{M-1} \leq m_{K,M} \leq n_{K,M} < p_{M+1}$ and $\|\sum_{i=m_{K,M}}^{n_{K,M}} x_K^{**}(f_i)x_i\| > \varepsilon$. As $[p_{M-1}, p_{M+1}]$ is finite, we may choose a sequence $(K_i)_{i=1}^{\infty} \in [\mathbb{N}]^{\omega}$ such that for every $M \in \mathbb{N}$ there exist $n_M, m_M \in \mathbb{N}$ such that $n_{K_i,M} = n_M$ and $m_{K_i,M} = m_M$ for all $i \geq M$. After passing to a further subsequence of $(K_i)_{i=1}^{\infty}$, we may assume that there exists $x^{**} \in X^{**}$ such that $x_{K_i}^{**}(f_j) \to x^{**}(f_j)$ for all $j \in \mathbb{N}$. Thus $\|\sum_{i=m_M}^{n_M} x^{**}(f_i)x_i\| \geq \varepsilon$. This contradicts the series $\sum_{i=1}^{\infty} x^{**}(f_i)x_i$ being norm convergent.

THEOREM 2.13. Let X be a Banach space with a shrinking Schauder frame $(x_i, f_i)_{i=1}^{\infty}$. Let Z be a Banach space with a basis $(z_i)_{i=1}^{\infty}$ such that $(x_i, f_i)_{i=1}^{\infty}$ is strongly shrinking relative to $(z_i)_{i=1}^{\infty}$. Let $(p_i)_{i=1}^{\infty} \in [\mathbb{N}]^{\omega}$ and $(\delta_i)_{i=1}^{\infty} \subset (0, 1)$ with $\delta_i \searrow 0$. Then there exist $(q_i)_{i=1}^{\infty}, (N_i)_{i=1}^{\infty} \in [\mathbb{N}]^{\omega}$ such that for any $(k_i)_{i=0}^{\infty} \in [\mathbb{N}]^{\omega}$ and $y^* \in S_{X^*}$, there exist $y_i^* \in X^*$ and $t_i \in$ $(N_{k_{i-1}-1}, N_{k_{i-1}})$ for all $i \in \mathbb{N}$ with $N_0 = 0$ and $t_0 = 0$ such that:

(a) $y^* = \sum_{i=1}^{\infty} y_i^*$

and for all $\ell \in \mathbb{N}$ we have

 $\begin{array}{ll} \text{(b)} & either \|y_{\ell}^{*}\| < \delta_{\ell}, \ or \ \sup_{p_{q_{t_{\ell-1}} \ge n \ge m}} \|\sum_{j=m}^{n} y_{\ell}^{*}(x_{j})f_{j}\| < \delta_{\ell}\|y_{\ell}^{*}\| \ and \\ & \sup_{n \ge m \ge p_{q_{t_{\ell}}}} \|\sum_{j=m}^{n} y_{\ell}^{*}(x_{j})f_{j}\| < \delta_{\ell}\|y_{\ell}^{*}\|, \\ \text{(c)} & \|P_{[p_{q_{N_{k_{\ell}}}}, p_{q_{N_{k_{\ell+1}}}})} \circ S^{*}(y_{\ell-1}^{*} + y_{\ell}^{*} + y_{\ell+1}^{*} - y^{*})\|_{Z^{*}} < \delta_{\ell}, \end{array}$

where P_I is the projection operator $P_I : [z_i^*] \to [z_i^*]$ given by $P_I(\sum a_i z_i^*) = \sum_{i \in I} a_i z_i^*$ for all $\sum a_i z_i^* \in [z_i^*]$ and all intervals $I \subseteq \mathbb{N}$.

Proof. By Theorems 2.1 and 2.5, $(f_i, x_i)_{i=1}^{\infty}$ is a boundedly complete frame for X^* with associated basis $(z_i^*)_{i=1}^{\infty}$. After renorming, we may assume without loss of generality that $(z_i)_{i=1}^{\infty}$ is bimonotone. We let K be the frame constant of $(f_i, x_i)_{i=1}^{\infty}$. Let $\varepsilon_i \searrow 0$ be such that $2\varepsilon_{i+1} < \varepsilon_i < \delta_i$ and $(1 + K)\varepsilon_i$ $< \delta_{i+1}^2$ for all $i \in \mathbb{N}$.

By repeatedly applying Lemma 2.8 to the frame $(x_i, f_i)_{i=1}^{\infty}$ of X, we may choose $(q_k)_{k=1}^{\infty} \in [\mathbb{N}]^{\omega}$ such that for all $k \in \mathbb{N}$,

(5)
$$\sup_{n \ge m \ge p_{q_{k+1}} > p_{q_k} \ge n_0 \ge m_0} \left\| \sum_{i=m_0}^{n_0} f_i \left(\sum_{j=m}^n f_j(x) x_j \right) x_i \right\| \le \varepsilon_k \|x\| \quad \text{for all } x \in X.$$

By Lemma 2.11, after possibly passing to a subsequence of $(q_k)_{k=1}^{\infty}$, we may assume that for all $k \in \mathbb{N}$,

(6)
$$\sup_{n \ge m \ge p_{q_{k+1}}} \left\| \sum_{i=1}^{p_{q_k}} \left(\sum_{j=m}^n x^*(x_j) f_j(x_i) \right) z_i^* \right\| \le \varepsilon_k \|x^*\| \quad \text{for all } x^* \in X^*.$$

By applying Lemma 2.7 to the frame $(x_i, f_i)_{i=1}^{\infty}$ of X, after possibly passing to a subsequence of $(q_k)_{k=1}^{\infty}$, we may assume that for all $k \in \mathbb{N}$,

(7)
$$\sup_{n \ge m \ge p_{q_{k+1}} > p_{q_k} \ge n_0 \ge m_0} \left\| \sum_{i=m}^n f_i \Big(\sum_{j=m_0}^{n_0} f_j(x) x_j \Big) x_i \right\| \le \varepsilon_{k+1} \|x\| \quad \text{for all } x \in X.$$

By Lemma 2.10, after possibly passing to a subsequence of $(q_k)_{k=1}^{\infty}$, we may assume that for all $k \in \mathbb{N}$,

(8)
$$\sup_{p_{q_k} > n \ge m \ge 1} \left\| \sum_{i=p_{q_{k+1}}}^{\infty} \left(\sum_{j=m}^n x^*(x_j) f_j(x_i) \right) z_i^* \right\| \le \varepsilon_{k+1} \|x^*\| \quad \text{for all } x^* \in X^*.$$

By Lemma 2.12, there exists $(N_i)_{i=0}^{\infty} \in [\mathbb{N}]^{\omega}$ such that $N_0 = 0$, and for all $x^* \in X^*$ and for all $k \in \mathbb{N}$ there exists $t_k \in \mathbb{N}$ such that $N_k < t_k < N_{k+1}$ and $\sup_{p_{q_{t_k-1}} \leq n \leq m < p_{q_{t_k+1}}} \|\sum_{i=n}^m x^*(x_i)f_i\| < \varepsilon_k \|x^*\|$.

Let $(k_i)_{i=0}^{\infty} \in [\mathbb{N}]^{\omega}$ and let $y^* \in S_{X^*}$. For each $i \in \mathbb{N}$, we choose $t_i \in (N_{k_i}, N_{k_{i+1}})$ with $t_0 = 1$ such that

(9)
$$\sup_{p_{q_{t_i+1}} > n \ge m \ge p_{q_{t_i-1}}} \left\| \sum_{j=m}^n y^*(x_j) f_j \right\| < \varepsilon_i.$$

We now set $y_i^* = \sum_{j=p_{q_{i-1}}}^{p_{q_{i-1}}} y^*(x_j) f_j$ for all $i \in \mathbb{N}$. We obtain

$$\sum_{i=1}^{\infty} y_i^* = \sum_{i=1}^{\infty} \sum_{j=p_{q_{t_i-1}}}^{p_{q_{t_i}}-1} y^*(x_j) f_j = \sum_{j=1}^{\infty} y^*(x_j) f_j = y^*.$$

Thus (a) is satisfied.

In order to prove (b), we let $\ell \in \mathbb{N}$ and assume that $||y_{\ell}^*|| > \delta_{\ell}$. Let $m, n \in \mathbb{N}$ be such that $n \geq m \geq p_{q_{t_{\ell}}}$. To prove property (b), we consider the inequalities

$$\begin{split} \left\|\sum_{j=m}^{n} y_{\ell}^{*}(x_{j})f_{j}\right\| &= \left\|\sum_{j=m}^{n}\sum_{i=p_{q_{t_{\ell}-1}}}^{p_{q_{t_{\ell}}-1}} y^{*}(x_{i})f_{i}(x_{j})f_{j}\right\| \\ &\leq \left\|\sum_{j=m}^{n}\sum_{i=p_{q_{t_{\ell}-1}}}^{p_{q_{t_{\ell}}-1}} y^{*}(x_{i})f_{i}(x_{j})f_{j}\right\| + \left\|\sum_{j=m}^{n}\sum_{i=p_{q_{t_{\ell}-1}}}^{p_{q_{t_{\ell}-1}}-1} y^{*}(x_{i})f_{i}(x_{j})f_{j}\right\| \\ &\leq K \left\|\sum_{i=p_{q_{t_{\ell}-1}}}^{p_{q_{t_{\ell}-1}}-1} y^{*}(x_{i})f_{i}\right\| + \left\|\sum_{j=m}^{n}\sum_{i=p_{q_{t_{\ell}-1}}}^{p_{q_{t_{\ell}-1}}-1} y^{*}(x_{i})f_{i}(x_{j})f_{j}\right\| \\ &< K\varepsilon_{t_{\ell}} + \varepsilon_{t_{\ell}} \quad \text{by (9) and (7)} \\ &< (1+K)\varepsilon_{t_{\ell}}\|y_{\ell}^{*}\|/\delta_{\ell} < (1+K)\varepsilon_{\ell}\|y_{\ell}^{*}\|/\delta_{\ell} < \delta_{\ell}\|y_{\ell}^{*}\|. \end{split}$$

Thus $\sup_{n \ge m \ge p_{q_{\ell_{\ell}}}} \|\sum_{j=m}^{n} y_{\ell}^*(x_j) f_j\| < \delta_{\ell} \|y_{\ell}^*\|$, proving the first of the two inequalities in (b). We now assume that $\ell > 1$, and let $m, n \in \mathbb{N}$ be such that $p_{q_{\ell_{\ell-1}}} \ge n \ge m$. To prove the remaining inequality in (b), we observe that

$$\begin{split} \left\| \sum_{j=m}^{n} y_{\ell}^{*}(x_{j}) f_{j} \right\| &= \left\| \sum_{j=m}^{n} \sum_{i=p_{q_{t_{\ell-1}}}}^{p_{q_{t_{\ell}}}-1} y^{*}(x_{i}) f_{i}(x_{j}) f_{j} \right\| \\ &\leq \left\| \sum_{j=m}^{n} \sum_{i=p_{q_{t_{\ell-1}+1}}}^{p_{q_{t_{\ell-1}+1}}} y^{*}(x_{i}) f_{i}(x_{j}) f_{j} \right\| + \left\| \sum_{j=m}^{n} \sum_{i=p_{q_{t_{\ell-1}}}}^{p_{q_{t_{\ell-1}+1}}-1} y^{*}(x_{i}) f_{i}(x_{j}) f_{j} \right\| \\ &\leq \left\| \sum_{j=m}^{n} \sum_{i=p_{q_{t_{\ell-1}+1}}}^{p_{q_{t_{\ell-1}+1}}} y^{*}(x_{i}) f_{i}(x_{j}) f_{j} \right\| + K \right\| \sum_{i=p_{q_{t_{\ell-1}}}}^{p_{q_{t_{\ell-1}}+1}-1} y^{*}(x_{i}) f_{i} \right\| \\ &< \varepsilon_{t_{\ell-1}+1} + K \varepsilon_{\ell-1} \quad \text{by (5) and (9)} \\ &< (\varepsilon_{t_{\ell-1}+1} + K \varepsilon_{\ell-1}) \| y_{\ell}^{*} \| / \delta_{\ell} < (1+K) \varepsilon_{\ell-1} \| y_{\ell}^{*} \| / \delta_{\ell} < \delta_{\ell} \| y_{\ell}^{*} \|. \end{split}$$

Thus $\sup_{p_{q_{\ell-1}} \ge n \ge m} \|\sum_{j=m}^n y_\ell^*(x_j) f_j\| < \delta_\ell \|y_\ell^*\|$, and hence all of (b) is satisfied.

To prove (c), we note that

$$\begin{split} \left\| P_{[p_{q_{N_{k_{\ell}}}}, p_{q_{N_{k_{\ell+1}}}}]} S^{*}(y_{\ell-1}^{*} + y_{\ell}^{*} + y_{\ell+1}^{*} - y^{*}) \right\|_{Z^{*}} \\ &= \left\| \sum_{i=p_{q_{N_{k_{\ell}}}}}^{p_{q_{N_{k_{\ell+1}}}}-1} (y_{\ell-1}^{*} + y_{\ell}^{*} + y_{\ell+1}^{*} - y^{*})(x_{i}) z_{i}^{*} \right\| \\ &= \left\| \sum_{i=p_{q_{N_{k_{\ell}}}}}^{p_{q_{N_{k_{\ell+1}}}}-1} (\sum_{j=1}^{p_{q_{t_{\ell-2}}}-1} y^{*}(x_{j}) f_{j}(x_{i}) + \sum_{j=p_{q_{t_{\ell+1}}}}^{\infty} y^{*}(x_{j}) f_{j}(x_{i})) z_{i}^{*} \right\| \end{split}$$

$$\leq \Big\| \sum_{i=p_{q_{N_{k_{\ell}+1}}}^{\infty} \Big(\sum_{j=1}^{p_{q_{t_{\ell-2}}}-1} y^{*}(x_{j}) f_{j}(x_{i}) \Big) z_{i}^{*} \Big\| + \Big\| \sum_{i=p_{q_{N_{\ell}}}}^{p_{q_{N_{k_{\ell+1}}}}-1} \Big(\sum_{j=p_{q_{t_{\ell+1}}}}^{\infty} y^{*}(x_{j}) f_{j}(x_{i}) \Big) z_{i}^{*} \Big\| \\ \leq \Big\| \sum_{i=p_{q_{N_{k_{\ell}}}}}^{\infty} \Big(\sum_{j=1}^{p_{q_{t_{\ell-2}}}-1} y^{*}(x_{j}) f_{j}(x_{i}) \Big) z_{i}^{*} \Big\| + \Big\| \sum_{i=1}^{p_{q_{N_{k_{\ell+1}}}}} \Big(\sum_{j=p_{q_{t_{\ell+1}}}}^{\infty} y^{*}(x_{j}) f_{j}(x_{i}) \Big) z_{i}^{*} \Big\| \\ < \varepsilon_{N_{k_{\ell}}-1} + \varepsilon_{N_{k_{\ell}+1}} < \varepsilon_{\ell} \quad \text{by (8) and (6).}$$

Thus (c) is satisfied. \blacksquare

Properties of coordinate systems for Banach spaces such as frames, bases, and FDDs can impose certain structure on infinite-dimensional subspaces. For our purposes, this structure can be intrinsically characterized in terms of even trees of vectors [OSZ1]. In order to index even trees, we define

$$T_{\infty}^{\text{even}} = \{ (n_1, \dots, n_{2\ell}) : n_1 < \dots < n_{2\ell} \text{ are in } \mathbb{N} \text{ and } \ell \in \mathbb{N} \}.$$

If X is a Banach space, an indexed family $(x_{\alpha})_{\alpha \in T_{\infty}^{\text{even}}} \subset X$ is called an *even tree*. Sequences of the form $(x_{(n_1,\ldots,n_{2\ell-1},k)})_{k=n_{2\ell-1}+1}^{\infty}$ are called *nodes*. This should not be confused with the more standard terminology where a node would refer to an individual member of the tree. Sequences of the form $(n_{2\ell-1}, x_{(n_1,\ldots,n_{2\ell})})_{\ell=1}^{\infty}$ are called *branches*. A *normalized tree*, i.e. one with $||x_{\alpha}|| = 1$ for all $\alpha \in T_{\infty}^{\text{even}}$, is called *weakly null* (resp. w^* -null) if every node is a weakly null (resp. w^* -null) sequence.

Given $1 > \varepsilon > 0$ and $A \subset (\mathbb{N} \times S_{X^*})^{\omega}$, we let $A_{\varepsilon} = \{(l_i, y_i^*) \in (\mathbb{N} \times S_{X^*})^{\omega}$: there exists $(k_i, x_i^*) \in A$ such that $k_i \leq \ell_i$, $||x_i^* - y_i^*|| < \varepsilon 2^{-i}$ for all $i \in \mathbb{N}\}$, and we let $\overline{A}_{\varepsilon}$ be the closure of A_{ε} in $(\mathbb{N} \times S_{X^*})^{\omega}$. We consider the following game between players S (subspace chooser) and P (point chooser). The game has an infinite sequence of moves; on the *n*th move S picks $k_n \in \mathbb{N}$ and a finite-codimensional w^* -closed subspace Z_n of X^* and P responds by picking an element $x_n^* \in S_{X^*}$ such that $d(x_n^*, Z_n) < \varepsilon 2^{-n}$. Now, S wins the game if the sequence $(k_i, x_i)_{i=1}^{\infty}$ the players generate is an element of $\overline{A}_{5\varepsilon}$, otherwise P is declared the winner. This is referred to as the (A, ε) -game and was introduced in [OSZ1]. The following proposition is essentially an extension of Proposition 2.6 in [FOSZ] from FDDs to frames, and relates properties of w^* -null even trees and winning strategies of the (A, ε) -game to blockings of a frame.

PROPOSITION 2.14. Let X be an infinite-dimensional Banach space with a shrinking Schauder frame $(x_i, f_i)_{i=1}^{\infty}$. Let $A \subseteq (\mathbb{N} \times S_{X^*})^{\omega}$. The following are equivalent:

(1) For all $\varepsilon > 0$ there exist $(K_i)_{i=1}^{\infty} \in [\mathbb{N}]^{\omega}$ and $\bar{\delta} = (\delta_i) \subset (0,1)$ with $\delta_i \searrow 0$ and $(p_i)_{i=1}^{\infty} \in [\mathbb{N}]^{\omega}$ such that if $(y_i^*)_{i=1}^{\infty} \subset S_{X^*}$ and

 $(r_i)_{i=0}^{\infty} \in [\mathbb{N}]^{\omega}$ satisfy $\sup_{p_{r_i-1}+1 \ge n \ge m \ge 1} \left\| \sum_{j=m}^n y_i^*(x_j) f_j \right\| < \delta_i$ and $\sup_{n\geq m\geq p_{r_i}} \left\|\sum_{j=m}^n y_i^*(x_j)f_j\right\| < \delta_i \text{ for all } i\in\mathbb{N}, \text{ then } (K_{r_{i-1}}, y_i^*)\in\overline{A}_{\varepsilon}.$ (2) For all $\varepsilon > 0$, \tilde{S} has a winning strategy for the (A, ε) -game.

- (3) For all $\varepsilon > 0$ every normalized w^{*}-null even tree in X^{*} has a branch in A_{ε} .

Proof. The equivalence $(2) \Leftrightarrow (3)$ is given in [FOSZ].

1

We assume (1) holds, and will prove (3). Let $(x^*_{(n_1,\ldots,n_{2\ell})})_{(n_1,\ldots,n_{2\ell})\in T_{\infty}^{\text{even}}}$ be a w^* -null even tree in X^* , and let $\varepsilon > 0$.

We shall construct by induction sequences $(r_i)_{i=0}^{\infty}, (n_i)_{i=1}^{\infty} \in [\mathbb{N}]^{\omega}$ such that $K_{r_i} = n_{2i+1}$ and $\sup_{p_{r_{i-1}+1} \ge n \ge m \ge 1} \left\| \sum_{j=m}^n x^*_{(n_1,\dots,n_{2i})}(x_j) f_j \right\| < \delta_i$ and $\sup_{n\geq m\geq p_{r_i}} \left\|\sum_{j=m}^n y_i^*(x_j)f_j\right\| < \delta_i$ for all $i \in \mathbb{N}$. To start, we let $r_0 = 1$ and $n_1 = K_1$. Now, if $\ell \in \mathbb{N}$ and $(r_i)_{i=0}^{\ell}$ and $(n_i)_{i=1}^{2\ell+1}$ have been chosen, then since $(x^*_{(n_1,...,n_{2\ell+1},k)})_{k=n_{2\ell+1}+1}^{\infty}$ is w^* -null, we may choose $n_{2\ell+2} > 0$ $n_{2\ell+1}$ such that $||x^*_{(n_1,\dots,n_{2\ell+1},n_{2\ell+2})}(x_j)f_j|| < (p_{r_\ell}+1)^{-1}\delta_{\ell+1}$. Thus, we have $\sup_{p_{r_{\ell}+1} \ge n \ge m \ge 1} \|\sum_{j=m}^{n} x_{(n_1,\dots,n_{2\ell+1},n_{2\ell+2})}^*(x_j)f_j\| < \delta_{\ell+1}. \text{ As } (x_j,f_j)_{i=1}^{\infty} \text{ is a Schauder frame, we may choose } r_{\ell+1} > r_{\ell} \text{ such that}$

$$\sup_{n \ge m \ge p_{r_{\ell+1}}} \left\| \sum_{j=m}^n x^*_{(n_1,\dots,n_{2\ell+1},n_{2\ell+2})}(x_j) f_j \right\| < \delta_{\ell+1}.$$

We then let $n_{2\ell+2} = K_{r_{\ell+1}}$. Thus, $(r_i)_{i=0}^{\infty}$ and $(n_i)_{i=1}^{\infty}$ may be constructed by induction to have the desired properties, giving $(n_{2i-1}, x^*_{(n_1,\dots,n_{2i})})_{i=1}^{\infty} =$ $(K_{r_{i-1}}, x^*_{(n_1,\dots,n_{2i})})_{i=1}^\infty \in \overline{A}_{\varepsilon}.$

We now assume (2) holds, and will prove (1). Let $\varepsilon > 0$. The fact that player S has a winning strategy for the (A, ε) -game means that there exist an indexed collection $(k_{(x_1^*,...,x_{\ell}^*)})_{(x_1^*,...,x_{\ell}^*)\in X^{<\mathbb{N}}}$ of natural numbers and an $\text{ indexed collection } (X^*_{(x^*_1,\ldots,x^*_\ell)})_{(x^*_1,\ldots,x^*_\ell)\in X^{<\mathbb{N}}} \text{ of cofinite-dimensional } w^*\text{-closed of the set of the set$ subsets of X^* such that if $(x_i^*)_{i=1}^{\infty} \subset S_{X^*}$ and $d(x_i^*, X_{(x_i^*, \dots, x_i^*)}^*) < \frac{1}{10} \varepsilon 2^{-i}$ for all $i \in \mathbb{N}$, then $(k_{(x_1^*, \dots, x_i^*)}, X_{(x_1^*, \dots, x_i^*)}^*)_{i=1}^{\infty} \in \overline{A}_{\varepsilon/2}$ and $(k_{(x_1^*, \dots, x_i^*)})_{i=1}^{\infty} \in [\mathbb{N}]^{\omega}$.

We construct by induction $(K_i)_{i=1}^{\infty}, (p_i)_{i=1}^{\infty} \in [\mathbb{N}]^{\omega}, (\delta_i)_{i=1}^{\infty} \in (0, 1)^{\omega}$, and a nested collection $(D_i)_{i=1}^{\infty} \subset [X^{<\omega}]^{\omega}$ such that D_i is $\frac{1}{20}\varepsilon^{2^{-i}}$ -dense in $[f_j]_{j=1}^{p_i}$ and if $(y_i^*)_{i=1}^{\infty} \subset S_{X^*}$ and $(r_i)_{i=0}^{\infty} \in [\mathbb{N}]^{\omega}$ are such that both $\sup_{p_{r_{i-1}+1} \ge n \ge m \ge 1} \left\| \sum_{j=m}^{n} y_i^*(x_j) f_j \right\| < \delta_i \text{ and } \sup_{n \ge m \ge p_{r_i}} \left\| \sum_{j=m}^{n} y_i^*(x_j) f_j \right\|$ $<\delta_i$ for all $i \in \mathbb{N}$, and $x_i^* \in D_{r_{i+1}}$ are such that $\|y_i^* - x_i^*\| < \frac{1}{20}\varepsilon^{2^{-i}}$ for all $i \in \mathbb{N}$, then $K_{r_{i-1}} \ge k_{(x_1^*, \dots, x_{i-1}^*)}$ and $d(x_i^*, X_{(x_1^*, \dots, x_i^*)}^*) < \frac{1}{10} \varepsilon 2^{-i}$. This would imply that $(k_{(x_1^*,\ldots,x_i^*)}, X_{(x_1^*,\ldots,x_i^*)}^*)_{i=1}^\infty \in \overline{A}_{\varepsilon/2}$. Hence, $(K_{r_{i-1}}, y_i^*)_{i=1}^\infty \in \overline{A}_{\varepsilon}$ as $K_{r_{i-1}} \geq k_{(x_1^*,\dots,x_{i-1}^*)}$ and $\|y_i^* - x_i^*\| < \frac{1}{20}\varepsilon 2^{-i}$ for all $i \in \mathbb{N}$. Thus all that remains is to show that $(K_i)_{i=1}^{\infty}, (p_i)_{i=1}^{\infty} \in [\mathbb{N}]^{\omega}$ and $(D_i)_{i=1}^{\infty} \subset [X^{<\omega}]^{\omega}$ may be constructed inductively with the desired properties.

We start by choosing $K_1 = k_{\emptyset}$. As $(x_i, f_i)_{i=1}^{\infty}$ is a shrinking Schauder frame for X and $X_{\emptyset}^* \subset X^*$ is cofinite-dimensional and w^* -closed, by Lemma 2.8 there exist $p_1 \in \mathbb{N}$ and $\delta_1 > 0$ such that if $\sup_{p_1 \ge n \ge m \ge 1} \|\sum_{j=m}^n y^*(x_j) f_j\|$ $< \delta_1$ for some $y^* \in S_{X^*}$, then $d(y^*, X_{\emptyset}^*) < \frac{1}{20}\varepsilon$. We then let D_1 be some finite $\frac{1}{20}\varepsilon$ -net in $[f_i]_{i=1}^{p_1}$. Now we assume $n \in \mathbb{N}$ and that $(K_i)_{i=1}^n \in [\mathbb{N}]^{<\omega}$, $(p_i)_{i=1}^n \in [\mathbb{N}]^{<\omega}$, $(\delta_i)_{i=1}^n \in (0, 1)^{<\omega}$, and $(D_i)_{i=1}^n \subset [X^{<\omega}]^{<\omega}$ have been suitably chosen. As $(x_i, f_i)_{i=1}^{\infty}$ is a shrinking Schauder frame for X, and $X_{(x_1^*, \dots, x_{\ell}^*)}^* \subset X^*$ is cofinite-dimensional and w^* -closed for all $(x_1^*, \dots, x_{\ell}^*)$ in $[D_n]^{<\omega}$, by Lemma 2.8 there exist $p_{n+1} \in \mathbb{N}$ and $\delta_{n+1} > 0$ such that if $\sup_{p_{n+1} \ge n \ge m \ge 1} \|\sum_{j=m}^n y^*(x_j) f_j\| < \delta_{n+1}$ for some $y^* \in S_{X^*}$, then $d(y^*, \bigcap_{(x_1^*, \dots, x_{\ell}^*) \in [D_n]^{<\omega}} X_{(x_1^*, \dots, x_{\ell}^*)}^*) < \frac{1}{20}\varepsilon 2^{-n-1}$. We then let

$$K_{n+1} = \max_{\substack{(x_1^*, \dots, x_{\ell}^*) \in [D_n] < \omega}} k_{(x_1^*, \dots, x_{\ell}^*)}$$

and let D_{n+1} be a finite $\frac{1}{20}\varepsilon 2^{-n-1}$ -net in $[f_i]_{i=1}^{p_{n+1}}$.

3. Upper and lower estimates. If $(x_i)_{i=1}^{\infty}$ and $(v_i)_{i=1}^{\infty}$ are two sequences in Banach spaces and $C \in \mathbb{R}$, then we say that $(x_i)_{i=1}^{\infty}$ is *C*-dominated by $(v_i)_{i=1}^{\infty}$ if $\|\sum a_i x_i\| \leq C \|\sum a_i v_i\|$ for all $(a_i) \in c_{00}$. Let *Z* be a Banach space with an FDD $(E_i)_{i=1}^{\infty}$, let $V = (v_i)_{i=1}^{\infty}$ be a normalized 1-unconditional basis, and let $1 \leq C < \infty$. We say that $(E_i)_{i=1}^{\infty}$ satisfies subsequential *C*-*V*-upper block estimates if every normalized block sequence $(z_i)_{i=1}^{\infty}$ of $(E_i)_{i=1}^{\infty}$ in *Z* is *C*-dominated by $(v_{m_i})_{i=1}^{\infty}$, where $m_i = \min \operatorname{supp}_E(z_i)$ for all $i \in \mathbb{N}$. We say that $(E_i)_{i=1}^{\infty}$ of $(E_i)_{i=1}^{\infty}$, where $m_i = \min \operatorname{supp}_E(z_i)$ for all $i \in \mathbb{N}$. We say that $(E_i)_{i=1}^{\infty}$ satisfies subsequential *C*-*V*-lower block estimates $(v_{m_i})_{i=1}^{\infty}$, where $m_i = \min \operatorname{supp}_E(z_i)$ for all $i \in \mathbb{N}$. We say that $(E_i)_{i=1}^{\infty}$ satisfies subsequential *C*-*V*-lower block estimates ($v_{m_i})_{i=1}^{\infty}$, where $m_i = \min \operatorname{supp}_E(z_i)$ for all $i \in \mathbb{N}$. We say that $(E_i)_{i=1}^{\infty}$ satisfies subsequential *C*-*V*-lower block estimates ($v_{m_i})_{i=1}^{\infty}$, where $m_i = \min \operatorname{supp}_E(z_i)$ for all $i \in \mathbb{N}$. We say that $(E_i)_{i=1}^{\infty}$ satisfies subsequential *V*-upper (resp. lower) block estimates if it satisfies subsequential *C*-*V*-upper (resp. lower) block estimates for some $1 \leq C < \infty$.

Note that if $(E_i)_{i=1}^{\infty}$ satisfies subsequential *C*-*V*-upper block estimates and $(z_i)_{i=1}^{\infty}$ is a normalized block sequence with $\max \operatorname{supp}_E(z_{i-1}) < k_i \leq \min \operatorname{supp}_E(z_i)$ for all i > 1, then $(z_i)_{i=1}^{\infty}$ is *C*-dominated by $(v_{k_i})_{i=1}^{\infty}$ (and a similar remark holds for lower estimates). The following well known lemma shows why upper estimates are useful for proving theorems about shrinking basic sequences, and why lower estimates are useful for proving theorems about boundedly complete basic sequences.

LEMMA 3.1. Let $(E_i)_{i=1}^{\infty}$ be an FDD and let $(v_i)_{i=1}^{\infty}$ be a normalized basic sequence. If $(v_i)_{i=1}^{\infty}$ is weakly null and $(E_i)_{i=1}^{\infty}$ satisfies subsequential V-upper block estimates, then $(E_i)_{i=1}^{\infty}$ is shrinking. If $(v_i)_{i=1}^{\infty}$ is boundedly complete and $(E_i)_{i=1}^{\infty}$ satisfies subsequential V-lower block estimates, then $(E_i)_{i=1}^{\infty}$ is boundedly complete.

Proof. We first assume that $(v_i)_{i=1}^{\infty}$ is weakly null and $(E_i)_{i=1}^{\infty}$ satisfies subsequential C-V-upper block estimates for some C > 0. Let $(z_i)_{i=1}^{\infty}$ be a normalized block sequence of $(E_i)_{i=1}^{\infty}$. We know that $(z_i)_{i=1}^{\infty}$ is C-dominated by $(v_{m_i})_{i=1}^{\infty}$ where $m_i = \min \operatorname{supp}_E(z_i)$ for all $i \in \mathbb{N}$. Let $\varepsilon > 0$. Because $(v_i)_{i=1}^{\infty}$ is weakly null, there exists $(\lambda_i)_{i=1}^n \subset (0,\infty)$ such that $\sum \lambda_i = 1$ and $\|\sum \lambda_i v_{m_i}\| < \varepsilon/C$. Hence, $\|\sum \lambda_i z_i\| \le C \|\sum \lambda_i v_i\| < \varepsilon$. We infer that 0 is contained in the closed convex hull of every normalized block sequence of (E_i) , and hence (E_i) is shrinking.

We now assume that $(v_i)_{i=1}^{\infty}$ is boundedly complete and $(E_i)_{i=1}^{\infty}$ satisfies subsequential V-lower block estimates. Let $(z_i)_{i=1}^{\infty}$ be a seminormalized block sequence of $(E_i)_{i=1}^{\infty}$. We see that $(v_{m_i})_{i=1}^{\infty}$ is dominated by $(z_i)_{i=1}^{\infty}$ where $m_i = \min \operatorname{supp}_E(z_i)$ for all $i \in \mathbb{N}$. As $(v_{m_i})_{i=1}^{\infty}$ is boundedly complete, $\operatorname{sup}_{n \in \mathbb{N}} \|\sum_{i=1}^n v_{m_i}\| = \infty$. Hence, $\operatorname{sup}_{n \in \mathbb{N}} \|\sum_{i=1}^n z_i\| = \infty$, and thus $(E_i)_{i=1}^{\infty}$ is boundedly complete as well.

Subsequential $V^{(*)}$ -upper block estimates and subsequential V-lower block estimates are dual properties, as shown in the following proposition from [OSZ1].

PROPOSITION 3.2 ([OSZ1, Proposition 2.14]). Assume that Z has an FDD $(E_i)_{i=1}^{\infty}$, and let $V = (v_i)_{i=1}^{\infty}$ be a normalized and 1-unconditional basic sequence. The following statements are equivalent:

- (a) (E_i)[∞]_{i=1} satisfies subsequential V-lower block estimates in Z.
 (b) (E^{*}_i)[∞]_{i=1} satisfies subsequential V^(*)-upper block estimates in Z^(*).

(Here subsequential $V^{(*)}$ -upper estimates are with respect to $(v_i^*)_{i=1}^{\infty}$, the sequence of biorthogonal functionals to $(v_i)_{i=1}^{\infty}$.)

Moreover, if $(E_i)_{i=1}^{\infty}$ is bimonotone in Z, then the equivalence holds true if one replaces, for some $C \ge 1$, V-lower estimates by C-V-lower estimates in (a) and $V^{(*)}$ -upper estimates by C- $V^{(*)}$ -upper estimates in (b).

Note that by duality, Proposition 3.2 holds if we interchange the words "upper" and "lower".

We are interested in creating associated spaces which have coordinate systems satisfying certain upper and lower block estimates. The particular Banach spaces that we start with do not themselves have such coordinate systems, and hence we need to consider intrinsic coordinate-free properties of a Banach space which characterize when a Banach space embeds into a different Banach space with a coordinate system having certain upper and lower block estimates. These intrinsic coordinate-free Banach space properties are defined in terms of even trees. Let X be a Banach space, $V = (v_i)_{i=1}^{\infty}$ be a normalized 1-unconditional basis, and $1 \leq C < \infty$. We say that X satisfies subsequential C-V-upper tree estimates if every weakly null even tree $(x_{\alpha})_{\alpha \in T_{\infty}^{\text{even}}}$ in X has a branch $(n_{2\ell-1}, x_{(n_1, \dots, n_{2\ell})})_{\ell=1}^{\infty}$ such that $(x_{(n_1,\ldots,n_{2\ell})})_{\ell=1}^{\infty}$ is C-dominated by $(v_{n_{2\ell-1}})_{\ell=1}^{\infty}$. We say that X satisfies subsequential V-upper tree estimates if it satisfies subsequential C-V-upper tree estimates for some $1 \leq C < \infty$. If X is a subspace of a dual space, we say that X satisfies subsequential C-V-lower w^* tree estimates if every w^* -null even tree $(x_{\alpha})_{\alpha\in T_{\infty}^{\text{even}}}$ in X has a branch $(n_{2\ell-1}, x_{(n_1,\ldots,n_{2\ell})})_{\ell=1}^{\infty}$ such that $(x_{(n_1,\ldots,n_{2\ell})})_{\ell=1}^{\infty}$ C-dominates $(v_{n_{2\ell-1}})_{\ell=1}^{\infty}$.

A basic sequence $V = (v_i)_{i=1}^{\infty}$ is called *C*-right dominant if for all sequences $m_1 < m_2 < \cdots$ and $n_1 < n_2 < \cdots$ of positive integers with $m_i \leq n_i$ for all $i \in \mathbb{N}$, the sequence $(v_{m_i})_{i=1}^{\infty}$ is *C*-dominated by $(v_{n_i})_{i=1}^{\infty}$. We say that $(v_i)_{i=1}^{\infty}$ is right dominant if for some $C \geq 1$ it is *C*-right dominant.

LEMMA 3.3 ([FOSZ, Lemma 2.7]). Let X be a Banach space with separable dual, and let $V = (v_i)_{i=1}^{\infty}$ be a normalized, 1-unconditional, right dominant basic sequence. If X satisfies subsequential V-upper tree estimates, then X^* satisfies subsequential V*-lower w* tree estimates.

Let $(x_i, f_i)_{i=1}^{\infty}$ be a shrinking Schauder frame for a Banach space X, and let $(v_i)_{i=1}^{\infty}$ be a normalized, 1-unconditional, block stable, 1-right dominant, and shrinking basic sequence. For any C > 0, we may apply Proposition 2.14 to the set $A = \{(n_i, x_i^*)_{i=1}^{\infty} \in (\mathbb{N} \times S_{X^*})^{\omega} : (x_i^*)_{i=1}^{\infty} C$ -dominates $(v_{n_i}^*)_{i=1}^{\infty}\}$ to obtain the following corollary.

COROLLARY 3.4. Let $(x_i, f_i)_{i=1}^{\infty}$ be a shrinking Schauder frame for a Banach space X, and let $V = (v_i)_{i=1}^{\infty}$ be a normalized, 1-unconditional, block stable, right dominant, and shrinking basic sequence. The following are equivalent:

- (1) There exist C > 0, $(K_i)_{i=1}^{\infty}, (p_i)_{i=1}^{\infty} \in [\mathbb{N}]^{\omega}$, and $\overline{\delta} = (\delta_i) \subset (0,1)$ with $\delta_i \searrow 0$ such that if $(y_i^*)_{i=1}^{\infty} \subset S_{X^*}$ and $(r_i)_{i=0}^{\infty} \in [\mathbb{N}]^{\omega}$ are sequences such that both $\sup_{1 \le n \le m \le p_{r_{i-1}+1}} \|\sum_{j=n}^m y_i^*(x_j)f_j\| < \delta_i$ and $\sup_{p_{r_i} \le n \le m} \|\sum_{j=n}^m y_i^*(x_j)f_j\| < \delta_i$ then $(y_i^*) \succeq_C (v_{K_{r_{i-1}}}^*)$.
- (2) X satisfies subsequential V-upper tree estimates.

Let Z be a Banach space with a basis $(z_i)_{i=1}^{\infty}$, let $(p_i)_{i=1}^{\infty} \in [\mathbb{N}]^{\omega}$ with $p_1 = 1$, and let $V = (v_i)_{i=1}^{\infty}$ be a normalized 1-unconditional basic sequence. The space $Z_V(p_i)$ is defined to be the completion of c_{00} with respect to the following norm $\|\cdot\|_{Z_V}$ where for $(a_i)_{i=1}^{\infty} \in c_{00}$:

$$\left\|\sum a_{i}z_{i}\right\|_{Z_{V}} = \max_{M \in \mathbb{N}, \ 1 \le r_{0} \le r_{1} < \dots < r_{M}} \left\|\sum_{i=1}^{M} \left\|\sum_{j=p_{r_{i}}}^{p_{r_{i+1}-1}} a_{j}z_{j}\right\|_{Z} v_{r_{i}}\right\|_{V}.$$

Note that $(z_i)_{i=1}^{\infty}$ is a Schauder basis for $Z_V(p_i)$ and $(\operatorname{span}_{j \in [p_i, p_{i+1})} z_j)_{i=1}^{\infty}$ is an FDD for $Z_V(p_i)$. The following proposition from [OSZ1] is what makes the space Z_V essential for us. Recall that in [OSZ1], a basis, $(v_i)_{i=1}^{\infty}$, is called *C*-block stable for some $C \ge 1$ if any two normalized block bases $(x_i)_{i=1}^{\infty}$ and $(y_i)_{i=1}^{\infty}$ with $\max(\operatorname{supp}(x_i), \operatorname{supp}(y_i)) < \min(\operatorname{supp}(x_{i+1}), \operatorname{supp}(y_{i+1}))$ for all

K. Beanland et al.

 $i \in \mathbb{N}$ are *C*-equivalent. We say that $(v_i)_{i=1}^{\infty}$ is *block stable* if it is *C*-block stable for some constant *C*. We will make use of the fact that the property of block stability dualizes. That is, if $(v_i)_{i=1}^{\infty}$ is a block stable basic sequence, then $(v_i^*)_{i=1}^{\infty}$ is also a block stable basic sequence. Another simple, though important, consequence of a normalized basic sequence $(v_i)_{i=1}^{\infty}$ being block stable is that there exists a constant $c \geq 1$ such that $(v_{n_i})_{i=1}^{\infty}$ is *c*-equivalent to $(v_{n_{i+1}})_{i=1}^{\infty}$ for all $(n_i)_{i=1}^{\infty} \in [\mathbb{N}]^{\omega}$. Block stability has been considered before in various forms and under different names. In particular, it has been called the blocking principle [CJT] and the shift property [CK] (see [FR] for alternative forms). The following proposition recalls some properties of $Z_V(p_i)$ which were shown in [OSZ1].

PROPOSITION 3.5 ([OSZ1, Corollary 3.2, Lemmas 3.3, 3.5, and 3.6]). Let $V = (v_i)_{i=1}^{\infty}$ be a normalized, 1-unconditional, and C-block stable basic sequence. If Z is a Banach space with a basis $(z_i)_{i=1}^{\infty}$ and $(p_i)_{i=1}^{\infty} \in [\mathbb{N}]^{\omega}$ with $p_1 = 1$, then $(\operatorname{span}_{j \in [p_i, p_{i+1})} z_j)_{i=1}^{\infty}$ satisfies subsequential 2C-V-lower block estimates in $Z_V(p_i)$. If the basis $(v_i)_{i=1}^{\infty}$ is boundedly complete, then $(z_i)_{i=1}^{\infty}$ is a boundedly complete basis for $Z_V(p_i)$. If the basis $(v_i)_{i=1}^{\infty}$ is shrinking and $(z_i)_{i=1}^{\infty}$ is shrinking in Z, then $(z_i)_{i=1}^{\infty}$ is a shrinking basis for $Z_V(p_i)$.

If $U = (u_i)_{i=1}^{\infty}$ is a normalized, 1-unconditional, and block stable basic sequence such that $(v_i)_{i=1}^{\infty}$ is dominated by $(u_i)_{i=1}^{\infty}$ and $(\operatorname{span}_{j \in [p_i, p_{i+1})} z_j)_{i=1}^{\infty}$ satisfies subsequential U-upper block estimates in Z, then $(\operatorname{span}_{j \in [p_i, p_{i+1})} z_j)_{i=1}^{\infty}$ also satisfies subsequential U-upper block estimates in $Z_V(p_i)$.

THEOREM 3.6. Let X be a Banach space with a shrinking Schauder frame $(x_i, f_i)_{i=1}^{\infty}$ which is strongly shrinking relative to some Banach space Z with basis $(z_i)_{i=1}^{\infty}$ and bounded operators $T: X \to Z$ and $S: Z \to X$ defined by $T(x) = \sum f_i(x)z_i$ for all $x \in X$ and $S(z) = \sum z_i^*(z)x_i$ for all $z \in Z$. Let $V = (v_i)_{i=1}^{\infty}$ be a normalized, 1-unconditional, block stable, right dominant, and shrinking basic sequence. If X satisfies subsequential V-upper tree estimates, then there exist $(n_i)_{i=1}^{\infty}, (K_i)_{i=1}^{\infty} \in [\mathbb{N}]^{\omega}$ such that $Z_{(v_{K_i})}^*(n_i)$ is an associated space of $(f_i, x_i)_{i=1}^{\infty}$ with bounded operators $S^*: X^* \to Z_{(v_{K_i})}^*(n_i)$ and $T^*: Z_{(v_{K_i})}^*(n_i) \to X$ given by $S^*(x^*) = \sum x^*(x_i)z_i^*$ for all $x^* \in X^*$ and $S^*(z^*) = \sum z^*(z_i)f_i$ for all $z^* \in Z_{(v_{K_i})}^*(n_i)$.

Proof. After renorming, we may assume that the basis $(z_i)_{i=1}^{\infty}$ is bimonotone. The sequence $(f_i, x_i)_{i=1}^{\infty}$ is a boundedly complete Schauder frame for X^* by Theorem 2.1, and we find that X^* satisfies subsequential V^* -lower w^* tree estimates by Lemma 3.3. By Theorem 2.5, the basis $(z_i^*)_{i=1}^{\infty}$ is an associated basis for $(f_i, x_i)_{i=1}^{\infty}$ with bounded operators $S^* : X^* \to Z^*$ and $T^* : Z^* \to X$ given by $S^*(x^*) = \sum x^*(x_i)z_i^*$ for all $x^* \in X^*$ and $S^*(z^*) = \sum z^*(z_i)f_i$ for all $z^* \in Z^*$. Let $\varepsilon > 0$. By Corollary 3.4, there exist C > 0, $(K_i)_{i=1}^{\infty}, (p_i)_{i=1}^{\infty} \in [\mathbb{N}]^{\omega}$, and $\overline{\delta} = (\delta_i) \subset (0,1)$ with $\delta_i \searrow 0$ and $\sum \delta_i < \varepsilon$ such that if $(y_i^*)_{i=1}^{\infty} \subset S_{X^*}$ and $(r_i)_{i=0}^{\infty} \in [\mathbb{N}]^{\omega}$ satisfy $\sup_{1 \le n \le m \le p_{r_{i-1}+1}} \|\sum_{j=n}^m y_i^*(x_j)f_j\| < \delta_i$ and $\sup_{p_{r_i} \le n \le m} \|\sum_{j=n}^m y_i^*(x_j)f_j\|$ $< \delta_i$, then $(y_i^*) \succeq_C (v_{K_{r_{i-1}}}^*)$. We apply Theorem 2.13 to $(x_i, f_i)_{i=1}^{\infty}, (p_i)_{i=1}^{\infty}$ $\in [\mathbb{N}]^{\omega}$, and $(\delta_i)_{i=1}^{\infty} \subset (0, 1)$ to obtain $(q_i)_{i=1}^{\infty}, (N_i)_{i=1}^{\infty} \in [\mathbb{N}]^{\omega}$ satisfying the conclusion of Theorem 2.13.

By Theorem 2.5, $(z_i^*)_{i=1}^{\infty}$ is an associated basis for $(f_i, x_i)_{i=1}^{\infty}$, and we denote the norm on Z^* by $\|\cdot\|_{Z^*}$. We block the basis $(z_i^*)_{i=1}^{\infty}$ into an FDD by setting $E_i = \operatorname{span}_{j \in [p_{q_{N_i}}, p_{q_{N_{i+1}}})} z_j^*$ for all $i \in \mathbb{N}$. We now define a new norm $\|\cdot\|_{\bar{Z}^*}$ on $\operatorname{span}(z_i^*)_{i=1}^{\infty}$ by setting, for all $(a_i) \in c_{00}$,

$$\left\|\sum a_{i}z_{i}^{*}\right\|_{\bar{Z}^{*}} = \max_{M \in \mathbb{N}, \ 1 \leq r_{0} \leq r_{1} < \dots < r_{M}} \left\|\sum_{i=1}^{M} \left\|\sum_{j=p_{q_{N_{r_{i}}}}}^{p_{q_{N_{r_{i+1}}}}-1} a_{j}z_{j}^{*}\right\|_{Z^{*}} v_{K_{N_{r_{i}}}}^{*}\right\|_{V^{*}}$$

We let \overline{Z}^* be the completion of $\operatorname{span}(z_i^*)_{i=1}^{\infty}$ under the norm $\|\cdot\|_{\overline{Z}^*}$. Note that $\|z^*\|_{Z^*} \leq \|z^*\|_{\overline{Z}^*}$ for all $z^* \in Z^*$. As $(v_i^*)_{i=1}^{\infty}$ is block stable, there exists a constant $c \geq 1$ such that $(v_{n_i}^*)_{i=1}^{\infty} \approx_c (v_{n_{i+1}}^*)_{i=1}^{\infty}$ for all $(n_i)_{i=1}^{\infty} \in [\mathbb{N}]^{\omega}$. We now show that $\|S^*(y^*)\|_{\overline{Z}^*} \leq (1+2\varepsilon)3cC\|S\| \|y^*\|$ for all $y^* \in X^*$.

Let $y^* \in X^*$ with $||y^*|| = 1$, $M \in \mathbb{N}$, and $1 \leq r_0 \leq r_1 < \cdots < r_M$. We will show that $(1+2\varepsilon)3cC||S|| ||y^*|| \geq ||\sum_{i=1}^M ||\sum_{j=p_{q_{N_{r_i}}}}^{p_{q_{N_{r_{i+1}}}}-1} y^*(x_j)z_j^*||v_{p_{N_{r_i}}}^*||_{V^*}$. By Theorem 2.13, there exist $y_i^* \in X^*$ and $t_i \in (N_{r_{i-1}-1}, N_{r_{i-1}})$ for all $i \in \mathbb{N}$ with $N_0 = 0$ and $t_0 = 0$ such that

(a) $y^* = \sum_{i=1}^{\infty} y_i^*$

and that for all $i \in \mathbb{N}$ we have

(b) either
$$||y_i^*|| < \delta_i$$
, or $\sup_{p_{q_{t_{i-1}}} \ge n \ge m} ||\sum_{j=m}^n y_i^*(x_j)f_j|| < \delta_i ||y_i^*||$ and
 $\sup_{n \ge m \ge p_{q_{t_i}}} ||\sum_{j=m}^n y_i^*(x_j)f_j|| < \delta_i ||y_i^*||,$
(c) $||P_{[p_{q_{N_{k_i}}}^*, p_{q_{N_{k_{i+1}}}})} \circ S^*(y_{i-1}^* + y_i^* + y_{i+1}^* - y^*)||_{Z^*} < \delta_i.$

We let $A = \{i \in \mathbb{N} : ||y_i^*|| > \delta_i\}$. By our choice of $(p_i)_{i=1}^{\infty}$, we deduce that $(y_i^*/||y_i^*||)_{i\in A} \succeq_C (v_{K_{r_{i-1}}}^*)_{i\in A}$. Thus, $C||\sum_{i\in A} y_i^*|| \ge ||\sum_{i\in A} ||y_i^*|| v_{K_{r_{i-1}}}^*||_{V^*}$. We now obtain the following lower estimate for $||y^*||$:

$$\begin{aligned} \|y^*\| &= \left\|\sum y_i^*\right\| \quad \text{by (a)} \\ &\geq \left\|\sum_{i \in A} y_i^*\right\| + \sum_{i \notin A} \|y_i^*\| - \varepsilon \quad \text{ as } \sum \delta_i < \varepsilon \\ &\geq \frac{1}{C} \left\|\sum_{i \in A} \|y_i^*\| v_{K_{r_{i-1}}}^*\right\|_{V^*} + \sum_{i \notin A} \|y_i^*\| - \varepsilon \ge \frac{1}{C} \left\|\sum \|y_i^*\| v_{K_{r_{i-1}}}^*\right\|_{V^*} - \varepsilon \end{aligned}$$

$$\begin{split} &\geq \frac{1}{3cC} \Big(\left\| \sum \|y_i^*\| v_{K_{r_{i-1}}}^* \right\|_{V^*} + \left\| \sum \|y_{i-1}^*\| v_{K_{r_{i-1}}}^* \right\|_{V^*} + \left\| \sum \|y_{i+1}^*\| v_{K_{r_{i-1}}}^* \right\|_{V^*} \Big) - \varepsilon \\ &\geq \frac{1}{3cC} \left\| \sum \|y_{i-1}^* + y_i^* + y_{i+1}^*\| v_{K_{r_{i-1}}}^* \right\|_{V^*} - \varepsilon \\ &\geq \frac{1}{3cC\|S\|} \left\| \sum \|S^*(y_{i-1}^* + y_i^* + y_{i+1}^*)\|_{Z^*} v_{K_{r_{i-1}}}^* \right\|_{V^*} - \varepsilon \\ &\geq \frac{1}{3cC\|S\|} \left\| \sum \|P_{[p_{q_{N_i}}, p_{q_{N_{i+1}}}]}^* S^*(y_{i-1}^* + y_i^* + y_{i+1}^*)\|_{Z^*} v_{K_{r_{i-1}}}^* \right\|_{V^*} - \varepsilon \\ &\geq \frac{1}{3cC\|S\|} \left\| \sum \|\sum_{j=p_{q_{N_i}}}^{p_{q_{N_{i+1}}} - 1} S^*(y_{i-1}^* + y_i^* + y_{i+1}^*)\|_{Z^*} v_{K_{r_{i-1}}}^* \right\|_{V^*} - \varepsilon \\ &\geq \frac{1}{3cC\|S\|} \left\| \sum \|\sum_{j=p_{q_{N_i}}}^{p_{q_{N_{i+1}}} - 1} y^*(x_j) z_j \| v_{K_{r_{i-1}}}^* \right\|_{V^*} - 2\varepsilon \quad \text{by (c).} \\ &\text{Thus } \|S^*y^*\|_{\bar{Z}^*} \leq (1+2\varepsilon) 3cC\|S\| \|y^*\| \text{ for all } y^* \in X^*. \text{ Furthermore, as } \\ \|z^*\|_{Z^*} \leq \|z^*\|_{\bar{Z}^*} \text{ for all } z^* \in Z^*, \text{ we have } \|y^*\| \leq \|T^*\| \|S^*y^*\|_{Z^*} \leq \|T^*\| \|S^*y^*\|_{\bar{Z}^*} \\ &\text{for all } y^* \in X^*. \text{ Hence, } S^* : X^* \to \bar{Z}^* \text{ is an isomorphism. We observe that } \\ &T^*: Z^* \to X^* \text{ is bounded, and hence } T^*: \bar{Z}^* \to X^* \text{ is bounded as well, since } \\ \|z^*\|_{Z^*} \leq \|z^*\|_{\bar{Z}^*} \text{ for all } z^* \in Z^*. \text{ Thus, } \bar{Z}^* \text{ is an associated space of } X^*. \end{array} \right. \end{split}$$

We now restate and prove Theorem 1.6, which is an extension of Theorem 1.1 in [FOSZ] to Schauder frames.

THEOREM 3.7. Let X be a Banach space with a shrinking Schauder frame $(x_i, f_i)_{i=1}^{\infty}$. Let $(v_i)_{i=1}^{\infty}$ be a normalized, 1-unconditional, block stable, right dominant, and shrinking basic sequence. If X satisfies subsequential $(v_i)_{i=1}^{\infty}$ -upper tree estimates, then there exist $(n_i)_{i=1}^{\infty}, (K_i)_{i=1}^{\infty} \in [\mathbb{N}]^{\omega}$ and an associated space Z with a shrinking basis $(z_i)_{i=1}^{\infty}$ such that $(\operatorname{span}_{j \in [n_i, n_{i+1})} z_i)_{i=1}^{\infty}$ is an FDD which satisfies subsequential $(v_{K_i})_{i=1}^{\infty}$ -upper block estimates.

Proof. As $(x_i, f_i)_{i=1}^{\infty}$ is a shrinking Schauder frame, it is strongly shrinking relative to some associated basis $(z_i)_{i=1}^{\infty}$ for a Banach space Z by Theorem 2.9. We thus have bounded operators $T: X \to Z$ and $S: Z \to X$ defined by $T(x) = \sum f_i(x)z_i$ for all $x \in X$ and $S(z) = \sum z_i^*(z)x_i$ for all $z \in Z$. By Theorem 3.6, there exist $(n_i)_{i=1}^{\infty}, (K_i)_{i=1}^{\infty} \in [\mathbb{N}]^{\omega}$ such that $Z_{(v_{K_i}^*)}^*(n_i)$ is an associated space of $(f_i, x_i)_{i=1}^{\infty}$ with bounded operators $S^*: X^* \to Z_{(v_{K_i}^*)}^*(n_i)$ and $T^*: Z_{(v_{K_i}^*)}^*(n_i) \to X$ given by $S^*(x^*) = \sum x^*(x_i)z_i^*$ for all $x^* \in X^*$ and $T^*(z^*) = \sum z^*(z_i)f_i$ for all $z^* \in Z_{(v_{K_i}^*)}^*(n_i)$. We define \overline{Z} as the completion of $[z_i]_{i=1}^{\infty}$ under the norm

$$\left\|\sum a_{i} z_{i}\right\|_{\bar{Z}} = \sup_{z^{*} \in B_{Z_{(v_{K_{i}}^{*})}^{*}(n_{i})}} z^{*} \left(\sum a_{i} z_{i}\right).$$

As $(z_i^*)_{i=1}^{\infty}$ is a boundedly complete basis of $Z_{(v_{K_i}^*)}^*(n_i)$ by Proposition 3.5, we see that $(z_i)_{i=1}^{\infty}$ is a shrinking basis for \bar{Z} and that the dual of \bar{Z} is $Z_{(v_{K_i}^*)}^*(n_i)$. We now prove that \bar{Z} is an associated space for $(x_i, f_i)_{i=1}^{\infty}$. If $(y_i^*)_{i=1}^{\infty} \subset X^*$ and $y_i^* \to_{w^*} 0$, then $(S^*(y_i^*))_{i=1}^{\infty}$ converges w^* to 0 as a sequence in Z^* . Thus $((S^*(y_i^*))(z_j))_{i=1}^{\infty}$ converges to 0 for all $j \in \mathbb{N}$. Furthermore, $(S^*(y_i^*))_{i=1}^{\infty}$ is bounded in $Z^*_{(v_{K_i}^*)}(n_i)$ as $Z^*_{(v_{K_i}^*)}(n_i)$ is an associated space of $(f_i, x_i)_{i=1}^{\infty}$. Proposition 3.5 shows that $(z_i^*)_{i=1}^{\infty}$ is a boundedly complete basis for $Z^*_{(v_{K_i}^*)}(n_i)$, and hence converging w^* to 0 in $Z^*_{(v_{K_i}^*)}(n_i)$ is equivalent to converging coordinatewise to 0 for bounded sequences. Hence, $(S^*(y_i^*))_{i=1}^{\infty}$ converges w^* to 0 in $Z^*_{(v_{K_i}^*)}(n_i)$. Thus $S^*: X^* \to Z^*_{(v_{K_i}^*)}(n_i)$ is w^* -to- w^* continuous, and hence is a dual operator. Thus, $S: \overline{Z} \to X$ is well defined and bounded.

If $(y_i^*)_{i=1}^{\infty} \subset Z_{(v_{K_i}^*)}^*(n_i)$ converges w^* to 0 in $Z_{(v_{K_i}^*)}^*(n_i)$, then $(y_i^*)_{i=1}^{\infty}$ is bounded and converges coordinatewise to 0. Hence, $(y_i^*)_{i=1}^{\infty}$ is bounded and converges coordinatewise to 0 in Z^* as $\|z^*\|_{Z^*} \leq \|z^*\|_{Z_{(v_{K_i}^*)}^*(n_i)}$ for all $z^* \in Z^*$. Hence, $(y_i^*)_{i=1}^{\infty}$ converges w^* to 0 in Z^* as $(z_i^*)_{i=1}^{\infty}$ is a boundedly complete basis for Z^* . Thus, $(T^*(y_i^*))_{i=1}^{\infty} \to w^*$ 0 in X^* . Hence $T^* : Z_{(v_{K_i}^*)}^*(n_i) \to X^*$ is w^* -to- w^* continuous, and so a dual operator. Thus, $T : X \to \overline{Z}$ is well defined and bounded. This tells us that \overline{Z} is an associated space for $(x_i, f_i)_{i=1}^{\infty}$. By Proposition 3.5, the FDD $(\operatorname{span}_{j\in[n_i,n_{i+1})} z_j^*)_{i=1}^{\infty}$ satisfies $(v_{K_i}^*)$ -lower block estimates in \overline{Z}^* , and hence $(\operatorname{span}_{j\in[n_i,n_{i+1})} z_j)_{i=1}^{\infty}$ satisfies $(v_{K_i})_{i=1}^{\infty}$ -upper block estimates in \overline{Z} . By Lemma 3.1 we conclude that $(z_i)_{i=1}^{\infty}$ is shrinking.

We now restate and prove Theorem 1.7, which is an extension of Theorem 4.6 in [OSZ1] to frames.

THEOREM 3.8. Let X be a Banach space with a shrinking and boundedly complete Schauder frame $(x_i, f_i)_{i=1}^{\infty}$. Let $(u_i)_{i=1}^{\infty}$ be a normalized, 1-unconditional, block stable, right dominant, and shrinking basic sequence, and let $(v_i)_{i=1}^{\infty}$ be a normalized, 1-unconditional, block stable, left dominant, and boundedly complete basic sequence such that (u_i) dominates (v_i) . Then X satisfies subsequential $(u_i)_{i=1}^{\infty}$ -upper tree estimates and subsequential $(v_i)_{i=1}^{\infty}$ -lower tree estimates if and only if there exist $(n_i)_{i=1}^{\infty}, (K_i)_{i=1}^{\infty} \in [\mathbb{N}]^{\omega}$ and a reflexive associated space Z with associated basis $(z_i)_{i=1}^{\infty}$ such that the FDD $(\operatorname{span}_{j \in [n_i, n_{i+1}]} z_j)_{i=1}^{\infty}$ satisfies subsequential $(u_{K_i})_{i=1}^{\infty}$ -upper block estimates and subsequential $(v_{K_i})_{i=1}^{\infty}$ -lower block estimates.

Proof. By Theorem 3.7, $(x_i, f_i)_{i=1}^{\infty}$ has an associated space Z with a shrinking basis $(z_i)_{i=1}^{\infty}$ such that there exist $(m_i)_{i=1}^{\infty}, (k_i)_{i=1}^{\infty} \in [\mathbb{N}]^{\omega}$ such that the FDD $(\operatorname{span}_{j \in [m_i, m_{i+1}]} z_j)_{i=1}^{\infty}$ satisfies subsequential $(u_{k_i})_{i=1}^{\infty}$ -upper block estimates. We notice that $(f_i, x_i)_{i=1}^{\infty}$ is a shrinking frame for X* which is strongly shrinking relative to the associated basis $(z_i^*)_{i=1}^{\infty}$ by Corollary 2.6. The space X satisfying subsequential $(v_i)_{i=1}^{\infty}$ -lower tree estimates implies

that X^* satisfies subsequential $(v_i^*)_{i=1}^{\infty}$ -upper tree estimates. Thus we may apply Theorem 3.6 to the space X^* , the frame $(f_i, x_i)_{i=1}^{\infty}$, and the associated basis $(z_i^*)_{i=1}^{\infty}$ to obtain $(n_i)_{i=1}^{\infty}, (K_i)_{i=1}^{\infty} \in [\mathbb{N}]^{\omega}$ such that $Z_{(v_{K_i})}(n_i)$ is an associated space for $(x_i, f_i)_{i=1}^{\infty}$. Furthermore, we may assume that $(n_i)_{i=1}^{\infty}$ is a subsequence of $(m_i)_{i=1}^{\infty}$ and that $(K_i)_{i=1}^{\infty}$ is a subsequence of $(k_i)_{i=1}^{\infty}$ as $(v_i)_{i=1}^{\infty}$ is left dominant. By Lemma 3.5, the FDD $(\operatorname{span}_{j\in[n_i,n_{i+1})} z_j)_{i=1}^{\infty}$ satisfies subsequential $(u_{K_i})_{i=1}^{\infty}$ -upper block estimates and subsequential $(v_{K_i})_{i=1}^{\infty}$ lower block estimates. By Lemma 3.1 we deduce that $(z_i)_{i=1}^{\infty}$ is shrinking and boundedly complete. Thus, Z is reflexive.

We now show that Theorem 1.5 follows immediately from Theorems 3.7 and 3.8.

Proof of Theorem 1.5. Let $(x_i, f_i)_{i=1}^{\infty}$ be a shrinking Schauder frame for a Banach space X. Then X must have separable dual by Theorem 2.1. We claim that X must satisfy subsequential $(v_i)_{i=1}^{\infty}$ -upper tree estimates for some normalized, 1-unconditional, block stable, right dominant, and shrinking basic sequence $(v_i)_{i=1}^{\infty}$. Indeed, as X has separable dual, X must have countable Szlenk index. In particular, $Sz(X) \leq \omega^{\alpha\omega}$ for some countable ordinal α . By [FOSZ, Theorem 1.3], there exists a constant $c \in (0, 1)$ such that X satisfies subsequential $T_{\alpha,c}$ -upper tree estimates, where $T_{\alpha,c}$ is Tsirelson's space of order α and constant c. The unit vector basis for $T_{\alpha,c}$ is a normalized, 1-unconditional, block stable, right dominant, and shrinking basic sequence (see [OSZ2, Proposition 3] as well as [CJT] and [LT] for a proof). Thus, we may apply Theorem 3.7 to obtain a shrinking associated basis for the shrinking Schauder frame $(x_i, f_i)_{i=1}^{\infty}$.

We now assume that $(x_i, f_i)_{i=1}^{\infty}$ is a shrinking and boundedly complete Schauder frame for a Banach space X. Then X must be reflexive by Theorem 2.2. We claim that X must satisfy subsequential $(u_i)_{i=1}^{\infty}$ -upper tree estimates and subsequential $(u_i^*)_{i=1}^{\infty}$ -lower tree estimates for some normalized, 1-unconditional, block stable, right dominant, and shrinking basic sequence $(u_i)_{i=1}^{\infty}$. Indeed, as X is separable and reflexive, both X and X* must have countable Szlenk index. In particular, $Sz(X), Sz(X^*) \leq \omega^{\alpha\omega}$ for some countable ordinal α . By [OSZ2, Theorem 21], there exists a constant $c \in (0, 1)$ such that X satisfies subsequential $T_{\alpha,c}$ -upper tree estimates and $T_{\alpha,c}^*$ -lower tree estimates. The unit vector basis for $T_{\alpha,c}$ is a normalized, 1-unconditional, block stable, right dominant, and shrinking basic sequence. Thus the unit vector basis for $T_{\alpha,c}^*$ is a normalized, 1-unconditional, block stable, right dominant, and shrinking basic sequence. Thus the unit vector basis for $T_{\alpha,c}^*$ is a normalized, stable, left dominant, and boundedly complete basic sequence. We apply Theorem 3.8 to obtain a reflexive associated space for the shrinking and boundedly complete Schauder frame $(x_i, f_i)_{i=1}^{\infty}$. Acknowledgements. The third author would like to express his gratitude to Edward Odell, who is greatly missed, for inviting him to visit the University of Texas at Austin in the fall of 2010 and spring of 2011.

The research of the second author was supported by grant DMS1332255 from the National Science Foundation. The third author was supported by the National Natural Science Foundation of China.

References

- [AD] S. A. Argyros and P. Dodos, Genericity and amalgamation of classes of Banach spaces, Adv. Math. 209 (2007), 666–748.
- [Bos] B. Bossard, Théorie descriptive des ensembles en géométrie des espaces de Banach, PhD thesis, Univ. Paris VI, 1994.
- [B] J. Bourgain, On separable Banach spaces, universal for all separable reflexive spaces, Proc. Amer. Math. Soc. 79 (1980), 241–246.
- [CL] D. Carando and S. Lassalle, Duality, reflexivity and atomic decompositions in Banach spaces, Studia Math. 191 (2009), 67–80.
- [CLS] D. Carando, S. Lassalle, and P. Schmidberg, The reconstruction formula for Banach frames and duality, J. Approx. Theory 163 (2011), 640–651.
- [CDOSZ] P. G. Casazza, S. J. Dilworth, E. Odell, Th. Schlumprecht, and A. Zsák, Coefficient quantization for frames in Banach spaces, J. Math. Anal. Appl. 348 (2008), 66–86.
- [CHL] P. G. Casazza, D. Han, and D. R. Larson, Frames for Banach spaces, in: The Functional and Harmonic Analysis of Wavelets and Frames, L. W. Baggett and D. R. Larson (eds.), Contemp. Math. 247, Amer. Math. Soc., Providence, RI, 1999, 149–182.
- [CJT] P. G. Casazza, W. B. Johnson, and L. Tzafriri, On Tsirelson's space, Israel J. Math. 47 (1984), 81–98.
- [CK] P. G. Casazza and N. J. Kalton, Unconditional bases and unconditional finitedimensional decompositions in Banach spaces, Israel J. Math. 95 (1996), 349– 373.
- [C1] R. Causey, Estimation of the Szlenk index of Banach spaces via Schreier spaces, Studia Math. 216 (2013), 149–178.
- [C2] R. Causey, Estimation of the Szlenk index of reflexive Banach spaces using generalized Baernstein spaces, Fund. Math. 228 (2015), 153–171.
- [DGM] I. Daubechies, A. Grossmann, and Y. Meyer, Painless nonorthogonal expansions, J. Math. Phys. 27 (1986), 1271–1283.
- [DF] P. Dodos and V. Ferenczi, Some strongly bounded classes of Banach spaces, Fund. Math. 193 (2007), 171–179.
- [DS] R. J. Duffin and A. C. Schaeffer, A class of nonharmonic Fourier series, Trans. Amer. Math. Soc. 72 (1952), 341–366.
- [FG] H. G. Feichtinger and K. Gröchenig, A unified approach to atomic decompositions via integrable group representations, in: Function Spaces and Applications, Lecture Notes in Math. 1302, Springer, Berlin, 1988, 52–73.
- [FR] V. Ferenczi and Ch. Rosendal, Ergodic Banach spaces, Adv. Math. 195 (2005), 259–282.
- [FOSZ] D. Freeman, E. Odell, Th. Schlumprecht, and A. Zsák, Banach spaces of bounded Szlenk index, II, Fund. Math. 205 (2009), 161–177.

188	K. Beanland et al.	
[FPWW]	D. Freeman, D. Poore, A. R. Wei, and M. Wyse, <i>Moving vector bundles</i> , Houston J. Math. 40 (2014), 817–832.	Parseval frames for
[G]	K. Gröchenig, <i>Describing functions: Atomic decomposi</i> Monatsh. Math. 112 (1991), 1–42.	tions versus frames,
[HL]	D. Han and D. R. Larson, Frames, bases and group re Amer. Math. Soc. 147 (2000), no. 697.	presentations, Mem.
[HLLL]	D. Han, D. R. Larson, B. Liu, and R. Liu, <i>Operator-valuea</i> and the theory of frames, Mem. Amer. Math. Soc. 229 (2	l measures, dilations, 2014), no. 1075.
[J]	W. B. Johnson, On quotients of L_p which are quotients of 34 (1977), 69–89.	$f \ell_p$, Compos. Math.
[LT]	D. H. Leung and WK. Tang, The Bourgain ℓ_1 -index of m. J. Funct. Anal. 199 (2003), 301–331.	nixed Tsirelson space,
[L]	R. Liu, On shrinking and boundedly complete Schaude spaces, J. Math. Anal. Appl. 365 (2010), 385–398.	r frames of Banach
[LZ]	R. Liu and B. Zheng, A characterization of Schauder fra Schauder bases, J. Fourier Anal. Appl. 16 (2010), 791–80	mes which are near- 3.
[OS1]	E. Odell and Th. Schlumprecht, <i>Trees and branches in B</i> Amer. Math. Soc. 354 (2002), 4085–4108.	anach spaces, Trans.
[OS2]	E. Odell and Th. Schlumprecht, A universal reflexive s uniformly convex Banach spaces, Math. Ann. 335 (2006).	pace for the class of 901–916.
[OSZ1]	E. Odell, Th. Schlumprecht, and A. Zsák, <i>A new infin spaces with applications</i> , in: Banach Spaces and Their App de Gruyter, Berlin, 2007, 147–182.	ite game in Banach lications in Analysis,
$\left[\mathrm{OSZ2}\right]$	E. Odell, Th. Schlumprecht, and A. Zsák, <i>Banach space index</i> , Studia Math. 183 (2007), 63–97.	es of bounded Szlenk
[Sz]	W. Szlenk, The non-existence of a separable reflexive Ba for all separable reflexive Banach spaces, Studia Math. 30	nach space universal 0 (1968), 53–61.
[Z]	M. Zippin, Banach spaces with separable duals, Trans. A (1988), 371–379.	mer. Math. Soc. 310
Kevin Bea	eanland	Daniel Freeman
Washingto	ton and Lee University Departm	nent of Mathematics

Washington and Lee University 204 W. Washington St. Lexington, VA 24450, U.S.A. E-mail: beanlandk@wlu.edu Daniel Freeman Department of Mathematics and Computer Science Saint Louis University St. Louis, MO 63103, U.S.A. E-mail: dfreema7@slu.edu

Rui Liu Department of Mathematics and LPMC Nankai University Tianjin 300071, P.R. China and Department of Mathematics Texas A&M University College Station, TX 77843, U.S.A. E-mail: ruiliu@nankai.edu.cn

> Received 6 January 2014; in revised form 3 April 2015