

Metric abstract elementary classes with perturbations

by

Åsa Hirvonen and Tapani Hyttinen (Helsinki)

Abstract. We define an abstract setting suitable for investigating perturbations of metric structures generalizing the notion of a metric abstract elementary class. We show how perturbation of Hilbert spaces with an automorphism and atomic Nakano spaces with bounded exponent fit into this framework, where the perturbations are built into the definition of the class being investigated. Further, assuming homogeneity and some other properties true in the example classes, we develop a notion of independence for this setting and show that it satisfies the usual independence axioms. Finally we define an isolation notion. Although it remains open whether this isolation gives any reasonable form of primeness, we prove that dominance works.

1. Introduction. In metric model theory there are examples of classes which come down in stability hierarchy if instead of the usual metric way of measuring stability one measures stability up to perturbations. Some come down as far as from non-superstability to omega-stability. So far, however, this phenomenon has mainly been observed but not used. This paper develops techniques to use the improvement in stability that perturbations offer, in particular it shows how ω -stability up to perturbation can be used to prove better behaviour of independence and constructible models than what one gets from usual metric stability or superstability.

The idea of perturbations was introduced by Ben Yaacov in [1] and [2]. We take a slightly different perspective. Instead of looking at later modifications of the structures and the mappings arising from such, we take as our starting point the mappings themselves and introduce classes of *generalized isomorphisms*. These are considered a form of isomorphisms and satisfy more or less the same properties as Ben Yaacov's perturbation mappings.

To allow for a natural treatment of generalized isomorphisms we use a syntax-free approach, that of *metric abstract elementary classes*. It was developed in [11] and is a very natural generalization of Shelah's abstract

2010 *Mathematics Subject Classification*: Primary 03C52; Secondary 03C45.

Key words and phrases: metric structures, perturbation.

elementary classes; basically the same definition was mentioned in [15]. The novelty of the approach in this paper is the adding of classes \mathbb{F}_ε of so-called ε -isomorphisms, which allow for the built-in treatment of perturbations. The abstract framework also gives a natural approach to unbounded structures, avoiding the somewhat cumbersome “emboundment” that Ben Yaacov introduces in [1] in order to be able to perturb the norm of Banach spaces.

As it seems that compactness (in the sense of metric model theory) is of limited value when everything is done only up to perturbations, we take homogeneity as our starting point and point out when stronger assumptions are needed. So as in [11] our basic framework is that of a metric abstract elementary class with countable Löwenheim–Skolem number (where size is measured by the density character), arbitrarily large models, joint embedding, amalgamation and homogeneity for Galois types (i.e. with respect to genuine isomorphisms). In addition we assume a form of amalgamation with respect to the ε -isomorphisms and a stronger perturbation property linking perturbation mappings to Galois types. The topology we consider on the type space is given by a metrizable uniformity \mathbf{d}^p , which resembles the distance arising in Ben Yaacov’s work by combining perturbation and the moving of realizations of types. In our setting it, however, also includes the moving of parameters, so our notion of ω -stability is weaker than “ ω -stability up to perturbation” in [2].

A natural example of a metric abstract elementary class with perturbations (although not presented here in further detail) is the class of all Banach spaces, where the perturbation mappings are Banach-space isomorphisms (i.e. linear homeomorphisms). Then the amalgamation property is given by Kislyakov’s construction, described in [8] (and in [11] in the setting of a metric abstract elementary class). It may be of interest to people in functional analysis that the monster model constructed using this form of amalgamation has the property that any Banach-space isomorphism between two small enough spaces and with any given isomorphism constant extends to a Banach-space automorphism of the monster with the same isomorphism constant. Further the perturbation property in this class arises from the fact that in the monster model finite tuples have the same almost isometric type if and only if they have the same Galois type.

Our main examples consist of on one hand Hilbert spaces with an automorphism and on the other atomic Nakano spaces with bounded exponent. Hilbert spaces with automorphisms and perturbations of them have been studied by Ben Yaacov, Berenstein, Usvyatsov and Zadka in [4] and [7], and they served as a first testing ground for additional assumptions on the perturbation mappings, needed for finding limits in constructions. Nakano spaces with bounded exponent have been studied by Poitevin and Ben Yaacov. The atomic case falls outside the compact range and thus offers a non-

compact example still satisfying the assumption of *complete type spaces*. This is a limit type assumption with a strong flavour of compactness and it is used in most of the constructions.

Having established a framework for structures with perturbation we develop an independence notion based on splitting and show that this independence notion satisfies the usual independence axioms (to the extent natural for a metric structure). Using this independence notion we also show that $\omega\text{-}\mathbf{d}^p$ -stable structures with complete type-spaces are $\lambda\text{-}\mathbf{d}^p$ -stable for all λ and further that $\omega\text{-}\mathbf{d}^p$ -stability and complete type spaces or good models imply ordinary stability in all $\lambda = \lambda^{N_0}$. We remind the reader that in stable classes there tends to be only one independence notion, i.e. any two are the same over the sets over which both behave well. One example of this phenomenon will be given.

Finally we introduce a notion of isolation. This enables us to define primary sets, but the question whether these are prime in any reasonable sense remains open. However, we do prove dominance for these primary sets.

In Section 2 we introduce the framework and present our notational conventions. Section 3 shows how Hilbert spaces with an automorphism fit into the framework. Section 4 studies atomic Nakano spaces with bounded exponent. In Section 5 we investigate the \mathbf{d}^p and present two ways of finding limit types. Section 6 presents the concept of $\omega\text{-}\mathbf{d}^p$ -saturation, which corresponds to Ben Yaacov's p -approximate N_0 -saturation. We show how to find $\omega\text{-}\mathbf{d}^p$ -saturated sets and prove that under various conditions true in the Hilbert example $\omega\text{-}\mathbf{d}^p$ -saturated models are almost isomorphic, a result obtained for continuous logic with perturbations by Ben Yaacov [1, Proposition 2.7]. Section 7 defines splitting and the independence notion based on it. Here we prove that the independence axioms hold and show how our stability and independence notions relate to the standard ones in homogeneous model theory. Finally Section 8 deals with isolation.

2. Generalizing metric abstract elementary classes. When reading the following definitions one may keep in mind the following example (in Sections 3 and 4 we give the main examples): Let \mathbb{K} be the class of all Banach spaces and let \preccurlyeq be the closed subspace relation. A map $f : \mathcal{A} \rightarrow \mathcal{B}$ is in \mathbb{F}_ε if f is an onto linear isomorphism in the functional analysis sense (i.e. a linear homeomorphism) with $\|f\|$ and $\|f^{-1}\|$ at most e^ε , where $e = \lim_{n \rightarrow \infty} (1 + 1/n)^n$. The exponential is used in order for the epsilons to behave nicely when composing functions (see Definition 2.2).

As in [11], we investigate a class of complete metric space structures of a fixed countable vocabulary. The structures are many-sorted and of the form

$$\mathcal{M} = \langle \mathcal{A}_0, \mathcal{A}_1, \dots, \mathbb{R}, d_0, d_1, \dots, c_0, c_1, \dots, R_0, R_1, \dots, F_0, F_1, \dots \rangle,$$

where

- (1) each \mathcal{A}_i is a complete metric space with metric d_i (with values in \mathbb{R}),
- (2) \mathbb{R} is an isomorphic copy of the ordered field of real numbers

$$(\mathbb{R}, +, \cdot, 0, 1, \leq, |\cdot|),$$

- (3) each c_i is a constant and each R_i a relation,
- (4) each F_i is a function $F_i : \mathcal{B}_0 \times \cdots \times \mathcal{B}_m \rightarrow \mathcal{B}_{m+1}$ where $\mathcal{B}_j \in \{\mathcal{A}_0, \mathcal{A}_1, \dots, \mathbb{R}\}$.

We will use some shorthand notation. When we talk about elements of a model \mathcal{M} we mean an element of some sort of \mathcal{M} , and $a \in \mathcal{M}$ refers to a finite tuple of elements of \mathcal{M} . For a set A , \bar{A} signifies, depending on whether we are inside a model or not, either the sortwise metric closure of A or some completion of A . Since size is measured with respect to density character, $|A| = \text{dens}(A)$. For \mathcal{M} a many-sorted model, $|\mathcal{M}|$ is the sum of the density characters of its sorts. The cardinality of a set A is denoted $\text{card}(A)$.

Both models and their domains will be denoted by \mathcal{A}, \mathcal{B} etc., A, B etc. refers to sets not necessary being models. After we construct a monster model \mathfrak{M} , models will be submodels of \mathfrak{M} with size small enough compared to $|\mathfrak{M}|$. Also by $A \subset \mathfrak{M}$ we mean that A is a small enough subset. (The notion of small enough will become apparent when we define \mathfrak{M} .)

We consider the distance $d(a, b)$ of two finite tuples a and b of the same length to be the maximum of their coordinatewise distances. When A and B are finite sets of the same cardinality we consider them to be ordered and hence by $d(A, B)$ mean this maximum of coordinatewise distances for a fixed ordering.

In [11] MAECs were defined as follows:

DEFINITION 2.1. We call a class $(\mathbb{K}, \preccurlyeq_{\mathbb{K}})$ of τ -structures for some fixed vocabulary τ a *metric abstract elementary class*, *MAEC*, if the following hold:

- (1) Both \mathbb{K} and the binary relation $\preccurlyeq_{\mathbb{K}}$ are closed under isomorphism.
- (2) If $\mathcal{A} \preccurlyeq_{\mathbb{K}} \mathcal{B}$ then \mathcal{A} is a substructure of \mathcal{B} (i.e. each sort of \mathcal{A} is a substructure of the corresponding sort of \mathcal{B}).
- (3) $\preccurlyeq_{\mathbb{K}}$ is a partial order on \mathbb{K} .
- (4) If $(\mathcal{A}_i)_{i < \delta}$ is a $\preccurlyeq_{\mathbb{K}}$ -increasing chain then there is a model $\overline{\bigcup_{i < \delta} \mathcal{A}_i}$, unique up to the choice of completion such that
 - (a) $\overline{\bigcup_{i < \delta} \mathcal{A}_i} \in \mathbb{K}$,
 - (b) for each $j < \delta$, $\mathcal{A}_j \preccurlyeq_{\mathbb{K}} \overline{\bigcup_{i < \delta} \mathcal{A}_i}$.

In addition if each $\mathcal{A}_i \preccurlyeq_{\mathbb{K}} \mathcal{B} \in \mathbb{K}$ and the completion is the metric closure in \mathcal{B} then $\overline{\bigcup_{i < \delta} \mathcal{A}_i} \preccurlyeq_{\mathbb{K}} \mathcal{B}$.

- (5) If $\mathcal{A}, \mathcal{B}, \mathcal{C} \in \mathbb{K}$, $\mathcal{A} \preccurlyeq_{\mathbb{K}} \mathcal{C}$, $\mathcal{B} \preccurlyeq_{\mathbb{K}} \mathcal{C}$ and $\mathcal{A} \subseteq \mathcal{B}$ then $\mathcal{A} \preccurlyeq_{\mathbb{K}} \mathcal{B}$.
- (6) There exists a Löwenheim–Skolem number $LS^d(\mathbb{K})$ such that if $\mathcal{A} \in \mathbb{K}$ and $A \subset \mathcal{A}$ then there is $\mathcal{B} \supseteq A$ such that $|\mathcal{B}| = |A| + LS^d(\mathbb{K})$ and $\mathcal{B} \preccurlyeq_{\mathbb{K}} \mathcal{A}$.

We generalize this by loosening the requirements of being an isomorphism and add a class \mathbb{F} of homeomorphisms to play the role of isomorphisms.

DEFINITION 2.2. $(\mathbb{K}, \preccurlyeq, \mathbb{F}_\varepsilon)_{\varepsilon \geq 0}$ is a *metric abstract elementary class with perturbations* if

- (1) $(\mathbb{K}, \preccurlyeq)$ is a MAEC.
 - (2) The \mathbb{F}_ε are collections of bijective mappings between members of \mathbb{K} such that:
 - (a) $\mathbb{F}_\delta \subseteq \mathbb{F}_\varepsilon$ for $\delta < \varepsilon$, $\mathbb{F}_0 = \bigcap_{\varepsilon > 0} \mathbb{F}_\varepsilon$ and \mathbb{F}_0 is the collection of genuine isomorphisms between models in \mathbb{K} ,
 - (b) for all $\varepsilon > 0$, all members of \mathbb{F}_ε have a common modulus of uniform continuity, i.e., there is a function $\Delta^\varepsilon : (0, \infty) \rightarrow (0, \infty)$ such that for any $f : \mathcal{A} \rightarrow \mathcal{B}$ with $f \in \mathbb{F}_\varepsilon$, any $x, y \in \mathcal{A}$, and any $\delta > 0$,
- $$\text{if } d(x, y) < \Delta^\varepsilon(\delta) \text{ then } d(f(x), f(y)) \leq \delta,$$
- (c) if $f \in \mathbb{F}_\varepsilon$ then $f^{-1} \in \mathbb{F}_\varepsilon$,
 - (d) if $f \in \mathbb{F}_\varepsilon$, $g \in \mathbb{F}_\delta$ and $\text{dom}(g) = \text{rng}(f)$ then $g \circ f \in \mathbb{F}_{\varepsilon+\delta}$.
 - (e) if $(f_i)_{i < \alpha}$ is an increasing chain of ε -isomorphisms, i.e. $f_i \in \mathbb{F}_\varepsilon$, $f_i : \mathcal{A}_i \rightarrow \mathcal{B}_i$, $\mathcal{A}_i \preccurlyeq \mathcal{A}_{i+1}$, $\mathcal{B}_i \preccurlyeq \mathcal{B}_{i+1}$ and $f_i \subseteq f_{i+1}$ for all $i < \alpha$, then there is an ε -isomorphism $f : \overline{\bigcup_{i < \alpha} \mathcal{A}_i} \rightarrow \overline{\bigcup_{i < \alpha} \mathcal{B}_i}$ such that $f|_{\mathcal{A}_i} = f_i$ for all $i < \alpha$.

DEFINITION 2.3 (ε -isomorphisms and ε -embeddings). If $\varepsilon \geq 0$, we refer to the mappings $f \in \mathbb{F}_\varepsilon$ as ε -isomorphisms. Correspondingly ε -automorphisms are ε -isomorphisms $f : \mathcal{A} \rightarrow \mathcal{A}$ for some \mathcal{A} . If $\mathcal{A}, \mathcal{B} \in \mathbb{K}$ and $f : \mathcal{A} \rightarrow \mathcal{B}$ is a mapping such that $f : \mathcal{A} \rightarrow f(\mathcal{A})$ is an ε -isomorphism and $f(\mathcal{A}) \preccurlyeq \mathcal{B}$, then f is called an ε -embedding. Here $f(\mathcal{A})$ means the image of \mathcal{A} with the structure induced by the inclusion mapping $f(\mathcal{A}) \hookrightarrow \mathcal{B}$.

It is worth noting that if $f : \mathcal{A} \rightarrow \mathcal{B}$ is an ε -isomorphism and $\mathcal{A}' \preccurlyeq \mathcal{A}$ then $f|_{\mathcal{A}'}$ need not be an ε -embedding.

DEFINITION 2.4 (Joint embedding property). \mathbb{K} is said to have the *joint embedding property* if for any $\mathcal{A}, \mathcal{B} \in \mathbb{K}$ there are $\mathcal{C} \in \mathbb{K}$ and 0-embeddings $f : \mathcal{A} \rightarrow \mathcal{C}$ and $g : \mathcal{B} \rightarrow \mathcal{C}$.

REMARK 2.5. Note that since a MAEC is closed under 0-isomorphisms, when applying the joint embedding property we may assume that one of the

embeddings is the identity mapping. However, with ε -embeddings we need to be more careful, since they do not necessarily preserve \preccurlyeq -submodels.

DEFINITION 2.6 (Amalgamation property). \mathbb{K} is said to have the *amalgamation property* if whenever $\mathcal{A}, \mathcal{B}, \mathcal{C} \in \mathbb{K}$, $\mathcal{A} \preccurlyeq \mathcal{B}$ and $f : \mathcal{A} \rightarrow \mathcal{C}$ is an ε -embedding then there are $\mathcal{B}', \mathcal{C}' \in \mathbb{K}$ with $\mathcal{B}' \succcurlyeq \mathcal{B}$, $\mathcal{C}' \succcurlyeq \mathcal{C}$ and an ε -embedding $g : \mathcal{B}' \rightarrow \mathcal{C}'$ with $g \supseteq f$.

REMARK 2.7. Note that, since 0-embeddings preserve \preccurlyeq -submodels, this version implies the ordinary amalgamation property: if $\mathcal{A} \preccurlyeq \mathcal{B}$ and $\mathcal{A} \preccurlyeq \mathcal{C}$ then there is $\mathcal{D} \in \mathbb{K}$ with $\mathcal{D} \succcurlyeq \mathcal{C}$ and a 0-embedding $f : \mathcal{B} \rightarrow \mathcal{D}$ with $f|_{\mathcal{A}} = \text{id}$.

We also have the following:

LEMMA 2.8. *Assuming the amalgamation property, if $\mathcal{A}', \mathcal{A} \in \mathbb{K}$, $\mathcal{A}' \preccurlyeq \mathcal{A}$ and $f : \mathcal{A}' \rightarrow \mathcal{A}$ is an ε -embedding, then there are $\mathcal{B}, \mathcal{D} \in \mathbb{K}$ with $\mathcal{A} \preccurlyeq \mathcal{B} \preccurlyeq \mathcal{D}$ and an ε -embedding $g : \mathcal{B} \rightarrow \mathcal{D}$ with $g \supseteq f$.*

Proof. By the amalgamation property there are $\mathcal{B} \succcurlyeq \mathcal{A}$ and $\mathcal{C} \succcurlyeq \mathcal{A}$ and an ε -embedding $g' : \mathcal{B} \rightarrow \mathcal{C}$ with $g' \supseteq f$. Then we may use standard amalgamation (with 0-embeddings) and amalgamate \mathcal{B} and \mathcal{C} over \mathcal{A} . So we get $\mathcal{D} \succcurlyeq \mathcal{B}$ and a 0-embedding $h : \mathcal{C} \rightarrow \mathcal{D}$ with $h|_{\mathcal{A}} = \text{id}$. Then $g = h \circ g'$ is an ε -embedding of \mathcal{B} into \mathcal{D} and $g|_{\mathcal{A}'} = h \circ g'|_{\mathcal{A}'} = h \circ f = f$. ■

DEFINITION 2.9 (Galois type in a model). For $\mathcal{A}, \mathcal{B} \in \mathbb{K}$ and $\{a_i : i < \alpha\} \subset \mathcal{A}$, $\{b_i : i < \alpha\} \subset \mathcal{B}$ we say that $(a_i)_{i < \alpha}$ and $(b_i)_{i < \alpha}$ have the same Galois type in \mathcal{A} and \mathcal{B} respectively,

$$\text{t}^g((a_i)_{i < \alpha}/\emptyset; \mathcal{A}) = \text{t}^g((b_i)_{i < \alpha}/\emptyset; \mathcal{B}),$$

if there are $\mathcal{C} \in \mathbb{K}$ and 0-embeddings $f : \mathcal{A} \rightarrow \mathcal{C}$ and $g : \mathcal{B} \rightarrow \mathcal{C}$ such that $f(a_i) = g(b_i)$ for every $i < \alpha$.

The amalgamation property ensures that having the same Galois type is a transitive relation. It is also true that elements have the same Galois type in a model and its \preccurlyeq -extensions and that 0-embeddings preserve Galois types.

DEFINITION 2.10 (0-homogeneity). We call \mathbb{K} 0-homogeneous or *Galois-homogeneous* if whenever $\mathcal{A}, \mathcal{B} \in \mathbb{K}$, $\{a_i : i < \alpha\} \subset \mathcal{A}$, $\{b_i : i < \alpha\} \subset \mathcal{B}$ and for all $n < \omega$, $i_0, \dots, i_{n-1} < \alpha$,

$$\text{t}^g((a_{i_0}, \dots, a_{i_{n-1}})/\emptyset; \mathcal{A}) = \text{t}^g((b_{i_0}, \dots, b_{i_{n-1}})/\emptyset; \mathcal{B})$$

then

$$\text{t}^g((a_i)_{i < \alpha}/\emptyset; \mathcal{A}) = \text{t}^g((b_i)_{i < \alpha}/\emptyset; \mathcal{B}).$$

With the properties defined so far we can construct a 0-homogeneous, ε -model-homogeneous monster model.

THEOREM 2.11. *Let $(\mathbb{K}, \preccurlyeq, \mathbb{F}_\varepsilon)_{\varepsilon \geq 0}$ be a metric abstract elementary class with perturbations satisfying the joint embedding property, the amalgamation property and 0-homogeneity. Let μ be a given infinite cardinal. Then there is $\mathfrak{M} \in \mathbb{K}$ satisfying:*

- (1) (μ -universality) *\mathfrak{M} is μ -universal, i.e. for all $\mathcal{A} \in \mathbb{K}$ with $|\mathcal{A}| < \mu$ there is a 0-embedding $f : \mathcal{A} \rightarrow \mathfrak{M}$.*
- (2) (μ -0-homogeneity) *If $(a_i)_{i < \alpha}$ and $(b_i)_{i < \alpha}$ are sequences of elements of \mathfrak{M} such that $\alpha < \mu$ and for all $n < \omega$ and $i_0, \dots, i_{n-1} < \alpha$,*

$$t^g((a_{i_0}, \dots, a_{i_{n-1}})/\emptyset; \mathfrak{M}) = t^g((b_{i_0}, \dots, b_{i_{n-1}})/\emptyset; \mathfrak{M})$$

then there is a 0-automorphism f of \mathfrak{M} such that $f(a_i) = b_i$ for all $i < \alpha$.

- (3) (μ - ε -model-homogeneity) *If $\mathcal{A} \preccurlyeq \mathfrak{M}$, $|\mathcal{A}| < \mu$ and there is an ε -embedding $f : \mathcal{A} \rightarrow \mathfrak{M}$ then there is an ε -automorphism of \mathfrak{M} extending f .*

Proof. Choose a regular $\kappa > \mu^{N_0}$. We define models \mathcal{A}_α for $\alpha < \kappa$ inductively as follows: When $\alpha = 0$, we use the joint embedding property to construct a μ -universal model \mathcal{A}_0 . When $\alpha = \delta$ is a limit, we let $\mathcal{A}_\delta = \overline{\bigcup_{\beta < \delta} \mathcal{A}_\beta}$. When $\alpha = \beta + 1$ we take care of properties (2) and (3) by an inner induction. Let $(f_i^\beta)_{i < \kappa_\beta}$ list all partial mappings $f : \mathcal{A}_\beta \rightarrow \mathcal{A}_\beta$ such that for all finite $b \in \text{dom}(f)$,

$$t^g(b/\emptyset; \mathcal{A}_\beta) = t^g(f(b)/\emptyset; \mathcal{A}_\beta).$$

Further let $(g_i^\beta)_{i < \kappa_\beta}$ list all mappings g such that $g : \mathcal{A}' \rightarrow \mathcal{A}_\beta$ is an ε -embedding for some $\varepsilon > 0$ and $\mathcal{A}' \preccurlyeq \mathcal{A}_\beta$. Then we define an increasing chain of models \mathcal{B}_i , $i < \kappa_\beta$, such that

- $\mathcal{B}_0 = \mathcal{A}_\beta$,
- $\mathcal{B}_\delta = \overline{\bigcup_{i < \delta} \mathcal{B}_i}$ when $\delta < \kappa_\beta$ is a limit,
- there is a 0-embedding $F_i^\beta : \mathcal{A}_\beta \rightarrow \mathcal{B}_{i+1}$ extending f_i^β ,
- if g_i^β is an ε -embedding $\mathcal{A}' \rightarrow \mathcal{A}_\beta$ then there is some \mathcal{B}' with $\mathcal{B}_i \preccurlyeq \mathcal{B}' \preccurlyeq \mathcal{B}_{i+1}$ and an ε -embedding $G_i^\beta : \mathcal{B}' \rightarrow \mathcal{B}_{i+1}$ extending g_i^β .

When constructing \mathcal{B}_{i+1} first note that by 0-homogeneity and the fact that Galois types are preserved under \preccurlyeq -extensions there are \mathcal{B}'_{i+1} and 0-embeddings $f : \mathcal{A}_\beta \rightarrow \mathcal{B}'_{i+1}$ and $g : \mathcal{B}_i \rightarrow \mathcal{B}'_{i+1}$ such that for all $b \in \text{dom}(f_i^\beta)$, $f(b) = g \circ f_i^\beta$ and as in 2.5 we may assume g is the identity. Hence f gives the required F_i^β . Next note that given g_i^β as above $\mathcal{A}' \preccurlyeq \mathcal{A}_\beta \preccurlyeq \mathcal{B}_i \preccurlyeq \mathcal{B}'_{i+1}$, so by Lemma 2.8 there are $\mathcal{B}_{i+1} \succ \mathcal{B}' \succ \mathcal{B}'_{i+1}$ and an ε -embedding $G_i^\beta : \mathcal{B}' \rightarrow \mathcal{B}_{i+1}$ extending g_i^β .

Hence we can define \mathcal{B}_i for all $i < \kappa_\beta$ and then let $\mathcal{A}_\alpha = \overline{\bigcup_{i < \kappa_\beta} \mathcal{B}_i}$.

Finally let $\mathfrak{M} = \bigcup_{i < \kappa} \mathcal{A}_i$ and note that since $\text{cf}(\kappa) > \omega$, $\mathfrak{M} = \overline{\mathfrak{M}}$. Then clearly \mathfrak{M} is μ -universal and to see that properties (2) and (3) hold note that inverses of type-preserving mappings (0-embeddings) are type-preserving and inverses of ε -embeddings are ε -embeddings. Hence any given embedding with $\text{dom} < \mu$ can be extended by a back-and-forth argument to an automorphism of \mathfrak{M} . Here we use property (2e) of Definition 2.2 to continue the back-and-forth construction through limits. ■

As usual, from now on we will work inside a monster model \mathfrak{M} as in 2.11 for some large enough μ . Then when we write $A \subset \mathfrak{M}$ or $\mathcal{A} \prec \mathfrak{M}$ this will include the assumption that A or \mathcal{A} has size (i.e. density character) less than μ .

NOTATION 2.12. We denote by $\text{Aut}_\varepsilon(\mathfrak{M}/A)$ the set of ε -automorphisms of \mathfrak{M} fixing A pointwise.

DEFINITION 2.13 (Galois type). We say that $(a_i)_{i < \alpha}$ and $(b_i)_{i < \alpha}$ have the same *Galois type over A*, $t^g((a_i)_{i < \alpha}/A) = t^g((b_i)_{i < \alpha}/A)$, if there is $f \in \text{Aut}_0(\mathfrak{M}/A)$ such that $f(a_i) = b_i$ for every $i < \alpha$.

Note that for $A = \emptyset$ the above definition coincides with the notion of Galois type in \mathfrak{M} in Definition 2.9.

In [11] we introduced the *perturbation property*, to achieve what the Perturbation Lemma gives in [10]. Here we will define an even stronger version of perturbation tying our ε -isomorphisms and Galois types together. The role of this property will be discussed in Section 5.

We define a nonnegative real-valued function \mathbf{d}^p on the set of pairs of Galois types of finite tuples over the empty set.

DEFINITION 2.14. For $a, b \in \mathfrak{M}$ and $\varepsilon > 0$ we write

$$\mathbf{d}^p(t^g(a/\emptyset), t^g(b/\emptyset)) \leq \varepsilon$$

if there are ε -automorphisms f and g of \mathfrak{M} such that $d(f(a), b) \leq \varepsilon$ and $d(g(b), a) \leq \varepsilon$.

The perturbation property then tells us that this function vanishes on (p, q) exactly when $p = q$:

DEFINITION 2.15 (perturbation property). Assume \mathbb{K} satisfies the joint embedding property, the amalgamation property and 0-homogeneity so that \mathfrak{M} can be constructed. Then \mathbb{K} is said to have the *perturbation property* if whenever $a, b \in \mathfrak{M}$ are such that $\mathbf{d}^p(t^g(a/\emptyset), t^g(b/\emptyset)) = 0$ (i.e. $\mathbf{d}^p(t^g(a/\emptyset), t^g(b/\emptyset)) \leq \varepsilon$ for all positive ε) then $t^g(a/\emptyset) = t^g(b/\emptyset)$.

We are now ready to list the assumptions we will make on \mathbb{K} .

ASSUMPTION 2.16. From now on we will assume that $(\mathbb{K}, \prec, \mathbb{F}_\varepsilon)_{\varepsilon \geq 0}$ is a metric abstract elementary class with perturbations satisfying the following:

- $\text{LS}^d(\mathbb{K}) = \omega$,
- \mathbb{K} has arbitrarily large models,
- \mathbb{K} has the joint embedding property 2.4,
- \mathbb{K} has the amalgamation property 2.6,
- \mathbb{K} is 0-homogeneous as defined in 2.10,
- \mathbb{K} has the perturbation property 2.15.

Hence we can construct a μ -universal, μ -0-homogeneous and μ - ε -model-homogeneous monster model \mathfrak{M} for some μ larger than any cardinality we will encounter and consider only \preccurlyeq -submodels of \mathfrak{M} .

Besides these assumptions we will mostly need to assume that the type space is complete with respect to \mathbf{d}^p (actually with respect to a metrizable uniformity defined by \mathbf{d}^p , see Section 5), more specifically:

DEFINITION 2.17. We say that \mathbb{K} has *complete type-spaces* if \mathbf{d}^p -Cauchy sequences of types over \emptyset have a limit, i.e. whenever $(a_i)_{i < \omega}$ is a sequence with the property that for all $\varepsilon > 0$ there is $n_0 < \omega$ such that for all $m, n \geq n_0$,

$$\mathbf{d}^p(t^g(a_m/\emptyset), t^g(a_n/\emptyset)) < \varepsilon,$$

then there exists some a with the property that for all $\varepsilon > 0$ there is $n_0 < \omega$ such that for all $n > n_0$,

$$\mathbf{d}^p(t^g(a/\emptyset), t^g(a_n/\emptyset)) < \varepsilon.$$

REMARK 2.18. Assuming perturbation, the definition above implies that \mathbf{d}^p -Cauchy sequences of types with finite parameter sets have limits, i.e. if $(t^g(a_n A/\emptyset))_{n < \omega}$ is a \mathbf{d}^p -Cauchy sequence then there is some a such that $t^g(a A/\emptyset)$ is a limit type of the sequence. To prove this, use the above property for the sequence $(t^g(a_n A/\emptyset))_{n < \omega}$. This has a limit type realized by some $a' A^+$. But $\mathbf{d}^p(t^g(A/\emptyset), t^g(A^+/\emptyset)) < \varepsilon$ for all $\varepsilon > 0$ so by perturbation they have the same Galois type and we may find $f \in \text{Aut}_0(\mathfrak{M})$ with $f(A^+) = A$. Then $a = f(a')$ is the desired limit element.

Completeness of type-spaces is our substitute for compactness. It, however, is a weaker property, as it holds in noncompact classes: In Section 4 we show that atomic Nakano spaces with bounded exponent satisfy completeness of type-spaces, although not even the unit balls of these spaces form a compact model class.

In some constructions the assumption of completeness of type-spaces can be replaced by the assumption of models being *good* (i.e. the pair $(\mathcal{A}, \mathcal{A})$ is good).

DEFINITION 2.19. If $a \in \mathcal{A}$ and $b \in \mathcal{B}$, we write

$$\mathbf{d}_\varepsilon(t^g(a/\emptyset; \mathcal{A}), t^g(b/\emptyset; \mathcal{B})) < \delta$$

if there are $\mathcal{A}' \succcurlyeq \mathcal{A}$ and $\mathcal{B}' \succcurlyeq \mathcal{B}$ and an ε -embedding $f : \mathcal{A}' \rightarrow \mathcal{B}'$ such that $d(f(a), b) < \delta$.

We then say that $f : A \rightarrow \mathcal{B}$ for $A \subset \mathcal{A}$ is a *weak ε -embedding* if for all $a \in A$, $\mathbf{d}_\varepsilon(t^g(a/\emptyset; \mathcal{A}), t^g(f(a)/\emptyset; \mathcal{B})) < \delta$ for all $\delta > 0$.

DEFINITION 2.20. Assume $A \subseteq \mathcal{A}$. We say that the pair (A, \mathcal{A}) is *good* if for all $\varepsilon > 0$ there is some $\delta > 0$ such that if $f : A \rightarrow \mathcal{B}$ is a weak δ -embedding then there are $\mathcal{A}' \succcurlyeq \mathcal{A}$, $\mathcal{B}' \succcurlyeq \mathcal{B}$ and an ε -embedding $g : \mathcal{A}' \rightarrow \mathcal{B}'$ such that $g \supseteq f$. (Note that $g|_{\mathcal{A}}$ need not be an ε -embedding.)

DEFINITION 2.21. When $A \subseteq \mathfrak{M}$, we say that A is *good in \mathfrak{M}* if for all $\varepsilon > 0$ there is some $\delta > 0$ such that for all weak δ -embeddings $f : A \rightarrow \mathfrak{M}$ there is $g \in \text{Aut}_\varepsilon(\mathfrak{M})$ such that $g \supseteq f$.

LEMMA 2.22. Let μ be as in the construction of \mathfrak{M} . Assume $A \subset \mathfrak{M}$ and $|A| < \mu$. Then A is good in \mathfrak{M} if and only if (A, \mathcal{A}) is good for some $\mathcal{A} \preccurlyeq \mathfrak{M}$ with $A \subseteq \mathcal{A}$ and $|\mathcal{A}| < \mu$. In particular, $\mathcal{A} \preccurlyeq \mathfrak{M}$ with $|\mathcal{A}| < \mu$ is good in \mathfrak{M} if and only if $(\mathcal{A}, \mathcal{A})$ is good.

Proof. First note that if $|\mathcal{A}| < \mu$ and a given mapping extends to an ε -embedding $\mathcal{A} \preccurlyeq \mathcal{A}' \rightarrow \mathcal{B}'$, we may always assume $|\mathcal{A}'|, |\mathcal{B}'| < \mu$, since by the assumption $\text{LS}^d = \omega$ and a back-and-forth construction we may find such small enough models (assuming $\mu > \aleph_0$). Hence if A is good in \mathfrak{M} , (A, \mathcal{A}) is good for any $\mathcal{A} \supset A$ in \mathfrak{M} . On the other hand, if (A, \mathcal{A}) is good for some small enough \mathcal{A} , and $f : \mathcal{A}' \rightarrow \mathcal{B}'$ is an ε -embedding extending some given weak embedding of A , we may assume $\mathcal{A}', \mathcal{B}'$ are small enough and hence inside \mathfrak{M} (by universality). Then f has an extension to an automorphism inside \mathfrak{M} . ■

In this context the natural notion of stability will be with respect to \mathbf{d}^p . However, for this we first need to generalize the definition.

DEFINITION 2.23. For $a, b \in \mathfrak{M}$ and $A \subset \mathfrak{M}$, we define

$$\mathbf{d}^p(t^g(a/A), t^g(b/A)) = \sup\{\mathbf{d}^p(t^g(ac/\emptyset), t^g(bc/\emptyset)) : c \in A \text{ finite}\}.$$

Then stability is defined in the natural way.

DEFINITION 2.24. We say that \mathbb{K} is λ - \mathbf{d}^p -stable if for any set A with $|A| \leq \lambda$, the set of types over A has density λ with respect to \mathbf{d}^p , i.e. looking at the situation inside \mathfrak{M} , there is a set A^* of cardinality λ such that for any $a \in \mathfrak{M}$ and $\varepsilon > 0$ there is some $a' \in A^*$ such that $\mathbf{d}^p(t^g(a/A), t^g(a'/A)) < \varepsilon$.

In Section 5 we will see that \mathbf{d}^p defines a metrizable uniformity on the type-space and that the above definition hence makes sense.

3. Hilbert spaces with an automorphism. We now turn to an example of a MAEC with perturbations investigated in [7] and [4] in the continuous setting as an example of a theory of continuous logic that is not

ω -stable but ω -stable up to perturbation (see e.g. [1]). The idea is that you allow arbitrarily small perturbations of parts of the language of the structure. In the setting of metric abstract elementary classes with perturbations, these perturbations are built into the \mathbb{F}_ε 's.

DEFINITION 3.1. Let \mathbb{K}_H be the class of complex Hilbert spaces equipped with an automorphism τ , i.e. τ is a surjective unitary operator. We then define $(H', \tau') \preccurlyeq (H, \tau)$ as the ordinary submodel relation, i.e. the \preccurlyeq -submodels of (H, τ) are closed subspaces closed under τ .

When $\varepsilon \geq 0$ we let $f : (H_1, \tau_1) \rightarrow (H_2, \tau_2)$ belong to \mathbb{F}_ε if f is an isometric isomorphism between the Hilbert spaces H_1 and H_2 and additionally

$$\|\tau_1 - f^{-1}\tau_2 f\| \leq \varepsilon \quad \text{and} \quad \|\tau_1^{-1} - f^{-1}\tau_2^{-1} f\| \leq \varepsilon$$

where $\|\cdot\|$ is the *operator norm* defined by $\|T\| = \sup_{\|x\|=1} \|T(x)\|$.

Ben Yaacov, Berenstein, Usvyatsov and Zadka consider generic automorphisms. Since considering all automorphisms or just the generic ones produces the same monster model, either choice constitutes an example class and we have chosen to let τ be any automorphism. Our demands on the ε -isomorphisms arise from the form of perturbation Ben Yaacov and Berenstein use in [4]: they define an *r-perturbation* of (H_0, τ_0) to (H_1, τ_1) to be an isometric isomorphism $U : H_0 \cong H_1$ satisfying $\|U\tau_0 U^{-1} - \tau_1\| \leq r$. This in turn is connected to a criterion by Ben Yaacov, Usvyatsov and Zadka for an automorphism to be generic and to the Weyl–von Neumann–Berg theorem. To state these results we need the definitions of some spectra of a bounded operator.

DEFINITION 3.2. Consider a Hilbert space H and a bounded operator T on H . Then we can define the *spectrum*, the *point spectrum* and the *essential spectrum* of T by

$$\begin{aligned} \sigma(T) &= \{\lambda \in \mathbb{C} : T - \lambda I \text{ is not invertible}\}, \\ \sigma_p(T) &= \{\lambda \in \mathbb{C} : \ker(T - \lambda I) \neq 0\}, \\ \sigma_e(T) &= \{\text{non-isolated points of } \sigma(T)\} \\ &\quad \cup \{\lambda \in \mathbb{C} : \dim \ker(T - \lambda I) = \infty\}. \end{aligned}$$

FACT 3.3 (Ben Yaacov, Usvyatov, Zadka [7]). *Let H be a Hilbert space and let τ be a unitary operator on H . Then (H, τ) is existentially closed as a model of the continuous first order theory of infinite-dimensional Hilbert spaces with an automorphism if and only if $\sigma(\tau) = S^1$ where S^1 is the unit circle.*

DEFINITION 3.4. Let H be a Hilbert space and let T_0, T_1 be bounded operators on H . Then T_0 and T_1 are *approximately unitarily equivalent* if there is a sequence $(U_n)_{n<\omega}$ of unitary operators such that $\|T_1 - U_n T_0 U_n^*\| \rightarrow 0$.

FACT 3.5 (Weyl–von Neumann–Berg Theorem). *Let H be a separable Hilbert space and let T_0, T_1 be normal operators on H . Then T_0 and T_1 are approximately unitarily equivalent if and only if*

- (1) $\sigma_e(T_0) = \sigma_e(T_1)$,
- (2) $\dim \ker(T_0 - \lambda I) = \dim \ker(T_1 - \lambda I)$ for all $\lambda \in \mathbb{C} \setminus \sigma_e(T_0)$.

This characterization of approximately unitarily equivalent spaces is the key point in the proof of ω -stability up to perturbation (see [4] and [7]).

FACT 3.6 (Ben Yaacov, Berenstein, Usvyatsov, Zadka). *The continuous first order theory of infinite-dimensional Hilbert spaces with a generic automorphism is ω -stable up to perturbation of the automorphism.*

REMARK 3.7. The above theory is not ω -stable in the ordinary metric sense, i.e. with respect to the distance defined as the infimum of distances of realizations of types. This is pointed out in [4, Remark 1.4], where the credit for the observation is given to Henson and Iovino.

We next turn to proving that the class defined in Definition 3.1 indeed satisfies the assumptions in 2.16.

LEMMA 3.8. *The class $(\mathbb{K}_H, \preccurlyeq)$ is a metric abstract elementary class with Löwenheim–Skolem number \aleph_0 , having arbitrarily large models and satisfying joint embedding and 0-homogeneity.*

Proof. It is trivial to see that $(\mathbb{K}_H, \preccurlyeq)$ forms a MAEC with $\text{LS}^d = \aleph_0$ and arbitrarily large models satisfying 0-homogeneity since the class is axiomatizable in continuous first order logic and joint embedding is taken care of by orthogonal direct sums. ■

LEMMA 3.9. *The collections \mathbb{F}_ε defined in 3.1 satisfy the properties of Definition 2.2.*

Proof. This is easily seen if we remember that in this example ε -isomorphisms are isometries of the underlying Hilbert spaces. ■

LEMMA 3.10. *The class defined in 3.1 has the amalgamation property.*

Proof. Assume that $(H', \tau') \preccurlyeq (H, \tau)$ and $f : (H', \tau') \rightarrow (H_1, \tau_1)$ is an ε -embedding. Then we may write $(H, \tau) = (H', \tau') \oplus (H'', \tau'')$ and embed (H, τ) into $(H_1, \tau_1) \oplus (H'', \tau'')$ with $f \oplus \text{id} \supseteq f$. ■

Since the amalgamation property holds, we have a monster model for the class and may consider our last property, perturbation.

LEMMA 3.11. *The class defined in 3.1 has the perturbation property.*

Proof. Let $a, b \in \mathfrak{M}$ be such that $\mathbf{d}^p(t^g(a/\emptyset), t^g(b/\emptyset)) = 0$, i.e. for all $\varepsilon > 0$ there are ε -automorphisms of \mathfrak{M} mapping a ε -close to b and vice versa.

Now consider the spaces spanned by $\{\tau^k a : k \in \mathbb{Z}\}$ and $\{\tau^k b : k \in \mathbb{Z}\}$ respectively. The assumption implies that any finite part $\{\tau^k a : -m < k < m\}$ can be mapped arbitrarily close to the corresponding set $\{\tau^k b : -m < k < m\}$ by an isometric isomorphism of the underlying Hilbert space. Hence the Hilbert spaces

$$\overline{\text{Span}\{\tau^k a : k \in \mathbb{Z}\}} \quad \text{and} \quad \overline{\text{Span}\{\tau^k b : k \in \mathbb{Z}\}}$$

are isometrically isomorphic, and since these spaces are closed under τ they are \preccurlyeq -submodels of \mathfrak{M} . Further, the function mapping each $\tau^k a$ to $\tau^k b$ makes no error in mapping τ , so the isomorphism is a 0-isomorphism of $(\mathbb{K}_H, \preccurlyeq, \mathbb{F}_\varepsilon)_{\varepsilon \geq 0}$ and hence extends to a 0-automorphism of \mathfrak{M} . ■

Collecting the results, we have:

THEOREM 3.12. *The class of Hilbert spaces with an automorphism, with \preccurlyeq defined as the ordinary submodel relation and the classes \mathbb{F}_ε defined as in 3.1, forms a metric abstract elementary class satisfying the assumptions in 2.16.*

We also have both completeness of type-spaces and goodness:

THEOREM 3.13. *\mathbb{K}_H has complete type-spaces.*

Proof. This follows from the fact that the class of Hilbert spaces with an automorphism is axiomatizable in continuous first order logic and hence closed under ultraproducts (see [5]). Now let $(a_i)_{i < \omega}$ be a sequence such that $(t^g(a_i/\emptyset))_{i < \omega}$ is \mathbf{d}^p -Cauchy. By switching to a subsequence if necessary, we may assume that $\mathbf{d}^p(t^g(a_j/\emptyset), t^g(a_i/\emptyset)) \leq 2^{-i}$ for all $i < j < \omega$. Then let f_{ji} be a 2^{-i} -automorphism witnessing this, i.e. satisfying $d(f_{ji}(a_j), a_i) \leq 2^{-i}$. Let $\{a_i : i < \omega\} \subset \mathcal{A} \preccurlyeq \mathfrak{M}$ be such that \mathcal{A} is closed under each f_{ji} . Further let D be an ultrafilter on ω extending the Fréchet filter and consider the ultrapower $(\mathcal{A})_D$. Then the diagonal embeddings of each a_i still form a \mathbf{d}^p -Cauchy sequence, witnessed by the functions $((f_{ji})_{n < \omega})/D$, which are 2^{-i} -automorphisms of $(\mathcal{A})_D$.

Now the sequence has a \mathbf{d}^p -limit, namely $a = ((a_i)_{i < \omega})/D$, and this is seen by considering the mappings

$$F_i = ((f_{ji})_{j < \omega})/D,$$

where we let $f_{ji} = \text{id}$ if $j \leq i$. Since the componentwise mappings of F_i are isometric automorphisms of the underlying Hilbert space of \mathcal{A} and the componentwise error in mapping τ is at most 2^{-i} , F_i is a 2^{-i} -automorphism of $(\mathcal{A})_D$. Also clearly the distance of $F_i(a)$ to the diagonal embedding of a_i is at most 2^{-i} for each $i < \omega$. ■

THEOREM 3.14. *If $\mathcal{A} \in \mathbb{K}_H$ then $(\mathcal{A}, \mathcal{A})$ is good as defined in 2.20.*

Proof. Let $\mathcal{A} \in \mathbb{K}_H$ and $\varepsilon > 0$ be given. We choose $\delta = \varepsilon/(2 + \sqrt{8})$ and show that if $f : \mathcal{A} \rightarrow \mathcal{B}$ is a weak δ -embedding, then there are $\mathcal{A}' \succcurlyeq \mathcal{A}$ and $\mathcal{B}' \succcurlyeq \mathcal{B}$ and an ε -embedding $g : \mathcal{A}' \rightarrow \mathcal{B}'$ extending f .

So let $f : \mathcal{A} \rightarrow \mathcal{B}$ be a weak δ -embedding, i.e. for all (finite tuples) $a \in \mathcal{A}$ and $\delta' > 0$ there are $\mathcal{A}_1 \succcurlyeq \mathcal{A}$ and $\mathcal{B}_1 \succcurlyeq \mathcal{B}$ and a δ -embedding $f_1 : \mathcal{A}_1 \rightarrow \mathcal{B}_1$ such that $d(f_1(a), f(a)) < \delta'$. Since δ -embeddings are isometric isomorphisms of the underlying Hilbert space, we see that weak embeddings must also be isometric isomorphisms of the Hilbert space. Further for any $a \in \mathcal{A}$ with $\|a\| = 1$ and any $\delta' > 0$ there is a δ -embedding f_1 for the tuple $(a, \tau_{\mathcal{A}}(a), \tau_{\mathcal{A}}^{-1}(a))$ as above and hence (noting that $\tau_{\mathcal{A}} = \tau_{\mathcal{A}_1}|_{\mathcal{A}}$ and similarly for $\tau_{\mathcal{B}}$)

$$\begin{aligned} \|(f\tau_{\mathcal{A}} - \tau_{\mathcal{B}}f)(a)\| &\leq \|(f\tau_{\mathcal{A}} - f_1\tau_{\mathcal{A}})(a)\| + \|(f_1\tau_{\mathcal{A}} - \tau_{\mathcal{B}_1}f_1)(a)\| \\ &\quad + \|(\tau_{\mathcal{B}_1}f_1 - \tau_{\mathcal{B}}f)(a)\| \\ &\leq \delta' + \delta + \delta', \end{aligned}$$

and similarly for τ^{-1} . Hence weak embeddings are isometric isomorphisms of the underlying Hilbert space satisfying the norm requirements for embeddings but whose images are not necessarily submodels of the target model, i.e. closed under τ . Since f is an isometric isomorphism of the Hilbert space, we know however that $f(\mathcal{A})$ must be a closed subspace of \mathcal{B} , so writing $\mathcal{B} = (H_{\mathcal{B}}, \tau_{\mathcal{B}})$ we may write $H_{\mathcal{B}} = H \oplus H'$ where $H = \text{rng}(f)$.

We now define $\mathcal{A}' = \mathcal{B}' = \mathcal{A} \oplus \mathcal{B}$, (i.e. $(H_{\mathcal{A}}, \tau_{\mathcal{A}}) \oplus (H_{\mathcal{B}}, \tau_{\mathcal{B}})$). Clearly $\mathcal{A} \preccurlyeq \mathcal{A}'$ and $\mathcal{B} \preccurlyeq \mathcal{B}'$ so we just need to find an ε -embedding $f' : \mathcal{A}' \rightarrow \mathcal{B}'$ extending f . But here we just define $f'|_{\mathcal{A}} = f$, $f'|_H = f^{-1}$ and $f'|_{H'} = \text{id}$. Then f' is an isometric isomorphism of the underlying Hilbert spaces and to see that it satisfies $\|\tau_{\mathcal{A}'} - f'^{-1}\tau_{\mathcal{B}'}f'\| \leq \varepsilon$ (we prove only the first condition since $\|\tau_{\mathcal{A}'}^{-1} - f'^{-1}\tau_{\mathcal{B}'}^{-1}f'\| \leq \varepsilon$ can be proved similarly) we first consider the parts separately:

CASE 1: If $a \in H_{\mathcal{A}}$ and $\|a\| = 1$ then since $f'|_{\mathcal{A}} = f$ we have

$$\|f'\tau_{\mathcal{A}'}(a) - \tau_{\mathcal{B}'}f'(a)\| = \|f\tau_{\mathcal{A}}(a) - \tau_{\mathcal{B}}f(a)\| \leq \delta.$$

CASE 2: If $b \in H$ and $\|b\| = 1$, then $b = f(a)$ for some $a \in \mathcal{A}$ with $\|a\| = 1$ and $\|f\tau_{\mathcal{A}}(a) - \tau_{\mathcal{B}}f(a)\| \leq \delta$. Hence

$$\begin{aligned} \|f'\tau_{\mathcal{A}'}(b) - \tau_{\mathcal{B}'}f'(b)\| &= \|f'\tau_{\mathcal{B}}(b) - \tau_{\mathcal{A}}f'(b)\| = \|f'\tau_{\mathcal{B}}f(a) - f'f\tau_{\mathcal{A}}(a)\| \\ &= \|\tau_{\mathcal{B}}f(a) - f\tau_{\mathcal{A}}(a)\| \leq \delta. \end{aligned}$$

CASE 3: If $b \in H'$ and $\|b\| = 1$, then since $\tau_{\mathcal{A}'}(b) = \tau_{\mathcal{B}}(b) \in \mathcal{B}$, there are $x \in H$ and $y \in H'$ such that $\tau_{\mathcal{A}'}(b) = x + y$. Then

$$\|f'\tau_{\mathcal{A}'}(b) - \tau_{\mathcal{B}'}f'(b)\| = \|f'(x + y) - (x + y)\| = \|f'(x) + x\| \leq \sqrt{2}\|x\|.$$

Now $\tau_{\mathcal{A}'}^{-1}(x + y) = b \in H'$, so $\text{Pr}_{H'}(\tau_{\mathcal{A}'}^{-1}(x)) + \text{Pr}_{H'}(\tau_{\mathcal{A}'}^{-1}(y)) = b$. Further $\|\text{Pr}_{H'}(\tau_{\mathcal{A}'}^{-1}(y))\| \leq \|y\|$ and since $\tau_{\mathcal{A}'}^{-1}(x)$ is $\delta\|x\|$ -close to something in H ,

$\|\text{Pr}_{H'}(\tau_{\mathcal{A}'}^{-1}(x))\| \leq \delta \|x\|$. Hence we have $1 = \|b\| \leq \delta \|x\| + \|y\|$ but by definition of x and y , $1 = \|b\|^2 = \|x\|^2 + \|y\|^2$. Hence

$$1 - \|x\|^2 = \|y\|^2 \geq (1 - \delta \|x\|)^2$$

and so

$$\|x\| \leq \frac{2\delta}{\delta^2 + 1} \leq 2\delta.$$

Combining this with the result above, we have

$$\|f'\tau_{\mathcal{A}'}(b) - \tau_{\mathcal{B}'}f'(b)\| \leq \sqrt{2}\|x\| \leq \sqrt{8}\delta.$$

Finally, if $a \in \mathcal{A}'$ is arbitrary, there are $x \in \mathcal{A}$, $y \in H$ and $z \in H'$ such that $a = x + y + z$. Then

$$\begin{aligned} \|(f'\tau_{\mathcal{A}'} - \tau_{\mathcal{B}'}f')(a)\| &\leq \|(f'\tau_{\mathcal{A}'} - \tau_{\mathcal{B}'}f')(x)\| + \|(f'\tau_{\mathcal{A}'} - \tau_{\mathcal{B}'}f')(y)\| \\ &\quad + \|(f'\tau_{\mathcal{A}'} - \tau_{\mathcal{B}'}f')(z)\| \\ &\leq \delta\|x\| + \delta\|y\| + \sqrt{8}\delta\|z\| \leq (2 + \sqrt{8})\delta\|a\| \leq \varepsilon\|a\|. \blacksquare \end{aligned}$$

THEOREM 3.15. \mathbb{K}_H is $\omega\text{-d}^p$ -stable. Actually, if $A \subset \mathfrak{M}$ is separable, then there is a countable set $A^* \subset \mathfrak{M}$ such that for any $a \in \mathfrak{M}$ and $\varepsilon > 0$ there is some $a' \in A^*$ and an ε -automorphism f of \mathfrak{M} such that $f|A = \text{id}$ and $d(f(a), a') < \varepsilon$.

Proof. This is proved by imitating Ben Yaacov's and Berenstein's proof of Fact 3.6. Let $A \subset \mathfrak{M}$ be separable. Then we may consider a separable $\mathcal{A} \preccurlyeq \mathfrak{M}$ containing A and by universality of \mathfrak{M} another separable $\mathcal{B} \preccurlyeq \mathfrak{M}$ such that $\tau_{\mathcal{B}}$ has full spectrum and $\mathcal{A} \cap \mathcal{B} = \{0\}$. Now consider $\mathcal{C} = \mathcal{A} \oplus \mathcal{B}$. We claim that any dense (countable) subset of \mathcal{C} will do as our A^* . So let $a \in \mathfrak{M}$. If $a \in \mathcal{A}$, there is nothing to prove. Otherwise consider $d = a - P_{\mathcal{A}}(a)$, where $P_{\mathcal{A}}$ is the orthogonal projection onto \mathcal{A} . By universality of \mathfrak{M} , let \mathcal{D} be a separable model containing d with $\mathcal{A} \cap \mathcal{D} = \{0\}$ and such that $\tau_{\mathcal{D}}$ has full spectrum. Fixing an isometric isomorphism of the Hilbert spaces of \mathcal{B} and \mathcal{D} and using Fact 3.5 we may for any $\varepsilon > 0$ find an ε -isomorphism $f : \mathcal{D} \rightarrow \mathcal{B}$. Then $\text{id}_{\mathcal{A}} \oplus f : \mathcal{A} \oplus \mathcal{D} \rightarrow \mathcal{A} \oplus \mathcal{B}$ belongs to \mathbb{F}_{ε} and extends to an ε -automorphism of \mathfrak{M} and $(\text{id}_{\mathcal{A}} \oplus f)(P_{\mathcal{A}}(a) + d)$ is the desired realization in \mathcal{C} . ■

REMARK 3.16. By considerations like the ones in the proof above we see that all (H, τ) where τ is generic (i.e. has full spectrum) are $\omega\text{-d}^p$ -saturated (see Definition 6.1).

4. Nakano spaces. In this section we study another example, of real-valued atomic Nakano spaces (variable exponent Lebesgue spaces) with bounded exponent. Nakano spaces with bounded exponent have been studied by Poitevin [13] and Ben Yaacov [3] but the purely atomic case falls outside the scope of continuous logic. When the exponent is unbounded, the

simple functions do not form a dense subset of the Nakano space and thus the spaces fail to form even a MAEC. However, they can be studied in a somewhat more general framework that will be presented in another paper by the present authors. The class is homogeneous giving rise to a nice model theory although ω -stability naturally fails.

Let (X, Σ, μ) be a measure space and let $L_0(X, \Sigma, \mu)$ be the space of all measurable functions $f : X \rightarrow \mathbb{R}$ up to equality almost everywhere. Further let $p : X \rightarrow [1, \infty)$ be a measurable function. Define the *convex modular* $\Theta_{p(\cdot)} : L_0(X, \Sigma, \mu) \rightarrow [0, \infty]$ by

$$\Theta_{p(\cdot)}(f) = \int_X |f(x)|^{p(x)} d\mu.$$

The corresponding *Nakano space* is defined as

$$L_{p(\cdot)}(X, \Sigma, \mu) = \{f \in L_0(X, \Sigma, \mu) : \Theta_{p(\cdot)}(\lambda f) < \infty \text{ for some } \lambda > 0\}.$$

This is a Banach space equipped with the *Luxemburg norm*

$$\|f\|_{p(\cdot)} = \inf\{\lambda > 0 : \Theta_{p(\cdot)}(f/\lambda) \leq 1\}.$$

In the purely atomic case this boils down to the following: Let I be an index set and $p : I \rightarrow [1, \infty)$. The modular $\Theta_{p(\cdot)} : \{(a_i)_{i \in I} : a_i \in \mathbb{R}\} \rightarrow [0, \infty]$ is defined by

$$\Theta_{p(\cdot)}((a_i)_{i \in I}) = \sum_{i \in I} |a_i|^{p(i)}.$$

Then the atomic Nakano space $\ell^{p(\cdot)}(I)$ is defined by

$$\ell^{p(\cdot)}(I) = \{(a_i)_{i \in I} : \Theta_{p(\cdot)}((\lambda a_i)_{i \in I}) < \infty \text{ for some } \lambda > 0\}$$

and the Luxemburg norm is

$$\|(a_i)_{i \in I}\|_{p(\cdot)} = \inf\{\lambda > 0 : \Theta_{p(\cdot)}((a_i/\lambda)_{i \in I}) \leq 1\}.$$

The modular and norm satisfy the following relations:

FACT 4.1 ([9, Corollary 1.1.14 and Lemma 2.4.2]). *In any Nakano space*

- (1) *if $\|x\| \leq 1$, then $\Theta(x) \leq \|x\|$,*
- (2) *if $\|x\| > 1$, then $\|x\| \leq \Theta(x)$.*

In addition, if the exponent is bounded,

- (3) *$\|x\| \leq 1$ if and only if $\Theta(x) \leq 1$,*
- (4) *$\|x\| < 1$ if and only if $\Theta(x) < 1$.*

We will study Nakano spaces as Banach lattices with the vocabulary

$$\mathcal{L}_{Bl} = \{0, -, +, \cdot, \|\cdot\|, \wedge, \vee\}$$

and with the pointwise interpretation of the lattice operators. The class will be determined by a real $r \geq 1$, the bound of the exponent of the spaces.

In [3] Ben Yaacov classifies Banach lattice isometries of Nakano spaces and proves that up to a measure density change they send characteristic functions to characteristic functions and preserve both measures and the exponent functions. In the atomic case we can prove a slightly stronger result, as no measure changes can occur.

THEOREM 4.2. *Let $\ell^{p(\cdot)}(I)$ and $\ell^{q(\cdot)}(J)$ be atomic Nakano spaces with bounded exponent and $|I| \geq 2$. Denote by $(e_i)_{i \in I}$ and $(e_j)_{j \in J}$ their standard bases. If U is a Banach lattice isometry from $\ell^{p(\cdot)}(I)$ to $\ell^{q(\cdot)}(J)$ then U is of the form*

$$U\left(\sum_{i \in I} a_i e_i\right) = \sum_{i \in I} a_i e_{\sigma(i)},$$

where σ is a bijection from I to J satisfying $q(\sigma(i)) = p(i)$ for all $i \in I$.

Proof. Let e_i be a basic vector of $\ell^{p(\cdot)}(I)$ and let $x = U(e_i) = \sum_{j \in J} a_j e_j$. Now

$$e_i = U^{-1}\left(\sum_{j \in J} a_j e_j\right) = \sum_{j \in J} a_j U^{-1}(e_j).$$

Since U^{-1} is a lattice isometry, it must preserve disjointness of supports, so

$$1 = |\text{supp}(e_i)| \geq \left| \text{supp} \left(\sum_{j \in J'} a_j e_j \right) \right|,$$

from which we deduce that $U(e_i) = a_j e_j$ for some $j \in J$. Taking into account that $\|e_i\| = 1 = \|e_j\|$ and $e_i \geq 0$ in the lattice order for any basic vector, we see that $a_j = 1$, i.e., $U(e_i) = e_j$.

To show that U preserves the exponents let $i, k \in I$ and $j, l \in J$ be such that $U(e_i) = e_j$, $U(e_k) = e_l$. For $t \in [0, 1]$ define

$$f_t = t^{1/p_i} e_i + (1 - t)^{1/p_k} e_k, \quad g_t = U(f_t) = t^{1/p_i} e_j + (1 - t)^{1/p_k} e_l.$$

Now

$$\Theta_{p(\cdot)}(f_t) = 1 \Rightarrow \|f_t\| = 1 \Rightarrow \|g_t\| = 1 \Rightarrow \Theta_{q(\cdot)}(g_t) = 1,$$

so

$$1 = \Theta_{q(\cdot)}(g_t) = t^{q_j/p_i} + (1 - t)^{q_l/p_k}, \quad \text{i.e. } (1 - t)^{q_l/p_k} = 1 - t^{q_j/p_i},$$

and since this has to hold for all $t \in [0, 1]$, we must have $q_l/p_k = q_j/p_i = 1$. ■

We wish to study Nakano spaces with respect to generalized isomorphisms. In [3] Ben Yaacov defines perturbation mappings as follows:

DEFINITION 4.3 (Ben Yaacov). Let (X, Σ, μ) be a measure space and $p, q : X \rightarrow [1, r]$ measurable. Define $\mathcal{E}_{p,q} : L_0(X, \Sigma, \mu) \rightarrow L_0(X, \Sigma, \mu)$ by

$$(\mathcal{E}_{p,q})(f) = \text{sgn}(f(x))|f(x)|^{p(x)/q(x)}.$$

FACT 4.4 (Ben Yaacov). *Let $(N, \Theta) = (L_{p(\cdot)}(X, \Sigma, \mu), \Theta_{p(\cdot)})$ and $(N', \Theta') = (L_{q(\cdot)}(X, \Sigma, \mu), \Theta_{q(\cdot)})$.*

- (1) *For each $f \in L_0(X, \Sigma, \mu)$ we have $\Theta(f) = \Theta'(\mathcal{E}_{p,q}f)$.*
- (2) *$\mathcal{E}_{p,q}$ is bijective and restricts to a bijection between N and N' .*
- (3) *$\mathcal{E}_{p,q}$ is uniformly continuous on the unit ball of N .*

We define our ε -isomorphisms based on Ben Yaacov's definition.

DEFINITION 4.5. We define the classes \mathbb{F}_ε to consist of all bijective map-pings

$$U : \ell^{p(\cdot)}(I) \rightarrow \ell^{q(\cdot)}(J)$$

given by

$$U\left(\sum_{i \in I} a_i e_i\right) = \sum_{i \in I} \text{sgn}(a_i) |a_i|^{p(i)/q(\sigma(i))} e_{\sigma(i)},$$

where σ is a bijection $I \rightarrow J$ satisfying $e^{-\varepsilon} \leq q(\sigma(i))/p(i) \leq e^\varepsilon$.

This definition, however, gives functions that are uniformly continuous only on bounded subsets of the space. In order to correct this we switch to a metric d which is equivalent to the one induced by the norm, thus giving the same notion of closure, but which moves attention to the unit ball, thus rendering our ε -isomorphisms uniformly continuous.

DEFINITION 4.6. Let $F_\lambda : \ell^{p(\cdot)}(I) \rightarrow \ell^{p(\cdot)}(I)$ be a scaling operator defined by

$$F_\lambda((a_i)_{i \in I}) = (\lambda^{-1/p(i)} a_i)_{i \in I}.$$

Let $F : \ell^{p(\cdot)}(I) \rightarrow \ell^{p(\cdot)}(I)$ be defined by

$$F(a) = \begin{cases} a & \text{if } \Theta(a) \leq 1, \\ F_{\Theta(a)}(a) & \text{otherwise.} \end{cases}$$

and $G : \ell^{p(\cdot)}(I) \rightarrow [1, \infty)$ by

$$G(a) = \max\{1, \Theta(a)\}.$$

Then define

$$d(a, b) = \|F(a) - F(b)\| + |G(a) - G(b)|.$$

It is straightforward to check that d is a metric on $\ell^{p(\cdot)}(I)$. We will show that it is also (topologically) equivalent to the norm metric $d_{\|\cdot\|}$. For this we need the following observation.

LEMMA 4.7. *For any $\lambda > 0$, F_λ is a bounded linear mapping satisfying:*

- if $\lambda \geq 1$ then $\lambda^{-1}\|a\| \leq \|F_\lambda(a)\| \leq \|a\|$,
- if $\lambda \leq 1$ then $\|a\| \leq \|F_\lambda(a)\| \leq \lambda^{-1}\|a\|$.

For $\eta, \lambda > 0$, $F_\lambda \circ F_\eta = F_{\lambda\eta}$.

Proof. Straightforward calculations. ■

PROPOSITION 4.8. *d is topologically equivalent to $d_{\|\cdot\|}$.*

Proof. We need to prove that every d -ball contains a $\|\cdot\|$ -ball and vice versa. So let $x \in \ell^{p(\cdot)}(I)$ and $r > 0$ be given.

First consider $B_d(x, r)$, the d -ball of radius r around x . Since the exponent is bounded, by [13, Lemma 2.1.5] the map $\Theta : (\ell^{p(\cdot)}(I), \|\cdot\|) \rightarrow [0, \infty)$ is uniformly continuous on bounded subsets. So there is a modulus of uniform continuity Δ for Θ on the $\|\cdot\|$ -ball of radius $\|x\| + 1$ around the origin. Now let $\varepsilon < \min\{r/3, r/(3\|x\|)\}$ and let $r' > 0$ be such that

$$r' < \min\{r/3, 1, \Delta(\varepsilon)\}$$

and additionally, if $\Theta(x) > 1$ (in which case also $\|x\| > 1$), we require

$$(4.1) \quad r' < \|x\| - 1.$$

Now let $y \in B_{\|\cdot\|}(x, r')$. Note that by the assumptions on r' this implies that $|\Theta(x) - \Theta(y)| \leq \varepsilon$. We wish to show that $d(x, y) < r$. We have three cases:

CASE 1: $\Theta(x), \Theta(y) \leq 1$. Then

$$d(x, y) = \|x - y\| + |1 - 1| < r' < r.$$

CASE 2: $\Theta(x), \Theta(y) > 1$. Then

$$\begin{aligned} d(x, y) &= \|F_{\Theta(x)}(x) - F_{\Theta(y)}(y)\| + |\Theta(x) - \Theta(y)| \\ &= \|F_{\Theta(y)}(F_{\Theta(x)/\Theta(y)}(x) - y)\| + |\Theta(x) - \Theta(y)| \\ &\leq \|F_{\Theta(x)/\Theta(y)}(x) - y\| + |\Theta(x) - \Theta(y)| \\ &\leq \|F_{\Theta(x)/\Theta(y)}(x) - x\| + \|x - y\| + |\Theta(x) - \Theta(y)| \\ &\leq \sup_{i \in I} \left| \left(\frac{\Theta(x)}{\Theta(y)} \right)^{-1/p(i)} - 1 \right| \|x\| + \|x - y\| + |\Theta(x) - \Theta(y)| \\ &\leq \left| \frac{\Theta(y)}{\Theta(x)} - 1 \right| \|x\| + \|x - y\| + |\Theta(x) - \Theta(y)| \\ &= \frac{|\Theta(y) - \Theta(x)|}{\Theta(x)} \|x\| + \|x - y\| + |\Theta(x) - \Theta(y)| \\ &\leq \varepsilon \|x\| + \|x - y\| + \varepsilon < r/3 + r/3 + r/3 = r. \end{aligned}$$

CASE 3: $\Theta(x) \leq 1, \Theta(y) > 1$. Then

$$\begin{aligned} d(x, y) &= \|x - F_{\Theta(y)}(y)\| + |1 - \Theta(y)| \\ &\leq \|x - F_{\Theta(y)}(x)\| + \|F_{\Theta(y)}(x) - F_{\Theta(y)}(y)\| + |\Theta(x) - \Theta(y)| \\ &\leq \sup_{i \in I} |1 - \Theta(y)^{-1/p(i)}| \|x\| + \|F_{\Theta(y)}(x - y)\| + |\Theta(x) - \Theta(y)| \\ &\leq \left| 1 - \frac{1}{\Theta(y)} \right| \|x\| + r' + \varepsilon \leq \varepsilon \|x\| + r' + \varepsilon < r. \end{aligned}$$

Note that the case $\Theta(x) > 1, \Theta(y) \leq 1$ does not occur, since if $\Theta(x) > 1$ then by (4.1), we have $\|y\| > \|x\| - r' > 1$ in which case also $\Theta(y) > 1$. We have proved that $B_{\|\cdot\|}(x, r') \subseteq B_d(x, r)$.

For the other direction consider $B_{\|\cdot\|}(x, r)$ and let

$$r'' < \min\{r/2, 1, r/(\|x\| + \Theta(x) + 1)\}$$

and again, if $\Theta(x) > 1$ we also require

$$(4.2) \quad r'' < \Theta(x) - 1.$$

We wish to show that $B_d(x, r'') \subseteq B_{\|\cdot\|}(x, r)$, so let $y \in B_d(x, r'')$. Now again, by (4.2), the case $\Theta(x) > 1, \Theta(y) \leq 1$ does not occur, so we have three cases:

CASE 1: $\Theta(x), \Theta(y) \leq 1$. Then

$$\|x - y\| = d(x, y) < r'' < r.$$

CASE 2: $\Theta(x), \Theta(y) > 1$. Then

$$\begin{aligned} d(x, y) &= \|F_{\Theta(x)}(x) - F_{\Theta(y)}(y)\| + |\Theta(x) - \Theta(y)| \\ &\geq \|F_{\Theta(x)}(x) - F_{\Theta(y)}(y)\| = \|F_{\Theta(y)}(F_{\Theta(x)/\Theta(y)}(x) - y)\| \\ &\geq \Theta(y)^{-1} \|F_{\Theta(x)/\Theta(y)}(x) - y\| \end{aligned}$$

and thus

$$\begin{aligned} \|x - y\| &\leq \|x - F_{\Theta(x)/\Theta(y)}(x)\| + \|F_{\Theta(x)/\Theta(y)}(x) - y\| \\ &\leq \|x - F_{\Theta(x)/\Theta(y)}(x)\| + \|F_{1/\Theta(y)}(F_{\Theta(x)}(x) - F_{\Theta(y)}(y))\| \\ &\leq \|x - F_{\Theta(x)/\Theta(y)}(x)\| + \Theta(y)d(x, y) \\ &\leq \sup_{i \in I} \left| 1 - \left(\frac{\Theta(x)}{\Theta(y)} \right)^{-1/p(i)} \right| \|x\| + (\Theta(x) + r'')d(x, y) \\ &\leq \left| 1 - \frac{\Theta(y)}{\Theta(x)} \right| \|x\| + (\Theta(x) + r'')r'' \\ &\leq r''\|x\| + (\Theta(x) + r'')r'' \leq r''(\|x\| + \Theta(x) + 1) < r. \end{aligned}$$

CASE 3: $\Theta(x) \leq 1, \Theta(y) > 1$. Then

$$d(x, y) = \|x - F_{\Theta(y)}(y)\| + |1 - \Theta(y)| \geq \|x - F_{\Theta(y)}(y)\|,$$

so

$$\begin{aligned} \|x - y\| &\leq \|x - F_{\Theta(y)}(y)\| + \|F_{\Theta(y)}(y) - y\| \leq d(x, y) + \|F_{\Theta(y)}(y) - y\| \\ &\leq d(x, y) + \sup_{i \in I} |\Theta(y)^{-1/p(i)} - 1| \|y\| \leq r'' + \left| \frac{1}{\Theta(y)} - 1 \right| \Theta(y) \\ &\leq r'' + |1 - \Theta(y)| \leq r'' + r'' < r. \end{aligned}$$

This finishes the proof that $B_d(x, r'') \subseteq B_{\|\cdot\|}(x, r)$. ■

PROPOSITION 4.9. *The metric d is definable in the sense that it is preserved by all automorphisms of $\ell^{p(\cdot)}(I)$. Thus, the introduction of it does not affect the automorphisms.*

Proof. Ben Yaacov [3, Theorem 3.1] has shown that the modular Θ is definable in Nakano spaces with bounded exponent and dimension at least two using the vocabulary for Banach lattices.

Further F_λ is the unique linear operator that preserves supports and satisfies

$$\Theta(F_\lambda(x)) = \frac{1}{\lambda} \Theta(x) \quad \text{for all } x \in \ell^{p(\cdot)}(I).$$

To see that ‘preserving supports’ is definable, note that for $\lambda \geq 1$, $F_\lambda(e_i) \wedge e_i = F_\lambda(e_i)$, thus $\text{supp}(F_\lambda(e_i)) \subseteq \text{supp}(e_i)$ and as $F_\lambda(e_i) \neq 0$ we must have $\text{supp}(F_\lambda(e_i)) = \text{supp}(e_i)$. Then the rest follows from $F_\lambda \circ F_{\lambda^{-1}} = \text{id}$.

Thus F and G are definable making the metric d definable. ■

In order to be able to treat our generalized isomorphisms, we expand the vocabulary to

$$\mathcal{L}_{Bl}^d = \{0, -, +, \cdot, \|\cdot\|, \wedge, \vee, d\}$$

and use d as the distinguished metric on our structures. As it is definable and (topologically) equivalent to the metric induced by the norm, we preserve isometries and closures. By Theorem 4.2, when given an atomic Nakano space $\ell^{p(\cdot)}(I)$ with bounded exponent, we can identify its standard basis $\mathbb{B}(\ell^{p(\cdot)}(I)) = (e_i)_{i \in I}$ as well as the partition thereof induced by p . Thus we can define:

DEFINITION 4.10. Let $(\mathbb{K}_r, \preccurlyeq)$ be the class of all structures Banach lattice isometrically isomorphic to some $\ell^{p(\cdot)}(I)$, with $p(i) \leq r$ (i.e. $p(i) \in [1, r]$) for all $i \in I$ and with $|I| \geq 2..$ (Strictly speaking, our structures are two-sorted with a copy of the reals as second sort.)

We then define $\mathcal{A} \preccurlyeq \mathcal{B}$ if

- $\mathbb{B}(\mathcal{A}) = \mathbb{B}(\mathcal{B}) \cap \mathcal{A}$, in which case we may write $\mathcal{A} = \ell^{p(\cdot)}(I)$ and $\mathcal{B} = \ell^{q(\cdot)}(J)$ with $I \subseteq J$,
- with the notation above $p(\cdot) = q(\cdot)|I$.

Note that as closures and strong submodels are the same whether we regard d or the norm as our metric, this definition gives the same model class as would be obtained without introducing d .

PROPOSITION 4.11. *For a given $r \in [1, \infty)$, $(\mathbb{K}_r, \preccurlyeq)$ is a MAEC with Löwenheim–Skolem number \aleph_0 .*

Proof. The first three items of the definition are trivial. For the fourth, note that if $\langle \mathcal{A}_i : i < \delta \rangle$ is an \preccurlyeq -increasing chain then we may write it as $\langle \ell^{p_i(\cdot)}(I_i) : i < \delta \rangle$ where $(I_i)_{i < \delta}$ is increasing and $p_i = p_j|I_i$ for all $i < j < \delta$.

As simple functions are dense in Nakano spaces with bounded exponent (see e.g. [9, Corollary 2.4.10]), the closure of the union is

$$\ell^{p(\cdot)}\left(\bigcup_{i<\delta} I_i\right)$$

where $p = \bigcup_{i<\delta} p_i$.

Also the fifth item of the definition is easy, and for the Löwenheim–Skolem number note that functions over I with finite support and rational values are dense in $\ell^{p(\cdot)}(I)$. ■

PROPOSITION 4.12. $(\mathbb{K}_r, \preccurlyeq, \mathbb{F}_\varepsilon)_{\varepsilon \geq 0}$ is a MAEC with perturbations.

Proof. We have already noted that $(\mathbb{K}_r, \preccurlyeq)$ is a MAEC, so what remains to be shown is that the ε -isomorphisms of Definition 4.5 satisfy the demands of Definition 2.2.

By Fact 4.4, ε -isomorphisms are bijections and they are uniformly continuous on the unit ball. As they map basic vectors to basic vectors, it is straightforward to see that a mapping in $\bigcap_{\varepsilon>0} \mathbb{F}_\varepsilon$ must preserve both basic vectors and their corresponding exponents, i.e. be a genuine isomorphism.

For uniform continuity we can use as modulus of uniform continuity $\Delta^\varepsilon(\cdot/2)$ where Δ^ε is the modulus arising from Ben Yaacov's result. If f is an ε -isomorphism, for any a , $\Theta(f(a)) = \Theta(a)$ and $F_\lambda(f(a)) = f(F_\lambda(a))$, so

$$F(f(a)) = \begin{cases} f(a) = f(F(a)) & \text{if } \Theta(f(a)) = \Theta(a) \leq 1, \\ F_{\Theta(f(a))}(f(a)) = f(F_{\Theta(a)}(a)) = f(F(a)) & \text{otherwise,} \end{cases}$$

i.e. F and f commute. Thus if $d(x, y) < \Delta^\varepsilon(\delta)$, we have both

$$\|F(x) - F(y)\| \leq d(x, y) < \Delta^\varepsilon(\delta)$$

and

$$|G(x) - G(y)| \leq d(x, y) < \Delta^\varepsilon(\delta) \leq \delta.$$

and also $\|F(x)\|, \|F(y)\| \leq 1$, so

$$\|f(F(x)) - f(F(y))\| \leq \delta$$

and thus

$$\begin{aligned} d(f(x), f(y)) &= \|F(f(x)) - F(f(y))\| + |G(f(x)) - G(f(y))| \\ &= \|f(F(x)) - f(F(y))\| + |G(x) - G(y)| \leq 2\delta. \end{aligned}$$

The rest of the demands on the ε -isomorphisms are straightforward to check. ■

PROPOSITION 4.13. $(\mathbb{K}_r, \preccurlyeq)$ has the joint embedding property and is 0-homogeneous.

Proof. If $\mathcal{A} \cong \ell^{p_1(\cdot)}(I)$ and $\mathcal{B} \cong \ell^{p_2(\cdot)}(J)$ (with I and J disjoint) then clearly they can be embedded into $\ell^{p_3(\cdot)}(I \cup J)$ where $p_3|I = p_1$ and $p_3|J = p_2$.

For homogeneity, assume that $(a_i)_{i<\alpha} \in \ell^{p(\cdot)}(I)$ and $(b_i)_{i<\alpha} \in \ell^{q(\cdot)}(J)$ and

$$(4.3) \quad t^g((a_{i_k})_{k< n}/\emptyset) = t^g((b_{i_k})_{k< n}/\emptyset) \quad \text{for each } n < \omega.$$

Since Nakano spaces with (essentially) bounded exponent are axiomatizable in continuous logic and we are working in a subclass of this class with essentially the same vocabulary, the sameness of Galois types is determined by the sameness of types in continuous logic. But (4.3) guarantees that $(a_i)_{i<\alpha}$ and $(b_i)_{i<\alpha}$ have the same syntactic type in continuous logic and hence they have the same Galois type. ■

PROPOSITION 4.14. $(\mathbb{K}_r \preccurlyeq, \mathbb{F}_\varepsilon)_{\varepsilon \geq 0}$ satisfies the amalgamation property (with respect to ε -isomorphisms).

Proof. Let $\ell^{p(\cdot)}(I') \preccurlyeq \ell^{p(\cdot)}(I)$ and $F : \ell^{p(\cdot)}(I') \rightarrow \ell^{q(\cdot)}(J)$ be an ε -embedding. Then I can be written as a disjoint union $I = I' \cup I''$. Since F maps basic vectors to basic vectors, we may write J as a disjoint union $J = J' \cup J''$ and note that for $i \in I'$, $e^{-\varepsilon} \leq q(i)/p(i) \leq e^\varepsilon$. Now consider $\mathcal{C} = \ell^{q(\cdot)}(J) + \ell^{p(\cdot)}(I'')$ (i.e. $\ell^{q^*(\cdot)}(I' \cup J' \cup I'')$ where $q^*|I' \cup J' = q$ and $q^*|I'' = p$, assuming the union is disjoint). Then $F + \text{id}_{\ell^{p(\cdot)}(I'')}$ is an ε -mapping $\ell^{p(\cdot)}(I) \rightarrow \mathcal{C}$. ■

PROPOSITION 4.15. $(\mathbb{K}_r \preccurlyeq, \mathbb{F}_\varepsilon)_{\varepsilon \geq 0}$ has the perturbation property.

Proof. Assume $a, b \in \mathfrak{M}$ satisfy $\mathbf{d}^p(t^g(a/\emptyset), t^g(b/\emptyset)) = 0$. We wish to show that $t^g(a/\emptyset) = t^g(b/\emptyset)$.

By assumption there is, for each $n < \omega$, a $1/n$ -automorphism of \mathfrak{M} satisfying $d(f_n(a), b) \leq 1/n$. Now let \mathcal{A} be a separable model containing a and let \mathcal{B} be a separable model containing b and $f_n(\mathcal{A})$ for each $n < \omega$. Let D be an ultrafilter on ω extending the Fréchet filter and define $\mathcal{C} = \prod_D \mathcal{B}$ (meaning the Banach space ultraproduct). As \mathcal{C} is not atomic, it is not in our class, but it is still a Nakano space (by Poitevin and Ben Yaacov). Now for each $c \in \mathcal{A}$, let

$$F(c) = \prod f_n(c)/D.$$

Then F is a Banach lattice isometry (considered in the class of all Nakano spaces with exponents bounded by r), but since these do not mix up atomic and atomless parts, we see that F is a 0-embedding of \mathcal{A} into the atomic part of \mathcal{C} . On the other hand the natural (diagonal) embedding $G : \mathcal{B} \rightarrow \mathcal{C}$ is a 0-embedding into the atomic part of \mathcal{C} and $F(a) = G(b)$ since

$$\{n < \omega : d(f_n(a), b) < 1/m\} \in D$$

for each $m < \omega$.

So the atomic part of \mathcal{C} together with the embeddings of \mathcal{A} and \mathcal{B} into it prove that $t^g(a/\emptyset; \mathcal{A}) = t^g(b/\emptyset; \mathcal{B})$, which implies $t^g(a/\emptyset) = t^g(b/\emptyset)$. ■

REMARK 4.16. The example of this section treats classes consisting of all atomic Nakano spaces with a given bound r for the exponent. Thus the range of the exponent of the monster model will always be the full interval $[1, r]$ and also each $p \in [1, r]$ will occur infinitely often. All work done so far could as such be generalized to the case where the exponent lies in a given bounded set. Then the next proposition would need the additional assumption that this range is closed.

PROPOSITION 4.17. *For a given $r \in [1, \infty)$, $(\mathbb{K}_r, \preccurlyeq, \mathbb{F}_\varepsilon)_{\varepsilon \geq 0}$ has complete type-spaces.*

Proof. Assume $\bar{a}_n \in \mathfrak{M}$ is such that $t^g(\bar{a}_n/\emptyset)$ forms a Cauchy sequence with respect to \mathbf{d}^p .

To make notation a bit easier we prove the claim for the case where $\lg(\bar{a}_n) = 1$. The generalization to the general case is straightforward.

The idea is to cut the elements up into their coordinates and find the limit element using real sequences formed by the coordinatewise coefficients. In making the real sequences we have to take into account that ε -isomorphisms permute the basis.

So we look at a sequence $(a_n)_{n < \omega}$ of elements of $\mathfrak{M} = \ell^{p(\cdot)}(I)$. First choose monotone sequences of positive reals $s_n < 1$ and $t_n > 1$ such that $\prod_{n < \omega} s_n > 1/2$ and $\prod_{n < \omega} t_n < 2$, i.e. (s_n) is increasing, (t_n) decreasing and both sequences tend to 1. Next choose a decreasing sequence of positive reals ε_n such that

$$\begin{aligned} \min\{r^{e^{-\varepsilon_n}}, r^{e^{\varepsilon_n}}\} - \varepsilon_n &> s_n r \quad \text{for all } r \geq 2^{-n}, \\ \max\{r^{e^{-\varepsilon_n}}, r^{e^{\varepsilon_n}}\} + \varepsilon_n &< t_n r \quad \text{for all } r \leq 2^n, \\ \sum_{i < \omega} \varepsilon_i &\leq 1. \end{aligned}$$

Now since $(t^g(a_n/\emptyset))_{n < \omega}$ is \mathbf{d}^p -Cauchy, for each $\varepsilon > 0$ there is some n_ε such that $\mathbf{d}^p(t^g(a_n/\emptyset), t^g(a_m/\emptyset)) < \varepsilon$ whenever $n, m \geq n_\varepsilon$. Switching to a subsequence and renumbering we may assume $a_k = a_{n_{\varepsilon_k}}$ for all $k < \omega$. We may also pick, for each $n < \omega$, an ε_n -automorphism f_n satisfying $d(f_n(a_n), a_{n+1}) \leq \varepsilon_n$.

Now each a_n is of the form $a_n = (a_n(i))_{i \in I}$. For each $i \in I$ we define a sequence $(a_n^i)_{n < \omega}$ as follows:

$$a_0^i = a_0(i), \quad a_1^i = a_1(j) \quad \text{where } f_0(e_i) = e_j,$$

and in general when $a_n^i = a_n(i')$ has been defined and $f_n(e_{i'}) = e_j$,

$$a_{n+1}^i = a_{n+1}(j).$$

Note that only countably many sequences have any nonzero elements. Also, as any a_n can be ε_0 -closely reached by an ε_0 -automorphism from a_0 , the

sequence $(\|a_n\|)_{n<\omega}$ is bounded and thus there is a universal bound M for the absolute value of all elements a_n^i , $i \in I$, $n < \omega$.

Next we show that these sequences converge. So let $i \in I$. If $(a_n^i)_{n<\omega}$ converges to zero, we are done, so we may assume it does not. Then there is some $\varepsilon > 0$ such that $|a_n^i| > \varepsilon$ cofinally often. Let n be large enough that $2^{-n} < \varepsilon$ and $M < 2^n$. Then let $m \geq n$ be such that $|a_m^i| > \varepsilon$.

Now

$$s_m|a_m^i| < |a_{m+1}^i| < t_m|a_m^i|$$

and we see that the elements of $(a_n^i)_{n<\omega}$ cannot switch signs after m . By induction we get, for any $m' \geq m$ and any $k < \omega$,

$$\prod_{j \geq m'} s_j |a_{m'}^i| < |a_{m'+k}^i| < \prod_{j \geq m'} t_j |a_{m'}^i|.$$

Since both $\prod_{j \geq m'} s_j$ and $\prod_{j \geq m'} t_j$ tend to 1 as $n \rightarrow \infty$, it is easy to deduce that $(a_n^i)_{n<\omega}$ is a Cauchy sequence and thus converges. Denote the limit by a^i .

Now we know what the coefficients of our limit element should be. Next we see in which coordinates to put the coefficients. So for each $i \in I$ consider the sequence of exponents p_n^i defined as for the a^i 's:

$$p_0^i = p(i)$$

and in general when $p_n^i = p(i')$ has been defined and $f_n(e_{i'}) = e_j$,

$$p_{n+1}^i = p(j).$$

Now for any $i \in I$ and $n < \omega$,

$$e^{-\varepsilon_n} \leq p_{n+1}^i / p_n^i \leq e^{\varepsilon_n},$$

and by induction for any $k < \omega$,

$$e^{-\sum_{j \geq n} \varepsilon_j} \leq p_{n+k}^i / p_n^i \leq e^{\sum_{j \geq n} \varepsilon_j};$$

as $\sum_{j \geq n} \varepsilon_j$ tends to 0, we see that $(p_n^i)_{n<\omega}$ converges, say to p^i .

Next let $I' = \{i \in I : a^i \neq 0\}$ and define $a \in \mathfrak{M}$ to be any element $\sum_{i \in I'} a^i e_{\sigma(i)}$ where σ is an injective mapping $I' \rightarrow I$ such that $p(\sigma(i)) = p^i$.

Finally, to show that the original sequence converges (in the sense of \mathbf{d}^p) to $\mathbf{t}^g(a/\emptyset)$, we need the following claim:

CLAIM. *For $m < \omega$ and $q > 0$ let $J_{<q}^m = \{i \in J : |a_m(i)| < q\}$. Then for every $\delta > 0$ there are $n < \omega$ and $q > 0$ such that for all $m \geq n$, $\|\sum_{i \in J_{<q}^m} a_m(i) e_i\| < \delta$.*

Proof. Assume this is not the case and $\delta > 0$ witnesses the fact. Let

$$\delta' < \Delta^\delta(\delta/2)/3$$

where Δ^δ denotes the modulus of uniform continuity for \mathbb{F}_δ . Let n be so large that $\mathbf{d}^p(t^g(a_n/\emptyset), t^g(a_m/\emptyset)) \leq \delta'$ for all $m \geq n$. As simple functions are dense in Nakano spaces with bounded exponents, we can find arbitrarily close finite support approximations of a_n and each such approximation can be chosen to consist of the coordinates where $|a_n(i)| \geq q$ for some $q > 0$. So we find a $q > 0$ and finite $J' \subset I$ such that

$$\left\| \sum_{i \in I} a_n(i) e_i - \sum_{i \in J'} a_n(i) e_i \right\| = \left\| \sum_{i \in J_{<q}^n} a_n(i) e_i \right\| < \delta'.$$

Let $k = |J'|$.

Next let

$$q' < \delta/(2k).$$

If our claim does not hold, there is some $m \geq n$ such that $\|\sum_{i \in J_{<q'}^m} a_m(i) e_i\| \geq \delta$. However, by our choice of n there is a δ' -automorphism f satisfying $d(f(a_m), a_n) \leq \delta'$. Since f maps basic vectors to basic vectors and is injective, it can only map k coordinates from $J_{<q'}^m$ into J' . The norm of this part is at most kq' as

$$\sum_{j=1}^k \left| \frac{a_m(i_j)}{kq'} \right|^p \leq k \cdot \left| \frac{q'}{kq'} \right|^1 = 1.$$

Thus the norm of the part over $J_{<q'}^m$ that must be mapped elsewhere (i.e. into $J_{<q}^n$) is by the reverse triangle inequality at least

$$\left\| \sum_{i \in J_{<q'}^m} a_m(i) e_i \right\| - kq' \geq \delta - kq' > \delta/2,$$

and when mapped with a δ' -isomorphism (as noted later in Fact 5.3) it gets mapped to an element of norm at least

$$\Delta^{\delta'}(\delta/2) \geq \Delta^\delta(\delta/2) > 3\delta'.$$

By our choice of m we now have

$$\begin{aligned} \delta' &\geq \|f(a_m) - a_n\| = \left\| \sum_{i \in I} (f(a_m)(i) - a_n(i)) e_i \right\| \\ &\geq \left\| \sum_{i \in J_{<q}^n} (f(a_m)(i) - a_n(i)) e_i \right\| \\ &\geq \left\| \left\| \sum_{i \in J_{<q}^n} f(a_m)(i) e_i \right\| - \left\| \sum_{i \in J_{<q}^n} a_n(i) e_i \right\| \right\| \geq 3\delta' - \delta' = 2\delta', \end{aligned}$$

a contradiction. Thus the claim must be true. ■

To show that $t^g(a_n/\emptyset) \rightarrow t^g(a/\emptyset)$ in the \mathbf{d}^p -metric, let $\varepsilon > 0$ be given and choose

$$\delta < \Delta^\varepsilon(\varepsilon/3) (\leq \varepsilon/3).$$

Then by the above claim find $n_1 < \omega$ and $q_1 > 0$ such that for $n \geq n_1$,

$$\left\| \sum_{J_i^n < q_1} a_n(i) e_i \right\| < \delta.$$

Then by density of simple functions find a finite subset J of the support of a such that

$$\left\| \sum_{j \in J} a(j) e_j - a \right\| < \delta.$$

As before we assume J consists of all coordinates where $|a(j)| \geq q_2$ for some $q_2 > 0$. Then let $q = \min\{q_1, q_2\}$.

Now only finitely many coordinates of a can have absolute value above $q/3$, so as $a_n^i \rightarrow a^i$ for all $i \in I$, there is some $n_2 \geq n_1$ such that from n_2 onwards all but finitely many of the sequences $(a_n^i)_{n < \omega}$ stay within the interval $(-q/2, q/2)$. As finitely many converging sequences have only finitely many accumulation points, this leaves gaps in the ranges. Thus there are q', q'' such that $q/2 < q' < q'' \leq q$ and $(\{|a_n^i| : n \geq n_2\} \cup \{|a^i|\}) \cap (q', q'') = \emptyset$. Now for some $n_3 \geq n_2$ none of the sequences occurring (after n_2) above $q/2$ can cross this gap, so from n_3 onwards we have a division of the sequences into a finite number, say k , occurring above q'' and the rest below. Denote the sets of the indices corresponding to these coefficients by $J_{\geq q''}^n$ and $J_{\leq q''}^n$.

As the corresponding p^i -sequences also converge, we can finally find some $n_4 \geq n_3$ such that for $n \geq n_4$ there is a δ -automorphism g_n mapping the coordinates corresponding to a_n^i to the ones corresponding to a^i for all sequences above q'' and making at most a δ/k -error in these coordinates. Regardless of what g_n does outside these coordinates, we have for $n \geq n_4$,

$$\begin{aligned} d(g_n(a_n), a) &\leq \left\| g_n(a_n) - \sum_{i \in J_{\geq q''}^n} g_n(a_n)(i) e_i \right\| \\ &\quad + \left\| \sum_{i \in J_{\geq q''}^n} g_n(a_n)(i) e_i - \sum_{i \in J_{\geq q''}^n} a(i) e_i \right\| + \left\| \sum_{i \in J_{\geq q''}^n} a(i) e_i - a \right\| \\ &\leq \left\| g_n(a_n) - g_n \left(\sum_{i \in J_{\geq q''}^n} a_n(i) e_i \right) \right\| \\ &\quad + \left\| \sum_{i \in J_{\geq q''}^n} \frac{\delta}{k} e_i \right\| + \left\| \sum_{i \in J_{\leq q''}^n} a(i) e_i \right\| \\ &\leq \varepsilon/3 + \delta + \delta < \varepsilon, \end{aligned}$$

where we have used that g_n is a δ -automorphism, hence an ε -isomorphism, and

$$\left\| a_n - \sum_{i \in J_{\geq q''}^n} a_n(i) e_i \right\| = \left\| \sum_{i \in J^n < q''} a_n(i) e_i \right\| \leq \left\| \sum_{i \in J^n < q_1} a_n(i) e_i \right\| < \delta. \blacksquare$$

As noted by Ben Yaacov in [3], we have

FACT 4.18. *The class is ω - \mathbf{d}^p -stable.*

5. Types and \mathbf{d}^p . Recall the definition of \mathbf{d}^p (the version below is what we get by the perturbation property):

DEFINITION 5.1. For $a, b \in \mathfrak{M}$ and $\varepsilon \geq 0$, $\mathbf{d}^p(t^g(a/\emptyset), t^g(b/\emptyset)) \leq \varepsilon$ if there are ε -automorphisms f and g of \mathfrak{M} such that $d(f(a), b) \leq \varepsilon$ and $d(g(b), a) \leq \varepsilon$.

For $a, b \in \mathfrak{M}$ and $A \subset \mathfrak{M}$,

$$\mathbf{d}^p(t^g(a/A), t^g(b/A)) = \sup\{\mathbf{d}^p(t^g(ac/\emptyset), t^g(bc/\emptyset)) : c \in A \text{ finite}\}.$$

It is easy to see that the definition only depends on the Galois types of a and b . Also, the perturbation property (together with 0-homogeneity) ensures that if $\mathbf{d}^p(t^g(a/A), t^g(b/A)) = 0$ then a and b do have the same Galois type over A (proved below). However, it is worth noting that when $\mathbf{d}^p(t^g(a/A), t^g(b/A))$ is strictly positive, we do not have a function fixing A pointwise that moves a near b .

LEMMA 5.2. *If $a, b \in \mathfrak{M}$, $A \subset \mathfrak{M}$ and $\mathbf{d}^p(t^g(a/A), t^g(b/A)) = 0$ then $t^g(a/A) = t^g(b/A)$.*

Proof. By 0-homogeneity it is enough to show that for any finite $c \in A$, $t^g(ac/\emptyset) = t^g(bc/\emptyset)$, which is equal to the demand $t^g(a/c) = t^g(b/c)$. But since our assumption asserts that $\mathbf{d}^p(t^g(a/A), t^g(b/A)) = 0$, i.e.

$$\sup\{\mathbf{d}^p(t^g(ac/\emptyset), t^g(bc/\emptyset)) : c \in A \text{ finite}\} = 0,$$

this is exactly what the perturbation property gives us. ■

Thus for any A , \mathbf{d}^p is a semimetric on $S(A)$. To investigate the behaviour of \mathbf{d}^p let us first have a look at the moduli of uniform continuity Δ^ε from Definition 2.2(2b). From the demands on the classes \mathbb{F}_ε we can easily deduce:

FACT 5.3.

- (1) *If $\delta < \varepsilon$ then $\Delta^\delta(x) \geq \Delta^\varepsilon(x)$ for all $x \in (0, \infty)$.*
- (2) *$\Delta^\varepsilon(x) \leq x$ for all $x \in (0, \infty)$.*
- (3) *If $f \in \mathbb{F}_\varepsilon$ and $d(x, y) > d$ then $d(f(x), f(y)) \geq \Delta^\varepsilon(d)$.*

Using the above we can easily see that if $\mathbf{d}^p(t^g(a/A), t^g(b/A)) < \Delta^\varepsilon(\varepsilon/2)$ and $\mathbf{d}^p(t^g(b/A), t^g(c/A)) < \Delta^\varepsilon(\varepsilon/2)$, then $\mathbf{d}^p(t^g(a/A), t^g(c/A)) < \varepsilon$.

Now for any set A the sets

$$D_\varepsilon^p = \{(p, q) \in S(A) : \mathbf{d}^p(p, q) \leq \varepsilon\}$$

form a base for a metrizable (diagonal) uniformity (see Chapter 9 of [16]). Thus, although \mathbf{d}^p is not itself a metric, it makes sense to talk about Cauchy sequences, limits and completeness with respect to \mathbf{d}^p .

To find limit types we use either goodness or completeness of type-spaces.

LEMMA 5.4. *Assume A is good in \mathfrak{M} . Then if $\varepsilon > 0$ and $\delta > 0$ are as in the definition of goodness (Definition 2.21), and if $F_n \in \text{Aut}_\delta(\mathfrak{M})$ for $n < \omega$ and for each $a \in A$ the sequence $(F_n(a))$ converges, then there is $F \in \text{Aut}_\varepsilon(\mathfrak{M})$ such that for each $a \in A$, $F(a) = \lim_{n \rightarrow \infty} F_n(a)$.*

Proof. By assumption, weak δ -embeddings $f : A \rightarrow \mathfrak{M}$ extend to ε -automorphisms of \mathfrak{M} . Hence it is enough to show that the function defined by $F(a) = \lim_{n \rightarrow \infty} F_n(a)$ for all $a \in A$ is a weak δ -embedding. But this is exactly what the functions F_n witness. ■

When using completeness of type-spaces (taking into account Remark 2.18) we can actually find limit types over directed systems. Note that as soon as we have types actually extending each other and not just being \mathbf{d}^p -close, 0-homogeneity gives these sorts of limit types:

FACT 5.5. *Assume (\mathcal{B}, \subseteq) is a directed system and $\{b_B : B \in \mathcal{B}\}$ are such that $t^g(b_{B'} / B) = t^g(b_B / B)$ when $B \subset B'$, $B, B' \in \mathcal{B}$. Then there is some b such that $t^g(b / B) = t^g(b_B / B)$ for all $B \in \mathcal{B}$.*

LEMMA 5.6. *Assume \mathbb{K} has complete type-spaces, (\mathcal{B}, \subseteq) is a directed system of finite sets and $\{t^g(b_B^n / B) : n < \omega, B \in \mathcal{B}\}$ is \mathbf{d}^p -coherent in the following sense: for every $B \in \mathcal{B}$ and $\varepsilon > 0$ there are $n_0 < \omega$ and $B_0 \in \mathcal{B}$ with $B_0 \supseteq B$ such that for all $n \geq m \geq n_0$ and $B_1, B_2 \in \mathcal{B}$ with $B_2 \supseteq B_1 \supseteq B_0$, $\mathbf{d}^p(t^g(b_{B_2}^n / B_1), t^g(b_{B_1}^m / B_1)) \leq \varepsilon$. Then there exists a limit type, i.e. there is some b such that for every $B \in \mathcal{B}$ and $\varepsilon > 0$ there are $n_0 < \omega$ and $B_0 \in \mathcal{B}$ with $B_0 \supseteq B$ such that for all $n \geq n_0$ and $B_1 \in \mathcal{B}$ with $B_1 \supseteq B_0$, $\mathbf{d}^p(t^g(b / B_1), t^g(b_{B_1}^n / B_1)) \leq \varepsilon$.*

Proof. For each $B \in \mathcal{B}$ we find a \mathbf{d}^p -limit b_B as follows: For every $m < \omega$ there are some $n_m < \omega$ and $B^m \supseteq B$ such that for all $n \geq n_m$ and $B' \supseteq B^m$,

$$(5.1) \quad \mathbf{d}^p(t^g(b_{B'}^n / B^m), t^g(b_{B^m}^{n_m} / B^m)) \leq 1/m,$$

and we may choose $n_{m+1} \geq n_m$ and $B^{m+1} \supseteq B^m$ for all m . Then $(t^g(b_{B^m}^{n_m} / B))_{m < \omega}$ is \mathbf{d}^p -Cauchy and by completeness of type-spaces has a \mathbf{d}^p -limit b_B .

Now if $B \subset C \in \mathcal{B}$ then $t^g(b_C / B) = t^g(b_B / B)$: For $\varepsilon > 0$ choose m large enough (and $> 1/\varepsilon$) such that

$$\mathbf{d}^p(t^g(b_B / B), t^g(b_{B^m}^{n_m} / B)) \leq \varepsilon$$

and similarly for b_C . Let $D \supset B^m \cup C^m$. Then by (5.1) for all large enough n ,

$$\mathbf{d}^p(t^g(b_D^n / B^m), t^g(b_{B^m}^{n_m} / B^m)) \leq 1/m < \varepsilon$$

and similarly for b_C . Since this can be done for any $\varepsilon > 0$, equality of the types $t^g(b_C / B)$ and $t^g(b_B / B)$ follows by perturbation. But then the claim follows by 0-homogeneity (Fact 5.5). ■

6. Saturation

DEFINITION 6.1. We say that A is $\omega\text{-}\mathbf{d}^p$ -saturated if for all finite $A' \subset A$, all $a \in \mathfrak{M}$ and all $\varepsilon > 0$ there is $a' \in A$ such that

$$\mathbf{d}^p(t^g(a/A'), t^g(a'/A')) \leq \varepsilon$$

(this is the same as demanding $\mathbf{d}^p(t^g(aA'/\emptyset), t^g(a'A'/\emptyset)) \leq \varepsilon$).

LEMMA 6.2. Assume \mathbb{K} is $\omega\text{-}\mathbf{d}^p$ -stable. If \mathcal{A} is $\omega\text{-}\mathbf{d}^p$ -saturated and $B \subset \mathcal{A}$ then there is an $\omega\text{-}\mathbf{d}^p$ -saturated $\mathcal{B} \preccurlyeq \mathcal{A}$ with $B \subseteq \mathcal{B}$ and $|\mathcal{B}| = |B| + \aleph_0$.

Proof. We build \mathcal{B} as a chain of models \bar{B}_n , where for each $n < \omega$:

- $\text{card}(B_n) \leq |B| + \aleph_0$,
- B_{n+1} realizes a \mathbf{d}^p -dense set of $S(A)$ for each finite $A \subset B_n$,
- $\bar{B}_n \preccurlyeq \mathcal{A}$.

We start by letting B'_0 be a dense subset of B of cardinality $|B|$ and then get B_0 by the assumption $\text{LS}^{\mathbf{d}} = \aleph_0$. When B_n has been defined we use $\omega\text{-}\mathbf{d}^p$ -stability and the $\text{LS}^{\mathbf{d}}$ assumption to get B_{n+1} . Finally we let $\mathcal{B} = \overline{\bigcup_{n < \omega} B_n}$. Then clearly $\mathcal{B} \preccurlyeq \mathcal{A}$ and $|\mathcal{B}| = |B| + \aleph_0$. It is also easy to see that \mathcal{B} is $\omega\text{-}\mathbf{d}^p$ -saturated, since parameters are allowed to move and we only need to realize types δ -closely. ■

REMARK 6.3. It is easy to see how to adapt the proof to show that any $\omega\text{-}\mathbf{d}^p$ -saturated set A with a given subset $A' \subset A$ contains an $\omega\text{-}\mathbf{d}^p$ -saturated set B with $A' \subseteq B$ and $|B| = |A'| + \aleph_0$. Moreover, if B is any $\omega\text{-}\mathbf{d}^p$ -saturated set, then \bar{B} is also $\omega\text{-}\mathbf{d}^p$ -saturated.

DEFINITION 6.4. We say that two models \mathcal{A} and \mathcal{B} are *almost isomorphic* if for each $\varepsilon > 0$ there is an ε -isomorphism $f : \mathcal{A} \rightarrow \mathcal{B}$.

We will prove that $\omega\text{-}\mathbf{d}^p$ -saturated separable models are unique up to almost isomorphism. In the continuous logic setting with perturbations this was proved by Ben Yaacov in [1, Proposition 2.7]. For our proof we need either goodness or the following continuity assumption for ε -isomorphisms:

DEFINITION 6.5. We say that \mathbb{K} is *weakly \mathbb{F} -homogeneous* if for every $\delta \geq 0$ if $f : \mathcal{A} \rightarrow \mathcal{B}$ is a weak δ -embedding (see Definition 2.19) which is onto then $f \in \mathbb{F}_\delta$.

REMARK 6.6. Note that \mathbb{K}_H from Section 3 is weakly \mathbb{F} -homogeneous.

THEOREM 6.7. Assume \mathcal{A} and \mathcal{B} are separable and $\omega\text{-}\mathbf{d}^p$ -saturated. Then if either \mathcal{A} and \mathcal{B} are good or \mathbb{K} is weakly \mathbb{F} -homogeneous, \mathcal{A} and \mathcal{B} are almost isomorphic.

Proof. First assume \mathcal{A} and \mathcal{B} are good. Fix $\varepsilon > 0$. We will construct an ε -isomorphism between \mathcal{A} and \mathcal{B} . So let $\delta > 0$ be such that weak δ -embeddings of both \mathcal{A} and \mathcal{B} extend to ε -embeddings, and choose for each

$n < \omega$, $\delta_n > 0$ such that $\delta_{n+1} \leq \delta_n$ for all n and $\sum_{n < \omega} \delta_n < \delta$. Denote $\varepsilon_n = \sum_{i < n} \delta_i$. Let $A \subset \mathcal{A}$ and $B \subset \mathcal{B}$ be countable and dense and enumerate them $A = \{a_n : n < \omega\}$, $B = \{b_n : n < \omega\}$. We will define finite sets $A_n \subset A$, $B_n \subset B$ and mappings f_n, g_n for $n < \omega$ such that:

- f_n is in $\text{Aut}_{\varepsilon_{2n}}(\mathfrak{M})$ and maps A_n $\Delta^{2\delta}(\delta_{2n-1})$ -close to B , i.e. there is $B'_n \subset B$ with $|B'_n| = |A_n|$ such that $d(f_n(A_n), B'_n) \leq \Delta^{2\delta}(\delta_{2n-1})$,
- g_n is in $\text{Aut}_{\varepsilon_{2n+1}}(\mathfrak{M})$ and maps B_n $\Delta^{2\delta}(\delta_{2n})$ -close to some $A'_n \subset A$,
- $A_{n+1} = A_n \cup \{a_i : i \leq n\} \cup A'_n$,
- $B_{n+1} = B_n \cup \{b_i : i \leq n\} \cup B'_{n+1}$.

We begin the construction by defining $A_0 = B_0 = \emptyset$ and $f_0 = \text{id}$. When A_n, B_n and f_n have been defined, consider the type $t^g(f_n^{-1}(B_n)/A_n)$. Since \mathcal{A} is ω -**d**^p-saturated, there is some $A'_n \subset A$ such that

$$\mathbf{d}^p(t^g(f_n^{-1}(B_n)/A_n), t^g(A'_n/A_n)) \leq \Delta^{2\delta}(\delta_{2n}).$$

Hence there is $g' \in \text{Aut}_{\delta_{2n}}(\mathfrak{M})$ such that $d(g'(f_n^{-1}(B_n)A_n), A'_n A_n) \leq \Delta^{2\delta}(\delta_{2n})$. Define $g_n = g' \circ f_n^{-1}$. Then $g_n \in \text{Aut}_{\varepsilon_{2n+1}}(\mathfrak{M})$ and g_n maps B_n $\Delta^{2\delta}(\delta_{2n})$ -close to $A'_n \subset A$.

Similarly define f_{n+1} and B'_{n+1} by considering $t^g(g_n^{-1}(A_{n+1})/B_n)$.

Now $g_n \circ f_n = g'$ (as defined above) so $d(g_n \circ f_n(A_n), A_n) \leq \Delta^{2\delta}(\delta_{2n})$ and similarly we see that $d(f_{n+1} \circ g_n(B_n), B_n) \leq \Delta^{2\delta}(\delta_{2n+1})$. Hence

$$\begin{aligned} d(f_{n+1}(A_n), f_n(A_n)) &\leq d(f_{n+1}(A_n), f_{n+1} \circ g_n \circ f_n(A_n)) \\ &\quad + d(f_{n+1} \circ g_n \circ f_n(A_n), f_{n+1} \circ g_n(B'_n)) \\ &\quad + d(f_{n+1} \circ g_n(B'_n), B'_n) + d(B'_n, f_n(A_n)) \\ &\leq 4\delta_{2n-1}. \end{aligned}$$

Hence we see that (f_n) converges (pointwise) on A , and since the mappings are uniformly continuous with the same modulus of uniform continuity and A is dense in \mathcal{A} , the sequence converges on \mathcal{A} . Since each f_n is a δ -automorphism, by Lemma 5.4 there is $F \in \text{Aut}_\varepsilon(\mathfrak{M})$ such that $F(a) = \lim_{n \rightarrow \infty} f_n(a)$ for each $a \in \mathcal{A}$. Similarly we obtain $G \in \text{Aut}_\varepsilon(\mathfrak{M})$ satisfying $G(b) = \lim_{n \rightarrow \infty} g_n(b)$ for each $b \in \mathcal{B}$.

To see that F maps \mathcal{A} into \mathcal{B} , note that for each $a \in \mathcal{A}$, $F(a)$ is the limit of a sequence of elements getting closer and closer to \mathcal{B} . Since \mathcal{B} is metrically closed, we must have $F(a) \in \mathcal{B}$. Similarly we see that $G(b) \in \mathcal{A}$ for each $b \in \mathcal{B}$. To see that F and G are onto, it is enough to show that for all $a \in \mathcal{A}$ and $b \in \mathcal{B}$, $d(G \circ F(a), a) \leq \delta$ and $d(F \circ G(b), b) \leq \delta$ for any positive δ . We prove the former claim since the latter is quite similar. So let $a \in \mathcal{A}$ and $\delta > 0$ be given and choose $\delta' < \Delta^{2\varepsilon}(\delta/7)$. Then we can find $a' \in \bigcup_{n < \omega} A_n$, $b' \in \bigcup_{n < \omega} B_n$ and $n < \omega$ such that:

- $d(a, a') \leq \delta'$,
- $d(F(a'), f_n(a')) \leq \delta'$,

- $d(f_n(a'), b') \leq \delta'$,
- $d(G(b'), g_n(b')) \leq \delta'$,
- $d(g_n \circ f_n(a'), a') \leq \delta'$.

Then

$$\begin{aligned} d(G \circ F(a), a) &\leq d(G \circ F(a), G \circ F(a')) + d(G \circ F(a'), G \circ f_n(a')) \\ &\quad + d(G \circ f_n(a'), G(b')) + d(G(b'), g_n(b')) \\ &\quad + d(g_n(b'), g_n \circ f_n(a')) + d(g_n \circ f_n(a'), a') + d(a', a) \\ &\leq \delta. \end{aligned}$$

Now if instead of goodness we assume weak \mathbb{F} -homogeneity then just choose the δ_n such that $\sum_{n<\omega} \delta_n < \varepsilon$ and do the construction as above. Then F and G are defined as the limit mappings on \mathcal{A} and \mathcal{B} respectively, giving the required approximability by ε -functions. The onto part is done in a similar fashion and then weak \mathbb{F} -homogeneity will ensure that F and G are in \mathbb{F}_ε . ■

REMARK 6.8. Note that if A is almost isomorphic to an ω - \mathbf{d}^p -saturated set then A is ω - \mathbf{d}^p -saturated.

7. Splitting and independence

DEFINITION 7.1. Assume $A \subset B \subset \mathfrak{M}$, A is finite and $a \in \mathfrak{M}$. We say that $t^g(a/B)$ ε -splits over A if for all $\delta > 0$ there are $b, c \in B$ such that $\mathbf{d}^p(t^g(b/A), t^g(c/A)) \leq \delta$ but $\mathbf{d}^p(t^g(ab/A), t^g(ac/A)) > \varepsilon$.

Splitting works nicely for ω - \mathbf{d}^p -stable classes \mathbb{K} , if we assume either goodness or completeness of type-spaces.

THEOREM 7.2. *If \mathbb{K} is ω - \mathbf{d}^p -stable then for all $a \in \mathfrak{M}$, all separable good $A \subset \mathfrak{M}$ and $\varepsilon > 0$ there is some finite $A' \subset A$ such that $t^g(a/A)$ does not ε -split over A' .*

Proof. Let A , a and $\varepsilon > 0$ be given. If there is no finite $A' \subset A$ such that $t^g(a/A)$ does not ε -split over A' then for all finite $A_n \subset A$ and all $\delta_n > 0$ there are $b_n, c_n \in A$ such that $\mathbf{d}^p(t^g(b_n/A_n), t^g(c_n/A_n)) \leq \delta_n$ but $\mathbf{d}^p(t^g(ab_n/A_n), t^g(ac_n/A_n)) > \varepsilon$. Let $\varepsilon' < \varepsilon/5$. Since A is good, there is some $\delta > 0$ such that weak δ -embeddings of A extend to ε' -automorphisms of \mathfrak{M} and we may assume $\delta \leq \Delta^\varepsilon(\varepsilon')$. Now there are A_n, b_n, c_n, f_n for $n < \omega$ such that:

- $A_0 = \emptyset$, $A_n \subset A_{n+1} \subset A$, $|A_n| < \aleph_0$ and $\overline{\bigcup_{n<\omega} A_n} \supset A$,
- $\delta_n > 0$ is such that $\sum_{n<\omega} \delta_n < \delta$,
- $\mathbf{d}^p(t^g(b_n/A_n), t^g(c_n/A_n)) \leq \Delta^\delta(\delta_n)$ and $\mathbf{d}^p(t^g(ab_n/A_n), t^g(ac_n/A_n)) > \varepsilon$,
- $f_n \in \text{Aut}_{\delta_n}(\mathfrak{M})$ and $d(f_n(b_n A_n), c_n A_n) \leq \Delta^\delta(\delta_n)$,
- $b_n, c_n \in A_{n+1}$.

Now we can define δ -automorphisms (of \mathfrak{M}) $F_{\eta \upharpoonright n}$ for all $\eta \in {}^\omega 2$ and $n < \omega$ as follows:

- $F_\emptyset = \text{id}$,
- $F_{\eta \upharpoonright n+1} = \begin{cases} F_{\eta \upharpoonright n} & \text{if } \eta(n) = 0, \\ F_{\eta \upharpoonright n} \circ f_n & \text{if } \eta(n) = 1. \end{cases}$

We have $F_{\eta \upharpoonright n} \in \text{Aut}_\delta(\mathfrak{M})$ for each $\eta \in {}^\omega 2$ and $n < \omega$. Further

$$d(F_{\eta \upharpoonright n+1}(A_n), F_{\eta \upharpoonright n}(A_n)) \leq d(F_{\eta \upharpoonright n} \circ f_n(A_n), F_{\eta \upharpoonright n}(A_n)) \leq \delta_n.$$

Hence for each $\eta \in {}^\omega 2$, $(F_{\eta \upharpoonright n})_{n < \omega}$ converges (pointwise) on $\bigcup_{n < \omega} A_n$ and since the functions $F_{\eta \upharpoonright n}$ have a common modulus of uniform continuity, on A . Thus by Lemma 5.4 there is $F_\eta \in \text{Aut}_{\varepsilon'}(\mathfrak{M})$ such that $F_\eta(c) = \lim_{n \rightarrow \infty} F_{\eta \upharpoonright n}(c)$ for each $c \in A$.

Now let

$$D = \bigcup \{F_{\eta \upharpoonright n}(A_n) : \eta \in {}^\omega 2, n < \omega\}.$$

Then D is countable. We wish to show that

$$\mathbf{d}^p(t^g(F_\eta(a)/D), t^g(F_\nu(a)/D)) > \Delta^\varepsilon(\varepsilon')$$

for all $\eta \neq \nu \in {}^\omega 2$, hence arriving at a contradiction. For this we need the following:

CLAIM. $\mathbf{d}^p(t^g(f_n(ab_n A_n)/\emptyset), t^g(ac_n A_n/\emptyset)) > \varepsilon - \delta_n$.

Proof. Otherwise there would be some $f \in \text{Aut}_{\varepsilon - \delta_n}(\mathfrak{M})$ with

$$d(f \circ f_n(ab_n A_n), ac_n A_n) \leq \varepsilon - \delta_n,$$

but then $f \circ f_n \in \text{Aut}_\varepsilon(\mathfrak{M})$ and $\mathbf{d}^p(t^g(ab_n/A_n), t^g(ac_n/A_n)) \leq \varepsilon$, a contradiction. ■

Now let $\eta, \nu \in {}^\omega 2$, $\eta \neq \nu$ and $n = \min\{n : \eta(n) \neq \nu(n)\}$. We may assume $\eta(n) = 0$ and then use the following:

- for any $\zeta \in {}^\omega 2$, $d(F_\zeta(A_{n+1}), F_{\zeta \upharpoonright n+1}(A_{n+1})) \leq \delta \leq \Delta^\varepsilon(\varepsilon')$,
- $d(F_{\nu \upharpoonright n+1}(b_n A_n), F_{\eta \upharpoonright n+1}(c_n A_n)) = d(F_{\nu \upharpoonright n} \circ f_n(b_n A_n), F_{\eta \upharpoonright n}(c_n A_n)) \leq \delta_n$.

Then we must have

$$\mathbf{d}^p(t^g(F_\nu(a)/F_{\eta \upharpoonright n+1}(A_{n+1})), t^g(F_\eta(a)/F_{\eta \upharpoonright n+1}(A_{n+1}))) > \Delta^\varepsilon(\varepsilon')$$

since otherwise we can find $g \in \text{Aut}_{\varepsilon'}$ with

$$d(g(F_\nu(a)F_{\eta \upharpoonright n+1}(A_{n+1})), F_\eta(a)F_{\eta \upharpoonright n+1}(A_{n+1})) \leq \Delta^\varepsilon(\varepsilon')$$

and then $F_\eta^{-1} \circ g \circ F_\nu \circ f_n^{-1} \in \text{Aut}_{4\varepsilon'}$ and

$$\begin{aligned}
& d(F_\eta^{-1} \circ g \circ F_\nu \circ f_n^{-1}(f_n(ab_n A_n)), ac_n A_n) \\
&= d(F_\eta^{-1} \circ g \circ F_\nu(ab_n A_n), ac_n A_n) \\
&\leq d(F_\eta^{-1} \circ g \circ F_\nu(ab_n A_n), F_\eta^{-1} \circ g(F_\nu(a) F_{\nu \upharpoonright n+1}(b_n A_n))) \\
&\quad + d(F_\eta^{-1} \circ g(F_\nu(a) F_{\nu \upharpoonright n+1}(b_n A_n)), F_\eta^{-1} \circ g(F_\nu(a) F_{\eta \upharpoonright n+1}(c_n A_n))) \\
&\quad + d(F_\eta^{-1} \circ g(F_\nu(a) F_{\eta \upharpoonright n+1}(c_n A_n)), F_\eta^{-1}(F_\eta(a) F_{\eta \upharpoonright n+1}(c_n A_n))) \\
&\quad + d(F_\eta^{-1}(F_\eta(a) F_{\eta \upharpoonright n+1}(c_n A_n)), ac_n A_n) \\
&\leq 4\varepsilon',
\end{aligned}$$

showing that $\mathbf{d}^p(t^g(f_n(ab_n A_n)/\emptyset), t^g(ac_n A_n/\emptyset)) \leq 4\varepsilon' < \varepsilon - \delta_n$, contradicting the claim (the other direction needed by the symmetry of \mathbf{d}^p is proved similarly). ■

THEOREM 7.3. *If \mathbb{K} is ω - \mathbf{d}^p -stable and all models are good (i.e. $(\mathcal{A}, \mathcal{A})$ is good for all $\mathcal{A} \in \mathbb{K}$) then for all $a \in \mathfrak{M}$, $\mathcal{A} \preccurlyeq \mathfrak{M}$ and $\varepsilon > 0$ there is some finite $A' \subset \mathcal{A}$ such that $t^g(a/\mathcal{A})$ does not ε -split over A' .*

Proof. Assume towards a contradiction that \mathcal{B} , a and $\varepsilon > 0$ are such that $t^g(a/\mathcal{B})$ ε -splits over every finite subset of \mathcal{B} . Then we may construct a separable $\mathcal{A} \in \mathcal{B}$ as the closure of a countable union of at most countable sets as follows:

First let $A_0 \subset \mathcal{B}$ be any finite set. When A_n has been defined and is at most countable, by assumption for any finite $A' \subset A_n$ and any rational $r > 0$ there are $b'_r, c'_r \in \mathcal{B}$ such that $\mathbf{d}^p(t^g(b'_r/A'), t^g(c'_r/A')) \leq r$ but $\mathbf{d}^p(t^g(ab'_r/A'), t^g(ac'_r/A')) > \varepsilon$. Then let A_{n+1} be a countable set containing $A_n \cup \{b'_r, c'_r : r \in \mathbb{Q}, r > 0, A' \subset A_n \text{ finite}\}$ and such that $\bar{A}_{n+1} \preccurlyeq \mathcal{B}$. In the end let

$$\mathcal{A} = \overline{\bigcup_{n<\omega} A_n} = \overline{\bigcup_{n<\omega} \bar{A}_n} \preccurlyeq \mathcal{B}.$$

Then \mathcal{A} is separable but $t^g(a/\mathcal{A})$ ε' -splits over all finite $A \subset \mathcal{A}$ when $\varepsilon' \leq \varepsilon/2$, contradicting Theorem 7.2. Namely let $A \subset \mathcal{A}$ be finite and $\delta > 0$ and we may assume $\delta \leq \varepsilon'$. Now choose some rational $r \leq \delta/2$ and a finite $A' \subset \bigcup_{n<\omega} A_n$ with $d(A, A') \leq \Delta^{\varepsilon'}(\delta/4)$. Then $A' \in A_n$ for some $n < \omega$ and there are $b'_r, c'_r \in \mathcal{A}$ such that

$$(7.1) \quad \mathbf{d}^p(t^g(b'_r/A'), t^g(c'_r/A')) \leq r$$

but

$$(7.2) \quad \mathbf{d}^p(t^g(ab'_r/A'), t^g(ac'_r/A')) > \varepsilon.$$

Now if f is an r -function witnessing (7.1), we have

$$d(f(A), A) \leq d(f(A), f(A')) + d(f(A'), A') + d(A', A) \leq \delta.$$

Hence $\mathbf{d}^p(t^g(b'_r/A), t^g(c'_r/A)) \leq \delta$. But $\mathbf{d}^p(t^g(ab'_r/A), t^g(ac'_r/A))$ must be at least ε' since if there were an ε' -function g witnessing the contrary, we would have $d(g(ab'_r), ac'_r) \leq \varepsilon'$ and $d(g(A'), A') \leq \delta/4 + \varepsilon' + \delta/4 < 2\varepsilon' \leq \varepsilon$, contradicting (7.2) ■

THEOREM 7.4. *If \mathbb{K} is ω - \mathbf{d}^p -stable and has complete type-spaces then for all $a \in \mathfrak{M}$, $A \subset \mathfrak{M}$ and $\varepsilon > 0$ there is some finite $A' \subset A$ such that $t^g(a/A)$ does not ε -split over A' .*

Proof. This is proved by a tree construction just as in Theorem 7.2, but here we may directly choose $\delta < \Delta^\varepsilon(\varepsilon/7)$ (and forget about ε') and we just need A_n to form an increasing sequence of finite subsets of A , without any demands on the union. Then the automorphisms $F_{\eta \upharpoonright n}$ are defined as before. Now for each $\eta \in {}^\omega 2$, $\{t^g(F_{\eta \upharpoonright n}(a)/F_{\eta \upharpoonright n}(A_n)) : n < \omega\}$ forms a coherent system in the sense that for each $\varepsilon > 0$ there is some $n_0 < \omega$ such that for $n > m > n_0$, $\mathbf{d}^p(t^g(F_{\eta \upharpoonright n}(a)/F_{\eta \upharpoonright m}(A_m)), t^g(F_{\eta \upharpoonright m}(a)/F_{\eta \upharpoonright m}(A_m))) < \varepsilon$. Hence by Lemma 5.6 there exists a limit type of the sequence

$$(t^g(F_{\eta \upharpoonright n}(a)/F_{\eta \upharpoonright n}(A_n)))_{n < \omega},$$

realized by some a_η .

Now let $\eta \neq \nu$ with $n = \min\{n : \eta(n) \neq \nu(n)\}$ and again we may assume $\eta(n) = 0$. By choosing N large enough we can find $g_\eta, g_\nu \in \text{Aut}_\delta$ such that

$$\begin{aligned} d(g_\nu(F_{\nu \upharpoonright N}(aA_{n+1})), a_\nu F_{\nu \upharpoonright N}(A_{n+1})) &< \delta, \\ d(g_\eta(a_\eta F_{\eta \upharpoonright N}(A_{n+1}), F_{\eta \upharpoonright N}(aA_{n+1}))) &< \delta. \end{aligned}$$

Then we can easily calculate that:

- for any $\zeta \in {}^\omega 2$ and $N > n$, $d(F_{\zeta \upharpoonright N}(A_{n+1}), F_{\zeta \upharpoonright n+1}(A_{n+1})) < \delta$,
- $d(F_{\nu \upharpoonright n+1}(b_n A_n), F_{\eta \upharpoonright n+1}(c_n A_n)) < \delta_n$.

As in Theorem 7.2 we can prove that

$$(7.3) \quad \mathbf{d}^p(t^g(f_n(ab_n A_n)/\emptyset), t^g(ac_n A_n/\emptyset)) > \varepsilon - \delta_n.$$

So, similarly to Theorem 7.2, we must have

$$(7.4) \quad \mathbf{d}^p(t^g(a_\nu/F_{\eta \upharpoonright n+1}(A_{n+1}), t^g(a_\eta/F_{\eta \upharpoonright n+1}(A_{n+1}))) > \delta,$$

since otherwise we can find $g \in \text{Aut}_\delta$ with

$$d(g(a_\nu F_{\eta \upharpoonright n+1}(A_{n+1})), a_\eta F_{\eta \upharpoonright n+1}(A_{n+1})) < \delta$$

and then

$$F_{\eta \upharpoonright N}^{-1} \circ g_\eta \circ g \circ g_\nu \circ F_{\nu \upharpoonright N} \circ f_n^{-1} \in \text{Aut}_{6\delta}$$

and

$$\begin{aligned}
& d(F_{\eta \upharpoonright N}^{-1} \circ g_\eta \circ g \circ g_\nu \circ F_{\nu \upharpoonright N} \circ f_n^{-1}(f_n(ab_n A_n)), ac_n A_n) \\
&= d(F_{\eta \upharpoonright N}^{-1} \circ g_\eta \circ g \circ g_\nu \circ F_{\nu \upharpoonright N}(ab_n A_n), ac_n A_n) \\
&\leq d(F_{\eta \upharpoonright N}^{-1} \circ g_\eta \circ g \circ g_\nu(F_{\nu \upharpoonright N}(ab_n A_n)), F_{\eta \upharpoonright N}^{-1} \circ g_\eta \circ g(a_\nu F_{\nu \upharpoonright N}(b_n A_n))) \\
&\quad + d(F_{\eta \upharpoonright N}^{-1} \circ g_\eta \circ g(a_\nu F_{\nu \upharpoonright N}(b_n A_n)), F_{\eta \upharpoonright N}^{-1} \circ g_\eta \circ g(a_\nu F_{\nu \upharpoonright n+1}(b_n A_n))) \\
&\quad + d(F_{\eta \upharpoonright N}^{-1} \circ g_\eta \circ g(a_\nu F_{\nu \upharpoonright n+1}(b_n A_n)), F_{\eta \upharpoonright N}^{-1} \circ g_\eta \circ g(a_\nu F_{\eta \upharpoonright n+1}(c_n A_n))) \\
&\quad + d(F_{\eta \upharpoonright N}^{-1} \circ g_\eta \circ g(a_\nu F_{\eta \upharpoonright n+1}(c_n A_n)), F_{\eta \upharpoonright N}^{-1} \circ g_\eta(a_\eta F_{\eta \upharpoonright n+1}(c_n A_n))) \\
&\quad + d(F_{\eta \upharpoonright N}^{-1} \circ g_\eta(a_\eta F_{\eta \upharpoonright n+1}(c_n A_n)), F_{\eta \upharpoonright N}^{-1} \circ g_\eta(a_\eta F_{\eta \upharpoonright N}(c_n A_n))) \\
&\quad + d(F_{\eta \upharpoonright N}^{-1} \circ g_\eta(a_\eta F_{\eta \upharpoonright N}(c_n A_n)), ac_n A_n) \\
&\leq 6\varepsilon/7 < \varepsilon - \delta_n,
\end{aligned}$$

contradicting (7.3). Now define $D = \bigcup\{F_{\eta \upharpoonright n}(A_n) : \eta \in {}^\omega 2, n < \omega\}$. Then D is countable and (7.4) contradicts the assumption of ω -**d^p**-stability. ■

Based on ε -splitting we can define a notion of independence.

DEFINITION 7.5. We write $a \downarrow_A^\varepsilon B$ if there is some finite $A' \subseteq A$ such that $t^g(a/A \cup B)$ does not ε -split over A' . We then define $a \downarrow_A^s B$ if for all $\varepsilon > 0$, $a \downarrow_A^\varepsilon B$.

We now set out to prove that this notion of independence satisfies the usual axioms for an independence notion. The axiom of finite character is replaced by countable character, as is to be expected in a metric setting.

THEOREM 7.6. *If \mathbb{K} is ω -**d^p**-stable and has complete type spaces then independence satisfies the following axioms:*

- (1) (Isomorphism invariance) *If $a \downarrow_A^s B$ and f is a 0-automorphism of \mathfrak{M} then $f(a) \downarrow_{f(A)}^s f(B)$.*
- (2) (Monotonicity) *If $A \subseteq B \subseteq C \subseteq D$ and $a \downarrow_A^s D$ then $a \downarrow_B^s C$.*
- (3) (Countable character of nonfreeness) *If A is ω -**d^p**-saturated and $a \not\downarrow_A^s B$ then there is a countable $A' \subseteq A$ and a finite $B' \subseteq B$ such that if $t^g(b/A'B') = t^g(a/A'B')$ then $b \not\downarrow_{A'}^s B'$.*
- (4) (Local character) *$a \downarrow_A^s A$ for all a and A , i.e. for every a , A and $\varepsilon > 0$ there is some finite $E \subseteq A$ such that $a \downarrow_E^\varepsilon A$.*
- (5) (Extension) *If $a \downarrow_A^s B$, B is ω -**d^p**-saturated and $A \subseteq B \subseteq C$ then there is b with $t^g(b/B) = t^g(a/B)$ satisfying $b \downarrow_A^s C$.*
- (6) (Stationarity) *If A is ω -**d^p**-saturated, $A \subseteq B$, $t^g(a/A) = t^g(b/A)$, $a \downarrow_A^s B$ and $b \downarrow_A^s B$, then $t^g(a/B) = t^g(b/B)$.*
- (7) (Transitivity) *If $A \subseteq B \subseteq C$, B is ω -**d^p**-saturated, $a \downarrow_A^s B$ and $a \downarrow_B^s C$ then $a \downarrow_A^s C$.*

- (8) (Symmetry) If A is ω - \mathbf{d}^p -saturated then $a \downarrow_A^s b$ if and only if $b \downarrow_A^s a$.
(9) (Reflexivity) If A is ω - \mathbf{d}^p -saturated and $a \notin A$ then $a \nmid_A^s a$.

REMARK 7.7. We will see that the assumption of complete type-spaces is used only in the proofs of local character and extension and then the extension property is used to prove transitivity and symmetry. Hence, we may exchange the assumption of complete type-spaces for the assumptions that models are good and extension holds. Then all the axioms hold with the minor change that local character is true only when A is a model.

Proof of Theorem 7.6. By the definition it is clear that independence is preserved under 0-automorphisms and is monotone. Local character is Theorem 7.4.

Countable character of nonfreeness. Let $A' \subset A$ be countable such that $\overline{A'}$ is ω - \mathbf{d}^p -saturated and $a \downarrow_{A'}^s A$. Let a' be such that $t^g(a'/A) = t^g(a/A)$ and $a' \downarrow_A^s B$. Then let $B' \subseteq B$ be such that $t^g(a/B') \neq t^g(a'/B')$. The claim follows by stationarity which is proved below.

Extension. Let, for each $n < \omega$, $A_n \subset A$ be finite and such that $t^g(a/B)$ does not $\frac{1}{n+1}$ -split over A_n . Then for each finite $c \in C$, by ω - \mathbf{d}^p -saturation of B , let $c_n^c \in B$ be such that $\mathbf{d}^p(t^g(c_n^c/A_n), t^g(c/A_n)) \leq \frac{1}{n+1}$ and let f_n^c be a $\frac{1}{n+1}$ -automorphism of \mathfrak{M} witnessing this. Then define $a_n^c = f_n^c(a)$. We claim that $\{t^g(a_n^c/A_n c) : n < \omega, c \in C \text{ finite}\}$ forms a \mathbf{d}^p -coherent system in the sense of Lemma 5.6. Namely, if $c \in C$ and $\varepsilon > 0$ are given, then choose n_1 large enough such that $1/n_1 < \Delta^\varepsilon(\varepsilon/3)$. Now $t^g(a/B)$ does not $1/n_1$ -split over A_{n_1} and there is some $\delta > 0$ witnessing this. Choose $n_0 \geq n_1$ such that $2/n_0 \leq \Delta^\varepsilon(\delta)$ and let $B_0 = A_{n_1} \cup c$. Then if $n \geq m \geq n_0$, for any finite $d, e \in C$, we need to show that $\mathbf{d}^p(t^g(a_n^{cd}/A_{n_1} cd), t^g(a_m^{cd}/A_{n_1} cd)) \leq \varepsilon$. But now

$$\begin{aligned} \mathbf{d}^p(t^g(a_n^{cd}/A_{n_1}), t^g(a, (cd)_n^{cd}/A_{n_1})) &\leq 1/n_1, \\ \mathbf{d}^p(t^g(a_m^{cd}/A_{n_1}), t^g(a, (cd)_m^{cd}/A_{n_1})) &\leq 1/n_1. \end{aligned}$$

Further

$$\mathbf{d}^p(t^g((cd)_n^{cd}/A_{n_1}), t^g((cd)cd_m/A_{n_1})) \leq \delta$$

and $(cd)_n^{cd}, (cd)_m^{cd} \in B$, so the claim follows by non- $1/n_1$ -splitting. Hence by Lemma 5.6 there is some b realizing the limit type of all a_n^c 's, i.e. for all finite $c \in C$ and $\varepsilon > 0$ there are $n_0 < \omega$ and $c_0 \in C$ such that for all $n > n_0$ and $d \in C$ with $c_0 \subseteq d$, $\mathbf{d}^p(t^g(b/A_nd), t^g(a_n^d/A_nd)) \leq \varepsilon$.

Next we claim $t^g(b/B) = t^g(a/B)$. For this it suffices to prove that $\mathbf{d}^p(t^g(bc/\emptyset), t^g(ac/\emptyset)) \leq \varepsilon$ for any $\varepsilon > 0$ and finite $c \in B$. So let $\varepsilon > 0$ and $c \in B$ be given and let n be large enough so that $t^g(a/B)$ does not $\varepsilon/3$ -split over A_n . Then let $\delta > 0$ witness this and be at most $\Delta^\varepsilon(\varepsilon/3)$. Choosing m large enough, we can ensure

- $\mathbf{d}^p(t^g(bc/A_n), t^g(a_m^c c/A_n)) \leq \delta$,
- $\mathbf{d}^p(t^g(a_m^c c/A_n), t^g(ac_m^c/A_n)) \leq \delta$.

Then by nonsplitting we get $\mathbf{d}^p(t^g(ac_m^c/A_n), t^g(ac/A_n)) \leq \varepsilon/3$. Combining these we get $\mathbf{d}^p(t^g(bc/A_n), t^g(ac/A_n)) \leq \varepsilon$.

Finally we need to show that $b \downarrow_A^s C$. We actually claim that $b \downarrow_{A_n}^{\frac{1}{n+1}} C$. But this is easy to see, since if for every $\delta > 0$ there were $c, d \in C$ witnessing $\frac{1}{n+1}$ -splitting, then for some m large enough, $\mathbf{d}^p(t^g(c_m^c/A_n), t^g(d_m^d/A_n))$ would be small enough to contradict the assumption $a \downarrow_{A_n}^{\frac{1}{n+1}} B$, since

$$\mathbf{d}^p(t^g(bc/A_n), t^g(a_m^c c/A_n))$$

can be made arbitrarily small with a large enough m and

$$\mathbf{d}^p(t^g(a_m^c c/A_n), t^g(ac_m^c/A_n)) \leq 1/m$$

for $m \geq n$ (and similarly for d).

Stationarity. To show that $t^g(a/B) = t^g(b/B)$, we need to show that for all finite $c \in B$, $t^g(ac/\emptyset) = t^g(bc/\emptyset)$. By perturbation, it is enough to show that for all finite $c \in B$ and all $\varepsilon > 0$, $\mathbf{d}^p(t^g(ac/\emptyset), t^g(bc/\emptyset)) \leq \varepsilon$. So fix some finite $c \in B$ and $\varepsilon > 0$. By assumption there is some finite $A' \subset A$ such that $t^g(a/B)$ and $t^g(b/B)$ do not $\Delta^\varepsilon(\varepsilon/2)$ -split over A' and there is some $\delta > 0$ witnessing this. Since A is ω - \mathbf{d}^p -saturated, there is $c' \in A$ such that $\mathbf{d}^p(t^g(c'/A'), t^g(c/A')) \leq \delta$. Then

- $\mathbf{d}^p(t^g(ac/\emptyset), t^g(ac'/\emptyset)) \leq \Delta^\varepsilon(\varepsilon/2)$,
- $\mathbf{d}^p(t^g(ac'/\emptyset), t^g(bc'/\emptyset)) = 0$,
- $\mathbf{d}^p(t^g(bc'/\emptyset), t^g(bc/\emptyset)) \leq \Delta^\varepsilon(\varepsilon/2)$.

Combining the above we get

$$\mathbf{d}^p(t^g(ac/\emptyset), t^g(bc/\emptyset)) \leq \varepsilon.$$

Transitivity. By the extension property let b be such that $t^g(b/B) = t^g(a/B)$ and $b \downarrow_A^s C$. Then $a \downarrow_B^s C$ and $b \downarrow_B^s C$ so by stationarity $t^g(a/C) = t^g(b/C)$, hence $a \downarrow_A^s C$.

Symmetry. The proof is postponed until Lemma 7.13.

Reflexivity. Let d be the distance of a to A , i.e. $d = \inf\{d(a, a') : a' \in A\}$. Since $a \notin \bar{A}$, we have $d > 0$. Now $t^g(a/Aa) \Delta^d(d)/3$ -splits over any finite $A' \subset A$. Namely let $A' \subset A$ be finite and $\delta > 0$. By ω - \mathbf{d}^p -saturation, let $a' \in A$ be such that $\mathbf{d}^p(t^g(a'/A'), t^g(a/A')) \leq \delta$. Then if $f \in \text{Aut}_\varepsilon$ for some $\varepsilon \leq d$, $d(f(aa'A'), aaA') \geq \Delta^\varepsilon(d)/2 \geq \Delta^d(d)/2$ and thus $\mathbf{d}^p(t^g(aa'/A'), t^g(aa/A')) > \Delta^d(d)/3$. ■

Although we have a weaker stability notion than in [11], the ω -stability notion at hand implies discrete λ -stability for all λ with $\lambda = \lambda^{\aleph_0}$.

LEMMA 7.8. *If \mathbb{K} is ω - \mathbf{d}^p -stable and either has complete type-spaces or good models (i.e. $(\mathcal{A}, \mathcal{A})$ is good for all $\mathcal{A} \in \mathbb{K}$) then for all $\lambda = \lambda^{\aleph_0}$, \mathbb{K} is λ -stable in the discrete sense, i.e. the number of types over a set of density character λ is λ (which then also is the cardinality of the set).*

Proof. Let $|B| \leq \lambda$ and let $a, b \in \mathfrak{M}$. By Lemma 6.2 we may assume B is an ω - \mathbf{d}^p -saturated model. By Theorem 7.4 or 7.3 there exists a separable $A \subset B$ such that $a \downarrow_A^s B$ and $b \downarrow_A^s B$, and again by Lemma 6.2 and monotonicity we may assume A is ω - \mathbf{d}^p -saturated and closed in B . Then by stationarity (Theorem 7.6) $t^g(a/B) = t^g(b/B)$ if and only if $t^g(a/A) = t^g(b/A)$. Hence to count the types over B it suffices to count the number of types over separable closed subsets of B . Since $\lambda^{\aleph_0} = \lambda$, there are only λ such subsets of B . Further types over separable sets are determined by their restrictions to some dense (countable) subset and there are only 2^{\aleph_0} types over countable sets (recall that the vocabulary is countable), so there are just λ types over B . ■

Since our stability notion in a sense considers weak types, our perturbation assumption gives a property that resembles the *metric homogeneity* assumption from [11]. As might be expected, we get similar stability results to [11] and below we prove that ω - \mathbf{d}^p -stability implies λ - \mathbf{d}^p -stability for all infinite λ . It is, however, worth noting that we do not assume metric homogeneity, which considers the orbits of Galois types over fixed parameter sets.

THEOREM 7.9. *If \mathbb{K} is ω - \mathbf{d}^p -stable and has complete type-spaces or good models (i.e. $(\mathcal{A}, \mathcal{A})$ is good for all $\mathcal{A} \in \mathbb{K}$) then it is λ - \mathbf{d}^p -stable for all infinite λ .*

Proof. Assume towards a contradiction that \mathbb{K} is not λ -stable. Then there is $B \subset \mathfrak{M}$ with $|B| = \lambda$ such that the density of $S(B)$ with respect to \mathbf{d}^p is at least λ^+ and by Lemma 6.2 we may assume B is an ω - \mathbf{d}^p -saturated model. Now there are a_i , $i < \lambda^+$, and some $\varepsilon > 0$ such that

$$\mathbf{d}^p(t^g(a_i/B), t^g(a_j/B)) \geq \varepsilon \quad \text{for all } i < j < \lambda^+.$$

Let $\varepsilon' = \Delta^\varepsilon(\varepsilon/6)$. For each $i < \lambda^+$, by either Theorem 7.3 or 7.4, choose a finite $A_i \subset B$ and a rational $\delta_i > 0$ such that $t^g(a_i/B)$ does not ε' -split over A_i and δ_i witnesses this. λ^+ of the pairs (A_i, δ_i) are the same, denote these by A and δ . Next, by Lemma 6.2 let $\mathcal{A} \preccurlyeq B$ be separable, ω - \mathbf{d}^p -saturated and contain A .

Now for $i \neq j$, since

$$\mathbf{d}^p(t^g(a_i/B), t^g(a_j/B)) = \sup\{\mathbf{d}^p(t^g(a_i c / \emptyset), t^g(a_j c / \emptyset)) : c \in B \text{ finite}\},$$

there is some finite $c_{ij} \in B$ such that

$$\mathbf{d}^p(t^g(a_i c_{ij} / \emptyset), t^g(a_j c_{ij} / \emptyset)) > \varepsilon/2.$$

For all $i \neq j$ such that $(A_i, \delta_i) = (A_j, \delta_j) = (A, \delta)$, let $c' \in \mathcal{A}$ be such that $\mathbf{d}^p(t^g(c_{ij}/A), t^g(c'/A)) \leq \delta$. Then we must have

$$(7.5) \quad \mathbf{d}^p(t^g(a_i/c'A), t^g(a_j/c'A)) > \varepsilon'$$

since by nonsplitting $\mathbf{d}^p(t^g(a_i c_{ij}/A), t^g(a_i c'/A)) \leq \varepsilon'$ (and similarly for a_j), and if (7.5) does not hold this would give

$$\mathbf{d}^p(t^g(a_i c_{ij}/A), t^g(a_j c_{ij}/A)) \leq 3\varepsilon/6,$$

a contradiction. But then we have $\mathbf{d}^p(t^g(a_i/\mathcal{A}), t^g(a_j/\mathcal{A})) > \varepsilon'$ for λ^+ many $i \neq j$, contradicting ω - \mathbf{d}^p -stability. ■

Next we will tie our notion of independence to the usual one in homogeneous model theory. In [11] it is shown how to put the abstract setting of homogeneous metric abstract elementary classes into a homogeneous first order context, hence gaining access to results from [12] (as long as we work in a stable class).

We recall some definitions from [12], as stated in a formula-free fashion in [11].

DEFINITION 7.10.

- (1) Let $A \subset B$ and $a \in \mathfrak{M}$. We say that $t^g(a/B)$ splits over A if there are $b, c \in B$ such that $t^g(b/A) = t^g(c/A)$ but $t^g(b/A \cup a) \neq t^g(c/A \cup a)$.
- (2) A type $t^g(a/B)$ is said to split strongly over $A \subset B$ if there are $b, c \in B$ and an infinite sequence I , indiscernible over A , with $b, c \in I$ such that $t^g(b/A \cup a) \neq t^g(c/A \cup a)$.
- (3) $\kappa(\mathbb{K})$ denotes the least cardinal such that there are no a, b_i and c_i for $i < \kappa(\mathbb{K})$ such that $t^g(a/\bigcup_{j \leq i}(b_j \cup c_j))$ splits strongly over $\bigcup_{j < i}(b_j \cup c_j)$ for each $i < \kappa(\mathbb{K})$.
- (4) $a \downarrow_A B$ if there is $C \subseteq A$ of cardinality $< \kappa(\mathfrak{M})$ such that for all $D \supseteq A \cup B$ there is b with $t^g(b/A \cup B) = t^g(a/A \cup B)$ such that $t^g(b/D)$ does not split strongly over C .
- (5) $\lambda(\mathfrak{M})$ is the least cardinality in which \mathfrak{M} is stable.

REMARK 7.11. Note that if $a \downarrow_A^s B$ then by perturbation $t^g(a/A \cup B)$ does not split over A and hence it does not split strongly over A .

Now the equivalence of \downarrow and \downarrow^s over saturated enough models can be proved just as in [11].

THEOREM 7.12. Assume \mathbb{K} is ω - \mathbf{d}^p -stable and has complete type-spaces. If $\mathcal{A} \subseteq B$, $a \in \mathfrak{M}$ and \mathcal{A} is 2^{\aleph_0} -saturated (i.e. realizes all Galois types over parameter sets of size $< 2^{\aleph_0}$) then

$$a \downarrow_A^s B \quad \text{if and only if} \quad a \downarrow_A B,$$

i.e. our notion of independence agrees with that in classical homogeneous model theory over 2^{\aleph_0} -saturated models.

Proof. First assume $a \downarrow_{\mathcal{A}} B$. By local character (Theorem 7.6), $a \downarrow_{\mathcal{A}}^s \mathcal{A}$. Let $\mathcal{B} \supseteq B$ be 2^{\aleph_0} -saturated and by extension (Theorem 7.6) find some b realizing $t^g(a/\mathcal{A})$ and satisfying $b \downarrow_{\mathcal{A}}^s \mathcal{B}$. Then $t^g(b/\mathcal{B})$ does not split strongly over \mathcal{A} , so by [12, Lemma 3.2(iii)], $b \downarrow_{\mathcal{A}} \mathcal{B}$, and by monotonicity $b \downarrow_{\mathcal{A}} B$. Then by stationarity of \downarrow ([12, Lemma 3.4]), $t^g(b/B) = t^g(a/B)$ so $a \downarrow_{\mathcal{A}}^s B$.

For the other direction assume $a \downarrow_{\mathcal{A}}^s B$. Again by [12, Lemma 3.2(iii)], $a \downarrow_{\mathcal{A}} \mathcal{A}$. Since \downarrow has built-in extensions, this implies the existence of some b realizing $t^g(a/\mathcal{A})$ and satisfying $b \downarrow_{\mathcal{A}} B$. Then by the previous direction $b \downarrow_{\mathcal{A}}^s B$, so by stationarity of \downarrow^s we are done. ■

Now using symmetry of \downarrow over so-called a-saturated (strongly $\kappa(\mathfrak{M})$ -saturated) models ([12, Lemma 3.6]) we can prove symmetry for \downarrow^s . We use the fact ([12, Lemma 1.9(iv)]) that $\lambda(\mathfrak{M})$ -saturated models are a-saturated and recall that by Lemma 7.8, $\lambda(\mathfrak{M})$ is at most 2^{\aleph_0} when \mathbb{K} is $\omega\text{-}\mathbf{d}^p$ -stable and has complete type-spaces.

LEMMA 7.13. *If \mathbb{K} is $\omega\text{-}\mathbf{d}^p$ -stable and has complete type-spaces, and if A is $\omega\text{-}\mathbf{d}^p$ -saturated and $a \downarrow_A^s b$, then $b \downarrow_A^s a$.*

Proof. Assume towards a contradiction that $a \downarrow_A^s b$ but $b \not\downarrow_A^s a$. Let $\mathcal{B} \supset A$ be 2^{\aleph_0} -saturated. Now $b \downarrow_A^s A$ so by the extension property (Theorem 7.6) there is some b' realizing $t^g(b/A)$ and satisfying $b' \downarrow_A^s \mathcal{B}$. We may assume $b' = b$ (since moving \mathcal{B} by a 0-automorphism does not affect its degree of saturation). Using the extension property and stationarity, we can find some a' realizing $t^g(a/Ab)$ and satisfying $a' \downarrow_A^s \mathcal{B}b$. Now by monotonicity we have $a' \downarrow_{\mathcal{B}}^s b$. But we cannot have $b \downarrow_{\mathcal{B}}^s a'$ since then by transitivity we would have $b \downarrow_A^s \mathcal{B}a'$, by monotonicity $b \downarrow_A^s a'$ and hence by invariance $b \downarrow_A^s a$. Hence $a' \downarrow_{\mathcal{B}}^s b$ and $b \not\downarrow_{\mathcal{B}}^s a'$, but since \mathcal{B} is 2^{\aleph_0} -saturated, by Lemma 7.12 this contradicts the symmetry of \downarrow ([12, Lemma 3.6]) over a-saturated models. ■

8. Isolation. In this section we develop a notion of isolation and build constructible sets. With our notion of isolation, dominance works, but whether our constructible sets are prime in the class of $\omega\text{-}\mathbf{d}^p$ -saturated models in any reasonable sense is an open question. We will discuss this matter at the end of the section.

DEFINITION 8.1. We say that $t^g(a/A)$ is ε -isolated if there are some $\delta > 0$ and finite $A' \subset A$ such that $\mathbf{d}^p(t^g(b/A'), t^g(a/A')) \leq \delta$ implies

$$\mathbf{d}^p(t^g(b/A), t^g(a/A)) \leq \varepsilon.$$

We say that $t^g(a/A)$ is isolated if it is ε -isolated for all $\varepsilon > 0$.

The next lemma also follows from [2, Proposition 3.7].

LEMMA 8.2. *If \mathbb{K} is ω - \mathbf{d}^p -stable and has complete type-spaces then for all A , $a, \varepsilon > 0$, $\delta > 0$ and finite $B \subset A$ there is some a' with*

$$\mathbf{d}^p(t^g(a'/B), t^g(a/B)) \leq \delta$$

such that $t^g(a'/A)$ is ε -isolated.

Proof. Assume towards a contradiction that the claim is false. Then for every a' with $\mathbf{d}^p(t^g(a'/B), t^g(a/B)) \leq \delta$ and for all $\delta' > 0$ and finite $A' \subset A$ there is some b such that $\mathbf{d}^p(t^g(b/A'), t^g(a'/A')) \leq \delta'$ but

$$\mathbf{d}^p(t^g(b/A), t^g(a'/A)) > \varepsilon.$$

Let $0 < \delta' \leq \min\{\delta/4, \Delta^\delta(\varepsilon/5)\}$. Then define, for $n < \omega$ and $\xi \in 2^{<\omega}$, $A_n \subset A$, $\delta_n > 0$, $A_\xi \subset A$ and a_ξ such that:

- $A_0 = B$, $A_n \subseteq A_{n+1}$, A_n is finite,
- $\delta_{n+1} \leq \delta_n$ and $\sum_{n < \omega} \delta_n < \Delta^{\delta'}(\delta')$,
- $a_\emptyset = a$ and $a_{\xi^\frown (0)} = a_\xi$,
- $\mathbf{d}^p(t^g(a_\xi/B), t^g(a/B)) \leq \sum_{i < \text{length}(\xi)} \delta_i < \delta'$,
- $\mathbf{d}^p(t^g(a_{\xi^\frown (0)}/A_{\text{length}(\xi)}), t^g(a_{\xi^\frown (1)}/A_{\text{length}(\xi)})) \leq \Delta^{\delta'}(\delta_{\text{length}(\xi)})$ but

$$\mathbf{d}^p(t^g(a_{\xi^\frown (0)}/A_\xi), t^g(a_{\xi^\frown (1)}/A_\xi)) > \varepsilon,$$
- $A_{n+1} = \bigcup_{\text{length}(\xi)=n} A_\xi$.

When $\text{length}(\xi) = n$ and A_n and a_ξ have been defined such that

$$\mathbf{d}^p(t^g(a_\xi/B), t^g(a/B)) \leq \sum_{i < n} \delta_i$$

then by our counter-assumption we may find some $a_{\xi^\frown (1)}$ satisfying

$$(8.1) \quad \mathbf{d}^p(t^g(a_\xi/A_n), t^g(a_{\xi^\frown (1)}/A_n)) \leq \Delta^{\delta'}(\delta_n),$$

$$(8.2) \quad \mathbf{d}^p(t^g(a_\xi/A), t^g(a_{\xi^\frown (1)}/A)) > \varepsilon.$$

Then we let $A_\xi \subset A$ be finite and witness (8.2), and note that (8.1) implies

$$\mathbf{d}^p(t^g(a_{\xi^\frown (1)}/B), t^g(a/B)) \leq \sum_{i < n+1} \delta_i \leq \delta',$$

keeping the induction going.

Next note that for each $\eta \in {}^\omega 2$, $\{t^g(a_{\eta \upharpoonright n}/A_n) : n < \omega\}$ forms a \mathbf{d}^p -coherent system in the sense of Lemma 5.6. Hence there is for each $\eta \in {}^\omega 2$ a limit element a_η . But now for $\eta \neq \nu$, $\eta, \nu \in {}^\omega 2$,

$$(8.3) \quad \mathbf{d}^p\left(t^g\left(a_\eta / \bigcup_{n < \omega} A_n\right), t^g\left(a_\nu / \bigcup_{n < \omega} A_n\right)\right) > \delta',$$

since otherwise, when $n = \min\{n : \eta(n) \neq \nu(n)\}$ and N is large enough,

- $\mathbf{d}^p(t^g(a_{\eta \upharpoonright n+1}/A_{n+1}), t^g(a_{\eta \upharpoonright N}/A_{n+1})) \leq \delta'$ (and similarly for ν) by construction,
- $\mathbf{d}^p(t^g(a_{\eta \upharpoonright N}/A_{n+1}), t^g(a_\eta/A_{n+1})) \leq \delta'$ (and similarly for ν) by choice of N and
- $\mathbf{d}^p(t^g(a_\eta/A_{n+1}), t^g(a_\nu/A_{n+1})) \leq \delta'$

and combining these gives

$$\mathbf{d}^p(t^g(a_{\eta \upharpoonright n+1}/A_{n+1}), t^g(a_{\nu \upharpoonright n+1}/A_{n+1})) \leq \varepsilon,$$

contradicting the way we chose the a_ξ s.

But (8.3) contradicts ω - \mathbf{d}^p -stability since $\bigcup_{n < \omega} A_n$ is countable. ■

THEOREM 8.3. *If \mathbb{K} is ω - \mathbf{d}^p -stable and has complete type-spaces then for all A , a , $\delta > 0$ and finite $B \subset A$ there is a' with $\mathbf{d}^p(t^g(a'/B), t^g(a/B)) \leq \delta$ such that $t^g(a'/A)$ is isolated.*

Proof. By the previous lemma we can, for each positive $n < \omega$ and δ'_n , find finite sets $A_n \subset A$, $\delta_n > 0$ and elements a_n such that A_n and δ_n $1/n$ -isolate $t^g(a_n/A)$ and $\mathbf{d}^p(t^g(a_{n+1}/A_n), t^g(a_n/A_n)) \leq \delta'_{n+1}$. We just need to make sure that all a_i for $i > n$ are within distance δ_n of a_n .

So we define:

- $A_0 = B$, $a_0 = a$, $\delta'_0 = \delta_0 = \delta/2$,
- $A_n \subset A_{n+1} \subset A$, A_n is finite and for $n > 0$, δ_n and A_n $1/n$ -isolate $t^g(a_n/A)$,
- $\delta'_{n+1} \leq \frac{1}{2} \min\{\Delta^\delta(\delta_n/2), \delta'_n\}$,
- $\mathbf{d}^p(t^g(a_{n+1}/A_n), t^g(a_n/A_n)) \leq \Delta^\delta(\delta'_{n+1})$.

Then $\{t^g(a_n/A_n) : n < \omega\}$ forms a \mathbf{d}^p -coherent system in the sense of Lemma 5.6 and hence there is a limit element a' such that

$$\mathbf{d}^p(t^g(a'/A_n), t^g(a_n/A_n)) \leq \Delta^\delta(\delta_n/2)$$

for each $n < \omega$. Hence in particular $\mathbf{d}^p(t^g(a'/B), t^g(a/B)) \leq \delta$ and it is fairly straightforward to see that if $1/n \leq \Delta^\varepsilon(\varepsilon/2)$ for a given $\varepsilon > 0$ then A_n and $\Delta^\delta(\delta_n/2)$ ε -isolate $t^g(a'/A)$. ■

DEFINITION 8.4. We say that A^* is *constructible over A* if $A^* = A \cup \{a_i : i < \alpha\}$ and for each $i < \alpha$, $t^g(a_i/A \cup \{a_j : j < i\})$ is isolated.

REMARK 8.5. Note that if B is constructible over A then \bar{B} is also constructible over A , since any element from B close enough to $b \in \bar{B}$ isolates the type of b over any set.

THEOREM 8.6. *If \mathbb{K} is ω - \mathbf{d}^p -stable and has complete type-spaces then for each A there is a complete ω - \mathbf{d}^p -saturated constructible set over A .*

Proof. This is done more or less in the standard way. If we have constructed $A_\alpha = A \cup \{a_i : i < \alpha\}$ and A_α is not ω - \mathbf{d}^p -saturated then there

is some finite $A' \subset A_\alpha$, some a and $\delta > 0$ such that no $a' \in A_\alpha$ satisfies $\mathbf{d}^p(t^g(a'/A'), t^g(a/A')) \leq \delta$. But by Theorem 8.3 we can find some a' such that $t^g(a'/A_\alpha)$ is isolated and $\mathbf{d}^p(t^g(a'/A'), t^g(a/A')) \leq \delta$ and then let $a_\alpha = a'$. The construction will eventually terminate at some α (essentially as in [14, Theorem IV.3.1]) giving a constructible ω - \mathbf{d}^p -saturated set A_α . However, A_α need not be complete, but by Remark 8.5 we may continue the construction until some $\alpha' \geq \alpha$ such that $A_{\alpha'} = \bar{A}_\alpha$, and by Remark 6.3, $A_{\alpha'}$ is still ω - \mathbf{d}^p -saturated. ■

DEFINITION 8.7. $B \supset A$ is *atomic over A* if for each $b \in B$, $t^g(b/A)$ is isolated.

THEOREM 8.8. *If B is constructible over A then B is atomic over A .*

Proof. This is done by a fairly straightforward induction on $i < \alpha$, where $B = A \cup \{a_i : i < \alpha\}$ and each $t^g(a_i/A \cup \{a_j : j < i\})$ is isolated. ■

THEOREM 8.9. *Assume \mathbb{K} is ω - \mathbf{d}^p -stable and has complete type-spaces. If A is ω - \mathbf{d}^p -saturated, $B \supseteq A$ and B^* is constructible over B then $a \downarrow_A^s B$ implies $a \downarrow_A^s B^*$.*

Proof. Assume towards a contradiction that $a \downarrow_A^s B$ but $a \not\downarrow_A^s B^*$. By countable character (Theorem 7.6) there is a finite $b \in B^*$ such that $a \not\downarrow_A^s b$. Further by symmetry $b \not\downarrow_A^s a$, so there is some $\varepsilon > 0$ such that $b \not\downarrow_A^\varepsilon a$, i.e. $t^g(b/Aa)$ ε -splits over every finite $A' \subset A$.

Let $\varepsilon' < \Delta^\varepsilon(\varepsilon/2)$ and $\varepsilon'' < \Delta^{\varepsilon'}(\varepsilon'/2)$.

By Theorem 8.8, B^* is atomic over B so we can find some finite $B' \subset B^*$ and $\delta' > 0$ with $\delta' \leq \varepsilon''$ such that B' and δ' witness the ε'' -isolation of $t^g(b/B)$. Then let $\delta'' < \Delta^\varepsilon(\delta'/2)$. Since $a \downarrow_A^s B$, we have $a \downarrow_A^s B'$ and again by symmetry $B' \downarrow_A^s a$. Hence there is some finite $A' \subset A$ such that $t^g(B'/Aa)$ does not δ'' -split over A' and there is some $\delta''' > 0$ witnessing this. Now since $b \not\downarrow_A^\varepsilon a$ there are $c, c' \in A \cup a$ such that $\mathbf{d}^p(t^g(c/A'), t^g(c'/A')) \leq \delta'''$ but

$$\mathbf{d}^p(t^g(bc/A'), t^g(bc'/A')) > \varepsilon.$$

Since δ''' witnesses the fact that $t^g(B'/Aa)$ does not δ'' -split over A' , we must have $\mathbf{d}^p(t^g(B'c/A'), t^g(B'c'/A')) \leq \delta''$. So there is $f \in \text{Aut}_{\delta''}$ such that

$$d(f(B'cA'), B'c'A') \leq \delta''.$$

CASE 1: $c' \in A$ (or $c \in A$, but that case is treated symmetrically). Then $\mathbf{d}^p(t^g(b/B'), t^g(f(b)/B')) \leq \delta''$ so by the ε'' -isolation of $t^g(b/B)$,

$$\mathbf{d}^p(t^g(b/B), t^g(f(b)/B)) \leq \varepsilon''.$$

So (since $A \subseteq B$) there is $g \in \text{Aut}_{\varepsilon''}$ satisfying $d(g(f(b)c'A'), bc'A') \leq \varepsilon''$ and as

$$d(g \circ f(bcA'), bc'A') \leq d(g \circ f(bcA'), g(f(b)c'A')) + d(g(f(b)c'A'), bc'A')$$

we have

$$\mathbf{d}^p(t^g(bc/A'), t^g(bc'/A')) \leq 2\varepsilon'' < \varepsilon,$$

contradicting the choice of c and c' .

CASE 2: $c, c' \notin A$. Since A is ω - \mathbf{d}^p -saturated, there is $d' \in A$ such that $\mathbf{d}^p(t^g(c'/A'), t^g(d'/A')) \leq \delta'''$, and again by non- δ''' -splitting of $t^g(B'/Aa)$ we may choose some $g \in \text{Aut}_{\delta''}$ with $d(g(B'c'A'), B'd'A') \leq \delta''$. Then $\mathbf{d}^p(t^g(g \circ f(b)/B'), t^g(b/B')) \leq \delta'$. So by ε'' -isolation, $\mathbf{d}^p(t^g(g \circ f(b)/B), t^g(b/B)) \leq \varepsilon''$. Also $\mathbf{d}^p(t^g(b/B), t^g(g(b)/B)) \leq \varepsilon''$ (as $\mathbf{d}^p(t^g(b/B'), t^g(g(b)/B')) \leq \delta'$). Hence $\mathbf{d}^p(t^g(g \circ f(b)/B), t^g(g(b)/B)) \leq \varepsilon'$, so letting $h \in \text{Aut}_{\varepsilon'}$ witness this we have $g^{-1} \circ h \circ g \circ f \in \text{Aut}_{2\varepsilon'}$ and

$$\begin{aligned} d(g^{-1} \circ h \circ g \circ f(bcA'), bc'A') \\ \leq d(g^{-1} \circ h \circ g \circ f(bcA'), g^{-1} \circ h \circ g(f(b)c'A')) \\ + d(g^{-1} \circ h \circ g(f(b)c'A'), g^{-1} \circ h(g \circ f(b)d'A')) \\ + d(g^{-1} \circ h(g \circ f(b)d'A'), g^{-1}(g(b)d'A')) \\ + d(g^{-1}(g(b)d'A'), bc'A') \\ \leq \varepsilon \end{aligned}$$

and thus $\mathbf{d}^p(t^g(bc/A'), t^g(bc'/A')) \leq \varepsilon$, again contradicting the choice of c and c' . ■

We now return to the question about the difficulties in proving the existence of prime models for the class of ω - \mathbf{d}^p -saturated models.

In classical model theory results like this are proved in the following way: Suppose $*$ -saturation is some notion of saturation and F is some reasonable isolation notion. To find prime models for the class of $*$ -saturated models one needs to prove the following two properties:

- (1) Every F -isolated type over A can be extended to an F -isolated type over $B \supset A$.
- (2) \mathcal{A} is $*$ -saturated if and only if \mathcal{A} is F -saturated, i.e. every F -isolated type over \mathcal{A} is realized in \mathcal{A} .

In our context we can prove \mathbf{d}^p -versions of (1) and (2) (see remark below) but we do not seem to be able to show that the properties imply the existence of prime models in any reasonable sense for the class of ω - \mathbf{d}^p -saturated models. Note that if \mathbf{d}^p is replaced by \mathbf{d} , the standard metric on types defined as the infimum of distances of realizations, this still works, as seen in [11].

REMARK 8.10. Assume \mathbb{K} is ω - \mathbf{d}^p -stable and has complete type-spaces.

- (1) If $t^g(a/A)$ is isolated and $B \supseteq A$ then for all $\varepsilon > 0$ there is b such that $t^g(b/B)$ is isolated and $\mathbf{d}^p(t^g(b/A), t^g(a/A)) \leq \varepsilon$.

- (2) If A is $\omega\text{-}\mathbf{d}^p$ -saturated and complete and $t^g(a/A)$ is isolated then $a \in A$.
- (3) If A realizes every isolated type over A then A is $\omega\text{-}\mathbf{d}^p$ -saturated and complete.

Proof. (1) and (3) are immediate consequences of Theorem 8.3. For (2) prove that the distance of a to the set A is less than any given ε , by using $\omega\text{-}\mathbf{d}^p$ -saturation over a suitably isolating set. ■

The problem here lies in the first item which forces us to “switch type” as we want to extend the isolation. Note that without this type-switching the item does not hold: otherwise $\omega\text{-}\mathbf{d}^p$ -saturated models would be ω -saturated. But considering the class of L^p -spaces with $\mathbb{F}_\varepsilon = \mathbb{F}_0$ for all ε , this would contradict Ben Yaacov’s and Usvyatsov’s observation in [6] that $L^p[0, 1]$ is not ω -saturated.

We finish this paper by a result exemplifying the kind of constructions made available by the theory developed in this paper. The proof is similar to the first order proof that nonunidimensional theories are not categorical. It is worth noting that the class \mathbb{K} in Lemma 8.12 may be unsuperstable in the ordinary metric sense and stability alone does not allow one to make constructions like the one in the lemma.

As an application of the lemma, one can prove that the class \mathbb{K}_H from Section 3 has the maximal number of models in density character ω_1 —a result that, however, has been known to functional analysts for almost a century. For this choose \mathcal{A} to be any existentially closed model and a_i for $i < \omega_1$ to be any nonzero element such that a_i is orthogonal to \mathcal{A} in the sense of Hilbert spaces and $\tau a_i = q_i a_i$, where $q_i, i < \omega_1$, are distinct complex numbers with norm 1.

DEFINITION 8.11. We say that $t^g(a/\emptyset)$ is *orthogonal* to $t^g(b/\emptyset)$ if for all $\omega\text{-}\mathbf{d}^p$ -saturated \mathcal{A} , a' and b' , if $t^g(a'/\emptyset) = t^g(a/\emptyset)$, $t^g(b'/\emptyset) = t^g(b/\emptyset)$, $a' \downarrow_\emptyset^s \mathcal{A}$ and $b' \downarrow_\emptyset^s \mathcal{A}$, then $a' \downarrow_\mathcal{A}^s b'$.

LEMMA 8.12. Suppose \mathbb{K} is $\omega\text{-}\mathbf{d}^p$ -stable, has complete type-spaces and every complete $\omega\text{-}\mathbf{d}^p$ -saturated set is a model. Assume that there are $\omega\text{-}\mathbf{d}^p$ -saturated \mathcal{A} and $a_i \notin \mathcal{A}$, $i < \omega_1$, such that $a_i \downarrow_\emptyset^s \mathcal{A}$ and for $i < j$, $t^g(a_i/\emptyset)$ is orthogonal to $t^g(a_j/\emptyset)$. Then the number of models in density character ω_1 is 2^{ω_1} .

Proof. Clearly we may assume that \mathcal{A} is separable. Let X_i , $i < 2^{\omega_1}$, be distinct nonempty subsets of ω_1 . For all $i < 2^\omega$ we construct $\mathcal{A}_i = \bigcup_{j < \omega_1} \mathcal{A}_i^j$ so that:

- (1) $\mathcal{A}_i^0 = \mathcal{A}$,
- (2) for limits j , \mathcal{A}_i^j is the completion of $\bigcup_{k < j} \mathcal{A}_i^k$,

- (3) \mathcal{A}_i^{j+1} is a separable, constructible, complete, ω - \mathbf{d}^p -saturated set over $a_i^j \mathcal{A}_i^j$, where a_i^j is such that $t^g(a_i^j / \mathcal{A}) = t^g(a_k / \mathcal{A})$ for some $k \in X_i$ and $a_i^j \downarrow_{\mathcal{A}}^s \mathcal{A}_i^j$,
- (4) for all $k \in X_i$, $|\{j < \omega_1 : t^g(a_i^j / \mathcal{A}) = t^g(a_k / \mathcal{A})\}| = \omega_1$.

By the proof of 8.6, it is easy to see that such models exist.

So it remains to prove that if $i < i'$, then \mathcal{A}_i is not isomorphic to $\mathcal{A}_{i'}$. For a contradiction suppose f is an isomorphism from \mathcal{A}_i to $\mathcal{A}_{i'}$. Without loss of generality we may assume that there is $k \in X_i - X_{i'}$. Clearly there is $j < \omega_1$ such that $f(\mathcal{A}_i^j) = \mathcal{A}_{i'}^j$. Now choose $j' \geq j$ so that $t^g(a_i^{j'} / \mathcal{A}) = t^g(a_k / \mathcal{A})$. An easy induction on $j' \leq j'' < \omega_1$ using dominance shows that $f(a_i^{j'}) \downarrow_{\emptyset}^s \mathcal{A}_{i'}^{j''}$ for all $j'' < \omega_1$. This contradicts the fact that $f(a_i^{j'}) \in \mathcal{A}_{i'}$. ■

Acknowledgements. The authors would like to thank Hans-Olav Tylli for many helpful discussions on Hilbert spaces and their operators.

Å. Hirvonen was supported by the Graduate School in Mathematical Logic and Algebra (MALJA) and the Finnish Academy of Science and Letters (Vilho, Yrjö and Kalle Väisälä foundation). T. Hyttinen was partially supported by the Academy of Finland, grant 1123110.

References

- [1] I. Ben Yaacov, *On perturbations of continuous structures*, J. Math. Logic 8 (2008), 225–249.
- [2] I. Ben Yaacov, *Topometric spaces and perturbations of metric structures*, Logic Anal. 1 (2008), 235–272.
- [3] I. Ben Yaacov, *Modular functionals and perturbations of Nakano spaces*, J. Logic Anal. 1 (2009), paper 1, 42 pp.
- [4] I. Ben Yaacov and A. Berenstein, *On perturbations of Hilbert spaces and probability algebras with a generic automorphism*, ibid., paper 7, 18 pp.
- [5] I. Ben Yaacov, A. Berenstein, C. W. Henson, and A. Usvyatsov, *Model theory for metric structures*, in: Model Theory with Applications to Algebra and Analysis, Vol. II, Z. Chatzidakis et al. (eds.), London Math. Soc. Lecture Note Ser. 350, Cambridge Univ. Press, Cambridge, 2008, 315–427.
- [6] I. Ben Yaacov and A. Usvyatsov, *On d-finiteness in continuous structures*, Fund. Math. 194 (2007), 67–88.
- [7] I. Ben Yaacov, A. Usvyatsov, and M. Zadka, *Generic automorphism of a Hilbert space*, preprint.
- [8] J. M. F. Castillo and M. González, *Three-Space Problems in Banach Space Theory*, Lecture Notes in Math. 1667, Springer, Berlin, 1997.
- [9] L. Diening, P. Harjulehto, P. Hästö, and M. Růžička, *Lebesgue and Sobolev spaces with variable exponents*, book manuscript, parts available at http://www.helsinki.fi/~pharjule/varsof/pdf/kirja_1_2_3_5_7_11.pdf.
- [10] C. W. Henson and J. Iovino, *Ultraproducts in analysis*, in: Analysis and Logic (Mons, 1997), London Math. Soc. Lecture Note Ser. 262, Cambridge Univ. Press, Cambridge, 2002, 1–110.

- [11] Å. Hirvonen and T. Hyttinen, *Categoricity in homogeneous complete metric spaces*, Arch. Math. Logic 48 (2009), 269–322.
- [12] T. Hyttinen and S. Shelah, *Strong splitting in stable homogeneous models*, Ann. Pure Appl. Logic 103 (2000), 201–228.
- [13] L. P. Poitevin, *Model theory of Nakano spaces*, PhD thesis, Univ. of Illinois at Urbana-Champaign, 2006; <http://www.salemstate.edu/~lpoitevin/thesis.pdf>.
- [14] S. Shelah, *Classification Theory and the Number of Nonisomorphic Models*, 2nd ed., Stud. Logic Found. Math. 92, North-Holland, Amsterdam, 1990.
- [15] S. Shelah and A. Usvyatsov, *Model theoretic stability and categoricity for complete metric spaces*, Israel J. Math. 182 (2011), 157–198.
- [16] S. Willard, *General Topology*, Dover Publ., Mineola, NY, 2004 (reprint of the 1970 original).

Åsa Hirvonen, Tapani Hyttinen
Department of Mathematics and Statistics
University of Helsinki
P.O. Box 68
00014 University of Helsinki, Finland
E-mail: asa.hirvonen@helsinki.fi
tapani.hyttinen@helsinki.fi

*Received 15 June 2010;
in revised form 16 May 2011 and 14 March 2012*