

The equivariant local ε -constant conjecture for unramified twists of $\mathbb{Z}_p(1)$

by

WERNER BLEY and ALESSANDRO COBBE (München)

Contents

1. Introduction	314
2. K-theory, determinants and refined Euler characteristics	317
3. The twisted local epsilon constant conjecture	319
3.1. The conjecture $C_{\text{EP}}^{\text{na}}(N/K, V)$	320
3.2. $C_{\text{EP}}^{\text{na}}(N/K, V)$ for unramified twists of $\mathbb{Z}_p(1)$	322
3.3. Organization of the paper	326
4. A representative for $RI(N, T)$	326
4.1. Some preliminary results	326
4.2. The construction of the complex	329
4.3. From $RI(N, \mathcal{F})$ to $RI(N, T)$	337
5. The term U_{cris}	341
5.1. The computation of U_{cris}	341
5.2. U_{cris} in the case $\chi^{\text{nr}} _{G_N} \neq 1$	343
5.3. U_{cris} in the case $\chi^{\text{nr}} _{G_N} = 1$	345
6. Epsilon constants of Weil–Deligne representations	352
6.1. Epsilon constants and Galois Gauß sums	352
6.2. Epsilon constants for unramified twists	354
7. The computation of the cohomological term	355
7.1. Some general results about $C_{N/K}$	355
7.2. The computation of $C_{N/K}$	360
7.3. The Euler characteristic when $\omega = 0$	367
7.4. The Euler characteristic when $\omega > 0$	369
8. Proof of Theorem 1	377
Appendix: A more detailed computation of U_{cris}	380
References	381

2010 *Mathematics Subject Classification*: Primary 11S23; Secondary 11S25.

Key words and phrases: local epsilon constant conjectures, equivariant Tamagawa number conjecture, weak ramification.

Received 7 July 2016; revised 20 October 2016.

Published online 26 April 2017.

1. Introduction. Let F/E denote a Galois extension of number fields and let $\Gamma := \text{Gal}(F/E)$. We fix a motive M which is defined over E and admits an action of the group algebra $\mathbb{Q}[\Gamma]$. Let $L(M, s)$ denote the equivariant complex L -function attached to M and F/E (see [BF01, Sec. 4]). We shall always assume that this L -function satisfies a functional equation of the following form relating the L -functions attached to M and its Kummer dual $M^*(1)$:

$$L(M, s) = \varepsilon(M, s) \frac{L_\infty(M^*(1), -s)}{L_\infty(M, s)} L(M^*(1), -s).$$

Here L_∞ is an Euler factor at infinity and the equivariant ε -factor $\varepsilon(M, s)$ decomposes into local factors:

$$\varepsilon(M, s) = \prod_v \varepsilon_v(M, s)$$

with v running through the set of finite places of E . For some more details see [BF01, Conj. 7].

In this paper we are interested in a local equivariant ε -constant conjecture which relates $\varepsilon_v(M, s)$ in a specific way to certain local cohomology groups of $RG(G_{E_v}, M_p)$ associated to the p -adic representation $V = M_p$. The validity of this ε -constant conjecture for all finite places v of E is closely related to the compatibility of the equivariant Tamagawa number conjectures for M and its Kummer dual $M^*(1)$ with the functional equation. Indeed, [BF01, Conj. 8] is a semilocal analogue of the local ε -constant conjecture and is directly related to the functional equation compatibility of the equivariant Tamagawa number conjecture (see [BF01, Cor. 1]).

So henceforth we fix a prime number p and let N/K denote a finite Galois extension of p -adic number fields with group $G := \text{Gal}(N/K)$. We write G_K (resp. G_N) for the absolute Galois group of K (resp. N) and let V denote a p -adic representation of G_K . Let $T \subseteq V$ be a G_K -stable \mathbb{Z}_p -sublattice such that $V = T \otimes_{\mathbb{Z}_p} \mathbb{Q}_p$.

Following seminal work of Fontaine and Perrin-Riou [FPR94] several authors have formulated ε -constant conjectures in this general context (see e.g. Benois and Berger [BB08] or Fukaya and Kato [FK06]). For the relations between the different formulations we refer the reader to [Izy12]. We will use the notation of [IV16, Conj. 5.1] and write $C_{\text{EP}}^{\text{na}}(N/K, V)$ for the equivariant local ε -constant conjecture. Here the subscript “EP” is short for Euler–Poincaré and refers to the fact that the conjecture is formulated in terms of certain Euler–Poincaré characteristics. The superscript “na” was added by Izychev and Venjakob for their non-abelian generalization. We will recall and formulate the conjecture in Section 3.

In a more specialized situation, Breuning [Bre04b] has formulated an ε -constant conjecture for $T = \mathbb{Z}_p(1)$ in terms of relative algebraic K -groups.

Extending ideas of [BB03] he proved his conjecture, amongst other cases, for tamely ramified extensions N/K . Breuning’s conjecture clearly has to be equivalent to $C_{\text{EP}}^{\text{na}}(N/K, V)$ for $V = \mathbb{Q}_p(1)$ (see e.g. [IV16, Appendix]). As a by-product of our work we will obtain a rigorous proof of this assertion.

Izychev and Venjakob [IV16] consider the case where $T = \mathbb{Z}_p(\chi^{\text{nr}})(1)$ is a one-dimensional unramified twist of $\mathbb{Z}_p(1)$. More precisely, if χ^{nr} is the restriction of an unramified character $\chi_{\mathbb{Q}_p}^{\text{nr}} : G_{\mathbb{Q}_p} \rightarrow \mathbb{Z}_p^\times$ with $\chi^{\text{nr}}|_{G_N} \neq 1$, they reformulate $C_{\text{EP}}^{\text{na}}(N/K, V)$ in the spirit of Breuning and adapt his proof of the tame case to show the validity of $C_{\text{EP}}^{\text{na}}(N/K, V)$ for tamely ramified extensions N/K and $V = \mathbb{Q}_p(\chi^{\text{nr}})(1)$. In [IV16, Appendix] they also indicate how to adapt Breuning’s arguments to prove the tame case for unramified characters as above with $\chi^{\text{nr}}|_{G_N} = 1$.

We recall that Benois and Berger [BB08, Th. 4.22] have proved $C_{\text{EP}}^{\text{na}}(N/K, V)$ for arbitrary crystalline representations V of G_K where K/\mathbb{Q}_p is unramified and N is a finite subfield of $K(\mu_{p^\infty})/K$. Here μ_{p^∞} denotes the group of all p -power roots of unity.

In this paper we will focus on weakly and wildly ramified extensions N/K of an unramified extension K/\mathbb{Q}_p . We recall that N/K is *weakly ramified* if the second ramification group in lower numbering is trivial. As above, we let χ^{nr} be an unramified character of G_K which is the restriction of an unramified character $\chi_{\mathbb{Q}_p}^{\text{nr}} : G_{\mathbb{Q}_p} \rightarrow \mathbb{Z}_p^\times$. We require no assumption on the restriction of χ^{nr} on G_N . The main result of our work is as follows.

THEOREM 1. *Let p be an odd prime and let K/\mathbb{Q}_p be the unramified extension of degree m . Let N/K be a weakly and wildly ramified finite abelian extension with cyclic ramification group. Let d denote the inertia degree of N/K and assume that m and d are relatively prime. Then the conjecture $C_{\text{EP}}^{\text{na}}(N/K, V)$ is true for N/K and $V = \mathbb{Q}_p(\chi^{\text{nr}})(1)$.*

REMARK 1. The assumptions of the theorem imply that the ramification group is cyclic of order p (cf. [PV13, Cor. 3.4]). More precisely, $|G| = pd$, $|G_0| = |G_1| = p$ and $|G_i| = 1$ for $i \geq 2$. Here G_i for $i \geq 0$ denotes a higher ramification subgroup.

REMARK 2. The proof of Theorem 1 is an adaptation of the methods in [BC16] where we prove Breuning’s conjecture for $T = \mathbb{Z}_p(1)$ and for extensions N/K as in the theorem. If $\chi^{\text{nr}}|_{G_N} = 1$, it is almost straightforward to extend this result to $T = \mathbb{Z}_p(\chi^{\text{nr}})(1)$ because then twisting commutes with taking G_N -cohomology.

REMARK 3. If $\chi^{\text{nr}}|_{G_N} \neq 1$, the proof of the above weakly and wildly ramified case is much more involved than the tame case of Izychev and Venjakob because of the following fact. In the tame case the cohomology groups of $R\Gamma(N, T)$ turn out to be perfect (see [IV16, Prop. 2.5]), whereas

this is in general no longer true in our weakly and wildly ramified case. This makes the computation of a certain refined Euler characteristic much more technically difficult.

We recall that by [Izy12, Ex. 5.20] the representation $T = \mathbb{Z}_p(\chi^{\text{nr}})(1)$ is naturally isomorphic to the p -adic Tate module $T_p\mathcal{F}$ of a one-dimensional Lubin–Tate formal group \mathcal{F} defined over the ring \mathbb{Z}_p . Conversely, every one-dimensional Lubin–Tate formal group \mathcal{F} defined over \mathbb{Z}_p gives rise to an unramified twist of $\mathbb{Z}_p(1)$ as above.

We let K_{nr} denote the maximal unramified extension of K and put $K_1 := K_{\text{nr}} \cap N$. We write N_{nr} for the maximal unramified extension of N . Let N_0 denote the completion of $N_{\text{nr}} = NK_{\text{nr}}$ and write $\widehat{N_0^\times}$ for the p -completion of N_0^\times . We set

$$\mathcal{I}_{N/K}(\chi^{\text{nr}}) := \text{Ind}_{\text{Gal}(N_{\text{nr}}/K_1)}^{\text{Gal}(N_{\text{nr}}/K)}(\widehat{N_0^\times}(\chi^{\text{nr}}))$$

and usually identify $\mathcal{I}_{N/K}(\chi^{\text{nr}})$ with $\prod_{i=1}^{d_{N/K}} \widehat{N_0^\times}$, where $d_{N/K}$ denotes the inertia degree of N/K . We will endow $\mathcal{I}_{N/K}(\chi^{\text{nr}})$ with a natural action of $\text{Gal}(K_{\text{nr}}/K) \times G$. If $\chi^{\text{nr}}|_{G_N} \neq 1$ we set $\omega_N := v_p(1 - \chi^{\text{nr}}(F_N))$ where F_N denotes the Frobenius of N . Let $F = F_K$ be the Frobenius of K .

In order to deal with the difficulties indicated in Remark 3 above we construct an exact sequence

$$(1) \quad 0 \rightarrow \mathcal{F}(\mathfrak{p}_N) \rightarrow \mathcal{I}_{N/K}(\chi^{\text{nr}}) \xrightarrow{(F-1) \times 1} \mathcal{I}_{N/K}(\chi^{\text{nr}}) \rightarrow \mathbb{Z}/p^{\omega_N}\mathbb{Z}(\chi^{\text{nr}}) \rightarrow 0$$

and show the following theorem, which may also be of independent interest.

THEOREM 2. *Let N/K be a finite Galois extension of p -adic fields. Let $\chi^{\text{nr}}: G_K \rightarrow \mathbb{Z}_p^\times$ be an unramified character as above and assume that $\chi^{\text{nr}}|_{G_N} \neq 1$. Then the complex*

$$C_{N,T}^\bullet := [\mathcal{I}_{N/K}(\chi^{\text{nr}}) \xrightarrow{(F-1) \times 1} \mathcal{I}_{N/K}(\chi^{\text{nr}})]$$

with non-trivial modules in degree 1 and 2 represents $R\Gamma(N, T)$.

REMARK 4. The exact sequence (1) should be seen as an analogue of the fundamental short exact sequence of [Ser79, Sec. XIII.5, Exercises] which by [Bre04a, Th. 4.20] is essentially a representative of $R\Gamma(G, \mathbb{Z}_p(1))$. Note that in the case $\chi^{\text{nr}}|_{G_N} \neq 1$ it does not seem to be possible to derive Theorem 2 from Serre’s result by a simple twisting argument.

In Section 2 we recall some preliminary results on K -theory, determinant functors and refined Euler characteristics which we will need throughout. After these preparations we formulate in Section 3 the conjecture $C_{\text{EP}}^{\text{na}}(N/K, V)$ following the exposition of [IV16] and then give a short description of the organization of the paper.

Notation. We fix some standard notation which will be used throughout. Given a finite field extension M/L , we will denote the norm and the trace by $\mathcal{N}_{M/L}$ and $\mathcal{T}_{M/L}$ respectively. If L/\mathbb{Q}_p is a finite extension we write L^c for the algebraic closure and set $G_L := \text{Gal}(L^c/L)$. We let L_{nr} denote the maximal unramified extension of L and write F_L for the arithmetic Frobenius of L . Note that F_L is a topological generator of $\text{Gal}(L_{\text{nr}}/L)$. As usual, we let \mathcal{O}_L denote the valuation ring of L and write $\mathfrak{p}_L = \pi_L \mathcal{O}_L$ for the maximal ideal.

We write d_L for the absolute inertia degree and e_L for the absolute ramification degree. If M/L is a finite extension we let $d_{M/L}$ and $e_{M/L}$ denote the relative inertia degree and ramification degree, respectively. Then we have $d_M = d_{M/L}d_L$, $e_M = e_{M/L}e_L$ and $F_M = F_L^{d_{M/L}}$. We will denote the inertia group by $I_{M/L}$.

We let L_0 be the completion of L_{nr} with respect to the p -adic topology. If A is an abelian group we denote the p -completion by \widehat{A} , i.e.,

$$\widehat{A} = \varprojlim_n A/p^n A.$$

Note that \widehat{A} is in a natural way a \mathbb{Z}_p -module. We will denote by

$$\nu_L: \widehat{L^\times} \otimes_{\mathbb{Z}_p} \mathbb{Q}_p \rightarrow \mathbb{Q}_p$$

the map induced by the normalized valuation of L^\times .

If Σ is any ring we denote its centre by $Z(\Sigma)$.

2. K-theory, determinants and refined Euler characteristics. Let A be any unital ring. Let $\text{PMod}(A)$ denote the category of finitely generated projective A -modules and write $\text{PMod}(A)^\bullet$ for the category of bounded complexes of such modules. We also write $D(A)$ for the derived category of complexes of A -modules and $D^p(A)$ for the full triangulated subcategory of $D(A)$ of perfect complexes. We recall that a complex C^\bullet of A -modules is *perfect* if there exists a complex $P^\bullet \in \text{PMod}(A)^\bullet$ and a quasi-isomorphism $P^\bullet \rightarrow C^\bullet$. We say that a A -module N is *perfect*, if the complex $N[0]$ belongs to $D^p(A)$.

Our main references for the theory of Picard categories, determinant functors and virtual objects are [BF01, Sec. 2] and [BB05]. For a more explicit approach we refer the reader to [FK06, Sec. 1]. By [FK06, Remark 1.2.10] both approaches should be equivalent.

Let $V(A)$ denote the Picard category of virtual objects associated to $\text{PMod}(A)$ and write $[\cdot]_A$ for the universal determinant functor

$$[\cdot]_A: (\text{PMod}(A), \text{is}) \rightarrow V(A),$$

where $(\text{PMod}(A), \text{is})$ denotes the subcategory of isomorphisms in $\text{PMod}(A)$.

By [BF01, Prop. 2.1] this functor extends to a functor

$$[\cdot]_\Lambda: (D^p(\Lambda), \text{is}) \rightarrow V(\Lambda).$$

We recall that the category $V(\Lambda)$ is equipped with a canonical bifunctor $(L, M) \mapsto L \otimes M$. We fix a unit object $\mathbf{1}_{V(\Lambda)}$ and for each object L an inverse L^{-1} with an isomorphism $L \otimes L^{-1} \cong \mathbf{1}_{V(\Lambda)}$. Each element of $V(\Lambda)$ is of the form $[P]_\Lambda \otimes [Q]_\Lambda^{-1}$ for modules $P, Q \in \text{PMod}(\Lambda)$. Furthermore, $[P]_\Lambda$ and $[Q]_\Lambda$ are isomorphic in $V(\Lambda)$ if and only if the classes of P and Q in $K_0(\Lambda)$ coincide.

For any Picard category \mathcal{P} we define $\pi_0(\mathcal{P})$ to be the group of isomorphism classes of objects of \mathcal{P} and set $\pi_1(\mathcal{P}) := \text{Aut}_{\mathcal{P}}(\mathbf{1}_{\mathcal{P}})$. The groups $\pi_0(V(\Lambda))$ and $\pi_1(V(\Lambda))$ are naturally isomorphic to $K_0(\Lambda)$ and $K_1(\Lambda)$, respectively.

In what follows we fix rings Λ and Σ and a ring homomorphism $\Lambda \rightarrow \Sigma$ such that Σ is flat as a right Λ -module. We assume that Σ is noetherian and *regular*, that is, every finitely generated Σ -module has finite projective dimension. We write $P_\Sigma = \Sigma \otimes_\Lambda P$ for the scalar extension map if P is a Λ -module or a complex of Λ -modules. By abuse of notation we will write $[P]_\Sigma$ in place of $[\Sigma \otimes_\Lambda P]_\Sigma$. We recall that there is a canonical exact sequence of algebraic K -groups

$$K_1(\Lambda) \rightarrow K_1(\Sigma) \xrightarrow{\partial_{\Lambda, \Sigma}^1} K_0(\Lambda, \Sigma) \xrightarrow{\partial_{\Lambda, \Sigma}^0} K_0(\Lambda) \rightarrow K_0(\Sigma),$$

where $K_0(\Lambda, \Sigma)$ is the relative algebraic K -group defined by Swan [Swa70, p. 215]. We also recall that elements in $K_0(\Lambda, \Sigma)$ are represented by triples $[P, g, Q]$ where $P, Q \in \text{PMod}(\Lambda)$ and $g: P_\Sigma \rightarrow Q_\Sigma$ is an isomorphism of Σ -modules.

If $\Sigma = L[G]$ for a finite group G and a finite field extension L/\mathbb{Q}_p , the reduced norm map induces an isomorphism

$$\text{Nrd}_\Sigma: K_1(\Sigma) \rightarrow Z(\Sigma)^\times$$

by [CR87, Th. (45.3)]. The same is true for algebraically closed fields L , in particular for $L = \mathbb{Q}_p^c$. We set $\hat{\partial}_{\Lambda, \Sigma}^1 := \partial_{\Lambda, \Sigma}^1 \circ \text{Nrd}_\Sigma^{-1}$.

The scalar extension functor $\text{PMod}(\Lambda) \rightarrow \text{PMod}(\Sigma)$ extends to a monoidal functor $V(\Lambda) \rightarrow V(\Sigma)$ and we again write $L \mapsto L_\Sigma$ for $L \in V(\Lambda)$. Still following closely the exposition of [BB05] we define a tensor category $V(\Lambda, \Sigma)$ as follows. Objects in $V(\Lambda, \Sigma)$ are pairs (L, λ) with $L \in V(\Lambda)$ and λ an isomorphism $L_\Sigma \rightarrow \mathbf{1}_{V(\Sigma)}$ in $V(\Sigma)$. A morphism $(L, \lambda) \rightarrow (M, \mu)$ is an isomorphism $\alpha: L \rightarrow M$ in $V(\Lambda)$ such that $\mu \circ \alpha_\Sigma = \lambda$. The product of (L, λ) and (M, μ) is $(L \otimes M, \nu)$ where $\nu: (L \otimes M)_\Sigma \rightarrow \mathbf{1}_{V(\Sigma)}$ is the isomorphism

$$(L \otimes M)_\Sigma \rightarrow L_\Sigma \otimes M_\Sigma \xrightarrow{\lambda \otimes \mu} \mathbf{1}_{V(\Sigma)} \otimes \mathbf{1}_{V(\Sigma)} \rightarrow \mathbf{1}_{V(\Sigma)}.$$

By [BB05, Lemma 5.1] the category $V(\Lambda, \Sigma)$ is a Picard category and there is a natural isomorphism

$$(2) \quad i_{\Lambda, \Sigma}: \pi_0(V(\Lambda, \Sigma)) \cong K_0(\Lambda, \Sigma).$$

For a complex $P \in \text{PMod}(\Lambda)^\bullet$ we define objects P^{ev} and P^{od} in $\text{PMod}(\Lambda)$ by

$$P^{\text{ev}} := \bigoplus_{i \text{ even}} P^i, \quad P^{\text{od}} := \bigoplus_{i \text{ odd}} P^i.$$

Similarly we define

$$H^{\text{ev}}(P) := \bigoplus_{i \text{ even}} H^i(P), \quad H^{\text{od}}(P) := \bigoplus_{i \text{ odd}} H^i(P), \quad H^{\text{all}}(P) := \bigoplus_{i \in \mathbb{Z}} H^i(P).$$

Let $P \in D^p(\Lambda)$ and let $t: H^{\text{ev}}(P_\Sigma) \rightarrow H^{\text{od}}(P_\Sigma)$ be an isomorphism of Σ -modules. We refer to t as a *trivialization* and to the pair (P, t) as a *trivialized complex*.

By [BB05, Def. 5.5] there is a (refined) Euler characteristic $\chi_{\Lambda, \Sigma}(P, t) \in K_0(\Lambda, \Sigma)$. We briefly recall the construction. The element $\chi_{\Lambda, \Sigma}(P, t)$ is defined by

$$\chi_{\Lambda, \Sigma}(P, t) = i_{\Lambda, \Sigma}(\left([U]_\Lambda, \kappa(P, t)\right)),$$

where $([U]_\Lambda, \kappa(P, t)) \in V(\Lambda, \Sigma)$ is defined as follows. Let $a: U \rightarrow P$ be a quasi-isomorphism with $U \in \text{PMod}(\Lambda)^\bullet$ and let $\kappa(P, t): [U]_\Sigma \rightarrow \mathbf{1}_{V(\Sigma)}$ be the isomorphism in $V(\Sigma)$ given by the composite

$$(3) \quad [U]_\Sigma \xrightarrow{\eta_{U_\Sigma}} [H(U_\Sigma)]_\Sigma \xrightarrow{H(a_\Sigma)} [H(P_\Sigma)]_\Sigma \\ \xrightarrow{\pi_{H(P_\Sigma)}} [H^{\text{ev}}(P_\Sigma)[0]]_\Sigma \otimes [H^{\text{od}}(P_\Sigma)[1]]_\Sigma \\ \xrightarrow{[t] \otimes \text{id}} [H^{\text{od}}(P_\Sigma)[0]]_\Sigma \otimes [H^{\text{od}}(P_\Sigma)[1]]_\Sigma \xrightarrow{\mu_{H^{\text{od}}(P_\Sigma)[0]}} \mathbf{1}_{V(\Sigma)}.$$

Here η_{U_Σ} is defined in [BB05, Prop. 3.1], $\pi_{H(P_\Sigma)}$ by [BB05, Prop. 4.4] and $\mu_{H^{\text{od}}(P_\Sigma)[0]}$ by [BB05, Lemma 2.3].

For computational applications it is often convenient to use an explicit construction of a refined Euler characteristic due to Burns [Bur04]. We still consider a ring homomorphism $\Lambda \rightarrow \Sigma$ as above, but now assume in addition that Σ is semisimple. We fix an object $P \in D^p(\Lambda)$ and a trivialization $t: H^{\text{ev}}(P_\Sigma) \rightarrow H^{\text{od}}(P_\Sigma)$. Following [BB05] we write $\chi_{\Sigma, \Lambda}^{\text{old}}(P, t^{-1}) \in K_0(\Lambda, \Sigma)$ for the refined Euler characteristic defined by Burns in [Bur04]. If we write $B^i(P_\Sigma) \subseteq P_\Sigma^i$ for the boundaries of P_Σ and set $B^{\text{od}}(P_\Sigma) := \bigoplus_{i \text{ odd}} B^i(P_\Sigma)$, then by [BB05, Th. 6.2] we have

$$(4) \quad -\chi_{\Lambda, \Sigma}^{\text{old}}(P, t^{-1}) = \chi_{\Lambda, \Sigma}(P, t) + \partial_{\Lambda, \Sigma}^1(B^{\text{od}}(P_\Sigma), -\text{id}).$$

3. The twisted local epsilon constant conjecture. In this section we formulate $C_{\text{EP}}^{\text{na}}(N/K, V)$ following the exposition of [IV16]. We recall the

necessary notation and results to the extent that we can relate $C_{\text{EP}}^{\text{na}}(N/K, V)$ to Breuning’s conjecture. For further details we refer the reader to [BB08] and [IV16].

3.1. The conjecture $C_{\text{EP}}^{\text{na}}(N/K, V)$. We will briefly recall the definition of local ε -constants following the expositions of [BB08] and [IV16].

Let E denote a large enough field of characteristic 0 such that E contains the p -power roots of unity μ_{p^∞} and the roots of unity of order $p - 1$. Let K/\mathbb{Q}_p be a finite extension. We write μ_K for the unique Haar measure on K normalized by $\mu_K(\mathcal{O}_K) = 1$. Let $\psi: K \rightarrow E^\times$ denote an additive continuous character.

We recall that the theory of Langlands and Deligne (cf. [Del73]) associates to each E -linear representation V of the Weil group of K a local constant $\varepsilon(V, \psi, \mu_K)$. The basic properties of these ε -constants are listed in [BB08, Sec. 2.3].

We fix a compatible system $\xi = (\xi_{p^n})_{n \geq 1}$ of p -power roots of unity and define an additive character $\psi_\xi: \mathbb{Q}_p \rightarrow E^\times$ by $\psi_\xi|_{\mathbb{Z}_p} = 1$ and $\psi_\xi(p^{-n}) = \xi_{p^n}$.

Let now N/K be a finite Galois extension with group G . We assume that E is large enough such that every irreducible character χ of G can be realized over E . For each character χ of G we fix an E -linear representation V_χ with character χ and write V_χ^* for the contragredient representation.

Following [IV16, Sec. 2.3] we define the element

$$\varepsilon_D(N/K, V) := \left(\varepsilon(D_{\text{pst}}(\text{Ind}_{K/\mathbb{Q}_p}(V \otimes_E V_\chi^*), \psi_\xi, \mu_{\mathbb{Q}_p})) \right)_{\chi \in \text{Irr}(G)}$$

in $\prod_{\chi \in \text{Irr}(G)} (\mathbb{Q}_p^c)^\times \cong Z(\mathbb{Q}_p^c[G])^\times$. For all unexplained notation we refer the reader to [IV16, Sec. 2.3].

We will apply the notation introduced and explained in [BB08, Sec. 1.1]. In particular, $B_{\text{cris}}, B_{\text{st}}$ and B_{dR} denote the p -adic period rings constructed by Fontaine. If V is a p -adic representation of G_K , we let

$$D_{\text{dR}}^K(V) := (B_{\text{dR}} \otimes_{\mathbb{Q}_p} V)^{G_K}, \quad D_{\text{cris}}^K(V) := (B_{\text{cris}} \otimes_{\mathbb{Q}_p} V)^{G_K}.$$

The K -vector space $D_{\text{dR}}^K(V)$ is finite-dimensional and filtered. The tangent space of V over K is defined by

$$t_V(K) := D_{\text{dR}}^K(V) / \text{Fil}^0 D_{\text{dR}}^K(V).$$

Finally, we write $\exp_V: t_V(K) \rightarrow H^1(K, V)$ for the exponential map of Bloch and Kato.

For any \mathbb{Q}_p -vector space W we write $W^* = \text{Hom}_{\mathbb{Q}_p}(W, \mathbb{Q}_p)$ for its \mathbb{Q}_p -linear dual. For convenience we usually write $t_V^*(K)$ instead of $t_V(K)^*$.

Let N/K be a finite Galois extension of p -adic fields with group G . We set $\Lambda := \mathbb{Z}_p[G], \Omega := \mathbb{Q}_p[G], \tilde{\Lambda} := \overline{\mathbb{Z}_p}^{\text{nr}}[G]$ and $\tilde{\Omega} := \overline{\mathbb{Q}_p}^{\text{nr}}[G]$, where $\overline{\mathbb{Z}_p}^{\text{nr}}$ denotes the ring of integers in the completion $\overline{\mathbb{Q}_p}^{\text{nr}}$ of \mathbb{Q}_p^{nr} .

We recall the fundamental 7-term exact sequence of $\mathbb{Q}_p[G]$ -modules of [BB08, (2.2)]:

$$(5) \quad 0 \rightarrow H^0(N, V) \rightarrow D_{\text{cris}}^N(V) \xrightarrow{1-\phi} D_{\text{cris}}^N(V) \oplus t_V(N) \xrightarrow{\text{exp}_V} H^1(N, V) \\ \xrightarrow{\text{exp}_V^*} D_{\text{cris}}^N(V^*(1))^* \oplus t_{V^*(1)}^*(N) \rightarrow D_{\text{cris}}^N(V^*(1))^* \rightarrow H^2(N, V) \rightarrow 0.$$

In the following we will freely use the properties of determinant functors as, for example, formulated in [BF01, Prop. 2.1]. As in [IV16, Sec. 3] we use

$$(6) \quad 0 \rightarrow t_{V^*(1)}^*(N) \rightarrow D_{\text{dR}}^N(V) \rightarrow t_V(N) \rightarrow 0,$$

and by [BF01, Prop. 2.1] we obtain an isomorphism

$$(7) \quad [D_{\text{dR}}^N(V)]_{\mathbb{Q}_p[G]} \otimes [R\Gamma(N, V)]_{\mathbb{Q}_p[G]} \\ \rightarrow [t_{V^*(1)}^*(N)]_{\mathbb{Q}_p[G]} \otimes [t_V(N)]_{\mathbb{Q}_p[G]} \otimes \bigotimes_i [H^i(N, V)]_{\mathbb{Q}_p[G]}^{(-1)^i}.$$

Using the commutativity constraint, we see that the right hand side of (7) is canonically isomorphic to

$$(8) \quad [H^0(N, V)]_{\mathbb{Q}_p[G]} \otimes [D_{\text{cris}}^N(V)]_{\mathbb{Q}_p[G]}^{-1} \otimes [D_{\text{cris}}^N(V)]_{\mathbb{Q}_p[G]} \otimes [t_V(N)]_{\mathbb{Q}_p[G]} \\ \otimes [H^1(N, V)]_{\mathbb{Q}_p[G]}^{-1} \otimes [t_{V^*(1)}^*(N)]_{\mathbb{Q}_p[G]} \otimes [D_{\text{cris}}^N(V^*(1))^*]_{\mathbb{Q}_p[G]} \\ \otimes [D_{\text{cris}}^N(V^*(1))^*]_{\mathbb{Q}_p[G]}^{-1} \otimes [H^2(N, V)]_{\mathbb{Q}_p[G]}.$$

Now the 7-term exact sequence (5) induces an isomorphism from the object in (8) to $\mathbf{1}_{V(\mathbb{Q}_p[G])}$, so that we obtain an isomorphism

$$(9) \quad \delta'(N/K, V): [D_{\text{dR}}^N(V)]_{\mathbb{Q}_p[G]} \otimes [R\Gamma(N, V)]_{\mathbb{Q}_p[G]} \rightarrow \mathbf{1}_{V(\mathbb{Q}_p[G])}.$$

We let

$$\Delta_{\text{EP}}^{\text{na}}(N/K, V) := [R\Gamma(N, V)]_{\mathbb{Q}_p[G]} \otimes [\text{Ind}_{N/\mathbb{Q}_p}(V)]_{\mathbb{Q}_p[G]} \in V(\mathbb{Q}_p[G]).$$

By [BB08, Lemme 2.13] we have an isomorphism

$$\psi: D_{\text{dR}}^N(V) \rightarrow D_{\text{dR}}^{\mathbb{Q}_p}(\text{Ind}_{N/\mathbb{Q}_p}(V)),$$

which induces a canonical comparison isomorphism

$$\text{comp}_V: B_{\text{dR}} \otimes_{\mathbb{Q}_p} D_{\text{dR}}^N(V) \rightarrow B_{\text{dR}} \otimes_{\mathbb{Q}_p} \text{Ind}_{N/\mathbb{Q}_p}(V), \quad c \otimes x \mapsto c\psi(x).$$

The morphism comp_V gives the isomorphism

$$\tilde{\alpha}(N/K, V): [D_{\text{dR}}^N(V)]_{B_{\text{dR}}[G]}^{-1} \otimes [\text{Ind}_{N/\mathbb{Q}_p}(V)]_{B_{\text{dR}}[G]} \rightarrow \mathbf{1}_{V(B_{\text{dR}}[G])}.$$

Following [BB08, Sec. 1.1] we define

$$h_i(V) := \dim_{\mathbb{Q}_p}(\text{Fil}^i D_{\text{dR}}^{\mathbb{Q}_p}(V) / \text{Fil}^{i+1} D_{\text{dR}}^{\mathbb{Q}_p}(V)) \quad \text{and} \quad t_H(V) := \sum_{i \in \mathbb{Z}} i h_i(V).$$

Furthermore, we set

$$\Gamma^*(V) := \prod_{i \in \mathbb{Z}} \Gamma^*(-i)^{h_i(V)}, \quad \text{where} \quad \Gamma^*(i) = \begin{cases} (i-1)! & \text{if } i > 0, \\ (-1)^i/(-i)! & \text{if } i \leq 0. \end{cases}$$

Multiplying $\tilde{\alpha}(N/K, V)$ by $t^{t_H(V)}$ where $t := \log[\xi] \in B_{\text{dR}}$ is the uniformizer of B_{dR}^+ associated with $\xi = (\xi_{p^n})_{n \geq 0}$, we get

$$(10) \quad \alpha(N/K, V) : [D_{\text{dR}}^N(V)]_{B_{\text{dR}}[G]}^{-1} \otimes [\text{Ind}_{N/\mathbb{Q}_p}(V)]_{B_{\text{dR}}[G]} \rightarrow \mathbf{1}_{V(B_{\text{dR}}[G])}.$$

Composing $\alpha(N/K, V)$ with the automorphism of $\mathbf{1}_{V(B_{\text{dR}}[G])}$ induced by the element $\Gamma^*(V)\varepsilon_D(N/K, V)$ we obtain

$$(11) \quad \beta(N/K, V) : [D_{\text{dR}}^N(V)]_{B_{\text{dR}}[G]}^{-1} \otimes [\text{Ind}_{N/\mathbb{Q}_p}(V)]_{B_{\text{dR}}[G]} \rightarrow \mathbf{1}_{V(B_{\text{dR}}[G])}.$$

We also define $\beta'(N/K, V)$ as the composite

$$(12) \quad \begin{aligned} \Delta_{\text{EP}}^{\text{na}}(N/K, V)_{B_{\text{dR}}[G]} &= [R\Gamma(N, V)]_{B_{\text{dR}}[G]} \otimes [\text{Ind}_{N/\mathbb{Q}_p}(V)]_{B_{\text{dR}}[G]} \\ &= [R\Gamma(N, V)]_{B_{\text{dR}}[G]} \otimes [D_{\text{dR}}^N(V)]_{B_{\text{dR}}[G]} \\ &\quad \otimes [D_{\text{dR}}^N(V)]_{B_{\text{dR}}[G]}^{-1} \otimes [\text{Ind}_{N/\mathbb{Q}_p}(V)]_{B_{\text{dR}}[G]} \\ &\xrightarrow{\delta'(N/K, V)} [D_{\text{dR}}^N(V)]_{B_{\text{dR}}[G]}^{-1} \otimes [\text{Ind}_{N/\mathbb{Q}_p}(V)]_{B_{\text{dR}}[G]} \xrightarrow{\alpha(N/K, V)} \mathbf{1}_{V(B_{\text{dR}}[G])}. \end{aligned}$$

Note that in order to apply $\delta'(N/K, V)$ in the above definition it is necessary to apply the commutativity constraint first.

The isomorphism $\delta(N/K, V)$ is defined in the same way, just writing $\beta(N/K, V)$ in place of $\alpha(N/K, V)$ in the above formula. Since according to [BB08, Lemme 2.15] the isomorphism $\beta(N/K, V)$ is actually defined over $\tilde{\Omega}$, the same is true for $\delta(N/K, V)$, so that we actually have an isomorphism

$$(13) \quad \delta(N/K, V) : \Delta_{\text{EP}}^{\text{na}}(N/K, V)_{\tilde{\Omega}} \rightarrow \mathbf{1}_{V(\tilde{\Omega})}.$$

We define

$$\Delta_{\text{EP}}^{\text{na}}(N/K, T) := [R\Gamma(N, T)]_{\mathbb{Z}_p[G]} \otimes [\text{Ind}_{N/\mathbb{Q}_p}(T)]_{\mathbb{Z}_p[G]} \in V(\mathbb{Z}_p[G]).$$

The element $\Delta_{\text{EP}}^{\text{na}}(N/K, T)$ actually does not depend on the choice of T and is therefore a canonical element in $V(\mathbb{Z}_p[G])$ whose base change gives $\Delta_{\text{EP}}^{\text{na}}(N/K, V)$. We can now finally formulate the epsilon constant conjecture (compare to [IV16, Conj. 5.1] or [BB08, Conj. 2.19]).

CONJECTURE 3.1.1. *If $V = \mathbb{Q}_p \otimes_{\mathbb{Z}_p} T$ is potentially semistable and N/K is a finite Galois extension, then the class $[(\Delta_{\text{EP}}^{\text{na}}(N/K, T), \delta(N/K, V))]$ $\in \pi_0(V(\mathbb{Z}_p[G], \tilde{\Omega}))$ is trivial in $\pi_0(V(\tilde{\Lambda}, \tilde{\Omega}))$.*

3.2. $C_{\text{EP}}^{\text{na}}(N/K, V)$ for unramified twists of $\mathbb{Z}_p(1)$. We now specialize to the situation considered in this paper.

Let p be a prime and let K be a finite extension of \mathbb{Q}_p . Let further $\chi_{\mathbb{Q}_p}^{\text{nr}} : G_{\mathbb{Q}_p} \rightarrow \mathbb{Z}_p^\times$ be a continuous unramified character and write $\chi^{\text{nr}} :=$

$\chi_{\mathbb{Q}_p}^{\text{nr}}|_{G_K}$ for its restriction to G_K . We write $\chi^{\text{cyc}}: G_K \rightarrow \mathbb{Z}_p^\times$ for the cyclotomic character and consider the p -adic representation

$$T := \mathbb{Z}_p(\chi^{\text{nr}})(\chi^{\text{cyc}}) = \mathbb{Z}_p(\chi^{\text{nr}})(1).$$

Explicitly, T is \mathbb{Z}_p as an abelian group equipped with the G_K -action defined by $gx := \chi^{\text{nr}}(g)\chi^{\text{cyc}}(g)x$ for $g \in G_K$ and $x \in \mathbb{Z}_p$. We set $V := \mathbb{Q}_p \otimes_{\mathbb{Z}_p} T$.

Let $\varphi = F_{\mathbb{Q}_p}$ denote the absolute Frobenius automorphism and set $u := \chi_{\mathbb{Q}_p}^{\text{nr}}(\varphi)$. Let \mathcal{F} denote the Lubin–Tate formal group associated with $\pi := up$. By [Izy12, Ex. 5.20] the representation T is the restriction to G_K of the p -adic Tate module $T_p\mathcal{F}$.

Let N/K be a finite Galois extension with group G . As usual, we write $\mathcal{F}(\mathfrak{p}_N)$ for the formal \mathbb{Z}_p -module \mathfrak{p}_N with the \mathbb{Z}_p -module structure induced by the formal group law \mathcal{F} .

We define the complex $R\Gamma(N, T)$ as the G_N -invariants of the continuous standard resolution of T (cf. [NSW08, Sec. I.2]). As usual, we write $H^i(N, T)$ for the cohomology groups of $R\Gamma(N, T)$. We recall some well-known facts about these cohomology groups. We write $v_p: \mathbb{Q}_p \rightarrow \mathbb{Z} \cup \{\infty\}$ for the normalized valuation map and set

$$(14) \quad \omega = \omega_N := v_p(1 - \chi^{\text{nr}}(F_N)).$$

Note that $\omega \in \mathbb{Z}$ if and only if $\chi^{\text{nr}}|_{G_N} \neq 1$.

LEMMA 3.2.1. *Assume the notation of Subsection 3.2.*

(a) *If $\chi^{\text{nr}}|_{G_N} \neq 1$, then*

- (i) $H^1(N, T) \cong (\widehat{N_0^\times}(\chi^{\text{nr}}))^{\text{Gal}(N_0/N)} \cong \mathcal{F}(\mathfrak{p}_N)$,
- (ii) $H^2(N, T) \cong (\mathbb{Z}_p/p^\omega\mathbb{Z}_p)(\chi^{\text{nr}})$,
- (iii) $H^i(N, T) = 0$ for $i \neq 1, 2$.

(b) *If $\chi^{\text{nr}}|_{G_N} = 1$, then*

- (i) $H^1(N, T) \cong \widehat{N^\times}(\chi^{\text{nr}})$,
- (ii) $H^2(N, T) \cong \mathbb{Z}_p(\chi^{\text{nr}})$,
- (iii) $H^i(N, T) = 0$ for $i \neq 1, 2$.

Proof. Part (a) is shown in [IV16, Prop. 2.5]. If $\chi^{\text{nr}}|_{G_N} = 1$, then taking cohomology commutes with twisting, i.e., $H^i(N, A(\chi^{\text{nr}})) = H^i(N, A)(\chi^{\text{nr}})$ for each G_N -module A . Hence part (b) is immediate from [Bre04a, Prop. 4.11]. ■

In Section 4 we will give an alternative proof of part (a) by explicitly constructing a representative for $R\Gamma(N, T)$. For details see Remark 4.3.2.

REMARK 3.2.2. In [BC16] and [Chi85] the symbol N_0 denotes the maximal unramified extension of N , which in the present manuscript is denoted by N_{nr} . In order to relate the results and computations of [BC16] to those

discussed here, it is worth noting that $\widehat{N_{\text{nr}}^\times} = \widehat{N_0^\times}$. Indeed, for each natural number n we have the canonical short exact sequence

$$0 \rightarrow N_{\text{nr}}^\times / (N_{\text{nr}}^\times \cap (N_0^\times)^{p^n}) \rightarrow N_0^\times / (N_0^\times)^{p^n} \rightarrow N_0^\times / N_{\text{nr}}^\times (N_0^\times)^{p^n} \rightarrow 0.$$

It is an easy exercise to show that $N_{\text{nr}}^\times (N_0^\times)^{p^n} = N_0^\times$ and $N_{\text{nr}}^\times \cap (N_0^\times)^{p^n} = (N_{\text{nr}}^\times)^{p^n}$. Hence we have a natural isomorphism $N_{\text{nr}}^\times / (N_{\text{nr}}^\times)^{p^n} \cong N_0^\times / (N_0^\times)^{p^n}$.

REMARK 3.2.3. In our situation we have

$$h_i(V) = \begin{cases} 1 & \text{if } i = -1, \\ 0 & \text{if } i \neq -1. \end{cases}$$

Indeed, by [FO10, p. 148] or [IV16, p. 499],

$$t_{V^*(1)}(\mathbb{Q}_p) \hookrightarrow \mathbb{C}_p((\chi^{\text{nr}})^{-1})(-1)^{G_N} = 0$$

and hence $t_{V^*(1)}(\mathbb{Q}_p) = 0$. Then the short exact sequence (6) implies that $D_{\text{dR}}^{\mathbb{Q}_p}(V) = t_V(\mathbb{Q}_p)$, i.e. $\text{Fil}^0 D_{\text{dR}}^{\mathbb{Q}_p}(V) = 0$. Furthermore the element $e_{\chi_{\mathbb{Q}_p,1}^{\text{nr}}} = t^{\text{nr}} \cdot t^{-1} \otimes (v \otimes \xi)$ defined in [IV16, p. 508] is a non-zero element in $\text{Fil}^{-1} D_{\text{dR}}^{\mathbb{Q}_p}(V)$. Since $\dim_{\mathbb{Q}_p}(D_{\text{dR}}^{\mathbb{Q}_p}(V)) = 1$, we deduce that $\text{Fil}^{-1} D_{\text{dR}}^{\mathbb{Q}_p}(V) = D_{\text{dR}}^{\mathbb{Q}_p}(V)$, so that there is only one jump in the filtration, namely at $i = -1$.

Hence we obtain $t_H(V) = -1$ and $\Gamma^*(V) = \Gamma^*(1) = 1$.

In this case we shall now reformulate Conjecture 3.1.1 in the language of relative algebraic K -groups and refined Euler characteristics.

Following [IV16] and [Bre04b] we define an element $\tilde{R}_{N/K} \in K_0(\mathbb{Z}_p[G], \tilde{\Omega})$ by

$$\tilde{R}_{N/K} = C_{N/K} + U_{\text{cris}} + \partial_{\mathbb{Z}_p[G], B_{\text{dR}}[G]}^1(t) + \partial_{\mathbb{Z}_p[G], B_{\text{dR}}[G]}^1(\varepsilon_D(N/K, V)),$$

where each of the terms is an element in $K_0(\mathbb{Z}_p[G], B_{\text{dR}}[G])$. After briefly recalling the definition of $C_{N/K}$ we give the formulation of the conjecture due to Izychev and Venjakob, and then, following the approach of Breuning, modify $\tilde{R}_{N/K}$ by the so-called unramified term in order to obtain an element $R_{N/K} \in K_0(\mathbb{Z}_p[G], \mathbb{Q}_p[G])$. In Section 6 we will write $\varepsilon_D(N/K, V)$ in terms of Galois Gauß sums in order to tie up to the notion of ε -constants as it is used in Breuning’s conjecture [Bre04b, Conj. 3.2].

The element $C_{N/K} \in K_0(\mathbb{Z}_p[G], \tilde{\Omega})$ is defined by

$$(15) \quad C_{N/K} = -\chi_{\mathbb{Z}_p[G], B_{\text{dR}}[G]}(M^\bullet, \lambda^{-1})$$

where the trivialized complex $(M^\bullet, \lambda^{-1})$ is given by

$$(16) \quad M^\bullet = R\Gamma(N, T) \oplus \text{Ind}_{N/\mathbb{Q}_p} T[0]$$

and

$$\lambda: H^{\text{od}}(M^\bullet)_{B_{\text{dR}}[G]} \rightarrow H^{\text{ev}}(M^\bullet)_{B_{\text{dR}}[G]},$$

$$\lambda = \begin{cases} \text{comp} \circ \exp_V^{-1} & \text{if } \chi^{\text{nr}}|_{G_N} \neq 1, \\ \text{comp} \circ \exp_V^{-1} + \nu_N & \text{if } \chi^{\text{nr}}|_{G_N} = 1. \end{cases}$$

Note that the minus sign in the definition of $C_{N/K}$ is due to a different convention about Euler characteristics in comparison to [IV16].

The explicit computation of $C_{N/K}$ is the technical heart of this paper. The computation is based on the construction of an explicit representative for $RF(N, T)$ (see Section 4), which is of independent interest. Our representative should be considered as the analogue of Serre’s explicit representative of the local fundamental class (see [Chi85, Prop. 6.1] or [Bre04a, Th. 4.20]).

The term U_{cris} will be defined in Section 5. In the case $\chi^{\text{nr}}|_{G_N} \neq 1$ our definition agrees with the one given in [IV16, Sec. 5]. In Section 5 we will also make the comparison to the correction term $M_{N/K}$ which occurs in (the generalization of) Breuning’s conjecture [Bre04b, Conj. 3.2].

CONJECTURE 3.2.4. *Under the current assumptions, $\tilde{R}_{N/K}$ equals 0 in $K_0(\tilde{\Lambda}, \tilde{\Omega})$.*

In the case $\chi^{\text{nr}}|_{G_N} \neq 1$ this is precisely the reformulation of Conjecture 3.1.1 as stated in [IV16, (5.2)].

We now follow the approach of [Bre04b] to define a modified element $R_{N/K}$ in $K_0(\mathbb{Z}_p[G], \mathbb{Q}_p[G])$. We write \mathcal{O}_p^t for the ring of integers in the maximal tamely ramified extension of \mathbb{Q}_p in \mathbb{Q}_p^c . Let $\iota: K_0(\mathbb{Z}_p[G], \mathbb{Q}_p^c[G]) \rightarrow K_0(\mathcal{O}_p^t[G], \mathbb{Q}_p^c[G])$ be the natural scalar extension map. We recall that by Taylor’s fixed point theorem the restriction of ι to $K_0(\mathbb{Z}_p[G], \mathbb{Q}_p[G])$ is injective. Let us define $U_{N/K}$ as in [Bre04b]; then by [Bre04b, Prop. 2.12] we have $\iota(U_{N/K}) = 0$. We define

$$(17) \quad R_{N/K} = C_{N/K} + U_{\text{cris}} + \partial_{\mathbb{Z}_p[G], B_{\text{dR}}[G]}^1(t) - U_{N/K} + \partial_{\mathbb{Z}_p[G], B_{\text{dR}}[G]}^1(\varepsilon_D(N/K, V)),$$

so that $R_{N/K} = \tilde{R}_{N/K} - U_{N/K}$. Using some of the properties of $U_{N/K}$ with respect to the action of $\text{Gal}(\mathbb{Q}_p^c/\mathbb{Q}_p)$ one can prove that $R_{N/K} \in K_0(\mathbb{Z}_p[G], \mathbb{Q}_p[G])$. We will do this in Subsection 7.1 and we will also prove that if $T = \mathbb{Z}_p(1)$, then the following conjecture is precisely Breuning’s conjecture [Bre04b, Conj. 3.2].

CONJECTURE 3.2.5. *Under the current assumptions, the element $R_{N/K}$ equals 0 in $K_0(\mathbb{Z}_p[G], \mathbb{Q}_p[G])$.*

PROPOSITION 3.2.6. *In the present setting, Conjectures 3.1.1, 3.2.4 and 3.2.5 are equivalent.*

The proof will be given in Subsection 7.1. In the rest of the article we write $C_{\text{EP}}^{\text{na}}(N/K, V)$ for the equivalent Conjectures 3.1.1, 3.2.4 and 3.2.5.

3.3. Organization of the paper. In Section 4 we construct an explicit representative of $R\Gamma(N, T)$. Section 5 is dedicated to the definition of the term U_{cris} and the comparison of U_{cris} to Breuning’s correction term $M_{N/K}$. In the following Section 6 we discuss epsilon constants and express them in terms of Galois Gauß sums. This will allow us to compare the term $\partial_{\mathbb{Z}_p[G], B_{\text{dR}}[G]}^1(\varepsilon_D(N/K, V))$ to Breuning’s element $T_{N/K}$. In Section 7 we express $C_{N/K}$ as a sum of a norm resolvent and the refined Euler characteristic of $R\Gamma(N, T)$. With these preparations at hand we study the Galois action of the individual terms in the definition of $R_{N/K}$, prove Proposition 3.2.6 and compute the refined Euler characteristic $C_{N/K}$. As a by-product we obtain the proof of the equivalence of Breuning’s conjecture [Bre04b, Conj. 3.2] and $C_{\text{EP}}^{\text{na}}(N/K, V)$ if $T = \mathbb{Z}_p(1)$. Finally, in Section 8 we provide the proof of Theorem 1.

4. A representative for $R\Gamma(N, T)$. In this section we construct an explicit representative for the complex $R\Gamma(N, T)$. We assume the situation described at the beginning of Subsection 3.2.

4.1. Some preliminary results

LEMMA 4.1.1. *Let L/K denote a finite Galois extension. Then we have a canonical exact sequence*

$$0 \rightarrow \mathcal{F}(\mathfrak{p}_L) \rightarrow (\widehat{L}_0^\times(\chi^{\text{nr}}))^{\text{Gal}(L_0/L)} \xrightarrow{\nu_L} (\mathbb{Z}_p(\chi^{\text{nr}}))^{\text{Gal}(L_0/L)}$$

of $I_{L/K}$ -modules. In particular, if $\chi^{\text{nr}}|_{G_L} \neq 1$, then there is an isomorphism

$$f_{\mathcal{F}, L}: \mathcal{F}(\mathfrak{p}_L) \rightarrow (\widehat{L}_0^\times(\chi^{\text{nr}}))^{\text{Gal}(L_0/L)}.$$

Explicitly $f_{\mathcal{F}, L}$ is given by a power series of the form $1 + \varepsilon^{-1}X + \text{deg} \geq 2$, where $\varepsilon \in \overline{\mathbb{Z}_p^{\text{nr}}}^\times$ is such that $u = \varepsilon^{\varphi-1}$.

Proof. We have $L_0^\times \cong \pi_L^{\mathbb{Z}} \times \kappa^\times \times U_{L_0}^{(1)}$, where κ denotes the residue class field of L_0 . Since any element of κ^\times has order coprime to p , and $U_{L_0}^{(1)}$ is p -complete, we obtain $\widehat{L}_0^\times \cong \mathbb{Z}_p \times U_{L_0}^{(1)}$, and hence a canonical short exact sequence

$$(18) \quad 0 \rightarrow U_{L_0}^{(1)} \rightarrow \widehat{L}_0^\times \rightarrow \mathbb{Z}_p \rightarrow 0.$$

From [LR78, Lemma, p. 237] we derive the existence of an isomorphism

$$\tilde{f}: \mathcal{F}(\mathfrak{p}_L) \rightarrow (U_{L_0}^{(1)}(\chi^{\text{nr}}))^{\text{Gal}(L_0/L)}.$$

Up to the obvious isomorphism $U_{L_0}^{(1)} \cong \mathbb{G}_m(\mathfrak{p}_{L_0})$, this isomorphism is given by the uniquely defined power series $\theta(X) = \varepsilon^{-1}X + \deg \geq 2$ of [Neu92, Koroll. V.2.3]. The result follows by twisting and taking $\text{Gal}(L_0/L)$ -fixed points in (18). ■

We set $\mathcal{G} := \text{Gal}(K_0/K) \times \text{Gal}(L/K)$. In analogy to [Chi85, (6.2)] we set

$$\mathcal{I}_{L/K}(\chi^{\text{nr}}) := \text{Ind}_{\text{Gal}(L_0/K)}^{\mathcal{G}}(\widehat{L}_0^\times(\chi^{\text{nr}}))$$

and we usually identify $\mathcal{I}_{L/K}(\chi^{\text{nr}})$ with $(\widehat{L}_0^\times)^{d_{L/K}}$. Recalling that $F = F_K$ denotes the Frobenius automorphism of K , note that the \mathcal{G} -module structure on $\mathcal{I}_{L/K}(\chi^{\text{nr}})$ is characterized by

$$(19) \quad (F \times 1) \cdot [x_1, \dots, x_{d_{L/K}}] = [x_{d_{L/K}}^{F_L \chi^{\text{nr}}(F_L)}, x_1, \dots, x_{d_{L/K}-1}],$$

$$(20) \quad (F^{-n} \times \sigma) \cdot [x_1, \dots, x_{d_{L/K}}] = [x_1^{\tilde{\sigma} \chi^{\text{nr}}(\tilde{\sigma})}, \dots, x_{d_{L/K}}^{\tilde{\sigma} \chi^{\text{nr}}(\tilde{\sigma})}],$$

where F^{-n} and $\sigma \in \text{Gal}(L/K)$ have the same restriction to $L \cap K_0$ and $\tilde{\sigma} \in \text{Gal}(L_0/K)$ is uniquely defined by $\tilde{\sigma}|_{K_0} = F^{-n}$ and $\tilde{\sigma}|_L = \sigma$.

We also have a G_K - and a $\text{Gal}(L/K)$ -action on $\mathcal{I}_{L/K}(\chi^{\text{nr}})$, via the natural maps $G_K \rightarrow \text{Gal}(L_0/K) \hookrightarrow \mathcal{G}$ and $\text{Gal}(L/K) \hookrightarrow \mathcal{G}$. Note that in the case $\chi^{\text{nr}}|_{G_L} = 1$ we have

$$(21) \quad \mathcal{I}_{L/K}(\chi^{\text{nr}}) = \text{Ind}_{\text{Gal}(L_0/K)}^{\mathcal{G}}(\widehat{L}_0^\times(\chi^{\text{nr}})) \cong (\text{Ind}_{\text{Gal}(L_0/K)}^{\mathcal{G}} \widehat{L}_0^\times)(\chi^{\text{nr}})$$

both as $\text{Gal}(L/K)$ - and G_K -modules. The following lemma clarifies the structure of $\mathcal{I}_{L/K}(\chi^{\text{nr}})$ as a $\text{Gal}(L/K)$ -module in the general situation.

LEMMA 4.1.2. *If we identify the inertia group $I_{L/K}$ with $\text{Gal}(L_0/K_0)$, then there is a $\text{Gal}(L/K)$ -isomorphism*

$$\tilde{f}_{\mathcal{F},L}: \text{Ind}_{I_{L/K}}^{\text{Gal}(L/K)} \widehat{L}_0^\times \rightarrow \mathcal{I}_{L/K}(\chi^{\text{nr}}).$$

In particular, $\mathcal{I}_{L/K}(\chi^{\text{nr}})$ is $\text{Gal}(L/K)$ -cohomologically trivial.

Proof. We identify $\text{Ind}_{I_{L/K}}^{\text{Gal}(L/K)} \widehat{L}_0^\times$ with $\mathbb{Z}_p[\text{Gal}(L/K)] \otimes_{\mathbb{Z}_p[I_{L/K}]} \widehat{L}_0^\times$ and set

$$\tilde{f}_{\mathcal{F},L}(\sigma \otimes x) := (1 \times \sigma) \cdot [x, 1, \dots, 1]$$

for $x \in \widehat{L}_0^\times$ and $\sigma \in \text{Gal}(L/K)$. To check that this is well-defined we take $\tau \in I_{L/K}$ and write $\tilde{\tau}$ for the corresponding element in $\text{Gal}(L_0/K_0)$. Then

$$\begin{aligned} \tilde{f}_{\mathcal{F},L}(\sigma\tau \otimes x) &= (1 \times \sigma\tau) \cdot [x, 1, \dots, 1] = (1 \times \sigma)(1 \times \tau) \cdot [x, 1, \dots, 1] \\ &= (1 \times \sigma) \cdot [x^{\tilde{\tau} \chi^{\text{nr}}(\tilde{\tau})}, 1, \dots, 1] = (1 \times \sigma) \cdot [x^{\tilde{\tau}}, 1, \dots, 1] \\ &= \tilde{f}_{\mathcal{F},L}(\sigma \otimes x^{\tilde{\tau}}). \end{aligned}$$

To prove the surjectivity of $\tilde{f}_{\mathcal{F},L}$, we check that all elements of the form $[1, \dots, 1, x, 1, \dots, 1]$ are in its image. Let $i + 1$ be the index of x and let us take $\sigma \in \text{Gal}(L/K)$ such that $\sigma|_{L \cap K_0} = F^{-i}$. Then

$$\begin{aligned} \tilde{f}_{\mathcal{F},L}(\sigma \otimes x^{\tilde{\sigma}^{-1}\chi^{\text{nr}}(\tilde{\sigma})^{-1}}) &= (1 \times \sigma) \cdot [x^{\tilde{\sigma}^{-1}\chi^{\text{nr}}(\tilde{\sigma})^{-1}}, 1, \dots, 1] \\ &= (F^i \times 1)(F^{-i} \times \sigma) \cdot [x^{\tilde{\sigma}^{-1}\chi^{\text{nr}}(\tilde{\sigma})^{-1}}, 1, \dots, 1] \\ &= (F^i \times 1) \cdot [x, 1, \dots, 1] = [1, \dots, 1, x, 1, \dots, 1]. \end{aligned}$$

It remains to prove injectivity. Let us take $\prod_i \sigma_i \otimes x_i \in \ker \tilde{f}_{\mathcal{F},L}$. Clearly we can assume that $0 \leq i \leq d_{L/K} - 1$ and $\sigma_i|_{L \cap K_0} = F^{-i}$. Then we calculate

$$\tilde{f}_{\mathcal{F},L} \left(\prod_{i=0}^{d_{L/K}-1} \sigma_i \otimes x_i \right) = [x_0^{\tilde{\sigma}_0\chi^{\text{nr}}(\tilde{\sigma}_0)}, x_1^{\tilde{\sigma}_1\chi^{\text{nr}}(\tilde{\sigma}_1)}, \dots, x_{d_{L/K}-1}^{\tilde{\sigma}_{d_{L/K}-1}\chi^{\text{nr}}(\tilde{\sigma}_{d_{L/K}-1})}].$$

Since we are assuming that the element is in the kernel, we deduce that all of the x_i are equal to 1. Hence $\tilde{f}_{\mathcal{F},L}$ is injective. The isomorphism is $\text{Gal}(L/K)$ -invariant by construction.

By [Ser79, Prop. XIII.5.14] the $I_{L/K}$ -module L_0^\times is $I_{L/K}$ -cohomologically trivial. By [Bre04a, Lemma 4.9] the same holds true for the p -completion $\widehat{L_0^\times}$. From Shapiro’s Lemma we finally derive that $\text{Ind}_{I_{L/K}}^{\text{Gal}(L/K)} \widehat{L_0^\times}$ is $\text{Gal}(L/K)$ -cohomologically trivial. ■

We now prove a slight generalization of a well-known lemma, see e.g. [Neu92, Lemma V.2.1].

LEMMA 4.1.3. *Given $c \in \widehat{U_{L_0}}$ there exists $x \in \widehat{U_{L_0}}$ with $x^{\chi^{\text{nr}}(F_L)F_L^{-1}} = c$. If x_n is such a solution modulo $U_{L_0}^{p^n}$, then we can assume that $x \equiv x_n \pmod{U_{L_0}^{p^n}}$. In particular, if $c \in U_{L_0}^{(1)}$, then we can assume $x \in U_{L_0}^{(1)}$.*

Proof. Let $n \in \mathbb{N}$, and $\psi_n \in \mathbb{Z}$ be such that $\psi_n \equiv \chi^{\text{nr}}(F_L) \pmod{p^n}$. First of all we want to determine an $x_n \in U_{L_0}/U_{L_0}^{p^n}$ such that $x_n^{\chi^{\text{nr}}(F_L)F_L^{-1}} \equiv x_n^{\psi_n F_L^{-1}} \equiv c \pmod{U_{L_0}^{p^n}}$. Let $c_n \in U_{L_0}$ be congruent to c modulo $U_{L_0}^{p^n}$. Then it is enough to find $x_n \in U_{L_0}$ such that $x_n^{\psi_n F_L^{-1}} = c_n$. The same proof as in [Neu92, Lemma V.2.1] works, simply writing $F_L \psi$ instead of φ everywhere. (Note that here we are using the fact that L_0 is complete.) Moreover, if we assume that we already have an $x_{n-1} \in U_{L_0}/U_{L_0}^{p^{n-1}}$ such that $x_{n-1}^{\chi^{\text{nr}}(F_L)F_L^{-1}} \equiv x_{n-1}^{\psi_{n-1} F_L^{-1}} \equiv c \pmod{U_{L_0}^{p^{n-1}}}$, then we can find x_n with the additional property that $x_n \equiv x_{n-1} \pmod{U_{L_0}^{p^{n-1}}}$.

By passing to the limit we obtain an element $x \in \widehat{U_{L_0}}$ which satisfies $x^{\chi^{\text{nr}}(F_L)F_L^{-1}} = c$. The remaining assertions are now straightforward. ■

4.2. The construction of the complex. We continue to assume the situation of Subsection 3.2.

THEOREM 4.2.1 (Serre). *If $\chi^{\text{nr}}(F_L) = 1$, then we have an exact sequence*

$$0 \rightarrow \widehat{L^\times}(\chi^{\text{nr}}) \rightarrow \mathcal{I}_{L/K}(\chi^{\text{nr}}) \rightarrow \mathcal{I}_{L/K}(\chi^{\text{nr}}) \rightarrow \mathbb{Z}_p(\chi^{\text{nr}}) \rightarrow 0$$

of $\mathbb{Z}_p[\text{Gal}(L/K)]$ -modules.

Proof. In view of the isomorphism (21) this follows directly from [Chi85, Prop. 6.1] by p -completing and twisting. ■

Now we want an analogous result for the case $\chi^{\text{nr}}(F_L) \neq 1$.

We set

$$\omega_L := v_p(1 - \chi^{\text{nr}}(F_L)),$$

which is finite when $\chi^{\text{nr}}(F_L) \neq 1$.

For $n > \omega_L$ we let

$$V_n = \{y \in L_0^\times / (L_0^\times)^{p^n} : \nu_L(y) \equiv 0 \pmod{p^{\omega_L}}\},$$

$$W_n = \{y \in L_0^\times / (L_0^\times)^{p^n} : y^{\chi^{\text{nr}}(F_L)F_L - 1} = 1\}.$$

Note that when $\omega_L = 0$ the congruence in the definition of V_n is always trivially satisfied, so that $V_n = L_0^\times / (L_0^\times)^{p^n}$.

LEMMA 4.2.2. *Assume $\chi^{\text{nr}}(F_L) \neq 1$. If $m \geq n \geq \omega_L$, then we have a commutative diagram of exact sequences:*

$$\begin{array}{ccccccc} 0 & \longrightarrow & W_m & \longrightarrow & L_0^\times / (L_0^\times)^{p^m} & \xrightarrow{\chi^{\text{nr}}(F_L)F_L - 1} & V_m \longrightarrow 0 \\ & & \downarrow \tau_{m,n} & & \downarrow \pi_{m,n} & & \downarrow \sigma_{m,n} \\ 0 & \longrightarrow & W_n & \longrightarrow & L_0^\times / (L_0^\times)^{p^n} & \xrightarrow{\chi^{\text{nr}}(F_L)F_L - 1} & V_n \longrightarrow 0 \end{array}$$

Here $\pi_{m,n}$ denotes the canonical projection and $\tau_{m,n}$ and $\sigma_{m,n}$ are induced by $\pi_{m,n}$. Moreover, $\pi_{m,n}$ and $\sigma_{m,n}$ are surjective and the projective system $(W_n)_n$ satisfies the Mittag-Leffler condition.

Proof. To prove the exactness of the rows we must show that

$$L_0^\times / (L_0^\times)^{p^n} \xrightarrow{\chi^{\text{nr}}(F_L)F_L - 1} V_n$$

is surjective. If $c \in V_n$, then $\nu_L(c) \in p^{\omega_L}\mathbb{Z}/p^n\mathbb{Z}$, so that there exists $k \in \mathbb{Z}$ such that

$$\nu_L(c) \equiv k(\chi^{\text{nr}}(F_L) - 1) \pmod{p^n\mathbb{Z}}.$$

We set

$$c' = \frac{c}{[\pi_L]_{\nu_L(c)}} \in U_{L_0} / (U_{L_0})^{p^n} \subseteq L_0^\times / (L_0^\times)^{p^n},$$

where $[\pi_L]$ denotes the class of π_L modulo $(L_0^\times)^{p^n}$. By Lemma 4.1.3 there exists $x' \in U_{L_0}/(U_{L_0})^{p^n}$ such that

$$(x')^{\chi^{\text{nr}}(F_L)F_L-1} = c'.$$

Then we set $x = x'[\pi_L]^k$ and obtain

$$x^{\chi^{\text{nr}}(F_L)F_L-1} = (x'[\pi_L]^k)^{\chi^{\text{nr}}(F_L)F_L-1} = c'[\pi_L]^k(\chi^{\text{nr}}(F_L)-1) = c'[\pi_L]^{\nu_L(c)} = c.$$

The commutativity of the diagram as well as the surjectivity of $\pi_{m,n}$ and $\sigma_{m,n}$ are evident. It remains to show that the projective system $(W_n)_n$ satisfies the Mittag–Leffler condition. To that end we prove that $\text{coker}(\tau_{m,n})$ stabilizes for $n \geq \omega_L$ as m tends to infinity.

By the Snake Lemma we obtain

$$(L_0^\times)^{p^n}/(L_0^\times)^{p^m} \xrightarrow{\chi^{\text{nr}}(F_L)F_L-1} \ker(\sigma_{m,n}) \rightarrow \text{coker}(\tau_{m,n}) \rightarrow 0.$$

Since $n > \omega_L$, it is easy to see that

$$\ker(\sigma_{m,n}) = (L_0^\times)^{p^n}/(L_0^\times)^{p^m},$$

so that we have to compute the cokernel of

$$(L_0^\times)^{p^n}/(L_0^\times)^{p^m} \xrightarrow{\psi} (L_0^\times)^{p^n}/(L_0^\times)^{p^m},$$

where ψ is induced by $\chi^{\text{nr}}(F_L)F_L - 1$. Clearly,

$$\text{im}(\psi) \subseteq (\pi_L^{p^{\omega_L+n}\mathbb{Z}} \times (U_{L_0})^{p^n})(L_0^\times)^{p^m}/(L_0^\times)^{p^m}.$$

We claim that we have equality. Let $c^{p^n} = \pi_L^{lp^{n+\omega_L}}v^{p^n}$ with $l \in \mathbb{Z}$ and $v \in U_{L_0}$ represent an element of the right hand side and set $c := \pi_L^{lp^{\omega_L}}v$. By the first part of the proof we can construct $x \in L_0^\times/(L_0^\times)^{p^m}$ such that $x^{\chi^{\text{nr}}(F_L)F_L-1} = [c]$. Then $(x^{p^n})^{\chi^{\text{nr}}(F_L)F_L-1} = [c^{p^n}]$, which proves our claim.

Finally we note that for $m \geq n + \omega_L$ one has $(L_0^\times)^{p^m} \subseteq \pi_L^{p^{\omega_L+n}\mathbb{Z}} \times (U_{L_0})^{p^n}$. Hence $\text{coker}(\tau_{m,n}) \cong (L_0^\times)^{p^n}/(\pi_L^{p^{\omega_L+n}\mathbb{Z}} \times (U_{L_0})^{p^n})$ stabilizes. ■

By the definition of V_n we have a commutative diagram of short exact sequences

$$\begin{array}{ccccccc} 0 & \longrightarrow & V_m & \longrightarrow & L_0^\times/(L_0^\times)^{p^m} & \xrightarrow{\nu_L} & \mathbb{Z}/p^{\omega_L}\mathbb{Z} \longrightarrow 0 \\ & & \downarrow \sigma_{m,n} & & \downarrow \pi_{m,n} & & \downarrow = \\ 0 & \longrightarrow & V_n & \longrightarrow & L_0^\times/(L_0^\times)^{p^n} & \xrightarrow{\nu_L} & \mathbb{Z}/p^{\omega_L}\mathbb{Z} \longrightarrow 0 \end{array}$$

Note that when $\omega_L = 0$ we simply interpret $\mathbb{Z}/p^{\omega_L}\mathbb{Z}$ as 0. By Lemma 4.2.2 the maps $\sigma_{m,n}$ are surjective, and hence the projective system $(V_n)_n$ satisfies the Mittag–Leffler condition. We define

$$W := \varprojlim_n W_n, \quad V := \varprojlim_n V_n,$$

and thus obtain by [Wei94, Prop. 3.5.7] two short exact sequences

$$0 \rightarrow W \rightarrow \widehat{L}_0^\times \rightarrow V \rightarrow 0, \quad 0 \rightarrow V \rightarrow \widehat{L}_0^\times \rightarrow \mathbb{Z}/p^{\omega_L}\mathbb{Z} \rightarrow 0.$$

Splicing together these two sequences we obtain

$$0 \rightarrow W \rightarrow \widehat{L}_0^\times \xrightarrow{\chi^{\text{nr}}(F_L)F_L-1} \widehat{L}_0^\times \xrightarrow{\nu_L} \mathbb{Z}/p^{\omega_L}\mathbb{Z} \rightarrow 0,$$

which we rewrite in the form

$$(22) \quad 0 \rightarrow W(\chi^{\text{nr}}) \rightarrow \widehat{L}_0^\times(\chi^{\text{nr}}) \xrightarrow{F_L-1} \widehat{L}_0^\times(\chi^{\text{nr}}) \xrightarrow{\nu_L} \mathbb{Z}/p^{\omega_L}\mathbb{Z}(\chi^{\text{nr}}) \rightarrow 0.$$

We clearly have $W(\chi^{\text{nr}}) \cong \widehat{L}_0^\times(\chi^{\text{nr}})^{\text{Gal}(L_0/L)}$, which by Lemma 4.1.1 is isomorphic to $\mathcal{F}(\mathfrak{p}_L)$.

Note that we also have a natural $\text{Gal}(L/K)$ -action on $(\widehat{L}_0^\times(\chi^{\text{nr}}))^{\text{Gal}(L_0/L)}$ and $\mathbb{Z}/p^{\omega_L}\mathbb{Z}(\chi^{\text{nr}})$ defined by extending an element $\sigma \in \text{Gal}(L/K)$ to an element $\tilde{\sigma} \in \text{Gal}(L_0/K)$.

So we can formulate the following theorem.

THEOREM 4.2.3. *Assume $\chi^{\text{nr}}(F_L) \neq 1$. We have an exact sequence*

$$0 \rightarrow (\widehat{L}_0^\times(\chi^{\text{nr}}))^{\text{Gal}(L_0/L)} \rightarrow \mathcal{I}_{L/K}(\chi^{\text{nr}}) \xrightarrow{(F-1) \times 1} \mathcal{I}_{L/K}(\chi^{\text{nr}}) \xrightarrow{w_L} \mathbb{Z}/p^{\omega_L}\mathbb{Z}(\chi^{\text{nr}}) \rightarrow 0$$

of $\mathbb{Z}_p[\text{Gal}(L/K)]$ -modules. Here w_L denotes the sum of the valuations ν_L .

REMARK 4.2.4. In the proof of Theorem 4.2.8 we will use this result together with the isomorphism $(\widehat{L}_0^\times(\chi^{\text{nr}}))^{\text{Gal}(L_0/L)} \cong \mathcal{F}(\mathfrak{p}_L)$ provided by Lemma 4.1.1.

Proof of Theorem 4.2.3. We start by showing that all the homomorphisms are $\text{Gal}(L/K)$ -invariant. Let $\sigma \in \text{Gal}(L/K)$ and let $\tilde{\sigma} \in \text{Gal}(L_0/K)$ be such that $\tilde{\sigma}|_L = \sigma$ and $\tilde{\sigma}|_{K_0} = F^{-n}$, where n is an integer. Let $x \in (\widehat{L}_0^\times(\chi^{\text{nr}}))^{\text{Gal}(L_0/L)}$. Then x maps diagonally to the element $[x, \dots, x] \in \mathcal{I}_{L/K}(\chi^{\text{nr}})$ and we have

$$\begin{aligned} (1 \times \sigma) \cdot [x, \dots, x] &= (F^n \times 1)(F^{-n} \times \sigma) \cdot [x, \dots, x] \\ &= (F^n \times 1) \cdot [x^{\tilde{\sigma}\chi^{\text{nr}}(\tilde{\sigma})}, \dots, x^{\tilde{\sigma}\chi^{\text{nr}}(\tilde{\sigma})}] \\ &= [x^{\tilde{\sigma}\chi^{\text{nr}}(\tilde{\sigma})F_L\chi^{\text{nr}}(F_L)}, \dots, x^{\tilde{\sigma}\chi^{\text{nr}}(\tilde{\sigma})F_L\chi^{\text{nr}}(F_L)}, x^{\tilde{\sigma}\chi^{\text{nr}}(\tilde{\sigma})}, \dots, x^{\tilde{\sigma}\chi^{\text{nr}}(\tilde{\sigma})}] \\ &= [x^{\tilde{\sigma}\chi^{\text{nr}}(\tilde{\sigma})}, \dots, x^{\tilde{\sigma}\chi^{\text{nr}}(\tilde{\sigma})}]. \end{aligned}$$

If we inject $\sigma \cdot x = x^{\tilde{\sigma}\chi^{\text{nr}}(\tilde{\sigma})}$ diagonally into $\mathcal{I}_{L/K}(\chi^{\text{nr}})$ we obviously get the same element, so that the first map is $\text{Gal}(L/K)$ -invariant. The second map is $\text{Gal}(L/K)$ -invariant because $(F-1) \times 1$ and $1 \times \sigma$ commute.

Let $[x_1, \dots, x_{d_{L/K}}] \in \mathcal{I}_{L/K}(\chi^{\text{nr}})$. Then

$$\begin{aligned} (1 \times \sigma) \cdot [x_1, \dots, x_{d_{L/K}}] &= (F^n \times 1)(F^{-n} \times \sigma) \cdot [x_1, \dots, x_{d_{L/K}}] \\ &= (F^n \times 1) \cdot [x_1^{\tilde{\sigma}\chi^{\text{nr}}(\tilde{\sigma})}, \dots, x_{d_{L/K}}^{\tilde{\sigma}\chi^{\text{nr}}(\tilde{\sigma})}] \\ &= [x_{d_{L/K}+1-n}^{\tilde{\sigma}\chi^{\text{nr}}(\tilde{\sigma})F_L\chi^{\text{nr}}(F_L)}, \dots, x_{d_{L/K}}^{\tilde{\sigma}\chi^{\text{nr}}(\tilde{\sigma})F_L\chi^{\text{nr}}(F_L)}, x_1^{\tilde{\sigma}\chi^{\text{nr}}(\tilde{\sigma})}, \dots, x_{d_{L/K}-n}^{\tilde{\sigma}\chi^{\text{nr}}(\tilde{\sigma})}]. \end{aligned}$$

Its valuation is

$$\begin{aligned} \chi^{\text{nr}}(\tilde{\sigma}) \left(\sum_{i=1}^{d_{L/K}-n} \nu_L(x_i) + \chi^{\text{nr}}(F_L) \sum_{i=d_{L/K}+1-n}^{d_{L/K}} \nu_L(x_i) \right) \\ \equiv \chi^{\text{nr}}(\tilde{\sigma})(\nu_L(x_1) + \dots + \nu_L(x_{d_{L/K}})) \pmod{p^{\omega_L}}, \end{aligned}$$

from which we deduce the $\text{Gal}(L/K)$ -invariance of w_L .

Next we show the exactness of the sequence. The first map is just the diagonal embedding.

Let $x = [x_1, \dots, x_{d_{L/K}}] \in \ker((F - 1) \times 1)$. Then

$$((F - 1) \times 1)(x) = \left[\frac{x_{d_{L/K}}^{F_L\chi^{\text{nr}}(F_L)}}{x_1}, \frac{x_1}{x_2}, \frac{x_2}{x_3}, \dots, \frac{x_{d_{L/K}-1}}{x_{d_{L/K}}} \right] = 1$$

implies that $x_1 = \dots = x_{d_{L/K}}$ and $x_1 = x_{d_{L/K}}^{F_L\chi^{\text{nr}}(F_L)} = x_1^{F_L\chi^{\text{nr}}(F_L)}$. It follows that the kernel of $(F - 1) \times 1$ is contained in the image of the first injection. Since the opposite inclusion is straightforward, we have exactness at the second non-trivial term of the sequence.

Let us consider exactness at the third non-trivial term. The following is just an adaptation of the arguments used to prove [BD13, Lemmas 11, 12].

Let $c = [c_1, \dots, c_{d_{L/K}}] \in \ker(w_L)$ for some $c_i \in \widehat{L}_0^\times(\chi^{\text{nr}})$. By assumption, $c_1 \dots c_{d_{L/K}}$ is in the kernel of ν_L , and hence by (22) there exists an element $y \in \widehat{L}_0^\times(\chi^{\text{nr}})$ such that $y\chi^{\text{nr}}(F_L)^{F_L-1} = c_1 \dots c_{d_{L/K}}$. Now we set

$$x = [c_2c_3 \dots c_{d_{L/K}}y, c_3 \dots c_{d_{L/K}}y, \dots, c_{d_{L/K}-1}c_{d_{L/K}}y, c_{d_{L/K}}y, y] \in \mathcal{I}_{L/K}(\chi^{\text{nr}}).$$

Then

$$\begin{aligned} ((F - 1) \times 1)(x) &= \left[\frac{x_{d_{L/K}}^{F_L\chi^{\text{nr}}(F_L)}}{x_1}, \frac{x_1}{x_2}, \frac{x_2}{x_3}, \dots, \frac{x_{d_{L/K}-1}}{x_{d_{L/K}}} \right] \\ &= \left[\frac{y^{F_L\chi^{\text{nr}}(F_L)}}{c_2c_3 \dots c_{d_{L/K}}y}, \frac{c_2c_3 \dots c_{d_{L/K}}y}{c_3c_4 \dots c_{d_{L/K}}y}, \frac{c_3c_4 \dots c_{d_{L/K}}y}{c_4c_5 \dots c_{d_{L/K}}y}, \dots, \frac{c_{d_{L/K}}y}{y} \right] \\ &= \left[\frac{y^{F_L\chi^{\text{nr}}(F_L)-1}}{c_2c_3 \dots c_{d_{L/K}}}, c_2, c_3, \dots, c_{d_{L/K}} \right] = [c_1, c_2, c_3, \dots, c_{d_{L/K}}]. \end{aligned}$$

The opposite inclusion is easy by the definition of ω_L . This proves exactness in the third term.

The surjectivity of the last map is clear. ■

We let $M/L/K$ be finite extensions such that both M/K and L/K are Galois. We have a canonical inclusion

$$\iota = \iota_{M/L}: \mathcal{I}_{L/K}(\chi^{\text{nr}}) \rightarrow \mathcal{I}_{M/K}(\chi^{\text{nr}}),$$

$$x \mapsto [x^{F_L^{d_{M/L}-1}} \chi^{\text{nr}}(F_L)^{d_{M/L}-1}, x^{F_L^{d_{M/L}-2}} \chi^{\text{nr}}(F_L)^{d_{M/L}-2}, \dots, x],$$

where $x = [x_1, \dots, x_{d_{L/K}}] \in (\widehat{L_0^\times})^{d_{L/K}}$ and the exponents of x have to be understood componentwise. There are three possibilities:

- (1) $\chi^{\text{nr}}(F_M) \neq 1$ and $\chi^{\text{nr}}(F_L) \neq 1$;
- (2) $\chi^{\text{nr}}(F_M) = 1$ and $\chi^{\text{nr}}(F_L) \neq 1$;
- (3) $\chi^{\text{nr}}(F_M) = 1$ and $\chi^{\text{nr}}(F_L) = 1$.

In case (1) we verify by a long but straightforward computation that the diagram of exact rows

$$\begin{array}{ccccccc} 0 \longrightarrow & (\widehat{L_0^\times}(\chi^{\text{nr}}))^{\text{Gal}(L_0/L)} & \longrightarrow & \mathcal{I}_{L/K}(\chi^{\text{nr}}) & \xrightarrow{(F-1)\times 1} & \mathcal{I}_{L/K}(\chi^{\text{nr}}) & \xrightarrow{w_L} & \mathbb{Z}/p^{\omega_L}\mathbb{Z}(\chi^{\text{nr}}) \longrightarrow 0 \\ & \downarrow \subseteq & & \downarrow \iota_{M/L} & & \downarrow \iota_{M/L} & & \downarrow \rho \\ 0 \longrightarrow & (\widehat{M_0^\times}(\chi^{\text{nr}}))^{\text{Gal}(M_0/M)} & \longrightarrow & \mathcal{I}_{M/K}(\chi^{\text{nr}}) & \xrightarrow{(F-1)\times 1} & \mathcal{I}_{M/K}(\chi^{\text{nr}}) & \xrightarrow{w_M} & \mathbb{Z}/p^{\omega_M}\mathbb{Z}(\chi^{\text{nr}}) \longrightarrow 0 \end{array}$$

commutes, where $\rho = e_{M/L} \sum_{i=0}^{d_{M/L}-1} F_L^i$.

In case (2) we have

$$\begin{aligned} w_M(\iota_{M/L}(x)) &= w_M(x) \sum_{i=0}^{d_{M/L}-1} \chi^{\text{nr}}(F_L)^i = w_M(x) \frac{\chi^{\text{nr}}(F_L)^{d_{M/L}} - 1}{\chi^{\text{nr}}(F_L) - 1} \\ &= w_M(x) \frac{\chi^{\text{nr}}(F_M) - 1}{\chi^{\text{nr}}(F_L) - 1} = 0. \end{aligned}$$

Therefore we obtain a commutative diagram

$$\begin{array}{ccccccc} 0 \longrightarrow & (\widehat{L_0^\times}(\chi^{\text{nr}}))^{\text{Gal}(L_0/L)} & \longrightarrow & \mathcal{I}_{L/K}(\chi^{\text{nr}}) & \xrightarrow{(F-1)\times 1} & \mathcal{I}_{L/K}(\chi^{\text{nr}}) & \xrightarrow{w_L} & \mathbb{Z}/p^{\omega_M}\mathbb{Z}(\chi^{\text{nr}}) \longrightarrow 0 \\ & \downarrow \subseteq & & \downarrow \iota_{M/L} & & \downarrow \iota_{M/L} & & \downarrow 0 \\ 0 \longrightarrow & (\widehat{M_0^\times}(\chi^{\text{nr}}))^{\text{Gal}(M_0/M)} & \longrightarrow & \mathcal{I}_{M/K}(\chi^{\text{nr}}) & \xrightarrow{(F-1)\times 1} & \mathcal{I}_{M/K}(\chi^{\text{nr}}) & \xrightarrow{w_M} & \mathbb{Z}_p(\chi^{\text{nr}}) \longrightarrow 0 \end{array}$$

Note that the bottom exact sequence comes from Theorem 4.2.1.

In case (3) our diagram takes the form

$$\begin{array}{ccccccc}
 0 \longrightarrow & \widehat{L}_0^\times(\chi^{\text{nr}})^{\text{Gal}(L_0/L)} & \longrightarrow & \mathcal{I}_{L/K}(\chi^{\text{nr}}) & \xrightarrow{(F-1)\times 1} & \mathcal{I}_{L/K}(\chi^{\text{nr}}) & \xrightarrow{w_L} \mathbb{Z}_p(\chi^{\text{nr}}) \longrightarrow 0 \\
 & \downarrow \subseteq & & \downarrow \iota_{M/L} & & \downarrow \iota_{M/L} & \downarrow [M:L] \\
 0 \longrightarrow & \widehat{M}_0^\times(\chi^{\text{nr}})^{\text{Gal}(M_0/M)} & \longrightarrow & \mathcal{I}_{M/K}(\chi^{\text{nr}}) & \xrightarrow{(F-1)\times 1} & \mathcal{I}_{M/K}(\chi^{\text{nr}}) & \xrightarrow{w_M} \mathbb{Z}_p(\chi^{\text{nr}}) \longrightarrow 0
 \end{array}$$

LEMMA 4.2.5. (a) *If $\chi^{\text{nr}}(F_L) \neq 1$ for every finite Galois extension L/K , then we have an exact sequence*

$$0 \rightarrow \varinjlim_L (\widehat{L}_0^\times(\chi^{\text{nr}}))^{\text{Gal}(L_0/L)} \rightarrow \varinjlim_L \mathcal{I}_{L/K}(\chi^{\text{nr}}) \xrightarrow{(F-1)\times 1} \varinjlim_L \mathcal{I}_{L/K}(\chi^{\text{nr}}) \rightarrow 0,$$

where the direct limit is taken over all finite Galois extensions L/K .

(b) *If $\chi^{\text{nr}}(F_L) = 1$ for some finite Galois extension L/K , then we have an exact sequence*

$$\begin{aligned}
 0 \rightarrow \varinjlim_L (\widehat{L}_0^\times(\chi^{\text{nr}}))^{\text{Gal}(L_0/L)} &\rightarrow \varinjlim_L \mathcal{I}_{L/K}(\chi^{\text{nr}}) \\
 &\xrightarrow{(F-1)\times 1} \varinjlim_L \mathcal{I}_{L/K}(\chi^{\text{nr}}) \rightarrow \mathbb{Q}_p(\chi^{\text{nr}}) \rightarrow 0.
 \end{aligned}$$

Proof. For the proof of (a) we use the diagram of case (1). Since the direct limit functor is exact, it suffices to show that $\varinjlim_L (\mathbb{Z}/p^{\omega_L}\mathbb{Z}(\chi^{\text{nr}}))$ is trivial.

We first note that

$$\begin{aligned}
 \omega_M &= v_p \left(\frac{1 - \chi^{\text{nr}}(F_M)}{1 - \chi^{\text{nr}}(F_L)} (1 - \chi^{\text{nr}}(F_L)) \right) \\
 &= \omega_L + v_p(1 + \chi^{\text{nr}}(F_L) + \dots + \chi^{\text{nr}}(F_L)^{d_{M/L}-1}).
 \end{aligned}$$

Let now $x \in \mathbb{Z}/p^{\omega_L}\mathbb{Z}(\chi^{\text{nr}})$. Then we let M/K be a finite Galois extension such that $L \subseteq M$ and $p^{\omega_L} | e_{M/L}$. So

$$e_{M/L}(1 + F_L + \dots + F_L^{d_{M/L}-1})x = e_{M/L}(1 + \chi^{\text{nr}}(F_L) + \dots + \chi^{\text{nr}}(F_L)^{d_{M/L}-1})x,$$

and since

$$\begin{aligned}
 v_p(e_{M/L}(1 + \chi^{\text{nr}}(F_L) + \dots + \chi^{\text{nr}}(F_L)^{d_{M/L}-1})) \\
 \geq \omega_L + v_p(1 + \chi^{\text{nr}}(F_L) + \dots + \chi^{\text{nr}}(F_L)^{d_{M/L}-1}) = \omega_M,
 \end{aligned}$$

we see that x becomes trivial in $\mathbb{Z}/p^{\omega_M}\mathbb{Z}(\chi^{\text{nr}})$.

The proof of (b) is achieved by taking the direct limit over the diagrams of cases (2) and (3). The result follows since it is straightforward to show that $\varinjlim_L \mathbb{Z}_p(\chi^{\text{nr}}) \cong \mathbb{Q}_p(\chi^{\text{nr}})$. ■

LEMMA 4.2.6. *We have*

$$(\mathcal{I}_{M/K}(\chi^{\text{nr}}))^{\text{Gal}(M/L)} = \iota_{M/L}(\mathcal{I}_{L/K}(\chi^{\text{nr}})).$$

Proof. For the proof we set $\tilde{d} := d_{M/L}$ and $d := d_{L/K}$. As usual, we identify $\mathcal{I}_{M/K}(\chi^{\text{nr}})$ and $(\widehat{M_0^\times})^{d\tilde{d}}$, the $\text{Gal}(K_0/K) \times \text{Gal}(M/K)$ -action being described by (19) and (20). Let us consider $[x_1, \dots, x_{\tilde{d}}] \in (\mathcal{I}_{M/K}(\chi^{\text{nr}}))^{\text{Gal}(M/L)}$ with $x_1, \dots, x_{\tilde{d}} \in (\widehat{M_0^\times})^d$.

Let $\tilde{\sigma} \in \text{Gal}(M_0/L_0)$ and $\sigma = \tilde{\sigma}|_M \in \text{Gal}(M/L)$. We have

$$(1 \times \sigma)[x_1, \dots, x_{\tilde{d}}] = [x_1^{\tilde{\sigma}\chi^{\text{nr}}(\tilde{\sigma})}, \dots, x_{\tilde{d}}^{\tilde{\sigma}\chi^{\text{nr}}(\tilde{\sigma})}].$$

Since $\tilde{\sigma}|_{K_0} = 1$, it follows that $\chi^{\text{nr}}(\tilde{\sigma}) = 1$. So from the above calculations, recalling that $[x_1, \dots, x_{\tilde{d}}]$ has to be fixed by the action of $1 \times \sigma$, we get $x_i^{\tilde{\sigma}} = x_i$ for $i = 1, \dots, \tilde{d}$. Since this holds for every $\tilde{\sigma} \in \text{Gal}(M_0/L_0)$, we conclude that $x_i \in (\widehat{L_0^\times})^d = \mathcal{I}_{L/K}(\chi^{\text{nr}})$ for $i = 1, \dots, \tilde{d}$.

Now let us take $\tilde{\tau} \in \text{Gal}(M_0/L)$ such that $\tilde{\tau}|_{L_0} = F_L$ and set $\tau := \tilde{\tau}|_M \in \text{Gal}(M/L)$. In particular the element $[x_1, \dots, x_{\tilde{d}}]$ must be fixed by the action of τ^{-1} . We have

$$\begin{aligned} [x_1, \dots, x_{\tilde{d}}] &= (1 \times \tau^{-1})[x_1, \dots, x_{\tilde{d}}] = (F^d \times 1)(F^{-d} \times \tau^{-1})[x_1, \dots, x_{\tilde{d}}] \\ &= (F^d \times 1)[x_1^{\tilde{\tau}^{-1}\chi^{\text{nr}}(\tilde{\tau})^{-1}}, \dots, x_{\tilde{d}}^{\tilde{\tau}^{-1}\chi^{\text{nr}}(\tilde{\tau})^{-1}}] \\ &= [x_{\tilde{d}}^{\tilde{\tau}^{-1}\chi^{\text{nr}}(\tilde{\tau})^{-1}F_M\chi^{\text{nr}}(F_M)}, x_1^{\tilde{\tau}^{-1}\chi^{\text{nr}}(\tilde{\tau})^{-1}}, \dots, x_{\tilde{d}-1}^{\tilde{\tau}^{-1}\chi^{\text{nr}}(\tilde{\tau})^{-1}}]. \end{aligned}$$

Since $x_i \in \mathcal{I}_{L/K}(\chi^{\text{nr}}) = (\widehat{L_0^\times})^d$ for $i = 1, \dots, \tilde{d}$, we can always replace $\tilde{\tau}$ by $\tilde{\tau}|_{L_0} = F_L$ and F_M by $F_M|_{L_0} = F_L^{\tilde{d}}$. Setting $x = x_{\tilde{d}}$ and comparing all the components in the above equality, except for the first one, we obtain

$$[x_1, \dots, x_{\tilde{d}}] = [x^{F_L^{\tilde{d}-1}\chi^{\text{nr}}(F_L)^{\tilde{d}-1}}, x^{F_L^{\tilde{d}-2}\chi^{\text{nr}}(F_L)^{\tilde{d}-2}}, \dots, x].$$

Thus we have proved that

$$[x_1, \dots, x_{\tilde{d}}] = \iota_{M/L}(x).$$

For the opposite inclusion we take $x \in \mathcal{I}_{L/K}(\chi^{\text{nr}})$ and have to show that $y := \iota_{M/L}(x)$ is fixed by all $\sigma \in \text{Gal}(M/L)$. As before, we let $\tilde{\tau} \in \text{Gal}(M_0/L)$ denote an element such that $\tilde{\tau}|_{L_0} = F_L$ and set $\tau := \tilde{\tau}|_M \in \text{Gal}(M/L)$. Then

$$\begin{aligned} (1 \times \tau^{-1})y &= (F^d \times 1)(F^{-d} \times \tau^{-1})[x^{F_L^{\tilde{d}-1}\chi^{\text{nr}}(F_L)^{\tilde{d}-1}}, x^{F_L^{\tilde{d}-2}\chi^{\text{nr}}(F_L)^{\tilde{d}-2}}, \dots, x] \\ &= (F^d \times 1)[x^{F_L^{\tilde{d}-2}\chi^{\text{nr}}(F_L)^{\tilde{d}-2}}, x^{F_L^{\tilde{d}-3}\chi^{\text{nr}}(F_L)^{\tilde{d}-3}}, \dots, x^{F_L^{-1}\chi^{\text{nr}}(F_L)^{-1}}] \\ &= [x^{F_L^{-1}\chi^{\text{nr}}(F_L)^{-1}F_M\chi^{\text{nr}}(F_M)}, x^{F_L^{\tilde{d}-2}\chi^{\text{nr}}(F_L)^{\tilde{d}-2}}, \dots, x] = \iota_{M/L}(x) = y. \end{aligned}$$

Let now $\sigma \in \text{Gal}(M/L)$ be arbitrary. Then there exists $i \in \mathbb{Z}$ such that $\sigma\tau^i|_{M \cap L_0} = \text{id}$. We then have $(1 \times \sigma\tau^i)y = y$ and the result follows from the above computation and $(1 \times \sigma) = (1 \times \sigma\tau^i)(1 \times \tau^{-i})$. ■

From Lemma 4.2.6 it is immediate that

$$(23) \quad \left(\varinjlim_L \mathcal{I}_{L/K}(\chi^{\text{nr}}) \right)^{G_N} = \mathcal{I}_{N/K}(\chi^{\text{nr}}),$$

where the direct limit is taken over all finite Galois extensions L/K . From now on, we will mainly be interested in the case $\chi^{\text{nr}}|_{G_N} \neq 1$, since the corresponding results for the case $\chi^{\text{nr}}|_{G_N} = 1$ are known. Under the assumption $\chi^{\text{nr}}|_{G_N} \neq 1$, we see that

$$(24) \quad (\mathbb{Q}_p(\chi^{\text{nr}}))^{G_N} = 0.$$

LEMMA 4.2.7. *The G_K -modules $\varinjlim_L \mathcal{I}_{L/K}(\chi^{\text{nr}})$ and $\mathbb{Q}_p(\chi^{\text{nr}})$ are acyclic with respect to the fixed point functor $A \mapsto A^{G_N}$.*

Proof. By [NSW08, Prop. 1.5.1], for $i > 0$ we have

$$H^i \left(N, \varinjlim_L \mathcal{I}_{L/K}(\chi^{\text{nr}}) \right) = \varinjlim_L H^i(\text{Gal}(L/N), \mathcal{I}_{L/K}(\chi^{\text{nr}})),$$

which is trivial by Lemma 4.1.2. The module $\mathbb{Q}_p(\chi^{\text{nr}})$ is cohomologically trivial since it is divisible (see [NSW08, Prop. 1.6.2]). ■

We can now state and prove the main result of this subsection.

THEOREM 4.2.8. *If $\chi^{\text{nr}}|_{G_N} \neq 1$, then the complex*

$$C_{N,\mathcal{F}}^\bullet := [\mathcal{I}_{N/K}(\chi^{\text{nr}}) \xrightarrow{(F-1) \times 1} \mathcal{I}_{N/K}(\chi^{\text{nr}})]$$

with non-trivial modules in degree 0 and 1 represents $R\Gamma(N, \mathcal{F})$.

Proof. We write X^\bullet for the standard resolution (as defined in [NSW08, Sec. I.2]) and I^\bullet for an injective resolution of $\varinjlim_L (\widehat{L}_0^\times(\chi^{\text{nr}}))^{\text{Gal}(L_0/L)}$. Let C^\bullet be the complex

$$\varinjlim_L \mathcal{I}_{L/K}(\chi^{\text{nr}}) \xrightarrow{(F-1) \times 1} \varinjlim_L \mathcal{I}_{L/K}(\chi^{\text{nr}}) \rightarrow \mathbb{Q}_p(\chi^{\text{nr}})$$

or the complex

$$\varinjlim_L \mathcal{I}_{L/K}(\chi^{\text{nr}}) \xrightarrow{(F-1) \times 1} \varinjlim_L \mathcal{I}_{L/K}(\chi^{\text{nr}}),$$

according to whether χ^{nr} becomes trivial for some G_L or not.

The G -modules $\text{Map}(G^n, \varinjlim_L (\widehat{L}_0^\times(\chi^{\text{nr}}))^{\text{Gal}(L_0/L)})$ appearing in X^\bullet are induced in the sense of [NSW08, Sec. I.3]. By [NSW08, Proposition 1.3.7] the standard resolution consists therefore of acyclic modules with respect to the fixed point functor. By [Lan02, Th. XX.6.2] there exists a morphism of complexes $X^\bullet \rightarrow I^\bullet$ inducing the identity on $\varinjlim_L (\widehat{L}_0^\times(\chi^{\text{nr}}))^{\text{Gal}(L_0/L)}$ and isomorphisms on cohomology. By Lemmas 4.2.5 and 4.2.7 we may apply [Lan02, Th. XX.6.2] once again and obtain a morphism of complexes $C^\bullet \rightarrow I^\bullet$, also

inducing the identity on $\varinjlim_L (\widehat{L}_0^\times(\chi^{\text{nr}}))^{\text{Gal}(L_0/L)}$ and isomorphisms on cohomology. By Lemma 4.1.1,

$$\left(\varinjlim_L (\widehat{L}_0^\times(\chi^{\text{nr}}))^{\text{Gal}(L_0/L)}\right)^{G_N} \cong (\widehat{N}_0^\times(\chi^{\text{nr}}))^{\text{Gal}(N_0/N)} \cong \mathcal{F}(\mathfrak{p}_N).$$

Therefore applying the G_N -fixed point functor together with (23) and (24) shows that $C_{N,\mathcal{F}}^\bullet = (C^\bullet)^{G_N}$ is quasi-isomorphic to $R\Gamma(N, \mathcal{F})$. ■

COROLLARY 4.2.9. *With the assumptions of Theorem 4.2.8 we have*

$$H^i(N, \mathcal{F}) = \begin{cases} \mathcal{F}(\mathfrak{p}_N) & \text{if } i = 0, \\ (\mathbb{Z}/p^\omega\mathbb{Z})(\chi^{\text{nr}}) & \text{if } i = 1, \\ 0 & \text{if } i \neq 0, 1. \end{cases}$$

Proof. This is an immediate consequence of Theorems 4.2.3 and 4.2.8. ■

In what follows, we always identify $R\Gamma(N, \mathcal{F})$ and $C_{N,\mathcal{F}}^\bullet$. We again set $G := \text{Gal}(N/K)$ and recall that a complex is said to be $\mathbb{Z}_p[G]$ -perfect if it is quasi-isomorphic to a bounded complex of finitely generated $\mathbb{Z}_p[G]$ -projectives.

COROLLARY 4.2.10. *If $\chi^{\text{nr}}|_{G_N} \neq 1$, then the complex $R\Gamma(N, \mathcal{F})$ is $\mathbb{Z}_p[G]$ -perfect. More precisely, there exists a complex*

$$P^\bullet := [P^{-1} \rightarrow P^0 \rightarrow P^1]$$

of finitely generated $\mathbb{Z}_p[G]$ -projectives together with a quasi-isomorphism $\eta: P^\bullet \rightarrow R\Gamma(N, \mathcal{F})$.

Proof. By [Lan02, Prop. XXI.1.1] and Theorem 4.2.8 there exists a quasi-isomorphism $f: K^\bullet \rightarrow R\Gamma(N, \mathcal{F})$, where $K^\bullet := [A \rightarrow B]$ is centered in degree 0 and 1 with B a $\mathbb{Z}_p[G]$ -projective module. As the cohomology groups $H^i(N, \mathcal{F})$ are finitely generated, the proof of [Lan02, Prop. XXI.1.1] actually shows that we can assume that A and B are both finitely generated. By our Lemma 4.1.2 the module $\mathcal{I}_{N/K}(\chi^{\text{nr}})$ is cohomologically trivial. We hence apply [Lan02, Prop. XXI.1.2] with \mathfrak{F} being the family of cohomologically trivial modules. We deduce that A is cohomologically trivial, and hence has a two-term resolution $0 \rightarrow P^{-1} \rightarrow P^0 \rightarrow A \rightarrow 0$ with finitely generated projective $\mathbb{Z}_p[G]$ -modules P^{-1} and P^0 . It follows that $R\Gamma(N, \mathcal{F})$ is quasi-isomorphic to $[P^{-1} \rightarrow P^0 \rightarrow P^1]$ with $P^1 = B$. ■

4.3. From $R\Gamma(N, \mathcal{F})$ to $R\Gamma(N, T)$. If C^\bullet is a complex and $p \in \mathbb{Z}$, then $C^\bullet[p]$ denotes the shifted complex, that is, $C^i[p] = C^{i+p}$. In this subsection we will show that the complex $P^\bullet[-1]$ of Corollary 4.2.10 is quasi-isomorphic to $R\Gamma(N, T)$. Together with Theorem 4.2.8 this will prove Theorem 2 whose formulation we recall once again.

THEOREM 4.3.1. *The complex*

$$C_{N,T}^\bullet := [\mathcal{I}_{N/K}(\chi^{\text{nr}}) \xrightarrow{(F-1)\times 1} \mathcal{I}_{N/K}(\chi^{\text{nr}})]$$

with non-trivial modules in degree 1 and 2 represents $R\Gamma(N, T)$.

REMARK 4.3.2. Theorem 4.3.1 combines with Theorem 4.2.3 to give a new proof of Lemma 3.2.1(a).

In the following we adapt the argument used to prove [Bre04a, Th. 4.20]. We will need the following lemma.

LEMMA 4.3.3. *Let $h: Y^\bullet \rightarrow W^\bullet$ be a map of cochain complexes. Suppose $h = 0$ in $D(\mathbb{Z}_p[G])$. Suppose also that Y^\bullet is bounded from above, comprising only projectives. Then h is homotopic to zero.*

Proof. It follows from [Wei94, Cor. 10.3.9] that there exists a quasi-isomorphism $t: Z^\bullet \rightarrow Y^\bullet$ such that $ht: Z^\bullet \rightarrow W^\bullet$ is null homotopic. By the dual of [Wei94, Lemma 10.4.6] there exists $s: Y^\bullet \rightarrow Z^\bullet$ such that ts is homotopy equivalent to the identity. Hence hts is homotopic to both 0 and h . ■

LEMMA 4.3.4. *There exist quasi-isomorphisms of complexes*

$$\varphi_n: P^\bullet/p^n \rightarrow R\Gamma(N, \mathcal{F}[p^n])[1]$$

which are compatible with the inverse systems.

Proof. The proof proceeds in two steps. First we construct morphisms of complexes φ_n , and then we modify the definition in such a way that the φ_n are compatible with the inverse systems.

By Kummer theory and [NSW08, Exercise I.3.1] we have a short exact sequence of complexes

$$0 \rightarrow R\Gamma(N, \mathcal{F}[p^n]) \rightarrow R\Gamma(N, \mathcal{F}) \xrightarrow{p^n} R\Gamma(N, \mathcal{F}) \rightarrow 0.$$

As a consequence of [Wei94, Ex. 10.4.9] we obtain an exact triangle

$$R\Gamma(N, \mathcal{F}[p^n]) \rightarrow R\Gamma(N, \mathcal{F}) \xrightarrow{p^n} R\Gamma(N, \mathcal{F}) \rightarrow .$$

Analogously we see that

$$P^\bullet \xrightarrow{p^n} P^\bullet \rightarrow P^\bullet/p^n \rightarrow$$

is an exact triangle.

Let $\eta: P^\bullet \rightarrow R\Gamma(N, \mathcal{F})$ be the quasi-isomorphism of Corollary 4.2.10. Then we obtain a commutative diagram in $D(\mathbb{Z}_p[G])$ of the following form:

$$\begin{array}{ccccccc} P^\bullet & \xrightarrow{p^n} & P^\bullet & \xrightarrow{\pi_n} & P^\bullet/p^n & \longrightarrow & \\ \downarrow \eta & & \downarrow \eta & & & & \\ R\Gamma(N, \mathcal{F}) & \xrightarrow{p^n} & R\Gamma(N, \mathcal{F}) & \xrightarrow{\lambda_n} & R\Gamma(N, \mathcal{F}[p^n])[1] & \longrightarrow & \end{array}$$

For clarity we use dotted arrows for morphisms in $D(\mathbb{Z}_p[G])$ which are not actual maps of complexes. As is common, we use $\xrightarrow{\sim}$ to indicate a quasi-isomorphism. By the definition of a morphism in the derived category there exists a complex Z^\bullet such that

$$R\Gamma(N, \mathcal{F}) \xrightarrow{\sim} Z^\bullet \rightarrow R\Gamma(N, \mathcal{F}[p^n])[1].$$

By [Mil80, Lemma VI.8.17] we construct a morphism of complexes

$$\psi_n : P^\bullet \rightarrow R\Gamma(N, \mathcal{F}[p^n])[1],$$

which makes the diagram

$$\begin{array}{ccccccc} P^\bullet & \xrightarrow{p^n} & P^\bullet & \xrightarrow{\pi_n} & P^\bullet/p^n & \longrightarrow & \\ \downarrow \eta & & \downarrow \eta & \searrow \psi_n & & & \\ R\Gamma(N, \mathcal{F}) & \xrightarrow{p^n} & R\Gamma(N, \mathcal{F}) & \cdots \xrightarrow{\lambda_n} & R\Gamma(N, \mathcal{F}[p^n])[1] & \longrightarrow & \end{array}$$

commutative in $D(\mathbb{Z}_p[G])$. As $R\Gamma(N, \mathcal{F}[p^n])$ is p^n -torsion, the map ψ_n factors through P^\bullet/p^n and we obtain morphisms φ_n of complexes such that

$$(25) \quad \begin{array}{ccccccc} P^\bullet & \xrightarrow{p^n} & P^\bullet & \xrightarrow{\pi_n} & P^\bullet/p^n & \longrightarrow & \\ \downarrow \eta & & \downarrow \eta & \searrow \psi_n & \downarrow \varphi_n & & \\ R\Gamma(N, \mathcal{F}) & \xrightarrow{p^n} & R\Gamma(N, \mathcal{F}) & \cdots \xrightarrow{\lambda_n} & R\Gamma(N, \mathcal{F}[p^n])[1] & \longrightarrow & \end{array}$$

commutes.

We want to show that (25) is actually a morphism of triangles. To that end we have to show that in

$$\begin{array}{ccccccc} P^\bullet & \xrightarrow{p^n} & P^\bullet & \xrightarrow{\pi_n} & P^\bullet/p^n & \xrightarrow{\omega} & P^\bullet[1] \\ \downarrow \eta & & \downarrow \eta & & \downarrow \varphi_n & & \downarrow \eta[1] \\ R\Gamma(N, \mathcal{F}) & \xrightarrow{p^n} & R\Gamma(N, \mathcal{F}) & \cdots \xrightarrow{\lambda_n} & R\Gamma(N, \mathcal{F}[p^n])[1] & \xrightarrow{\nu} & R\Gamma(N, \mathcal{F})[1] \end{array}$$

the right hand rectangle commutes. Since $P^\bullet \xrightarrow{\pi_n} P^\bullet/p^n$ is surjective, it suffices to show that $\eta[1] \circ \omega \circ \pi_n = \nu \circ \varphi_n \circ \pi_n$. By the commutativity of the central square this is equivalent to $\eta[1] \circ \omega \circ \pi_n = \nu \circ \lambda_n \circ \eta$ which holds true by [Wei94, Sec. 10.2, (TR3)].

We can therefore apply the 5-Lemma [Wei94, Exercise 10.2.2] and derive that the morphism φ_n is a quasi-isomorphism.

In a second step we now show that the maps φ_n can be chosen so that they are compatible with the inverse systems. We recall the identification $C_{N, \mathcal{F}}^\bullet = R\Gamma(N, \mathcal{F})$ and we set $Q_n^\bullet := R\Gamma(N, \mathcal{F}[p^n])[1]$. We write $\pi_n^Q : Q_n^\bullet \rightarrow Q_{n-1}^\bullet$ for the canonical map induced by the epimorphism $\mathcal{F}[p^n] \rightarrow \mathcal{F}[p^{n-1}]$.

We view P^i/p^n , Q_n^{i-1} and Q_{n-1}^{i-1} as $\mathbb{Z}/p^n\mathbb{Z}[G]$ -modules and observe that P^i/p^n is $\mathbb{Z}/p^n\mathbb{Z}[G]$ -projective. Hence there exists a lift \tilde{h}_n^i of h_n^i such that

$$\begin{array}{ccc} P^i/p^n & & \\ \downarrow \tilde{h}_n^i & \searrow h_n^i & \\ Q_n^{i-1} & \xrightarrow{\pi_n^{Q,i-1}} & Q_{n-1}^{i-1} \longrightarrow 0 \end{array}$$

commutes. We now replace each φ_n^i by

$$\varphi_n^i + d_{Q_n}^{i-1} \tilde{h}_n^i + \tilde{h}_n^{i+1} d_{P,n}^i.$$

Since we have only changed φ_n by a homotopy, diagram (25) still commutes in $D(\mathbb{Z}_p[G])$. By a straightforward computation we see that $\pi_n^Q \varphi_n = \varphi_{n-1} \pi_n^P$. This concludes the proof of the inductive step. ■

Proof of Theorem 4.3.1. From Lemma 4.3.4 we obtain a map $\varphi := \varprojlim_n \varphi_n$ of complexes,

$$\varphi: P^\bullet \rightarrow R\Gamma(N, T)[1].$$

Here we have naturally identified $\varprojlim_n P^\bullet/p^n$ with P^\bullet and $\varprojlim_n R\Gamma(N, \mathcal{F}[p^n])$ with $R\Gamma(N, T)$. Since each φ_n is a quasi-isomorphism, we obtain for each i an isomorphism

$$H^i(\varphi): \varprojlim_n H^i(P^\bullet/p^n) \rightarrow \varprojlim_n H^{i+1}(N, \mathcal{F}[p^n]).$$

By [BF98, Lemma 9] we conclude that $\varprojlim_n H^i(P^\bullet/p^n) = H^i(P^\bullet)$ and also $\varprojlim_n H^{i+1}(N, \mathcal{F}[p^n]) = H^{i+1}(N, T)$, so φ is a quasi-isomorphism. ■

5. The term U_{cris} . The term U_{cris} arises from the comparison of the trivialization λ used to define the refined Euler characteristic

$$C_{N/K} = -\chi_{\mathbb{Z}_p[G], B_{\text{dR}}[G]}(M^\bullet, \lambda^{-1})$$

in (15) and the trivialization $\beta'(N/K, V)$ defined in (12). In this section we will make this comparison explicit, and in the case $\chi^{\text{ur}}|_{G_N} = 1$ also compare U_{cris} to the generalization of Breuning’s correction term $M_{N/K}$.

5.1. The computation of U_{cris} . We define U_{cris} by the equation

$$(26) \quad \begin{aligned} i_{\mathbb{Z}_p[G], B_{\text{dR}}[G]}(\Delta_{\text{EP}}^{\text{na}}(N/K, T), \beta'(N/K, V)) \\ = C_{N/K} + \partial_{\mathbb{Z}_p[G], B_{\text{dR}}[G]}^1(t) + U_{\text{cris}}, \end{aligned}$$

where $i_{\mathbb{Z}_p[G], B_{\text{dR}}[G]}$ denotes the natural isomorphism from (2).

LEMMA 5.1.1. *Conjectures 3.1.1 and 3.2.4 are equivalent.*

Proof. Since $\dim_{\mathbb{Q}_p}(V) = 1$ for $V = \mathbb{Q}_p \otimes_{\mathbb{Z}_p} T$, the representation V is obviously potentially semistable. Conjecture 3.1.1 states that

$$i_{\mathbb{Z}_p[G], B_{\text{dR}}[G]}(\Delta_{\text{EP}}^{\text{na}}(N/K, T), \delta(N/K, V))$$

becomes 0 in $K_0(\tilde{A}, B_{\text{dR}}[G])$, while Conjecture 3.2.4 states that $\tilde{R}_{N/K}$ becomes 0 in $K_0(\tilde{A}, B_{\text{dR}}[G])$. Recalling the definition of $\tilde{R}_{N/K}$ and U_{cris} , Conjecture 3.2.4 is therefore equivalent to the assertion that

$$i_{\mathbb{Z}_p[G], B_{\text{dR}}[G]}(\Delta_{\text{EP}}^{\text{na}}(N/K, T), \beta'(N/K, V)) + \partial_{\mathbb{Z}_p[G], B_{\text{dR}}[G]}^1(\varepsilon_D(N/K, V))$$

becomes trivial. Recall that in our special situation we have $\Gamma^*(V) = 1$. Hence the definition of $\delta(N/K, V)$ in (13) implies

$$\delta(N/K, V) = \varepsilon_D(N/K, V)\beta'(N/K, V).$$

The equivalence of the two conjectures follows easily. ■

We proceed now to explicitly compute U_{cris} .

We write $H_e^1(N, V)$ and $H_f^1(N, V)$ for the exponential and finite part of $H^1(N, V)$ as defined by Bloch and Kato. For basic properties as used in our context we refer the reader to [BB08, §1.4].

For our calculation of U_{cris} we need to recall the fundamental exact sequences from [BB08, (1.1) and (1.2)]:

$$(27) \quad 0 \rightarrow H^0(N, V) \rightarrow D_{\text{cris}}^N(V) \xrightarrow{\phi=1} t_V(N) \xrightarrow{\text{exp}_V} H_e^1(N, V) \rightarrow 0$$

and

$$(28) \quad 0 \rightarrow H^0(N, V) \rightarrow D_{\text{cris}}^N(V) \xrightarrow{1-\phi} D_{\text{cris}}^N(V) \oplus t_V(N) \xrightarrow{\varepsilon_V} H_f^1(N, V) \rightarrow 0.$$

Note that $\varepsilon_V|_{t_V(N)} = \text{exp}_V$. We will also need the dual of (28) with $V^*(1)$ in place of V :

$$(29) \quad 0 \rightarrow H_f^1(N, V^*(1))^* \rightarrow D_{\text{cris}}^N(V^*(1))^* \oplus t_{V^*(1)}^*(N) \rightarrow D_{\text{cris}}^N(V^*(1))^* \rightarrow H^0(N, V^*(1))^* \rightarrow 0.$$

By local duality we have

$$(30) \quad H^0(N, V^*(1))^* \cong H^2(N, V), \quad H_f^1(N, V^*(1))^* \cong \frac{H^1(N, V)}{H_f^1(N, V)}.$$

Splicing together (28) and (29) we obtain the fundamental 7-term exact sequence (5) which induces the trivialization $\delta'(N/K, V)$ from (9).

Recall that $V = \mathbb{Q}_p(\chi^{\text{nr}})(1)$ and $V^*(1) = \mathbb{Q}_p((\chi^{\text{nr}})^{-1})$.

LEMMA 5.1.2. *We have:*

- (a) $t_{V^*(1)}(N) = 0$.
- (b) $\dim_{\mathbb{Q}_p} H_f^1(N, V^*(1)) = \begin{cases} 0 & \text{if } \chi^{\text{nr}}|_{G_N} \neq 1, \\ 1 & \text{if } \chi^{\text{nr}}|_{G_N} = 1. \end{cases}$

- (c) $H_f^1(N, V) = H_e^1(N, V) \cong \begin{cases} H^1(N, V) & \text{if } \chi^{\text{nr}}|_{G_N} \neq 1, \\ U_N^{(1)}(\chi^{\text{nr}}) \otimes_{\mathbb{Z}_p} \mathbb{Q}_p & \text{if } \chi^{\text{nr}}|_{G_N} = 1. \end{cases}$
- (d) $H^1(N, V) = \widehat{(N_0^\times(\chi^{\text{nr}}))}^{\text{Gal}(N_0/N)} \otimes_{\mathbb{Z}_p} \mathbb{Q}_p$. In particular, if $\chi^{\text{nr}}|_{G_N} = 1$, then $H^1(N, V) = \widehat{N^\times}(\chi^{\text{nr}}) \otimes_{\mathbb{Z}_p} \mathbb{Q}_p$.

Proof. (a) By [FO10, p. 148] or [IV16, p. 499] we have

$$t_{V^*(1)}(N) \hookrightarrow \mathbb{C}_p((\chi^{\text{nr}})^{-1})(-1)^{G_N} = 0,$$

and (a) follows.

(b) Using (29) and (a) we deduce

$$\dim_{\mathbb{Q}_p} H_f^1(N, V^*(1)) = \dim_{\mathbb{Q}_p} H^0(N, V^*(1)).$$

Clearly

$$H^0(N, V^*(1)) = \mathbb{Q}_p((\chi^{\text{nr}})^{-1})^{G_N} = \begin{cases} 0 & \text{if } \chi^{\text{nr}}|_{G_N} \neq 1, \\ \mathbb{Q}_p((\chi^{\text{nr}})^{-1}) & \text{if } \chi^{\text{nr}}|_{G_N} = 1. \end{cases}$$

(c) The proof of [IV16, Lemma 6.3] shows that $1-\phi: D_{\text{cris}}^N(V) \rightarrow D_{\text{cris}}^N(V)$ is an isomorphism. Hence $H_f^1(N, V) = H_e^1(N, V)$ is immediate from [BB08, Lemme 1.3].

From (30) we deduce

$$\dim_{\mathbb{Q}_p} H_f^1(N, V) + \dim_{\mathbb{Q}_p} H_f^1(N, V^*(1)) = \dim_{\mathbb{Q}_p} H^1(N, V).$$

Combining this with (b) we obtain the equality

$$H_f^1(N, V) = H^1(N, V)$$

in the case $\chi^{\text{nr}}|_{G_N} \neq 1$. For $\chi^{\text{nr}}|_{G_N} = 1$, the isomorphism

$$H_f^1(N, V) \cong U_N^{(1)}(\chi^{\text{nr}}) \otimes_{\mathbb{Z}_p} \mathbb{Q}_p$$

is well-known (see for example [BK90, Ex. 3.9]).

(d) Immediate by Lemma 3.2.1. ■

5.2. U_{cris} in the case $\chi^{\text{nr}}|_{G_N} \neq 1$. By Lemmas 5.1.2, 3.2.1(a) and the duality results (30), the 7-term exact sequence (5) degenerates to the two exact sequences

$$\begin{aligned} 0 \rightarrow D_{\text{cris}}^N(V) \xrightarrow{1-\phi} D_{\text{cris}}^N(V) \oplus t_V(N) \rightarrow H^1(N, V) \rightarrow 0, \\ 0 \rightarrow D_{\text{cris}}^N(V^*(1))^* \xrightarrow{1-\phi^*} D_{\text{cris}}^N(V^*(1))^* \rightarrow 0. \end{aligned}$$

By (27), Lemma 5.1.2 and the fact that $\varepsilon_V|_{t_V(N)} = \exp_V$, the first sequence is split by \exp_V^{-1} . Recall the definition of $i_{\mathbb{Z}_p[G], B_{\text{dR}}[G]}$ and the definition of the trivialization in (3). If we compare the trivializations $\kappa = \kappa(M^\bullet, \lambda^{-1})$ and $\beta'(N/K, V)$, then we see that (26) holds if and only if

$$(31) \quad U_{\text{cris}} = \partial_{\mathbb{Z}_p[G], \mathbb{Q}_p[G]}^1(\kappa \circ \beta'^{-1}) - \partial_{\mathbb{Z}_p[G], B_{\text{dR}}[G]}^1(t),$$

where we interpret $\kappa \circ \beta^{t-1} \in \pi_1(V(\mathbb{Q}_p[G]))$ as an element in $K_1(\mathbb{Q}_p[G])$. It follows that

$$(32) \quad U_{\text{cris}} = \partial_{\mathbb{Z}_p[G], \mathbb{Q}_p[G]}^1([D_{\text{cris}}^N(V), 1 - \phi]) - \partial_{\mathbb{Z}_p[G], \mathbb{Q}_p[G]}^1([D_{\text{cris}}^N(V^*(1))^*, 1 - \phi^*]).$$

Note that the sign comes from the fact that the first term in the first sequence is in odd degree, whereas the first term in the second sequence is in even degree.

We introduce the following notation. If $x \in Z(\mathbb{Q}_p[G])$ we let $*x \in Z(\mathbb{Q}_p[G])^\times$ denote the invertible element which on the Wedderburn decomposition $Z(\mathbb{Q}_p[G]) = \bigoplus_{i=1}^r F_i$ for suitable finite extensions F_i/\mathbb{Q}_p is given by $(*x_i)$ with $*x_i = 1$ if $x_i = 0$ and $*x_i = x_i$ otherwise.

We now compute U_{cris} explicitly. Recall that $F = F_K = \varphi^{d_K}$.

LEMMA 5.2.1. *If $\chi^{\text{nr}}|_{G_N} \neq 1$, the endomorphism $1 - \phi^*$ of $D_{\text{cris}}^N(V^*(1))^*$ is an isomorphism. Furthermore,*

$$\partial_{\mathbb{Z}_p[G], \mathbb{Q}_p[G]}^1([D_{\text{cris}}^N(V^*(1))^*, 1 - \phi^*]) = \hat{\delta}_{\mathbb{Z}_p[G], \mathbb{Q}_p[G]}^1(*((1 - u^{d_K} F^{-1})e_I))$$

in $K_0(\mathbb{Z}_p[G], \mathbb{Q}_p[G])$.

Proof. We let v^* be a \mathbb{Q}_p -basis of $V^*(1)$. In particular, $\sigma v^* = \chi^{\text{nr}}(\sigma)^{-1} v^*$ for all $\sigma \in G_K$.

We consider the \mathbb{Z}_p -module $(\overline{\mathbb{Z}_p^{\text{nr}}}(\chi_{\mathbb{Q}_p}^{\text{nr}}))^{G_{\mathbb{Q}_p}}$. For $s \in \overline{\mathbb{Z}_p^{\text{nr}}}$ one has

$$s \in (\overline{\mathbb{Z}_p^{\text{nr}}}(\chi_{\mathbb{Q}_p}^{\text{nr}}))^{G_{\mathbb{Q}_p}} \Leftrightarrow u\varphi(s) - s = 0.$$

Adapting the proof of [Neu92, Lemma V.2.1] we construct $t^{\text{nr}} \in \overline{\mathbb{Z}_p^{\text{nr}}}$ such that $(t^{\text{nr}})^{p-1} \equiv u^{-1} \pmod{p}$ and $u\varphi(t^{\text{nr}}) = t^{\text{nr}}$. It follows that $t^{\text{nr}} \in \overline{\mathbb{Z}_p^{\text{nr}}}^\times$, and furthermore it is easy to show that $(\overline{\mathbb{Z}_p^{\text{nr}}}(\chi^{\text{nr}}))^{G_{\mathbb{Q}_p}} = \mathbb{Z}_p t^{\text{nr}}$. Note that t^{nr} is unique up to multiplication by units of \mathbb{Z}_p .

By [FO10, Th. 6.14] we know that $\overline{\mathbb{Z}_p^{\text{nr}}} \subseteq B_{\text{cris}}$, so that the element

$$e_0^* := (t^{\text{nr}})^{-1} \otimes v^*$$

is contained in $B_{\text{cris}} \otimes_{\mathbb{Q}_p} V^*(1)$. Since $\sigma(t^{\text{nr}}) = \chi_{\mathbb{Q}_p}^{\text{nr}}(\sigma)^{-1} t^{\text{nr}}$ for all $\sigma \in G_{\mathbb{Q}_p}$, we derive that $e_0^* \in (B_{\text{cris}} \otimes_{\mathbb{Q}_p} V^*(1))^{G_K} \subseteq D_{\text{cris}}^N(V^*(1))$. Let $N_1 \subseteq N$ denote the maximal unramified subextension of N/\mathbb{Q}_p . So we have $[N_1 : \mathbb{Q}_p] = d_N$. Since $\dim_{N_1} D_{\text{cris}}^N(V^*(1)) \leq \dim_{\mathbb{Q}_p}(V^*(1)) = 1$, the element e_0^* generates $D_{\text{cris}}^N(V^*(1))$ as an N_1 -vector space, i.e., $D_{\text{cris}}^N(V^*(1)) = N_1 e_0^*$. By the φ -semilinearity of ϕ we obtain $\phi(e_0^*) = \chi_{\mathbb{Q}_p}^{\text{nr}}(\varphi) e_0^* = u e_0^*$.

We now fix a normal basis element θ of N_1/\mathbb{Q}_p . Then $w_j := \varphi^{-j} \theta e_0^*$, $j = 0, \dots, d_K - 1$, is a $\mathbb{Q}_p[G/I]$ -basis of $D_{\text{cris}}^N(V^*(1))$. Let $\psi_l \in D_{\text{cris}}^N(V^*(1))^*$,

$l = 0, \dots, d_K - 1$, be the dual basis. Then

$$D_{\text{cris}}^N(V^*(1))^* = \bigoplus_{l=0}^{d_K-1} \mathbb{Q}_p[G/I]\psi_l$$

and for $0 < j < d_K$ we have

$$\phi^*(\psi_l)(F^i w_j) = \psi_l(\phi(F^i \varphi^{-j} \theta e_0^*)) = \psi_l(u F^i \varphi^{-j+1} \theta e_0^*) = u \psi_l(F^i w_{j-1}),$$

which is u for $i = 0$ and $l = j - 1$, and 0 otherwise. For $j = 0$ we get

$$\phi^*(\psi_l)(F^i w_0) = \psi_l(\phi(F^i \theta e_0^*)) = \psi_l(u F^{i+1} \varphi^{-d_K+1} \theta e_0^*) = u \psi_l(F^{i+1} w_{d_K-1}),$$

which is u for $i = -1$ and $l = d_K - 1$, and 0 otherwise. Hence $\phi^*(\psi_l) = u \psi_{l+1}$ for $0 \leq l < d_K - 1$ and $\phi^*(\psi_{d_K-1}) = u F^{-1} \psi_0$.

With respect to the $\mathbb{Q}_p[G/I]$ -basis $\psi_0, \dots, \psi_{d_K-1}$ the matrix associated to $1 - \phi^*$ is given by

$$A = \begin{pmatrix} 1 & 0 & 0 & \cdots & 0 & -u F^{-1} \\ -u & 1 & 0 & \cdots & 0 & 0 \\ 0 & -u & 1 & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & \cdots & 1 & 0 \\ 0 & 0 & 0 & \cdots & -u & 1 \end{pmatrix}.$$

Since $\det(A) = 1 - u^{d_K} F^{-1}$ the lemma follows. Note that $\det(A)$ (and hence also $1 - \phi^*$) is invertible if and only if $\chi^{\text{nr}}|_{G_N} \neq 1$. ■

LEMMA 5.2.2. *The endomorphism $1 - \phi$ of $D_{\text{cris}}^N(V)$ is an isomorphism and we have*

$$\partial_{\mathbb{Z}_p[G], \mathbb{Q}_p[G]}^1([D_{\text{cris}}^N(V), 1 - \phi]) = \hat{\partial}_{\mathbb{Z}_p[G], \mathbb{Q}_p[G]}^1(*((1 - p^{-d_K} u^{-d_K} F) e_I))$$

in $K_0(\mathbb{Z}_p[G], \mathbb{Q}_p[G])$.

Proof. The proof is analogous to the previous lemma, following the computations in [IV16, Lemma 6.3]. ■

5.3. U_{cris} in the case $\chi^{\text{nr}}|_{G_N} = 1$. By Lemmas 5.1.2, 5.2.2 (note that the latter holds also for $\chi^{\text{nr}}|_{G_N} = 1$), (27)–(30) and Lemma 3.2.1(b), the 7-term exact sequence (5) degenerates to

$$(33) \quad 0 \rightarrow D_{\text{cris}}^N(V) \xrightarrow{1-\phi} D_{\text{cris}}^N(V) \rightarrow 0,$$

$$(34) \quad 0 \rightarrow t_V(N) \xrightarrow{\text{exp}_V} H^1(N, V) \rightarrow \frac{H^1(N, V)}{H_f^1(N, V)} \rightarrow 0$$

and

$$(35) \quad 0 \rightarrow \frac{H^1(N, V)}{H_f^1(N, V)} \rightarrow D_{\text{cris}}^N(V^*(1))^* \xrightarrow{1-\phi^*} D_{\text{cris}}^N(V^*(1))^* \rightarrow H^2(N, V) \rightarrow 0.$$

In the case $\chi^{\text{nr}}|_{G_N} = 1$, taking cohomology commutes with twisting, so that we may and will assume that $V = \mathbb{Q}_p(1)$ (see Remark 5.3.3 below). By [BK90, Ex. 3.10.1] and Lemma 5.1.2, the short exact sequence (34) fits into the diagram

$$(36) \quad \begin{array}{ccccccc} 0 & \longrightarrow & t_V(N) & \xrightarrow{\text{exp}_V} & H^1(N, V) & \longrightarrow & \frac{H^1(N, V)}{H_f^1(N, V)} \longrightarrow 0 \\ & & \downarrow & & \downarrow \partial_{Ku}^{-1} & & \downarrow \\ 0 & \longrightarrow & N & \xrightarrow{\text{exp}} & \widehat{N^\times} \otimes_{\mathbb{Z}_p} \mathbb{Q}_p & \longrightarrow & \frac{\widehat{N^\times} \otimes_{\mathbb{Z}_p} \mathbb{Q}_p}{U_N^{(1)} \otimes_{\mathbb{Z}_p} \mathbb{Q}_p} \longrightarrow 0 \\ & & \downarrow = & & \downarrow = & & \downarrow \nu_N \\ 0 & \longrightarrow & N & \xrightarrow{\text{exp}} & \widehat{N^\times} \otimes_{\mathbb{Z}_p} \mathbb{Q}_p & \xrightarrow{\nu_N} & \mathbb{Q}_p \longrightarrow 0 \end{array}$$

where exp is induced by the p -adic exponential map and ∂_{Ku} denotes the connecting homomorphism induced by the Kummer sequence. We have a morphism γ_1 in $V(\mathbb{Q}_p[G])$:

$$\begin{aligned} \gamma_1: \left[\frac{H^1(N, V)}{H_f^1(N, V)} \right]_{\mathbb{Q}_p[G]}^{-1} &\otimes [H^2(N, V)]_{\mathbb{Q}_p[G]} \\ &\rightarrow \left[\frac{H^1(N, V)}{H_f^1(N, V)} \right]_{\mathbb{Q}_p[G]}^{-1} \otimes [D_{\text{cris}}^N(V^*(1))^*]_{\mathbb{Q}_p[G]} \\ &\quad \otimes [D_{\text{cris}}^N(V^*(1))^*]_{\mathbb{Q}_p[G]}^{-1} \otimes [H^2(N, V)]_{\mathbb{Q}_p[G]} \\ &\rightarrow \mathbf{1}_{V(\mathbb{Q}_p[G])}, \end{aligned}$$

where the last arrow is induced by the exact sequence (35). From Lemma 5.1.2 together with the isomorphism $\text{inv}_N: H^2(N, V) \rightarrow \mathbb{Q}_p$ we deduce a morphism γ_2 in $V(\mathbb{Q}_p[G])$:

$$\gamma_2: \left[\frac{H^1(N, V)}{H_f^1(N, V)} \right]_{\mathbb{Q}_p[G]}^{-1} \otimes [H^2(N, V)]_{\mathbb{Q}_p[G]} \rightarrow \mathbf{1}_{V(\mathbb{Q}_p[G])},$$

which is induced by the valuation map ν_N . We set

$$\gamma := \gamma_2 \circ \gamma_1^{-1} \in \pi_1(V(\mathbb{Q}_p[G])) \cong K_1(\mathbb{Q}_p[G]).$$

The comparison of the trivializations $\kappa = \kappa(M^\bullet, \lambda^{-1})$ and $\beta'(N/K, V)$ shows

(as explained in some more detail in the Appendix) that

$$(37) \quad U_{\text{cris}} = \partial_{\mathbb{Z}_p[G], \mathbb{Q}_p[G]}^1([D_{\text{cris}}^N(V), 1 - \phi]) + \partial_{\mathbb{Z}_p[G], \mathbb{Q}_p[G]}^1(\gamma).$$

We now proceed to compute U_{cris} explicitly. The first summand is computed by Lemma 5.2.2. For non-trivial characters the same computation as in the proof of Lemma 5.2.1 shows that

$$\begin{aligned} \partial_{\mathbb{Z}_p[G], \mathbb{Q}_p[G]}^1([(1 - e_G)D_{\text{cris}}^N(V^*(1))^*, 1 - \phi^*]) \\ = \hat{\partial}_{\mathbb{Z}_p[G], \mathbb{Q}_p[G]}^1(*((1 - F^{-1})(e_I - e_G))) = \hat{\partial}_{\mathbb{Z}_p[G], \mathbb{Q}_p[G]}^1(*((1 - F^{-1})e_I)). \end{aligned}$$

It therefore remains to compute the trivial component of γ . For this we will need some preparations.

We recall that the exact sequence (35) is derived by dualizing and then using local duality from

$$(38) \quad 0 \rightarrow H^0(N, V^*(1)) \xrightarrow{\iota} D_{\text{cris}}^N(V^*(1)) \xrightarrow{1-\phi} D_{\text{cris}}^N(V^*(1)) \rightarrow H_f^1(N, V^*(1)) \rightarrow 0,$$

which is (27) with V replaced by $V^*(1)$. Recall that N_1 is the maximal unramified subextension of N/\mathbb{Q}_p . The sequence (38) is explicitly given by

$$0 \rightarrow \mathbb{Q}_p \xrightarrow{\iota} N_1 \xrightarrow{1-\varphi} N_1 \xrightarrow{\xi} H_f^1(N, V^*(1)) \rightarrow 0,$$

where ξ is the connecting homomorphism. Explicitly, for $x \in N_1$ and $\sigma \in G_N$ we have $\xi_x(\sigma) = \sigma(y) - y$, if $y \in N_0$ is such that $(1 - \varphi)(y) = x$. By [BK90, Ex. 3.9], $H_f^1(N, V^*(1)) \subseteq \text{Hom}_{\text{cont}}(G_N, \mathbb{Q}_p) = H^1(N, V^*(1))$ are the unramified homomorphisms, hence

$$H_f^1(N, V^*(1)) = \text{Hom}_{\text{cont}}(\overline{\langle F_N \rangle}, \mathbb{Q}_p).$$

We will always identify $H_f^1(N, V^*(1))$ with \mathbb{Q}_p via

$$\mu: H_f^1(N, V^*(1)) \rightarrow \mathbb{Q}_p, \quad \eta \mapsto \eta(F_N).$$

Recall that $\mathcal{T}_{N_1/\mathbb{Q}_p}$ denotes the trace map. Since

$$\begin{aligned} \xi_x(F_N) &= F_N(y) - y = \varphi^{d_N}(y) - y \\ &= \sum_{i=0}^{d_N-1} (\varphi^{i+1}(y) - \varphi^i(y)) = - \sum_{i=0}^{d_N-1} \varphi^i(x) = -\mathcal{T}_{N_1/\mathbb{Q}_p}(x), \end{aligned}$$

we have the commutative diagram

$$(39) \quad \begin{array}{ccccccccc} 0 & \longrightarrow & \mathbb{Q}_p & \xrightarrow{\iota} & N_1 & \xrightarrow{1-\varphi} & N_1 & \xrightarrow{\xi} & H_f^1(N, V^*(1)) & \longrightarrow & 0 \\ & & \downarrow = & & \downarrow = & & \downarrow = & & \downarrow \mu & & \\ 0 & \longrightarrow & \mathbb{Q}_p & \xrightarrow{\iota} & N_1 & \xrightarrow{1-\varphi} & N_1 & \xrightarrow{-\mathcal{T}_{N_1/\mathbb{Q}_p}} & \mathbb{Q}_p & \longrightarrow & 0 \end{array}$$

By dualizing we obtain

$$\begin{array}{ccccccccc}
 0 & \longrightarrow & \mathbb{Q}_p^* & \xrightarrow{(-\mathcal{T}_{N_1/\mathbb{Q}_p})^*} & N_1^* & \xrightarrow{(1-\varphi)^*} & N_1^* & \xrightarrow{\iota^*} & \mathbb{Q}_p^* & \longrightarrow & 0 \\
 & & \downarrow \mu^* & & \downarrow = & & \downarrow = & & \downarrow = & & \\
 0 & \longrightarrow & H_f^1(N, V^*(1))^* & \xrightarrow{\xi^*} & N_1^* & \xrightarrow{(1-\varphi)^*} & N_1^* & \xrightarrow{\iota^*} & \mathbb{Q}_p^* & \longrightarrow & 0 \\
 (40) & & \downarrow s_1 & & & & & & \downarrow s_2 & & \\
 & & H^1(N, V)/H_f^1(N, V) & & & & & & H^2(N, V) & & \\
 & & \downarrow \cong & & & & & & \downarrow \text{inv}_N & & \\
 & & \frac{\widehat{N^\times} \otimes \mathbb{Q}_p}{U_N^{(1)} \otimes \mathbb{Q}_p} & \xrightarrow{\nu_N} & & & & & \mathbb{Q}_p & &
 \end{array}$$

where s_1 and s_2 denote the duality maps from (30).

LEMMA 5.3.1.

- (a) Let $\eta_1 \in \mathbb{Q}_p^* = \text{Hom}_{\mathbb{Q}_p}(\mathbb{Q}_p, \mathbb{Q}_p)$ be defined by $\eta_1(1) = 1$. Then $s_1(\mu^*(\eta_1)) = [\pi_N]$ where $[\pi_N]$ denotes the class of π_N in $\frac{\widehat{N^\times} \otimes \mathbb{Q}_p}{U_N^{(1)} \otimes \mathbb{Q}_p}$.
- (b) Let $\eta_2 \in \mathbb{Q}_p^*$ be defined by $\eta_2(1) = 1$. Then $\text{inv}_N(s_2(\eta_2)) = 1$.

Proof. (a) We want to prove that $\langle \pi_N, \cdot \rangle = \mu^*(\eta_1)$, where

$$(41) \quad \langle \cdot, \cdot \rangle : H^1(N, V) \times H^1(N, V^*(1)) \rightarrow H^2(N, \mathbb{Q}_p(1)) \xrightarrow{\text{inv}_N} \mathbb{Q}_p$$

denotes the duality pairing. As in the proof of Lemma 5.2.1 we let $\theta \in N_1$ be a normal basis element. In addition, we require the property $\mathcal{T}_{N_1/\mathbb{Q}_p}(\theta) = -1$. Since $H_f^1(N, V^*(1)) = \mathbb{Q}_p \xi_\theta$, it is enough to prove that $\langle \pi_N, \xi_\theta \rangle = \mu^*(\eta_1)(\xi_\theta)$. By the defining property of θ we obtain

$$\mu^*(\eta_1)(\xi_\theta) = \eta_1(\mu(\xi_\theta)) = \eta_1(-\mathcal{T}_{N_1/\mathbb{Q}_p}(\theta)) = \eta_1(1) = 1.$$

By [NSW08, Cor. 7.2.13] we obtain

$$\langle \pi_N, \xi_\theta \rangle = \text{inv}_N(\pi_N \cup \xi_\theta) = \xi_\theta(\text{rec}_N(\pi_N)) = \xi_\theta(F_N) = -\mathcal{T}_{N_1/\mathbb{Q}_p}(\theta) = 1.$$

(b) If

$$\langle \cdot, \cdot \rangle : H^2(N, V) \times H^0(N, V^*(1)) \rightarrow H^2(N, \mathbb{Q}_p(1)) \xrightarrow{\text{inv}_N} \mathbb{Q}_p$$

denotes the local duality pairing, then by definition $\langle s_2(\eta_2), \cdot \rangle = \eta_2$, and therefore

$$\text{inv}_N(s_2(\eta_2)) = \langle s_2(\eta_2), 1 \rangle = \eta_2(1) = 1. \quad \blacksquare$$

As explained in [BF01, Sec. 2.5], we can and will identify here $V(\mathbb{Q}_p)$ with the category of graded invertible modules with the commutativity constraint

ψ as in [BF01, (5)]. Explicitly, for objects (L, α) , (M, β) and $l \in L$, $m \in M$ one has

$$\psi(l \otimes m) = (-1)^{\alpha\beta} m \otimes l.$$

We also point out that for an object (L, α) we fix a right inverse by setting $(L, \alpha)^{-1} := (\text{Hom}_{\mathbb{Q}_p}(L, \mathbb{Q}_p), -\alpha)$, i.e. $(L, \alpha) \otimes (L, \alpha)^{-1} = (\mathbb{Q}_p, 0)$.

LEMMA 5.3.2. *Under our assumption that $\chi^{\text{nr}} = 1$,*

$$\partial_{\mathbb{Z}_p[G], \mathbb{Q}_p[G]}^1(\gamma) = \frac{*(d_{N/K}e_G)}{*((1 - F^{-1})e_I)}.$$

Proof. As already mentioned, the computation of the $(1 - e_G)$ -component is analogous to the proof of Lemma 5.2.1.

For the computation of the trivial component, by Lemma 5.3.1 we can identify $\eta_1^{-1} \otimes \eta_2$ with $[\pi_N]^{-1} \otimes 1$, so that we need to compute $\gamma_1(\eta_1^{-1} \otimes \eta_2)$ and $\gamma_2([\pi_N]^{-1} \otimes 1)$, where by abuse of notation we have denoted the e_G -component of γ_1 and γ_2 again by the same symbol.

Using the commutativity constraint we obtain

$$(42) \quad \gamma_2([\pi_N]^{-1} \otimes 1) = -1.$$

For the computation of the trivial component it remains by Lemma 5.3.1 to compute the image of $\eta_1^{-1} \otimes \eta_2$ under the morphism γ_1 . By (42) we have to show

$$\gamma_1(\eta_1^{-1} \otimes \eta_2) = -\frac{1}{d_{N/K}}.$$

The e_G -component of the top sequence in diagram (40) is given by

$$0 \rightarrow \mathbb{Q}_p^* \xrightarrow{(-\mathcal{T}_{N_1/\mathbb{Q}_p})^*} e_G N_1^* \xrightarrow{(1-\varphi)^*} e_G N_1^* \xrightarrow{\iota^*} \mathbb{Q}_p^* \rightarrow 0.$$

The elements $e_G \psi_0, \dots, e_G \psi_{d_K-1}$ constitute a \mathbb{Q}_p -basis of $e_G N_1^*$, where $\psi_1, \dots, \psi_{d_K-1}$ are defined as in the proof of Lemma 5.2.1. We set

$$\psi'_i := e_G \psi_i, \quad i = 0, \dots, d_K - 1.$$

Since $(1 - \varphi)^*(\psi'_i) = \psi'_i - \psi'_{i+1}$, we set $z_i := \psi'_i - \psi'_{i+1}$, so that

$$Z := \langle z_0, \dots, z_{d_K-2} \rangle_{\mathbb{Q}_p}$$

is the image of $(1 - \varphi)^*$. We consider the two short exact sequences

$$(43) \quad 0 \rightarrow \mathbb{Q}_p^* \xrightarrow{(-\mathcal{T}_{N_1/\mathbb{Q}_p})^*} e_G N_1^* \xrightarrow{(1-\varphi)^*} Z \rightarrow 0,$$

$$(44) \quad 0 \rightarrow Z \xrightarrow{\subseteq} e_G N_1^* \xrightarrow{\iota^*} \mathbb{Q}_p^* \rightarrow 0$$

and compute the image of $\eta_1^{-1} \otimes \psi' \otimes (\psi')^{-1} \otimes \eta_2$ with $\psi' := \psi'_0 \wedge \dots \wedge \psi'_{d_K-1}$ under the natural map

$$[\mathbb{Q}_p^*]_{\mathbb{Q}_p}^{-1} \otimes [e_G N_1^*]_{\mathbb{Q}_p} \otimes [e_G N_1^*]_{\mathbb{Q}_p}^{-1} \otimes [\mathbb{Q}_p^*]_{\mathbb{Q}_p} \rightarrow \mathbf{1}_{V(\mathbb{Q}_p)},$$

which is induced by the top sequence of (40).

We express $(-\mathcal{T}_{N_1/\mathbb{Q}_p})^*(\eta_1)$ in terms of the \mathbb{Q}_p -basis $\psi_0, \dots, \psi_{d_N-1}$ of N_1^* , which is dual to the basis w_0, \dots, w_{d_N-1} of N_1 defined by $w_j = \varphi^{-j}\theta$. As usual, θ is a normal basis element with the additional property that $\mathcal{T}_{N_1/\mathbb{Q}_p}(\theta) = -1$. One has

$$(-\mathcal{T}_{N_1/\mathbb{Q}_p})^*(\eta_1)(w_i) = \eta_1(-\mathcal{T}_{N_1/\mathbb{Q}_p}(w_i)) = \eta_1(1) = 1$$

for all $i = 0, \dots, d_N - 1$. We set $b_1 := (-\mathcal{T}_{N_1/\mathbb{Q}_p})^*(\eta_1)$ and obtain

$$b_1 = \sum_{i=0}^{d_N-1} \psi_i = \left(\sum_{j=0}^{d_{N/K}-1} \varphi^{d_K j} \right) \sum_{i=0}^{d_K-1} \psi_i = d_{N/K} \sum_{i=0}^{d_K-1} \psi'_i.$$

Hence $(b_1, \psi'_0, \dots, \psi'_{d_K-2}) = (\psi'_0, \dots, \psi'_{d_K-1})A$ with the matrix $A \in \text{Gl}_{d_K}(\mathbb{Q}_p)$ given by

$$A = \begin{pmatrix} d_{N/K} & 1 & 0 & \cdots & 0 \\ d_{N/K} & 0 & 1 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ d_{N/K} & 0 & 0 & \cdots & 1 \\ d_{N/K} & 0 & 0 & \cdots & 0 \end{pmatrix}.$$

A splitting τ of $(1 - \varphi)^*$ is given by $\tau(z_i) := \psi'_i, i = 0, \dots, d_K - 2$. Therefore, under the isomorphism

$$[e_G N_1^*]_{\mathbb{Q}_p} \cong [\mathbb{Q}_p^*]_{\mathbb{Q}_p} \otimes [Z]_{\mathbb{Q}_p}$$

induced by (43) we obtain

$$\psi' \mapsto \det(A)^{-1}(\eta_1 \otimes z),$$

where $z := z_0 \wedge \cdots \wedge z_{d_K-2}$.

We define

$$\sigma: \mathbb{Q}_p^* \rightarrow N_1^*, \quad \eta_2 \mapsto -\frac{1}{d_N} \sum_{i=0}^{d_N-1} \psi_i.$$

Then σ is a splitting of ι^* because

$$\iota^*(\sigma(\eta_2))(1) = -\frac{1}{d_N} \sum_{i=0}^{d_N-1} \psi_i(1) = \frac{1}{d_N} \sum_{i=0}^{d_N-1} \psi_i \left(\sum_{j=0}^{d_N-1} w_j \right) = 1.$$

We set $b_2 := \sigma(\eta_2)$. Then

$$b_2 = -\frac{1}{d_N} \sum_{i=0}^{d_N-1} \psi_i = -\frac{1}{d_K} \sum_{i=0}^{d_K-1} \psi'_i.$$

Hence $(z_0, \dots, z_{d_K-2}, b_2) = (\psi'_0, \dots, \psi'_{d_K-1})B$ with the matrix $B \in \text{Gl}_{d_K}(\mathbb{Q}_p)$

given by

$$B = \begin{pmatrix} 1 & 0 & \cdots & 0 & 0 & -1/d_K \\ -1 & 1 & \cdots & 0 & 0 & -1/d_K \\ 0 & -1 & \cdots & 0 & 0 & -1/d_K \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots \\ 0 & 0 & \cdots & -1 & 1 & -1/d_K \\ 0 & 0 & \cdots & 0 & -1 & -1/d_K \end{pmatrix}.$$

Therefore

$$\psi' \mapsto \det(B)^{-1}(z \otimes \eta_2)$$

under the isomorphism induced by (44), resp. $(\psi')^{-1} \mapsto \det(B)(\eta_2^{-1} \otimes z^{-1})$. Note that

$$\det(A) = (-1)^{d_K+1}d_{N/K}, \quad \det(B) = -1.$$

Putting things together we obtain

$$\begin{aligned} \eta_1^{-1} \otimes \psi' \otimes (\psi')^{-1} \otimes \eta_2 &\mapsto (-1)^{d_K+1} \frac{1}{d_{N/K}} (\eta_1^{-1} \otimes (\eta_1 \otimes z) \otimes (\psi')^{-1} \otimes \eta_2) \\ &\mapsto (-1)^{d_K} \frac{1}{d_{N/K}} (z \otimes (\psi')^{-1} \otimes \eta_2) \\ &\mapsto (-1)^{d_K+1} \frac{1}{d_{N/K}} (z \otimes (\eta_2^{-1} \otimes z^{-1}) \otimes \eta_2) \\ &\mapsto (-1)^{d_K+1+d_K-1} \frac{1}{d_{N/K}} (z \otimes z^{-1} \otimes \eta_2^{-1} \otimes \eta_2) \\ &\mapsto \frac{1}{d_{N/K}} (\eta_2^{-1} \otimes \eta_2) \mapsto -\frac{1}{d_{N/K}}. \blacksquare \end{aligned}$$

REMARK 5.3.3. In the case $\chi^{\text{nr}}|_{G_N} = 1$ but $\chi^{\text{nr}} \neq 1$ the restriction of χ^{nr} to N defines an abelian character of $G = \text{Gal}(N/K)$. We write

$$e_{\chi^{\text{nr}}} = \frac{1}{|G|} \sum_{\sigma \in G} \chi^{\text{nr}}(\sigma) \sigma^{-1}$$

for the associated idempotent. The calculations in this case are basically the same as for $V = \mathbb{Q}_p(1)$. Diagram (39) becomes

$$\begin{array}{ccccccc} 0 \longrightarrow & \mathbb{Q}_p((\chi^{\text{nr}})^{-1}) & \xrightarrow{\iota} & N_1((\chi^{\text{nr}})^{-1}) & \xrightarrow{f} & N_1((\chi^{\text{nr}})^{-1}) & \xrightarrow{\xi} H_f^1(N, V^*(1)) \longrightarrow 0 \\ & \downarrow = & & \downarrow = & & \downarrow = & \downarrow \mu \\ 0 \longrightarrow & \mathbb{Q}_p((\chi^{\text{nr}})^{-1}) & \xrightarrow{\iota} & N_1((\chi^{\text{nr}})^{-1}) & \xrightarrow{f} & N_1((\chi^{\text{nr}})^{-1}) & \xrightarrow{-\tau_{N_1/\mathbb{Q}_p}} \mathbb{Q}_p((\chi^{\text{nr}})^{-1}) \longrightarrow 0 \end{array}$$

Note that the maps in (39) and also their splittings do not change; this is the reason why we write f instead of $1 - \varphi$ which otherwise could falsely

be interpreted as $x \mapsto x - \chi^{\text{nr}}(\varphi)\varphi(x)$. For the rest of the proof of the second part of the next proposition one just has to take into account that only the action of G on the modules changes by the twist so that we obtain the non-trivial contribution from non-torsion cohomology for the character χ^{nr} .

We summarize the main results of Section 5.

PROPOSITION 5.3.4.

(a) If $\chi^{\text{nr}}|_{G_N} \neq 1$, then $U_{\text{cris}} = \partial(m_{N/K})$ with

$$m_{N/K} = \frac{*((1 - p^{-d_K} u^{-d_K} F)e_I)}{*((1 - u^{d_K} F^{-1})e_I)}.$$

(b) If $\chi^{\text{nr}}|_{G_N} = 1$, then $U_{\text{cris}} = \partial(m_{N/K})$ with

$$m_{N/K} = \frac{*((1 - p^{-d_K} u^{-d_K} F)e_I) \cdot *(d_{N/K} e_{\chi^{\text{nr}}})}{*((1 - u^{d_K} F^{-1})e_I)}.$$

Proof. Combine (32) and (37) with Lemmas 5.2.1, 5.2.2, 5.3.2 and Remark 5.3.3. ■

REMARK 5.3.5. In the case $\chi^{\text{nr}} = 1$ we precisely recover the correction term $M_{N/K}^{\text{Breu}}$ in Breuning’s conjecture [Bre04b, Conj. 3.2].

6. Epsilon constants of Weil–Deligne representations. In this section we study the epsilon constant defined in [IV16, Sec. 2.3], and in the situation described in Subsection 3.2 we rewrite it in terms of Galois Gauß sums.

6.1. Epsilon constants and Galois Gauß sums. If L/\mathbb{Q}_p is a finite extension and W is an Artin representation of G_L , we let $\mathfrak{f}(W)$ denote the Artin conductor of W . Let $\psi = \psi_\xi$ be the standard additive character, as defined in Subsection 3.1. By [IV16, Lemma 4.1] we have

(45)
$$\varepsilon_D(N/K, V) = (\varepsilon(\text{Ind}_{K/\mathbb{Q}_p} \chi, -\psi, \mu_{\mathbb{Q}_p})^{-1} \chi_{\mathbb{Q}_p}^{\text{nr}} (\mathfrak{f}(\text{Ind}_{K/\mathbb{Q}_p} V_\chi)^{-1}))_{\chi \in \text{Irr}(G)}.$$

Here we view $\chi_{\mathbb{Q}_p}^{\text{nr}}$ as a character $\mathbb{Q}_p^\times \xrightarrow{\text{rec}} G_{\mathbb{Q}_p}^{\text{ab}} \xrightarrow{\chi_{\mathbb{Q}_p}^{\text{nr}}} \mathbb{Q}_p^\times$ via the local reciprocity map.

REMARK 6.1.1. Contrary to the convention in [IV16] we normalize reciprocity maps by sending a uniformizer π_L to the arithmetic Frobenius map F_L .

LEMMA 6.1.2. *Assume that M/L is an abelian extension of p -adic number fields. Let η be an irreducible character of $\text{Gal}(M/L)$. Let $\mathfrak{D}_L = \pi_L^{s_L} \mathcal{O}_L$*

be the different of L/\mathbb{Q}_p . Then

$$\varepsilon(\eta, \psi, \mu_L) = p^{d_L s_L} \tau_L(\eta),$$

where $\tau_L(\eta)$ is a Galois Gauß sum (see e.g. [PV13, p. 1184]).

Proof. We will apply some of properties (1)–(6) listed in [BB08, Sec. 2.3]. If η is unramified, we derive from property (1) of loc. cit. the equality

$$\varepsilon(\eta, \psi, \mu_L) = \eta(\pi_L^{-s_L}) p^{d_L s_L}.$$

We point out that we have $\eta(\pi_L^{-s_L})$ (and not $\eta(\pi_L^{s_L})$ as in [BB08] or [IV16]) due to our different normalization of the reciprocity map. By the definition of Galois Gauß sums one has $\tau_L(\eta) = \eta(\pi_L^{-s_L})$ and the lemma follows for unramified characters η .

Let now η be a ramified character. We write $\mathfrak{f}(\eta) = \pi_L^{m_\eta} \mathcal{O}_L$ for the conductor of η . By property (1) of [BB08] and [Del73, (3.4.3.2)] we have

$$\begin{aligned} \varepsilon(\eta, \psi, \mu_L) &= \int_{\{v_L(x) = -s_L - m_\eta\}} \eta(x) \psi(x) d\mu_L \\ &= \int_{x \in \mathcal{O}_L^\times} \frac{1}{\mu_L(\mathfrak{p}_L^{s_L + m_\eta})} \eta\left(\frac{x}{\pi_L^{s_L + m_\eta}}\right) \psi\left(\frac{x}{\pi_L^{s_L + m_\eta}}\right) d\mu_L \\ &= \frac{1}{\mu_L(\mathfrak{p}_L^{s_L + m_\eta})} \int_{x \in \mathcal{O}_L^\times} \eta\left(\frac{x}{\pi_L^{s_L + m_\eta}}\right) \psi\left(\frac{x}{\pi_L^{s_L + m_\eta}}\right) d\mu_L \\ &= \frac{1}{\mu_L(\mathfrak{p}_L^{s_L + m_\eta})} \mu_L(U_L^{(m_\eta)}) \sum_{x \in U_L/U_L^{(m_\eta)}} \eta\left(\frac{x}{\pi_L^{s_L + m_\eta}}\right) \psi\left(\frac{x}{\pi_L^{s_L + m_\eta}}\right) \\ &= \frac{\mu_L(U_L^{(m_\eta)})}{\mu_L(\mathfrak{p}_L^{s_L + m_\eta})} \tau_L(\eta) = p^{d_L s_L} \tau_L(\eta), \end{aligned}$$

where the next-to-last equality results from the definition of Galois Gauß sums. ■

We will now use Brauer induction in degree 0 to deduce the following proposition.

PROPOSITION 6.1.3. *Let χ be an arbitrary character of $G_{\mathbb{Q}_p}$ with open kernel. Then*

$$\varepsilon(\chi, \psi, \mu_{\mathbb{Q}_p}) = \tau_{\mathbb{Q}_p}(\chi).$$

Proof. By [Ser77, Exercise 10.6] there exist integers a_ϕ such that

$$\chi - \chi(1)1_{G_{\mathbb{Q}_p}} = \sum_{\phi} a_\phi \text{Ind}_{K_\phi/\mathbb{Q}_p}(\phi - 1)$$

with linear characters ϕ of $G_{\mathbb{Q}_p}$ and $K_\phi = (\mathbb{Q}_p^c)^{\ker(\phi)}$. We will again apply some of properties (1)–(6) listed in [BB08, Sec. 2.3]. By property (2), local

constants are additive, and Lemma 6.1.2 applied for $M = L = \mathbb{Q}_p$ together with the fact that $\tau_K(1_{G_{\mathbb{Q}_p}}) = 1$ shows that $\varepsilon(1_{G_{\mathbb{Q}_p}}, \psi, \mu_{\mathbb{Q}_p}) = 1$. Therefore we conclude

$$\begin{aligned} \varepsilon(\chi, \psi, \mu_{\mathbb{Q}_p}) &= \varepsilon(\chi - \chi(1)1_{G_{\mathbb{Q}_p}}, \psi, \mu_{\mathbb{Q}_p})\varepsilon(1_{G_{\mathbb{Q}_p}}, \psi, \mu_{\mathbb{Q}_p})^{\chi(1)} \\ &= \varepsilon(\chi - \chi(1)1_{G_{\mathbb{Q}_p}}, \psi, \mu_{\mathbb{Q}_p}). \end{aligned}$$

We now use properties (2) and (4) of loc. cit. in order to derive

$$\varepsilon(\chi, \psi, \mu_{\mathbb{Q}_p}) = \prod_{\phi} \varepsilon(\text{Ind}_{K_{\phi}/\mathbb{Q}_p}(\phi - 1), \psi, \mu_{\mathbb{Q}_p})^{a_{\phi}} = \prod_{\phi} \varepsilon(\phi - 1, \psi_{\phi}, \mu_{K_{\phi}})^{a_{\phi}},$$

where we write $\psi_{\phi}: K_{\phi} \rightarrow (\mathbb{Q}_p^{\times})^{\times}$ for the standard additive character of K_{ϕ} . Because of $\dim(\text{Ind}_{K_{\phi}/\mathbb{Q}_p}(\phi - 1)) = 0$ the λ -constants which arise in property (4) of loc. cit. do not show up.

We again apply additivity followed by our Lemma 6.1.2 and obtain

$$\begin{aligned} \varepsilon(\chi, \psi, \mu_{\mathbb{Q}_p}) &= \prod_{\phi} \varepsilon(\phi, \psi_{\phi}, \mu_{K_{\phi}})^{a_{\phi}} \cdot \prod_{\phi} \varepsilon(1, \psi_{\phi}, \mu_{K_{\phi}})^{-a_{\phi}} \\ &= \prod_{\phi} (p^{d_{K_{\phi}} s_{K_{\phi}}} \tau_{K_{\phi}}(\phi))^{a_{\phi}} \cdot \prod_{\phi} (p^{d_{K_{\phi}} s_{K_{\phi}}} \tau_{K_{\phi}}(1))^{-a_{\phi}} \\ &= \prod_{\phi} (\tau_{K_{\phi}}(\phi) \tau_{K_{\phi}}(1)^{-1})^{a_{\phi}}. \end{aligned}$$

Now we recall from [Frö83] that Galois Gauß sums are additive, inductive in degree zero and equal to 1 for the trivial character. It follows that

$$\begin{aligned} \varepsilon(\chi, \psi, \mu_{\mathbb{Q}_p}) &= \prod_{\phi} \tau_{K_{\phi}}(\phi - 1)^{a_{\phi}} = \prod_{\phi} \tau_{\mathbb{Q}_p}(\text{Ind}_{K_{\phi}/\mathbb{Q}_p}(\phi - 1))^{a_{\phi}} \\ &= \tau_{\mathbb{Q}_p}(\chi - \chi(1)1_{G_{\mathbb{Q}_p}}) = \tau_{\mathbb{Q}_p}(\chi). \quad \blacksquare \end{aligned}$$

6.2. Epsilon constants for unramified twists. We apply the results of this section to unramified twists of $\mathbb{Z}_p(1)$.

PROPOSITION 6.2.1. *Assume that N/K is an arbitrary Galois extension of p -adic number fields with Galois group G . Let $\chi_{\mathbb{Q}_p}^{\text{nr}}$ be as usual and $\chi_{\mathbb{Q}_p}^{\text{nr}}(\varphi) = u$. Let $\chi \in \text{Irr}(G)$ and write $\mathfrak{f}(\chi) = \pi_K^{m_{\chi}} \mathcal{O}_K$ for the Artin conductor. Then*

$$\varepsilon_D(N/K, V)_{\chi} = (\det(\text{Ind}_{K/\mathbb{Q}_p} \chi))(-1) \cdot u^{-d_K(s_K \chi(1) + m_{\chi})} \cdot \tau_{\mathbb{Q}_p}(\text{Ind}_{K/\mathbb{Q}_p} \chi)^{-1}.$$

Proof. By property (3) of [BB08, Sec. 2.3] we have

$$\varepsilon(\text{Ind}_{K/\mathbb{Q}_p} \chi, -\psi, \mu_{\mathbb{Q}_p}) = \det(\text{Ind}_{K/\mathbb{Q}_p} \chi)(-1) \varepsilon(\text{Ind}_{K/\mathbb{Q}_p} \chi, \psi, \mu_{\mathbb{Q}_p}).$$

By Proposition 6.1.3 we deduce further

$$(46) \quad \varepsilon(\text{Ind}_{K/\mathbb{Q}_p} \chi, -\psi, \mu_{\mathbb{Q}_p}) = \det(\text{Ind}_{K/\mathbb{Q}_p} \chi)(-1) \tau_{\mathbb{Q}_p}(\text{Ind}_{K/\mathbb{Q}_p} \chi).$$

Finally, by [Neu92, VII.11.7], we get

$$f(\text{Ind}_{K/\mathbb{Q}_p} \chi) = \mathfrak{d}_K^{\chi(1)} N_{K/\mathbb{Q}_p}(f(\chi)) = p^{d_K s_K \chi(1)} p^{d_K m_\chi},$$

where \mathfrak{d}_K denotes the discriminant of K/\mathbb{Q}_p . Since $\chi_{\mathbb{Q}_p}^{\text{nr}}(p) = \chi_{\mathbb{Q}_p}^{\text{nr}}(\varphi) = u$, we obtain

$$(47) \quad \chi_{\mathbb{Q}_p}^{\text{nr}}(f(\text{Ind}_{K/\mathbb{Q}_p} \chi)) = u^{d_K(s_K \chi(1) + m_\chi)}.$$

Combining (45)–(47) leads to the assertion. ■

For later reference we record the following lemma.

LEMMA 6.2.2. *Assume that N/K is an arbitrary Galois extension of p -adic number fields with Galois group G . If $\chi \in \text{Irr}(G)$, then*

$$\partial_{\mathbb{Z}_p[G], \mathbb{Q}_p[G]}^1((\det(\text{Ind}_{K/\mathbb{Q}_p} \chi))(a)) = 0$$

for all $a \in \mathbb{Q}_p^\times$.

Proof. We write $\sigma \in G_{\mathbb{Q}_p}$ for any lift of $\text{rec}_{\mathbb{Q}_p}(a) \in G_{\mathbb{Q}_p}^{\text{ab}}$. Let us denote by $\text{Ver}: G_{\mathbb{Q}_p}^{\text{ab}} \rightarrow G_K^{\text{ab}}$ the transfer homomorphism (see e.g. [NSW08, Sec. I.5]). By [Del73, §1] we have the rule

$$\det(\text{Ind}_{K/\mathbb{Q}_p} \chi) = \det(\chi) \circ \text{Ver}.$$

Therefore $\det(\text{Ind}_{K/\mathbb{Q}_p} \chi)(a) = (\det \chi)(\tau)$ with $\tau := \text{Ver}(\sigma)$. The linear character $\det \chi$ factors through the quotient $\text{Gal}(K^{\text{ab}} \cap N/K)$. If $\sigma' \in G$ denotes any lift of $\tau|_{K^{\text{ab}} \cap N}$, then $(\det \chi)(\tau) = (\det \chi)(\sigma')$. It follows that $\det(\text{Ind}_{K/\mathbb{Q}_p} \chi)(a)$ is the image of $\sigma' \in K_1(\mathbb{Z}_p[G])$ under the canonical homomorphism $K_1(\mathbb{Z}_p[G]) \rightarrow K_1(\mathbb{Q}_p[G])$. The result now follows from the localization sequence. ■

7. The computation of the cohomological term. This section contains the more technical part of the paper, concerning the computation of the cohomological term $C_{N/K}$ in the special case we are considering. This will allow us to prove Theorem 1 in the last section.

7.1. Some general results about $C_{N/K}$. We start with some general results about the cohomological term, for which we do not need the restrictive assumptions of Theorem 1. We will use these results to prove the rationality of $R_{N/K}$ and the equivalence of Conjectures 3.1.1, 3.2.4 and 3.2.5 as stated in Proposition 3.2.6. Furthermore we will show that Breuning’s conjecture [Bre04b, Conj. 3.2] is a special case of Conjecture 3.2.5.

We recall that $T = \mathbb{Z}_p(\chi^{\text{nr}})(1)$ with an unramified character χ^{nr} which is the restriction to G_K of an unramified character $\chi_{\mathbb{Q}_p}^{\text{nr}}: G_{\mathbb{Q}_p} \rightarrow \mathbb{Z}_p^\times$. Then $T \cong T_p \mathcal{F}$ where \mathcal{F} is the Lubin–Tate formal group associated with $\pi = up$ and $u = \chi_{\mathbb{Q}_p}^{\text{nr}}(\varphi)$.

Let $\mathcal{L} \subseteq \mathcal{O}_N$ be a full projective $\mathbb{Z}_p[G]$ -sublattice such that the exponential map $\exp_{\mathcal{F}}: \mathbb{G}_a \rightarrow \mathcal{F}$ converges on \mathcal{L} . We first assume that $\chi^{\text{nr}}|_{G_N} \neq 1$. We set $X(\mathcal{L}) := \exp_{\mathcal{F}}(\mathcal{L})$ and obtain in this way a full $\mathbb{Z}_p[G]$ -projective sublattice of $\mathcal{F}(\mathfrak{p}_N)$, which by Lemma 3.2.1(a) we identify with $H^1(N, T)$. If $\chi^{\text{nr}}|_{G_N} = 1$, we note that $\exp_{\mathbb{G}_m} = \exp - 1$ where \exp denotes the p -adic exponential map. In this case we define $X(\mathcal{L}) := \exp(\mathcal{L})$ and obtain a full $\mathbb{Z}_p[G]$ -projective sublattice of $U_N \subseteq \widehat{N^\times}$. Recall that by Lemma 3.2.1(b) we identify $H^1(N, T)$ with $\widehat{N^\times}(\chi^{\text{nr}})$.

So in either case we have a natural embedding $X(\mathcal{L}) \hookrightarrow H^1(N, T)$. As in the proof of Corollary 4.2.10 we fix once and for all a representative $[A \rightarrow B]$ of $R\Gamma(N, T)$, i.e., A and B are finitely generated cohomologically trivial $\mathbb{Z}_p[G]$ -modules centered in degrees 1 and 2. In addition, we will always assume that B is $\mathbb{Z}_p[G]$ -projective. Thus we have a perfect 2-extension

$$0 \rightarrow H^1(N, T) \rightarrow A \rightarrow B \rightarrow H^2(N, T) \rightarrow 0.$$

The embedding $X(\mathcal{L}) \hookrightarrow H^1(N, T)$ induces an injective map of complexes

$$X(\mathcal{L})[-1] \rightarrow M^\bullet,$$

where $M^\bullet = R\Gamma(N, T) \oplus \text{Ind}_{N/\mathbb{Q}_p} T[0]$ was defined in (16).

We set

$$\begin{aligned} K^\bullet(\mathcal{L}) &:= \text{Ind}_{N/\mathbb{Q}_p}(T)[0] \oplus X(\mathcal{L})[-1], \\ M^\bullet(\mathcal{L}) &:= [A/X(\mathcal{L}) \rightarrow B] \quad \text{with modules in degrees 1 and 2,} \end{aligned}$$

and have thus constructed an exact sequence of complexes

$$0 \rightarrow K^\bullet(\mathcal{L}) \rightarrow M^\bullet \rightarrow M^\bullet(\mathcal{L}) \rightarrow 0.$$

By the additivity of Euler characteristics (see [BB05, Th. 5.7]) and the definition of $C_{N/K}$ in (15) we obtain

$$(48) \quad C_{N/K} = -\chi_{\mathbb{Z}_p[G], B_{\text{dR}}[G]}(K^\bullet(\mathcal{L}), \lambda_1^{-1}) - \chi_{\mathbb{Z}_p[G], B_{\text{dR}}[G]}(M^\bullet(\mathcal{L}), \lambda_2^{-1})$$

with $\lambda_1 = \text{comp}_V \circ \exp_V^{-1}$ and

$$\lambda_2: (H^1(N, T)/X(\mathcal{L})) \otimes_{\mathbb{Z}_p[G]} B_{\text{dR}}[G] \rightarrow H^2(N, T) \otimes_{\mathbb{Z}_p[G]} B_{\text{dR}}[G]$$

given by

$$\lambda_2 = \begin{cases} \nu_N & \text{if } \chi^{\text{nr}}|_{G_N} = 1, \\ 0 & \text{if } \chi^{\text{nr}}|_{G_N} \neq 1. \end{cases}$$

From (4) we obtain

$$(49) \quad \begin{aligned} &\chi_{\mathbb{Z}_p[G], B_{\text{dR}}[G]}(K^\bullet(\mathcal{L}), \lambda_1^{-1}) \\ &= -\chi_{\mathbb{Z}_p[G], B_{\text{dR}}[G]}^{\text{old}}(K^\bullet(\mathcal{L}), \lambda_1) - \partial_{\mathbb{Z}_p[G], B_{\text{dR}}[G]}^1((B^{\text{od}}(K^\bullet(\mathcal{L})), -\text{id})). \end{aligned}$$

By the definition of $\chi_{\mathbb{Z}_p[G], B_{\text{dR}}[G]}^{\text{old}}$ we derive

$$(50) \quad \chi_{\mathbb{Z}_p[G], B_{\text{dR}}[G]}^{\text{old}}(K^\bullet(\mathcal{L}), \lambda_1) = [X(\mathcal{L}), \text{comp}_V \circ \exp_V^{-1}, \text{Ind}_{N/\mathbb{Q}_p} T].$$

Furthermore, by [BB05, Lemma 6.3], we have

$$(51) \quad \begin{aligned} \partial_{\mathbb{Z}_p[G], B_{\text{dR}}[G]}^1((B^{\text{od}}(K^\bullet(\mathcal{L})), -\text{id})) \\ = \partial_{\mathbb{Z}_p[G], B_{\text{dR}}[G]}^1((X(\mathcal{L}) \otimes_{\mathbb{Z}_p[G]} B_{\text{dR}}[G], -\text{id})). \end{aligned}$$

Since $X(\mathcal{L}) \otimes_{\mathbb{Z}_p[G]} \mathbb{Q}_p[G] \cong \mathbb{Q}_p[G]^{[K:\mathbb{Q}_p]}$ is a free $\mathbb{Q}_p[G]$ -module, the right hand side is trivial, so we deduce from (48)–(51) that

$$(52) \quad \begin{aligned} C_{N/K} = [X(\mathcal{L}), \text{comp}_V \circ \exp_V^{-1}, \text{Ind}_{N/\mathbb{Q}_p} T] \\ - \chi_{\mathbb{Z}_p[G], B_{\text{dR}}[G]}(M^\bullet(\mathcal{L}), \lambda_2^{-1}). \end{aligned}$$

Similarly, it can be shown that

$$(53) \quad \chi_{\mathbb{Z}_p[G], B_{\text{dR}}[G]}(M^\bullet(\mathcal{L}), \lambda_2^{-1}) = -\chi_{\mathbb{Z}_p[G], B_{\text{dR}}[G]}^{\text{old}}(M^\bullet(\mathcal{L}), \lambda_2).$$

REMARK 7.1.1. If \mathfrak{p}_N^n is $\mathbb{Z}_p[G]$ -projective for some $n \in \mathbb{N}$ and n is large enough so that $\exp_{\mathcal{F}}$ converges on \mathfrak{p}_N^n , then we can use $\mathcal{L} = \mathfrak{p}_N^n$. In this case we obtain $X(\mathcal{L}) = \mathcal{F}(\mathfrak{p}_N^n)$ resp. $X(\mathcal{L}) = U_N^{(n)}$. For example, if N/K is at most tamely ramified, every ideal \mathfrak{p}_N^n is $\mathbb{Z}_p[G]$ -projective. If N/K is weakly ramified, then \mathfrak{p}_N^n is $\mathbb{Z}_p[G]$ -projective if and only if $n \equiv 1 \pmod{|G_1|}$. Both properties are implied by [Köc04, Th. 1.1].

The computation of $\chi_{\mathbb{Z}_p[G], B_{\text{dR}}[G]}(M^\bullet(\mathcal{L}), \lambda_2^{-1})$ in the special case of Theorem 1 will be postponed to the next subsections. For the purpose of this subsection we just remark that

$$(54) \quad \chi_{\mathbb{Z}_p[G], B_{\text{dR}}[G]}(M^\bullet(\mathcal{L}), \lambda_2^{-1}) \in K_0(\mathbb{Z}_p[G], \mathbb{Q}_p[G]).$$

The term $[X(\mathcal{L}), \text{comp}_V \circ \exp_V^{-1}, \text{Ind}_{N/\mathbb{Q}_p}(T)]$ can be made more explicit if we let \mathcal{L} be of the following special form. Let $b \in N$ be a normal basis element of N/K , i.e. $N = K[G]b$, and suppose that $\mathcal{L} = \mathcal{O}_K[G]b$ is such that $\exp_{\mathcal{F}}$ converges on \mathcal{L} . Let

$$\rho = (\rho_\chi)_{\chi \in \text{Irr}(G)} \in Z(\mathbb{Q}_p^c[G])^\times = \prod_{\chi \in \text{Irr}(G)} (\mathbb{Q}_p^c)^\times$$

be defined by

$$\rho_\chi = \mathfrak{d}_K^{\chi(1)} \mathcal{N}_{K/\mathbb{Q}_p}(b|\chi),$$

where \mathfrak{d}_K denotes the discriminant of K/\mathbb{Q}_p and $\mathcal{N}_{K/\mathbb{Q}_p}(b|\chi)$ the usual norm resolvent (see e.g. [PV13, Sec. 2.2]).

LEMMA 7.1.2. *Let $\mathcal{L} = \mathcal{O}_K[G]b$ be as above. Then*

$$[X(\mathcal{L}), \text{comp}_V \circ \exp_V^{-1}, \text{Ind}_{N/\mathbb{Q}_p} T] + \partial_{\mathbb{Z}_p[G], B_{\text{dR}}[G]}^1(t) = \hat{\partial}_{\mathbb{Z}_p[G], B_{\text{dR}}[G]}^1(\rho)$$

in $K_0(\mathbb{Z}_p[G], B_{\text{dR}}[G])$.

Proof. The proof is similar to that of [IV16, Lemma 6.1]. ■

Combining (52) and Lemma 7.1.2 we obtain

$$(55) \quad C_{N/K} = -\partial_{\mathbb{Z}_p[G], B_{\text{dR}}[G]}^1(t) + \partial_{\mathbb{Z}_p[G], B_{\text{dR}}[G]}^1(\rho) - \chi_{\mathbb{Z}_p[G], B_{\text{dR}}[G]}(M^\bullet(\mathcal{L}), \lambda_2^{-1}).$$

We recall the basic properties of the so-called *unramified term* $U_{N/K}$ defined by Breuning in [Bre04b, Prop. 2.12]. We write \mathcal{O}_p^t for the ring of integers in the maximal tamely ramified extension of \mathbb{Q}_p in \mathbb{Q}_p^c . For subrings $\mathbb{Z}_p[G] \subseteq \Lambda_1 \subseteq \Lambda_2 \subseteq \mathbb{Q}_p^c[G]$ we write

$$j_{\Lambda_1, \Lambda_2}: K_0(\Lambda_1, \mathbb{Q}_p^c[G]) \rightarrow K_0(\Lambda_2, \mathbb{Q}_p^c[G])$$

for the natural scalar extension map. We write

$$\iota: K_0(\mathbb{Z}_p[G], \mathbb{Q}_p^c[G]) \rightarrow K_0(\mathcal{O}_p^t[G], \mathbb{Q}_p^c[G])$$

for $j_{\mathbb{Z}_p[G], \mathcal{O}_p^t[G]}$. We recall that by Taylor’s fixed point theorem the restriction of ι to the subgroup $K_0(\mathbb{Z}_p[G], \mathbb{Q}_p[G])$ is injective.

Let \mathbb{Q}_p^{ab} be the maximal abelian extension of \mathbb{Q}_p inside \mathbb{Q}_p^c . Then we recall that $\mathbb{Q}_p^{\text{ab}} = \mathbb{Q}_p^{\text{nr}}\mathbb{Q}_p^{\text{ram}}$, where \mathbb{Q}_p^{nr} denotes the maximal unramified extension of \mathbb{Q}_p and $\mathbb{Q}_p^{\text{ram}} := \mathbb{Q}_p(\mu_{p^\infty})$. For each $\omega \in G_{\mathbb{Q}_p}$ we define elements $\omega^{\text{nr}}, \omega^{\text{ram}} \in \text{Gal}(\mathbb{Q}_p^{\text{ab}}/\mathbb{Q}_p)$ by

$$\omega^{\text{nr}}|_{\mathbb{Q}_p^{\text{nr}}} = \omega|_{\mathbb{Q}_p^{\text{nr}}}, \quad \omega^{\text{nr}}|_{\mathbb{Q}_p^{\text{ram}}} = 1, \quad \omega^{\text{ram}}|_{\mathbb{Q}_p^{\text{nr}}} = 1, \quad \omega^{\text{ram}}|_{\mathbb{Q}_p^{\text{ram}}} = \omega|_{\mathbb{Q}_p^{\text{ram}}}.$$

Then $U_{N/K}$ is the unique element in $K_0(\mathbb{Z}_p[G], \mathbb{Q}_p^c[G])$ satisfying the following two properties (see [Bre04b, Prop. 2.12]):

- (1) $\iota(U_{N/K}) = 0$.
- (2) If $(\alpha_\chi)_{\chi \in \text{Irr}(G)} \in \prod_{\chi \in \text{Irr}(G)} (\mathbb{Q}_p^c)^\times$ is any preimage of $U_{N/K}$ under $\hat{\partial}_{\mathbb{Z}_p[G], \mathbb{Q}_p^c[G]}^1$, then

$$(56) \quad \omega(\alpha_{\omega^{-1} \circ \chi}) = \alpha_\chi \det_{\text{Ind}_{K/\mathbb{Q}_p}(\chi)}(\omega^{\text{nr}})$$

for all $\omega \in G_{\mathbb{Q}_p}$.

Recall the definition (17):

$$R_{N/K} = C_{N/K} + U_{\text{cris}} + \partial_{\mathbb{Z}_p[G], B_{\text{dR}}[G]}^1(t) - U_{N/K} + \partial_{\mathbb{Z}_p[G], B_{\text{dR}}[G]}^1(\varepsilon_D(N/K, V)).$$

We are now able to prove the following result.

PROPOSITION 7.1.3. *The element $R_{N/K}$ is rational, i.e. $R_{N/K}$ belongs to $K_0(\mathbb{Z}_p[G], \mathbb{Q}_p[G])$.*

Proof. For elements $x, y \in K_0(\mathbb{Z}_p[G], \mathbb{Q}_p^c[G])$ we use the notation $x \equiv y$ when $x - y \in K_0(\mathbb{Z}_p[G], \mathbb{Q}_p[G])$. Since U_{cris} is rational, we derive from Proposition 6.2.1, (54) and (55) that

$$R_{N/K} \equiv \partial_{\mathbb{Z}_p[G], B_{\text{dR}}[G]}^1(\rho) - U_{N/K} - T_{N/K},$$

where $T_{N/K} := \partial_{\mathbb{Z}_p[G], B_{\text{dR}}[G]}^1(\tau_{\mathbb{Q}_p}(\text{Ind}_{K/\mathbb{Q}_p} \chi)_{\chi \in \text{Irr}(G)})$ is precisely the element defined by Breuning [Bre04b, Sec. 2.3].

By [Bre04b, Lemma 2.4] we find that for all preimages

$$(\beta_\chi)_{\chi \in \text{Irr}(G)} \in \prod_{\chi \in \text{Irr}(G)} (\mathbb{Q}_p^\times)$$

of $T_{N/K}$ under $\hat{\partial}_{\mathbb{Z}_p[G], \mathbb{Q}_p^\times[G]}^1$ and all $\omega \in G_{\mathbb{Q}_p}$,

$$(57) \quad \omega(\beta_{\omega^{-1} \circ \chi}) = \beta_\chi \det_{\text{Ind}_{K/\mathbb{Q}_p}(\chi)}(\omega^{\text{ram}}).$$

From the proof of [Bre04b, Lemma 2.8] we derive

$$(58) \quad \omega(\rho_{\omega^{-1} \circ \chi}) = \rho_\chi \det_{\text{Ind}_{K/\mathbb{Q}_p}(\chi)}(\omega).$$

The result now follows by combining (56)–(58). ■

Proof of Proposition 3.2.6. We have to show the equivalence of the following assertions:

- (i) $j_{\mathbb{Z}_p[G], \tilde{\Lambda}}([\Delta_{\text{EP}}^{\text{na}}(N/K, T), \delta(N/K, V)]) = 0$ in $\pi_0(V(\tilde{\Lambda}, \tilde{\Omega})) \cong K_0(\tilde{\Lambda}, \tilde{\Omega})$.
- (ii) $\tilde{R}_{N/K} = 0$ in $K_0(\tilde{\Lambda}, \tilde{\Omega})$.
- (iii) $R_{N/K} = 0$ in $K_0(\mathbb{Z}_p[G], \tilde{\Omega})$.

By Taylor’s fixed point theorem, the restriction of the map $j_{\mathbb{Z}_p[G], \tilde{\Lambda}}$ to $K_0(\mathbb{Z}_p[G], \mathbb{Q}_p[G])$ is injective. In order to prove the equivalence of (ii) and (iii) it suffices by Proposition 7.1.3 to show that $j_{\mathbb{Z}_p[G], \tilde{\Lambda}}(R_{N/K}) = \tilde{R}_{N/K}$. For this purpose we just remark that the proof of [Bre04b, Prop. 2.12] actually shows that $j_{\mathbb{Z}_p[G], \tilde{\Lambda}}(U_{N/K}) = 0$.

The equivalence of (i) and (ii) was already verified in Lemma 5.1.1. ■

To conclude this subsection we prove the following proposition.

PROPOSITION 7.1.4. *If $T = \mathbb{Z}_p(1)$, then Conjecture 3.2.5 and [Bre04b, Conj. 3.2] are equivalent.*

Proof. [Bre04b, Conj. 3.2] states that

$$R_{N/K}^{\text{Breu}} = T_{N/K}^{\text{Breu}} + C_{N/K}^{\text{Breu}} + U_{N/K}^{\text{Breu}} - M_{N/K}^{\text{Breu}} = 0,$$

where we write the index Breu to refer to the terms defined by Breuning. The terms $U_{N/K}^{\text{Breu}}$ and $U_{N/K}$ are defined in the same way. By Remark 5.3.5, $M_{N/K}^{\text{Breu}} = U_{\text{cris}}$. By Proposition 6.2.1 together with Lemma 6.2.2 we have

$$T_{N/K}^{\text{Breu}} = \partial_{\mathbb{Z}_p[G], B_{\text{dR}}[G]}^1 \tau_{\mathbb{Q}_p}(\text{Ind}_{K/\mathbb{Q}_p} \chi) = -\partial_{\mathbb{Z}_p[G], B_{\text{dR}}[G]}^1 \varepsilon_D(N/K, V)_\chi.$$

Recalling the definition of $E(X)$ in [BB03, (19)], the proof of [BB03, Lemma 3.7] and equation (53), we see that

$$\begin{aligned}
 (59) \quad E(\exp(\mathcal{L}))_p &= \chi_{\mathbb{Z}_p[G], B_{\text{dR}}[G]}^{\text{old}}(M^\bullet(\mathcal{L})[1], \lambda_2^{-1}) \\
 &= -\chi_{\mathbb{Z}_p[G], B_{\text{dR}}[G]}^{\text{old}}(M^\bullet(\mathcal{L}), \lambda_2) \\
 &= \chi_{\mathbb{Z}_p[G], B_{\text{dR}}[G]}(M^\bullet(\mathcal{L}), \lambda_2^{-1}).
 \end{aligned}$$


Using [Bre04b, Prop. 2.6, Lemma 2.7], (59), Lemma 7.1.2 and (52) we obtain

$$\begin{aligned}
 C_{N/K}^{\text{Breu}} &= E(\exp(\mathcal{L}))_p - [\mathcal{L}, \rho_L, H_L] \\
 &= \chi_{\mathbb{Z}_p[G], B_{\text{dR}}[G]}(M^\bullet(\mathcal{L}), \lambda_2^{-1}) - \hat{\partial}_{\tilde{\Lambda}, B_{\text{dR}}[G]}^1(\rho) \\
 &= \chi_{\mathbb{Z}_p[G], B_{\text{dR}}[G]}(M^\bullet(\mathcal{L}), \lambda_2^{-1}) \\
 &\quad - [X(\mathcal{L}), \text{comp}_V \circ \exp_V^{-1}, \text{Ind}_{N/\mathbb{Q}_p}(T)] - \partial_{\tilde{\Lambda}, B_{\text{dR}}[G]}^1(t) \\
 &= -C_{N/K} - \partial_{\tilde{\Lambda}, B_{\text{dR}}[G]}^1(t).
 \end{aligned}$$

Substituting all those terms in $R_{N/K}^{\text{Breu}}$ we get $-R_{N/K}$. ■

7.2. The computation of $C_{N/K}$. Our aim now is to make formula (55) more explicit in the special situation of Theorem 1 by calculating the Euler characteristic $\chi_{\mathbb{Z}_p[G], B_{\text{dR}}[G]}(M^\bullet(\mathcal{L}), \lambda_2^{-1})$. We will use several preliminary results and constructions of [BC16], and therefore stick closely to the notation used there.

We will consider local field extensions as follows:



Here K/\mathbb{Q}_p is the unramified extension of degree m , i.e. $d_K = m = [K : \mathbb{Q}_p]$. Furthermore, K'/K is the maximal unramified subextension of N/K and we set $d := d_{N/K} = [K' : K]$. We assume throughout that $(m, d) = 1$. Finally, M/K is a weakly and wildly ramified cyclic extension of degree p and $N = MK'$. Since $(m, d) = 1$, there exists \tilde{K}'/\mathbb{Q}_p of degree d such that $K' = K\tilde{K}'$.

Let $F := F_K \in \text{Gal}(N_{\text{nr}}/M) \cong \text{Gal}(K_{\text{nr}}/K)$ be the Frobenius automorphism; we will keep the notation F_N for the element $F^d \in \text{Gal}(N_{\text{nr}}/N) \cong$

$\text{Gal}(K_{\text{nr}}/K')$, which in [BC16] used to be called F_0 . We set $q = p^m$, $b = F^{-1}$ and consider an element $a \in \text{Gal}(N_{\text{nr}}/K)$ such that $\text{Gal}(M/K) = \langle a|_M \rangle$, $a|_{K_{\text{nr}}} = 1$. Since there will be no ambiguity, we will denote by the same letters a, b their restrictions to N . Then $\text{Gal}(N/K) = \langle a, b \rangle$ and $\text{ord}(a) = p$, $\text{ord}(b) = d$. We also define $\mathcal{T}_a := \sum_{i=0}^{p-1} a^i$.

Any irreducible character ψ of G decomposes as $\psi = \chi\phi$ where χ is an irreducible character of $\langle a \rangle$ and ϕ an irreducible character of $\langle b \rangle$.

We will always identify $K_0(\mathbb{Z}_p[G], \mathbb{Q}_p^c[G])$ with $\mathbb{Q}_p^c[G]^\times / \mathbb{Z}_p[G]^\times$ where the isomorphism is induced by $\hat{\delta}_{\mathbb{Z}_p[G], \mathbb{Q}_p^c[G]}^1$. We say that $\alpha \in \mathbb{Q}_p^c[G]^\times$ represents $c \in K_0(\mathbb{Z}_p[G], \mathbb{Q}_p^c[G])$ if $\hat{\delta}_{\mathbb{Z}_p[G], \mathbb{Q}_p^c[G]}^1(\alpha) = c$.

If $\chi^{\text{nr}}|_{G_N} = 1$, the Euler characteristic $\chi_{\mathbb{Z}_p[G], B_{\text{dR}}[G]}(M^\bullet(\mathcal{L}), \lambda_2^{-1})$ is just a twist of the element $E(\exp(\mathcal{L}))_p$ which we computed in [BC16, Prop. 4.3.1].

From now on we assume that $\chi^{\text{nr}}|_{G_N} \neq 1$. We will have to study separately the cases $\omega = 0$ and $\omega > 0$, where $\omega = \omega_N = v_p(1 - \chi^{\text{nr}}(F_N))$ is as in (14). We start with some preliminary results and constructions which will be used in both cases.

LEMMA 7.2.1. *For $n \geq 2$ one has*

$$\mathcal{F}(\mathfrak{p}_N^n) \text{ is } \mathbb{Z}_p[G]\text{-projective} \Leftrightarrow n \equiv 1 \pmod{p}.$$

Moreover,

$$\mathcal{F}(\mathfrak{p}_N) \text{ is } \mathbb{Z}_p[G]\text{-projective} \Leftrightarrow \omega = 0.$$

Proof. For $n \geq 2$ the formal logarithm induces an isomorphism $\mathcal{F}(\mathfrak{p}_N^n) \cong \mathfrak{p}_N^n$ of $\mathbb{Z}_p[G]$ -modules by [Sil09, Th. IV.6.4]. Hence the first assertion follows from [Köc04, Th. 1.1 and Prop. 1.3].

We henceforth assume $n = 1$. By Lemmas 4.1.1, 4.1.2 and Theorem 4.2.3 we already know that $\mathcal{F}(\mathfrak{p}_N)$ is cohomologically trivial if and only if $\omega = 0$. Hence it suffices to prove that if $\omega = 0$ then $\mathcal{F}(\mathfrak{p}_N)$ is torsionfree. By [Sil09, Th. IV.6.4], $\log_{\mathcal{F}}$ converges on $\mathcal{F}(\mathfrak{p}_N)$, so that we may set $J := \log_{\mathcal{F}}(\mathcal{F}(\mathfrak{p}_N)) \subseteq N$. We consider the commutative diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & \mathcal{F}(\mathfrak{p}_N^2) & \longrightarrow & \mathcal{F}(\mathfrak{p}_N) & \longrightarrow & \mathcal{F}(\mathfrak{p}_N)/\mathcal{F}(\mathfrak{p}_N^2) \longrightarrow 0 \\ & & \downarrow \log_{\mathcal{F}} & & \downarrow \log_{\mathcal{F}} & & \downarrow \lambda_{\mathcal{F}} \\ 0 & \longrightarrow & \mathfrak{p}_N^2 & \longrightarrow & J & \longrightarrow & J/\mathfrak{p}_N^2 \longrightarrow 0 \end{array}$$

where $\lambda_{\mathcal{F}}(x + \mathcal{F}(\mathfrak{p}_N^2)) = \log_{\mathcal{F}}(x) + \mathfrak{p}_N^2$. Recall that in Subsection 3.2 we introduced the Lubin–Tate formal group \mathcal{F} associated with $\pi = up$. By [Neu92, Sekt. V.4, Aufgabe 4] we may choose \mathcal{F} such that

$$\log_{\mathcal{F}}(x) = \sum_{j=0}^{\infty} \frac{x^{p^j}}{\pi^j}.$$

We recall that $v_N(\pi) = p$. If $v_N(x) = 1$, then x^p/π is a unit and so also is $x + x^p/\pi$. Since for $j \geq 2$,

$$v_N\left(\frac{x^{p^j}}{\pi^j}\right) = p^j - jp \geq 2,$$

we deduce that $v_N(\log_{\mathcal{F}}(x)) = 0$ for $x \in \mathcal{F}(\mathfrak{p}_N) \setminus \mathcal{F}(\mathfrak{p}_N^2)$, so that $\lambda_{\mathcal{F}}$ is injective. By the Snake Lemma the middle vertical map is therefore also injective and we have an isomorphism $\log_{\mathcal{F}}: \mathcal{F}(\mathfrak{p}_N) \rightarrow J$ of \mathbb{Z}_p -modules. Obviously, $J \subseteq N$ is torsionfree. ■

By the above lemma we can use $\mathcal{L} = \mathfrak{p}_N^{p+1}$ in our constructions (see also Remark 7.1.1). In this case we have $X(\mathcal{L}) = \mathcal{F}(\mathfrak{p}_N^{p+1})$.

We now recall some results and constructions from [BC16]. Let $\theta_1 \in M$ be such that $\mathcal{O}_K[\text{Gal}(M/K)]\theta_1 = \mathfrak{p}_M$ and $\mathcal{T}_{M/K}\theta_1 = p$ (see [BC16, Lemma 3.1.2]). Let θ_2 (resp. A) be a normal integral basis generator of trace one for the extension \tilde{K}'/\mathbb{Q}_p (resp. K/\mathbb{Q}_p). Such elements exist by [BC16, Lemma 3.1.1].

Since $a \in G_1 \setminus G_2$, where G_i is the i th ramification group of $G = \text{Gal}(N/K)$, we know by [Ser79, Prop. IV.2.5] that $\theta_1^{a-1} \equiv 1 - \alpha_1\theta_1 \pmod{\mathfrak{p}_M^2}$ for some unit $\alpha_1 \in \mathcal{O}_M^\times$. Since α_1 can be replaced by any element in the same residue class in $\mathcal{O}_M/\mathfrak{p}_M = \mathcal{O}_K/\mathfrak{p}_K$, we can assume that $\alpha_1 \in \mathcal{O}_K^\times$.

By our choice of A , we know that $A, A^\varphi, \dots, A^{\varphi^{m-1}}$ is a basis of \mathcal{O}_K over \mathbb{Z}_p , where φ denotes as before the Frobenius automorphism of $K_{\text{nr}}/\mathbb{Q}_p$. Since $1 = \mathcal{T}_{K/\mathbb{Q}_p}A = \sum_{i=0}^{m-1} A^{\varphi^i}$ and $\alpha_1 \in \mathcal{O}_K^\times$, it easily follows that also

$$(60) \quad \alpha_1, \alpha_2 = \alpha_1 A, \alpha_3 = \alpha_1 A^\varphi, \dots, \alpha_m = \alpha_1 A^{\varphi^{m-2}}$$

constitute a basis of \mathcal{O}_K over \mathbb{Z}_p . In particular, $A = \alpha_2/\alpha_1$.

We recall from [BC16, Lemma 3.1.3] that the polynomial $X^p - X + A\theta_2$ divides $X^{q^d} - X + 1$ in $\mathcal{O}_{K'}/\mathfrak{p}_{K'}[X]$. As in [BC16], we choose $x_2 \in \mathcal{O}_{K_{\text{nr}}}$ such that $x_2/\alpha_1 \pmod{\mathfrak{p}_{K'}}$ is a root of $X^p - X + A\theta_2$. The following lemma is analogous to [BC16, Lemma 3.1.4].

LEMMA 7.2.2. *Assume $\omega > 0$. If we consider $1 + x_2\theta_1$ as an element of $\widehat{N_0^\times}$, then*

$$(1 + x_2\theta_1)^{\chi^{\text{nr}}(F_N)^{F_N-1}} \equiv \theta_1^{a-1} \equiv 1 - \alpha_1\theta_1 \pmod{\mathfrak{p}_{N_0}^2}.$$

Proof. From $(x_2/\alpha_1)^{q^d} - x_2/\alpha_1 \equiv -1 \pmod{\mathfrak{p}_{K'}}$ we obtain $x_2^{q^d} - x_2 \equiv -\alpha_1 \pmod{\mathfrak{p}_{N_0}}$. Since $\omega > 0$, we have $u^{dm} \equiv 1 \pmod{p}$, and hence

$$(61) \quad u^{dm}x_2^{q^d} - x_2 \equiv -\alpha_1 \pmod{\mathfrak{p}_{N_0}}.$$

Recalling $\chi^{\text{nr}}(F_N) = u^{dm} \equiv 1 \pmod{p}$ and $v_N(\theta_1) = 1$ we conclude

$$\begin{aligned} (1 + x_2\theta_1)^{\chi^{\text{nr}}(F_N)F_N-1} &= (1 + x_2^{F_N}\theta_1)^{\chi^{\text{nr}}(F_N)}(1 + x_2\theta_1)^{-1} \\ &\equiv (1 + u^{dm}x_2^{q^d}\theta_1)(1 - x_2\theta_1) \pmod{\mathfrak{p}_{N_0}^2} \\ &\equiv 1 + u^{dm}x_2^{q^d}\theta_1 - x_2\theta_1 \pmod{\mathfrak{p}_{N_0}^2} \\ &\equiv 1 - \alpha_1\theta_1 \pmod{\mathfrak{p}_{N_0}^2}, \end{aligned}$$

where the last congruence follows from (61). ■

As in [BC16, Lemma 3.1.5], we can use Lemma 4.1.3 to find an element $\gamma \in U_{N_0}^{(1)}$ such that $\gamma^{\chi^{\text{nr}}(F_N)F_N-1} = \theta_1^{a-1}$. If $\omega > 0$, using also Lemma 7.2.2, we can assume that $\gamma \equiv 1 + x_2\theta_1 \pmod{\mathfrak{p}_{N_0}^2}$.

Let

$$W' = \mathbb{Z}_p[G]z_1 \oplus \mathbb{Z}_p[G]z_2, \quad W_{\geq n} = \bigoplus_{j=n}^{p-1} \bigoplus_{k=1}^m \mathbb{Z}_p[G]v_{k,j}, \quad W = W_{\geq 0}.$$

Note that these are the same as the p -completions of \mathcal{F}' , $\mathcal{F}_{\geq n}$, $\mathcal{F} = \mathcal{F}_{\geq 0}$ in [BC16], but here we prefer to change the notation to avoid confusion with the notation for the Lubin–Tate formal group.

Note that the assignment $v_{k,j} \mapsto \alpha_k w_j$ induces an isomorphism

$$(62) \quad W \xrightarrow{\cong} \bigoplus_{j=0}^{p-1} \bigoplus_{k=1}^m \mathbb{Z}_p[G]\alpha_k w_j = \bigoplus_{j=0}^{p-1} \mathcal{O}_K[G]w_j$$

of free $\mathbb{Z}_p[G]$ -modules. We will always identify W with $\bigoplus_{j=0}^{p-1} \mathcal{O}_K[G]w_j$.

We set

$$(63) \quad E = \begin{cases} 1 & \text{if } \omega = 0, \\ \sum_{i=0}^{dm-1} u^{-i}(A\theta_2)^{\varphi^i} & \text{if } \omega > 0, \end{cases}$$

and

$$(64) \quad \tilde{u} = \begin{cases} 1 & \text{if } \omega = 0, \\ u & \text{if } \omega > 0. \end{cases}$$

Recall that in Lemma 4.1.1 we defined $\varepsilon \in \overline{\mathbb{Z}_p^{\text{nr}}}^\times$ such that $u = \varepsilon^{\varphi-1}$.

LEMMA 7.2.3. *We have $E \in \mathcal{O}_{K'}^\times$, and $E^\varphi \equiv \tilde{u}E \pmod{p\mathcal{O}_{K'}}$. In particular, if $\omega > 0$ then we can also assume $\varepsilon \equiv E \pmod{p\mathcal{O}_{K'}}$.*

Proof. For $\omega = 0$ there is nothing to prove. So let us assume $\omega > 0$.

Since $A\theta_2$ is an integral normal basis generator for the unramified extension K'/\mathbb{Q}_p , the elements $(A\theta_2)^{\varphi^i}$ are linearly independent modulo $p\mathcal{O}_{K'}$. Hence $E \not\equiv 0 \pmod{p\mathcal{O}_{K'}}$ and we deduce $E \in \mathcal{O}_{K'}^\times$.

Moreover,

$$E^\varphi = u \sum_{i=0}^{dm-1} u^{-(i+1)} (A\theta_2)^{\varphi^{i+1}} \equiv uE \pmod{p\mathcal{O}_{K'}}$$

because $u^{dm} \equiv 1 \pmod{p\mathcal{O}_{K'}}$ as a consequence of $0 < \omega = v_p(1 - \chi^{\text{nr}}(F_N))$ and $\chi^{\text{nr}}(F_N) = u^{dm}$. ■

We define

$$\tilde{f}_{3,W} : W \rightarrow \mathcal{F}(\mathfrak{p}_N) \quad \text{by} \quad \tilde{f}_{3,W}(v_{k,j}) = E\alpha_k(a-1)^j\theta$$

for all k and j , where $\theta = \theta_1\theta_2$. We denote by $f_{3,W}$ the composition of $\tilde{f}_{3,W}$ with the projection to $\mathcal{F}(\mathfrak{p}_N)/\mathcal{F}(\mathfrak{p}_N^{p+1})$.

The following lemma is the analogue of [BC16, Lemma 4.1.3].

LEMMA 7.2.4. *The map $f_{3,W}$ is surjective. More precisely, for $j \geq 0$,*

$$f_{3,W}(W_{\geq j}) = \mathcal{F}(\mathfrak{p}_N)^{j+1}/\mathcal{F}(\mathfrak{p}_N^{p+1}).$$

Proof. For $j = p$ we have $W_{\geq p} = \{0\}$ and $\mathcal{F}(\mathfrak{p}_N^{p+1})/\mathcal{F}(\mathfrak{p}_N^{p+1}) = \{0\}$, so the result is trivial.

We assume the result for $j + 1$ and proceed by descending induction. Recall that by [BC16, Lemma 3.2.6], $\mathfrak{p}_N^{j+1} = (p, (a-1)^j)\theta$. Hence if $x \in \mathcal{F}(\mathfrak{p}_N)^{j+1}$, then

$$E^{-1}x \equiv \mu p\theta + \nu(a-1)^j\theta \pmod{\mathfrak{p}_N^{j+2}}$$

for some $\mu, \nu \in \mathcal{O}_K[G]$. We write $\nu = \sum_{k,\ell} \nu_{k,\ell} a^k b^\ell$. Let $\tilde{\nu} = \sum_{k,\ell} \tilde{u}^{m\ell} \nu_{k,\ell} a^k b^\ell$. By Lemma 7.2.3 we have

$$\begin{aligned} \tilde{u}^{ml} a^k b^\ell (E(a-1)^j\theta) &= \tilde{u}^{ml} E^{\varphi^{-ml}} \cdot a^k b^\ell (a-1)^j\theta \\ &\equiv E \cdot a^k b^\ell (a-1)^j\theta \pmod{\mathfrak{p}_N^{p+1}}. \end{aligned}$$

Since $\mu p\theta \in \mathfrak{p}_N^{p+1}$, we further conclude that

$$x \equiv E\nu(a-1)^j\theta \equiv \tilde{\nu}E(a-1)^j\theta \pmod{\mathfrak{p}_N^{p+1}}.$$

By the analogue of [BC16, Lemma 4.1.7] we have

$$\tilde{\nu}E(a-1)^j\theta \equiv \tilde{f}_{3,W}(\tilde{\nu}w_j) \pmod{\mathfrak{p}_N^{j+2}}$$

and thus $x \equiv \tilde{f}_{3,W}(\tilde{\nu}w_j) \pmod{\mathfrak{p}_N^{j+2}}$. This means that $\pi(x)$ is the sum of an element in the image of $W_{\geq j}$ and an element in $\mathcal{F}(\mathfrak{p}_N)^{j+2}/\mathcal{F}(\mathfrak{p}_N^{p+1})$, which is by assumption in the image of $W_{\geq j+1} \subseteq W_{\geq j}$. ■

As in [BC16], we will need to construct some particular elements in the kernel of $f_{3,W}$. We will proceed as in [BC16, Lemmas 4.2.4–4.2.6].

LEMMA 7.2.5. *Let $0 \leq j \leq p - 1$ and $1 \leq k \leq m$. Then there exists $\mu_{j,k} \in W_{\geq j+2}$ such that the element*

$$s_{j,k} = \alpha_k(a - 1)w_j - \alpha_k w_{j+1} + \mu_{j,k}$$

is in the kernel of $f_{3,W}$. Here w_p should be interpreted as 0.

Proof. The formal subtraction of X and Y takes the form

$$X -_{\mathcal{F}} Y = X - Y + AXY + BX^2 + CY^2 + \text{deg} \geq 3$$

with $A, B, C \in \mathbb{Z}_p$. Taking $Y = 0$ we see that $B = 0$, taking $X = Y$ we then get $C = -A$. Hence

$$X -_{\mathcal{F}} Y = X - Y + AXY - AY^2 + \text{deg} \geq 3.$$

For

$$x := E\alpha_k(a - 1)^j a\theta, \quad y := E\alpha_k(a - 1)^j \theta, \quad z := x - y = E\alpha_k(a - 1)^{j+1} \theta,$$

from [BC16, Lemma 3.2.5] and Lemma 7.2.3 we obtain

$$v_N(x) = v_N(y) = j + 1, \quad v_N(z) = v_N(x - y) \geq j + 2.$$

In the following computation all congruences are modulo \mathfrak{p}_N^{j+3} :

$$\begin{aligned} x -_{\mathcal{F}} y -_{\mathcal{F}} z &\equiv x - y + Axy - Ay^2 - z + A(x - y + Axy - Ay^2)z - Az^2 \\ &\equiv z + Ayz - z + A(z + Ayz)z - Az^2 \equiv 0. \end{aligned}$$

Therefore

$$\begin{aligned} &\tilde{f}_{3,W}(\alpha_k(a - 1)w_j - \alpha_k w_{j+1}) \\ &= E\alpha_k(a - 1)^j a\theta -_{\mathcal{F}} E\alpha_k(a - 1)^j \theta -_{\mathcal{F}} E\alpha_k(a - 1)^{j+1} \theta \equiv 0 \pmod{\mathcal{F}(\mathfrak{p}_N^{j+3})}. \end{aligned}$$

Using Lemma 7.2.4, by the same arguments used in the proof of [BC16, Lemma 4.2.4] the result follows. ■

Recall that $(m, d) = 1$. Let \tilde{m} denote an integer such that $m\tilde{m} \equiv 1 \pmod{d}$. The next lemma is the analogue of [BC16, Lemmas 4.2.5, 4.2.6].

LEMMA 7.2.6. *The elements*

$$\begin{aligned} r_1 &= \alpha_1 \mathcal{T}_a w_0 + (u^{-1} \tilde{u}^{1-m\tilde{m}} b^{-\tilde{m}} \alpha_1 - \alpha_1) w_{p-1}, \\ r_k &= \alpha_k \mathcal{T}_a w_0 + (u^{-1} \tilde{u}^{1-m\tilde{m}} b^{-\tilde{m}} \alpha_{k+1} - \alpha_k) w_{p-1} \quad \text{for } 1 < k < m, \\ r_m &= \alpha_m \mathcal{T}_a w_0 + \left(u^{-1} \tilde{u}^{1-m\tilde{m}} b^{-\tilde{m}} \alpha_1 - u^{-1} \tilde{u}^{1-m\tilde{m}} b^{-\tilde{m}} \sum_{i=2}^m \alpha_i - \alpha_m \right) w_{p-1} \end{aligned}$$

are in the kernel of $f_{3,W}$.

Proof. Recall the isomorphism

$$f_{\mathcal{F},N}: \mathcal{F}(\mathfrak{p}_N) \rightarrow (\widehat{N_0^\times}(\chi^{\text{nr}}))^{\text{Gal}(N_0/N)}$$

from Lemma 4.1.1. By [Ser79, Prop. V.6.8], we know that $\mathcal{N}_{N_0/K_0} U_{N_0}^{(2)} \subseteq U_{K_0}^{(2)} \subseteq U_{N_0}^{(2p)} \subseteq U_{N_0}^{(p+1)}$. This implies that

$$\mathcal{N}_{N_0/K_0}(f_{\mathcal{F},N}(E\alpha_k\theta)) \equiv \mathcal{N}_{N_0/K_0}(1 + \varepsilon^{-1}E\alpha_k\theta) \pmod{\mathfrak{p}_{N_0}^{p+1}}.$$

Now we calculate $\tilde{f}_{3,W}(\alpha_k \mathcal{T}_a w_0)$, using the results and calculations of [BC16] and the fact that $\theta_2^p \equiv \theta_2^{p^{m\tilde{m}}} \equiv \theta_2^{q^{\tilde{m}}} \equiv \theta_2^{b^{-\tilde{m}}} \pmod{\mathfrak{p}_{\tilde{K}'}}$:

$$\begin{aligned} \tilde{f}_{3,W}(\alpha_k \mathcal{T}_a w_0) &= \mathcal{T}_a E\alpha_k\theta = f_{\mathcal{F},N}^{-1}(\mathcal{T}_a \cdot (f_{\mathcal{F},N}(E\alpha_k\theta))) \\ &= f_{\mathcal{F},N}^{-1}(\mathcal{N}_{N_0/K_0}(f_{\mathcal{F},N}(E\alpha_k\theta))) \\ &\equiv f_{\mathcal{F},N}^{-1}(\mathcal{N}_{N_0/K_0}(1 + \varepsilon^{-1}E\alpha_k\theta)) \pmod{\mathfrak{p}_{N_0}^{p+1}} \\ &\equiv f_{\mathcal{F},N}^{-1}\left(1 + \left(\frac{\varepsilon^{-1}E\alpha_k\theta_2}{\alpha_1} - \left(\frac{\varepsilon^{-1}E\alpha_k\theta_2}{\alpha_1}\right)^p\right)\alpha_1 p\right) \pmod{\mathfrak{p}_{N_0}^{p+1}} \\ &\equiv \left(\frac{\alpha_k\theta_2}{\alpha_1} - \varepsilon^{1-p}E^{p-1}\left(\frac{\alpha_k\theta_2}{\alpha_1}\right)^p\right)E\alpha_1 p \pmod{\mathfrak{p}_{N_0}^{p+1}} \\ &\equiv \left(\alpha_k\theta_2 - u^{-1}\tilde{u}\alpha_1\left(\frac{\alpha_k}{\alpha_1}\right)^p\theta_2^{b^{-\tilde{m}}}\right)Ep \pmod{\mathfrak{p}_{N_0}^{p+1}}. \end{aligned}$$

By (60) we have

$$\left(\frac{\alpha_k}{\alpha_1}\right)^p = (A^{\varphi^{k-2}})^p \equiv A^{p^{k-1}} \equiv \frac{\alpha_{k+1}}{\alpha_1} \pmod{\mathfrak{p}_K} \quad \text{for } 1 < k < m,$$

and

$$\left(\frac{\alpha_m}{\alpha_1}\right)^p = (A^{\varphi^{m-2}})^p \equiv A^{\varphi^{m-1}} = 1 - \sum_{i=0}^{m-2} A^{\varphi^i} = 1 - \sum_{i=0}^{m-2} \frac{\alpha_{i+2}}{\alpha_1} \pmod{\mathfrak{p}_K}.$$

Hence we obtain the following congruences modulo \mathfrak{p}_N^{p+1} :

$$\tilde{f}_{3,W}(\alpha_k \mathcal{T}_a w_0) \equiv \begin{cases} (\alpha_1\theta_2 - u^{-1}\tilde{u}\alpha_1\theta_2^{b^{-\tilde{m}}})Ep & \text{if } k = 1, \\ (\alpha_k\theta_2 - u^{-1}\tilde{u}\alpha_{k+1}\theta_2^{b^{-\tilde{m}}})Ep & \text{if } 1 < k < m, \\ \left(\alpha_m\theta_2 + u^{-1}\tilde{u}\left(-\alpha_1 + \sum_{i=2}^m \alpha_i\right)\theta_2^{b^{-\tilde{m}}}\right)Ep & \text{if } k = m. \end{cases}$$

Recalling Lemma 7.2.3 and [BC16, Lemma 3.2.4], for $1 \leq i \leq m$ we calculate

$$\begin{aligned} \tilde{f}_{3,W}(u^{-1}\tilde{u}^{1-m\tilde{m}}b^{-\tilde{m}}\alpha_i w_{p-1}) &= u^{-1}\tilde{u}^{1-m\tilde{m}}b^{-\tilde{m}}(E\alpha_i(a-1)^{p-1}\theta) \\ &= u^{-1}\tilde{u}^{1-m\tilde{m}}(Eb^{-\tilde{m}}\alpha_i^{b^{-\tilde{m}}}(a-1)^{p-1}\theta^{b^{-\tilde{m}}}) \\ &= u^{-1}\tilde{u}^{1-m\tilde{m}}E\varphi^{m\tilde{m}}\alpha_i\theta_2^{b^{-\tilde{m}}}(a-1)^{p-1}\theta_1 \\ &\equiv u^{-1}\tilde{u}E\alpha_i\theta_2^{b^{-\tilde{m}}}p \pmod{\mathfrak{p}_N^{p+1}} \end{aligned}$$

and

$$\tilde{f}_{3,W}(\alpha_i w_{p-1}) = E\alpha_i(a-1)^{p-1}\theta \equiv E\alpha_i\theta_2 p \pmod{\mathfrak{p}_N^{p+1}}.$$

For $x, y \in \mathfrak{p}_N^p$ we see that $x - \mathcal{F}y \equiv x - y + Axy - Ay^2 \equiv x - y \pmod{\mathfrak{p}_N^{p+1}}$, so that we easily deduce the lemma by the above calculations. ■

From now on we will need to distinguish between the cases $\omega = 0$ and $\omega > 0$.

7.3. The Euler characteristic when $\omega = 0$. As already mentioned, we use $\mathcal{L} = \mathfrak{p}_N^{p+1}$ and thus have $X(\mathcal{L}) = \mathcal{F}(\mathfrak{p}_N^{p+1})$. Since $\omega = 0$, the second cohomology group of $M^\bullet(\mathcal{L})$ vanishes and we have the short exact sequence

$$0 \rightarrow \mathcal{F}(\mathfrak{p}_N)/\mathcal{F}(\mathfrak{p}_N^{p+1}) \rightarrow A/X(\mathcal{L}) \rightarrow B \rightarrow 0.$$

Recall that for $\omega = 0$ both $\mathcal{F}(\mathfrak{p}_N)$ and $\mathcal{F}(\mathfrak{p}_N^{p+1})$ are $\mathbb{Z}_p[G]$ -projective by Lemma 7.2.1. Hence, recalling also (53), we derive

$$\chi_{\mathbb{Z}_p[G], B_{\text{dR}}[G]}(M^\bullet(\mathcal{L}), \lambda_2^{-1}) = [\mathcal{F}(\mathfrak{p}_N^{p+1}), \text{id}, \mathcal{F}(\mathfrak{p}_N)].$$

By Lemma 7.2.4, the map $f_{3,W}: W \rightarrow \mathcal{F}(\mathfrak{p}_N)/\mathcal{F}(\mathfrak{p}_N^{p+1})$ is onto. Thus we obtain the short exact sequence

$$0 \rightarrow \ker(f_{3,W}) \rightarrow W \xrightarrow{f_{3,W}} \mathcal{F}(\mathfrak{p}_N)/\mathcal{F}(\mathfrak{p}_N^{p+1}) \rightarrow 0,$$

which implies

$$[\mathcal{F}(\mathfrak{p}_N^{p+1}), \text{id}, \mathcal{F}(\mathfrak{p}_N)] = [\ker(f_{3,W}), \text{id}, W].$$

Recall the definition of the elements $s_{j,k}$ and r_k in Lemmas 7.2.5 and 7.2.6.

LEMMA 7.3.1. *The pm elements $r_k, s_{j,k}$ for $0 \leq j \leq p - 2, 1 \leq k \leq m$ constitute a $\mathbb{Z}_p[G]$ -basis of $\ker(f_{3,W})$.*

Proof. At the end we want to adapt the proofs of [BC16, Lemmas 4.2.7 and 4.2.9], but first we need some preparations. We write the coefficients of the $\alpha_i w_{p-1}$ -components, $i = 1, \dots, m$, of the elements $r_j, j = 1, \dots, m$, into the columns of an $m \times m$ matrix which we call \mathcal{M} ,

$$\mathcal{M} = \begin{pmatrix} u^{-1}b^{-\tilde{m}} - 1 & 0 & 0 & \cdots & 0 & 0 & u^{-1}b^{-\tilde{m}} \\ 0 & -1 & 0 & \cdots & 0 & 0 & -u^{-1}b^{-\tilde{m}} \\ 0 & u^{-1}b^{-\tilde{m}} & -1 & \cdots & 0 & 0 & -u^{-1}b^{-\tilde{m}} \\ 0 & 0 & u^{-1}b^{-\tilde{m}} & \cdots & 0 & 0 & -u^{-1}b^{-\tilde{m}} \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots \\ 0 & 0 & 0 & \cdots & u^{-1}b^{-\tilde{m}} & -1 & -u^{-1}b^{-\tilde{m}} \\ 0 & 0 & 0 & \cdots & 0 & u^{-1}b^{-\tilde{m}} & -1 - u^{-1}b^{-\tilde{m}} \end{pmatrix}.$$

By an easy induction argument we see that each $n \times n$ matrix

$$M_{C,n} = \begin{pmatrix} -1 & 0 & \cdots & 0 & 0 & -C \\ C & -1 & \cdots & 0 & 0 & -C \\ 0 & C & \cdots & 0 & 0 & -C \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots \\ 0 & 0 & \cdots & C & -1 & -C \\ 0 & 0 & \cdots & 0 & C & -1 - C \end{pmatrix}$$

has determinant $(-1)^n \sum_{i=0}^n C^i$. Setting $C := u^{-1}b^{-\tilde{m}}$ we obtain

$$\begin{aligned} \det(\mathcal{M}) &= (u^{-1}b^{-\tilde{m}} - 1)(-1)^{m-1} \sum_{i=0}^{m-1} (u^{-1}b^{-\tilde{m}})^i \\ &= (-1)^{m-1}(u^{-m}b^{-\tilde{m}m} - 1) = (-1)^{m-1}(u^{-m}b^{-1} - 1). \end{aligned}$$

We claim that $\det(\mathcal{M})$ is a unit in $\mathbb{Z}_p[G]$. To prove this it is sufficient to show that $\det(\mathcal{M})$ is a unit in the maximal \mathbb{Z}_p -order A' in $\mathbb{Q}_p[G]$ because $(A')^\times \cap \mathbb{Z}_p[G] = \mathbb{Z}_p[G]^\times$. For any irreducible character ψ of G we have $\psi(\det(\mathcal{M})) = (-1)^{m-1}(u^{-m}\zeta_d - 1)$ for some d th root of unity ζ_d (depending on ψ). If now $\psi(\det(\mathcal{M}))$ were not a unit, we would conclude that $u^{dm} = \chi^{\text{nr}}(F_N) \equiv 1 \pmod{p}$, contradicting the assumption $\omega = 0$.

We are now prepared to apply the arguments used to prove [BC16, Lemmas 4.2.7 and 4.2.9]. Following the proof of [BC16, Lemma 4.2.7], we can demonstrate that the $pm + m$ elements $r_k, s_{j,k}$ for $0 \leq j \leq p - 1, 1 \leq k \leq m$ generate $\ker(f_{3,W})$ as a $\mathbb{Z}_p[G]$ -module. Since the determinant of \mathcal{M} is a unit, it can be shown as in the proof of [BC16, Lemma 4.2.9] that for $k = 1, \dots, m$ we have $s_{p-1,k} \in \langle r_1, \dots, r_m \rangle_{\mathbb{Z}_p[G]}$. Therefore the elements listed in the statement of our lemma generate $\ker(f_{3,W})$. We actually have a basis since $\ker(f_{3,W})$ is free of $\mathbb{Z}_p[G]$ -rank $\text{rk}_{\mathbb{Z}_p[G]}(W) = pm$. ■

PROPOSITION 7.3.2. *Assume that $\omega = 0$. For $\mathcal{L} = \mathfrak{p}_N^{p+1}$ the element*

$$\chi_{\mathbb{Z}_p[G], B_{\text{dR}}[G]}(M^\bullet(\mathcal{L}), \lambda_2^{-1}) \in K_0(\mathbb{Z}_p[G], B_{\text{dR}}[G])$$

is contained in $K_0(\mathbb{Z}_p[G], \mathbb{Q}_p[G])$ and represented by $\varepsilon \in \mathbb{Q}_p[G]^\times$ where

$$\varepsilon_{\chi\phi} = \begin{cases} p^m & \text{if } \chi = \chi_0, \\ (-1)^{m-1}(u^{-m}\phi(b)^{-1} - 1)(\chi(a) - 1)^{m(p-1)} & \text{if } \chi \neq \chi_0. \end{cases}$$

Proof. We choose the elements r_k and $s_{j,k}$ of Lemma 7.3.1 as a $\mathbb{Z}_p[G]$ -basis of $\ker(f_{3,W})$ and fix the canonical $\mathbb{Z}_p[G]$ -basis of W . Then

$$\chi_{\mathbb{Z}_p[G], B_{\text{dR}}[G]}(M^\bullet(\mathcal{L}), \lambda_2^{-1}) = [\ker(f_{3,W}), \text{id}, W]$$

is represented by the determinant of

$$\mathfrak{M} = \begin{pmatrix} \mathcal{T}_a I & (a-1)I & 0 & \cdots & 0 & 0 \\ 0 & -I & (a-1)I & \cdots & 0 & 0 \\ 0 & * & -I & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & * & * & \cdots & -I & (a-1)I \\ \mathcal{M} & * & * & \cdots & * & -I \end{pmatrix}.$$

Recalling that p is odd, we get

$$\begin{aligned} \det(\chi\phi(\mathfrak{M})) &= \begin{cases} p^m(-1)^{m(p-1)} & \text{if } \chi = \chi_0, \\ (-1)^{m^2(p-1)} \det(\chi\phi(\mathcal{M}))(\chi(a) - 1)^{m(p-1)} & \text{if } \chi \neq \chi_0 \end{cases} \\ &= \begin{cases} p^m & \text{if } \chi = \chi_0, \\ (-1)^{m-1}(u^{-m}\phi(b)^{-1} - 1)(\chi(a) - 1)^{m(p-1)} & \text{if } \chi \neq \chi_0. \blacksquare \end{cases} \end{aligned}$$

7.4. The Euler characteristic when $\omega > 0$. Recall the definitions of E and \tilde{u} in (63) and (64) respectively.

LEMMA 7.4.1. *There is a commutative diagram of $\mathbb{Z}_p[G]$ -modules with exact rows*

$$\begin{array}{ccccccccc} 0 & \longrightarrow & X(2) \oplus W & \longrightarrow & W' \oplus W & \xrightarrow{\delta_2} & \mathbb{Z}_p[G]z_0 & \xrightarrow{\pi} & \mathbb{Z}_p/p^\omega\mathbb{Z}_p(\chi^{\text{nr}}) & \longrightarrow & 0 \\ & & \downarrow \tilde{f}_4 & & \downarrow \tilde{f}_3 & & \downarrow \tilde{f}_2 & & \downarrow = & & \\ 0 & \longrightarrow & \mathcal{F}(\mathfrak{p}_N) & \xrightarrow{f_{\mathcal{F},N}} & \mathcal{I}_{N/K}(\chi^{\text{nr}}) & \xrightarrow{(F-1)\times 1} & \mathcal{I}_{N/K}(\chi^{\text{nr}}) & \longrightarrow & \mathbb{Z}_p/p^\omega\mathbb{Z}_p(\chi^{\text{nr}}) & \longrightarrow & 0 \end{array}$$

where

$$\begin{aligned} \delta_2(z_1) &= (u^m b - 1)z_0, & \tilde{f}_3(z_1) &= [\theta_1, 1, \dots, 1], \\ \delta_2(z_2) &= (a - 1)z_0, & \tilde{f}_3(z_2) &= [\gamma, \dots, \gamma], \\ \delta_2(v_{j,k}) &= 0 \text{ for all } k \text{ and } j, & \tilde{f}_3(v_{k,j}) &= f_{\mathcal{F},N}(\tilde{f}_{3,W}(v_{k,j})), \\ \tilde{f}_2(z_0) &= [\theta_1, 1, \dots, 1], & \pi(z_0) &= 1. \end{aligned}$$

Further, $X(2) = \ker(\delta_2|_{W'})$ and \tilde{f}_4 is the restriction of \tilde{f}_3 to $X(2) \oplus W$.

Proof. We first remark that the natural G -structure on $\mathcal{I}_{N/K}(\chi^{\text{nr}})$ given by the inclusion $G \hookrightarrow \text{Gal}(K_{\text{nr}}/K) \times G$ induces on $\mathbb{Z}_p/p^\omega\mathbb{Z}_p(\chi^{\text{nr}})$ the G -structure characterized by

$$a \cdot 1 = 1, \quad b \cdot 1 = \chi^{\text{nr}}(F^{-1}).$$

Note that this is well-defined because $\chi^{\text{nr}}(F)^d = \chi^{\text{nr}}(F_N) \equiv 1 \pmod{p^\omega}$ by the definition of ω .

The exactness of the bottom 2-extension is shown in Theorem 4.2.3. For the exactness of the top row we have to verify exactness at $\mathbb{Z}_p[G]z_0$; the rest

is clear by the definitions. If we set $H := \langle (a - 1)z_0, (u^mb - 1)z_0 \rangle$, then we must prove $H = \ker(\pi)$. The inclusion $H \subseteq \ker(\pi)$ is implied by

$$\pi((a - 1)z_0) = (a - 1) \cdot 1 = 0, \quad \pi((u^mb - 1)z_0) = (\chi^{\text{nr}}(F)b - 1) \cdot 1 = 0.$$

Let $\lambda z_0 \in \ker(\pi)$ with $\lambda \in \mathbb{Z}_p[G]$. Then clearly $\lambda z_0 \equiv xz_0 \pmod{H}$ for some $x \in \mathbb{Z}_p$ with $v_p(x) \geq \omega$. Since $\omega = v_p(1 - \chi^{\text{nr}}(F_N))$ and $\chi^{\text{nr}}(F_N) = u^{dm}$, we can write $x = (u^{dm} - 1)y$ with $y \in \mathbb{Z}_p$. Hence $xz_0 = ((u^mb)^d - 1)y z_0 \in H$.

The proof of commutativity is straightforward using $\theta_1^{a-1} = \gamma^{\chi^{\text{nr}}(F_N)F_N-1}$, which holds by the definition of γ . For the convenience of the reader we give the computations:

$$\begin{aligned} \tilde{f}_2(\delta_2(z_1)) &= \tilde{f}_2((\chi^{\text{nr}}(F)b - 1)z_0) = \frac{(F \times 1)(F^{-1} \times b)[\theta_1^{\chi^{\text{nr}}(F)}, 1, \dots, 1]}{[\theta_1, 1, \dots, 1]} \\ &= \frac{(F \times 1)[\theta_1, 1, \dots, 1]}{[\theta_1, 1, \dots, 1]} = ((F - 1) \times 1)\tilde{f}_3(z_1), \\ \tilde{f}_2(\delta_2(z_2)) &= \tilde{f}_2((a - 1)z_0) = \frac{(1 \times a)[\theta_1, 1, \dots, 1]}{[\theta_1, 1, \dots, 1]} = [\theta_1^{a-1}, 1, \dots, 1] \\ &= [\gamma^{\chi^{\text{nr}}(F_N)F_N-1}, 1, \dots, 1] = \frac{[\gamma^{\chi^{\text{nr}}(F_N)F_N}, \gamma, \dots, \gamma]}{[\gamma, \dots, \gamma]} \\ &= ((F - 1) \times 1)[\gamma, \dots, \gamma] = ((F - 1) \times 1)\tilde{f}_3(z_2). \blacksquare \end{aligned}$$

We let $f_4: X(2) \oplus W \rightarrow \mathcal{F}(\mathfrak{p}_N)/\mathcal{F}(\mathfrak{p}_N^{p+1})$ denote the composite of \tilde{f}_4 with the canonical projection.

LEMMA 7.4.2. *The complex*

$$F^\bullet := [\ker(f_4) \rightarrow W' \oplus W \rightarrow \mathbb{Z}_p[G]z_0]$$

with modules in degrees 0, 1 and 2 is a representative of $M^\bullet(\mathcal{L})$ for $\mathcal{L} = \mathfrak{p}_N^{p+1}$.

Proof. By Lemmas 7.4.1 and 7.2.4 we have a quasi-isomorphism of complexes

$$F^\bullet \rightarrow [\mathcal{I}_{N/K}(\chi^{\text{nr}})/f_{\mathcal{F},N}(\mathcal{F}(\mathfrak{p}_N^{p+1})) \rightarrow \mathcal{I}_{N/K}(\chi^{\text{nr}})].$$

By Theorem 4.3.1 the right hand side represents $M^\bullet(\mathcal{L})$. \blacksquare

Our aim is to use Lemma 7.4.2 to compute the Euler characteristic

$$\chi_{\mathbb{Z}_p[G], B_{\text{dR}}[G]}(M^\bullet(\mathcal{L}), \lambda_2^{-1}).$$

We therefore must compute $\ker(f_4)$ explicitly. A first step in this direction is the following lemma which is similar to [BC16, Lemma 4.1.5].

LEMMA 7.4.3. *We have*

$$X(2) = \langle \mathcal{T}_a z_2, (a - 1)z_1 - (u^mb - 1)z_2 \rangle_{\mathbb{Z}_p[G]}.$$

Proof. The inclusion “ \supseteq ” is clear from the definition of δ_2 . For the reverse inclusion we let

$$x = \sum_{i=0}^{p-1} \sum_{j=0}^{d-1} \alpha_{i,j} a^i b^j z_1 + \sum_{i=0}^{p-1} \sum_{j=0}^{d-1} \beta_{i,j} a^i b^j z_2$$

be an element in $X(2)$ with $\alpha_{i,j}, \beta_{i,j} \in \mathbb{Z}_p$. From $\delta_2(x) = 0$ we derive

$$(65) \quad u^m \alpha_{i,j-1} - \alpha_{i,j} + \beta_{i-1,j} - \beta_{i,j} = 0$$

for all $0 \leq i < p$ and $0 \leq j < d$. Here and in the following we regard all indices as integers modulo p and d respectively.

Taking the sum over the index i we deduce

$$(66) \quad u^m \sum_{i=0}^{p-1} \alpha_{i,j-1} = \sum_{i=0}^{p-1} \alpha_{i,j}$$

for all $0 \leq j < d$. Applying (66) repeatedly we get

$$\sum_{i=0}^{p-1} \alpha_{i,0} = u^m \sum_{i=0}^{p-1} \alpha_{i,d-1} = \dots = u^{dm} \sum_{i=0}^{p-1} \alpha_{i,0}.$$

As $u^{dm} = \chi^{\text{nr}}(F_N) \neq 1$ by assumption, we obtain $\sum_{i=0}^{p-1} \alpha_{i,0} = 0$. Using (66) again, we find that

$$(67) \quad \sum_{i=0}^{p-1} \alpha_{i,j} = 0$$

for all $0 \leq j < d$. So in particular the above sum does not depend on the choice of j , hence we can continue our proof exactly as in [BC16, Lemma 4.1.5]. Note that with the notation of [BC16] we would obtain $\mu_0 = 0$, so that in the present situation two generators are enough. We include the computations for the convenience of the reader.

We want to find $\gamma_{i,j}$ and ν_j in \mathbb{Z}_p such that

$$\begin{aligned} x &= \sum_{i=0}^{p-1} \sum_{j=0}^{d-1} \gamma_{i,j} a^i b^j ((a-1)z_1 - (u^m b - 1)z_2) + \sum_{j=0}^{d-1} \nu_j b^j \mathcal{T}_a z_2 \\ &= \sum_{i=0}^{p-1} \sum_{j=0}^{d-1} (\gamma_{i-1,j} - \gamma_{i,j}) a^i b^j z_1 + \sum_{i=0}^{p-1} \sum_{j=0}^{d-1} (-u^m \gamma_{i,j-1} + \gamma_{i,j} + \nu_j) a^i b^j z_2. \end{aligned}$$

So we need to solve

$$(68) \quad \gamma_{i-1,j} - \gamma_{i,j} = \alpha_{i,j},$$

$$(69) \quad -u^m \gamma_{i,j-1} + \gamma_{i,j} + \nu_j = \beta_{i,j}.$$

We set $\gamma_{i,j} := -\sum_{1 \leq \ell \leq i} \alpha_{\ell,j}$ and $\nu_j = \beta_{0,j}$. Then equation (68) is clearly satisfied for $i = 1, \dots, p-1$, and for $i = 0$ it is immediately implied by (67). Equation (69) is proved by an easy induction on i using (65). ■

Still following the conventions in [BC16] we use $[x]$ to denote the element $[x, \dots, x] \in \mathcal{I}_{N/K}(\chi^{\text{nr}})$ or even in $\mathcal{I}_{N/K}(\chi^{\text{nr}})/f_{\mathcal{F},N}(\mathcal{F}(\mathfrak{p}_N^{p+1}))$.

Now we need to evaluate f_4 (or equivalently \tilde{f}_3) at the generators of $W(2)$ determined in the previous lemma, as we did in [BC16, Lemma 4.1.6].

LEMMA 7.4.4. *We have*

$$\tilde{f}_3((a-1)z_1 - (u^m b - 1)z_2) = [\gamma^{1-b}], \quad \tilde{f}_3(\mathcal{T}_a z_2) = [\gamma^{\mathcal{T}_a}].$$

Proof. Using (19), (20), $\theta_1^{a-1} = \gamma^{\chi^{\text{nr}}(F_N)F_N^{-1}}$, $\theta_1^b = \theta_1$ and $\chi^{\text{nr}}(F^{-1}) = u^{-m}$ we obtain

$$\begin{aligned} \tilde{f}_3((a-1)z_1 - (u^m b - 1)z_2) &= \frac{[\theta_1^{a-1}, 1, \dots, 1][\gamma, \dots, \gamma]}{(F \times 1)(F^{-1} \times b)[\gamma^{u^m}, \dots, \gamma^{u^m}]} \\ &= \frac{[\theta_1^{a-1}\gamma, \gamma, \dots, \gamma]}{(F \times 1)[\gamma^b, \dots, \gamma^b]} = \frac{[\theta_1^{a-1}\gamma, \gamma, \dots, \gamma]}{[\gamma^{\chi^{\text{nr}}(F_N)F_N^b}, \gamma^b, \dots, \gamma^b]} \\ &= \frac{[\theta_1^{a-1}\gamma, \gamma, \dots, \gamma]}{[\theta_1^{(a-1)b}\gamma^b, \gamma^b, \dots, \gamma^b]} = [\gamma^{1-b}, \dots, \gamma^{1-b}]. \end{aligned}$$

The second equality is obvious by the definitions. ■

The following results are the analogues of [BC16, Lemmas 4.2.1 and 4.2.2] and [BC16, Lemma 4.2.2].

LEMMA 7.4.5. *Let \tilde{m} be an integer such that $m\tilde{m} \equiv 1 \pmod{d}$. Set*

$$\begin{aligned} \tilde{t}_1 &:= (a-1)z_1 - (u^m b - 1)z_2 \\ &+ \left(\sum_{i=2}^m \alpha_i (u^m b)^{1-(i-2)\tilde{m}} + \left(\alpha_1 - \sum_{i=2}^m \alpha_i \right) (u^m b)^{\tilde{m}} \right) w_0. \end{aligned}$$

Then there exists $y_1 \in W_{\geq 1}$ such that $t_1 := \tilde{t}_1 + y_1 \in \ker(f_4)$.

Proof. Recalling that γ has been chosen so that $\gamma \equiv 1 + x_2\theta_1 \pmod{\mathfrak{p}_{N_0}^2}$ and following the calculations in the proof of [BC16, Lemma 4.2.1] we get

$$\gamma^{1-b} \equiv (1 + x_2\theta_1)^{1-b} = 1 - \sum_{i=2}^m \alpha_i \theta^{b^{1-(i-2)\tilde{m}}} - \alpha_1 \theta^{b^{\tilde{m}}} + \sum_{i=2}^m \alpha_i \theta^{b^{\tilde{m}}} \pmod{\mathfrak{p}_{N_0}^2}.$$

By Lemmas 4.1.1 and 7.2.3,

$$(70) \quad f_{\mathcal{F},N}(X) \equiv 1 + E^{-1}X + \deg \geq 2 \pmod{p\mathcal{O}_{K'}},$$

and so by Lemma 7.4.4 we deduce

$$\begin{aligned} & \tilde{f}_4((a-1)z_1 - (u^m b - 1)z_2) \\ & \equiv - \sum_{i=2}^m \alpha_i E \theta^{b^{1-(i-2)\tilde{m}}} - \alpha_1 E \theta^{b^{\tilde{m}}} + \sum_{i=2}^m \alpha_i E \theta^{b^{\tilde{m}}} \pmod{\mathfrak{p}_{N_0}^2}. \end{aligned}$$

Since by Lemma 7.2.3, for any integer i ,

$$(u^m b)^i \cdot (E\theta) = u^{mi} E b^i \theta^{b^i} \equiv u^{mi} u^{-mi} E \theta^{b^i} \equiv E \theta^{b^i} \pmod{\mathfrak{p}_{N_0}^2},$$

we have

$$\begin{aligned} & \tilde{f}_4\left(\sum_{i=2}^m \alpha_i (u^m b)^{1-(i-2)\tilde{m}} + \left(\alpha_1 - \sum_{i=2}^m \alpha_i\right) (u^m b)^{\tilde{m}}\right) w_0 \\ & \equiv \left(\sum_{i=2}^m \alpha_i (u^m b)^{1-(i-2)\tilde{m}} + \left(\alpha_1 - \sum_{i=2}^m \alpha_i\right) (u^m b)^{\tilde{m}}\right) \cdot (E\theta) \pmod{\mathfrak{p}_{N_0}^2} \\ & \equiv \sum_{i=2}^m \alpha_i E \theta^{b^{1-(i-2)\tilde{m}}} + \alpha_1 E \theta^{b^{\tilde{m}}} - \sum_{i=2}^m \alpha_i E \theta^{b^{\tilde{m}}} \pmod{\mathfrak{p}_{N_0}^2} \end{aligned}$$

and we can conclude that $\tilde{f}_4(\tilde{t}_1) \equiv 0 \pmod{\mathfrak{p}_{N_0}^2}$. Therefore

$$f_4(\tilde{t}_1) \in \mathcal{F}(\mathfrak{p}_N^2)/\mathcal{F}(\mathfrak{p}_N^{p+1}),$$

and by Lemma 7.2.4 there exists $y_1 \in W_{\geq 1}$ such that $f_4(-y_1) = f_4(\tilde{t}_1)$, i.e. $\tilde{t}_1 + y_1 \in \ker(f_4)$. ■

Note that t_1 is not exactly the same as in [BC16] since there is a u^m factor which appears at all the occurrences of b .

LEMMA 7.4.6. *The element*

$$t_2 := \mathcal{T}_a z_2 - \beta w_{p-1} \quad \text{with} \quad \beta = \begin{cases} \alpha_1 & \text{if } m = 1, \\ \alpha_2 & \text{if } m > 1 \end{cases}$$

is in the kernel of f_4 .

Proof. As in the previous lemma, we can follow the calculations in the proof of [BC16, Lemma 4.2.2]:

$$\gamma^{T_a} \equiv \mathcal{N}_{N_{nr}/K_{nr}}(1 + x_2 \theta_1) \equiv 1 + x_2 p - x_2^p \alpha_1^{1-p} p \equiv 1 + \beta \theta_2 p \pmod{\mathfrak{p}_{N_0}^{p+1}}.$$

As above, we deduce that

$$\tilde{f}_4(\mathcal{T}_a z_2) \equiv E \beta \theta_2 p \pmod{\mathfrak{p}_{N_0}^{p+1}}.$$

Next using [BC16, Lemma 3.2.4] we calculate

$$\tilde{f}_4(\beta w_{p-1}) \equiv E \beta (a-1)^{p-1} \theta \equiv E \beta \theta_2 p \pmod{\mathfrak{p}_{N_0}^{p+1}}.$$

Hence we easily conclude since $\mathcal{F}(\mathfrak{p}_{N_0}^p)/\mathcal{F}(\mathfrak{p}_{N_0}^{p+1}) \cong \mathfrak{p}_{N_0}^p/\mathfrak{p}_{N_0}^{p+1}$. ■

LEMMA 7.4.7. *The element*

$$r_1 = \mathcal{T}_a t_1 + (u^m b - 1)t_2$$

belongs to $\ker(f_4) \cap W$ and its $\alpha_1 w_0$ -component is $(u^m b)^{\tilde{m}} \mathcal{T}_a$.

Proof. Straightforward calculations: see [BC16, proof of Lemma 4.2.5]. ■

Following the strategy in [BC16] we consider the $\alpha_i w_{p-1}$ -components, $i = 1, \dots, m$, of t_2 and r_2, \dots, r_m . We write these components as the columns of an $m \times m$ matrix \mathcal{M} . If $m > 1$, then we obtain

$$\mathcal{M} = \begin{pmatrix} 0 & 0 & 0 & \cdots & 0 & 0 & (u^m b)^{-\tilde{m}} \\ -1 & -1 & 0 & \cdots & 0 & 0 & -(u^m b)^{-\tilde{m}} \\ 0 & (u^m b)^{-\tilde{m}} & -1 & \cdots & 0 & 0 & -(u^m b)^{-\tilde{m}} \\ 0 & 0 & (u^m b)^{-\tilde{m}} & \cdots & 0 & 0 & -(u^m b)^{-\tilde{m}} \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots \\ 0 & 0 & 0 & \cdots & (u^m b)^{-\tilde{m}} & -1 & -(u^m b)^{-\tilde{m}} \\ 0 & 0 & 0 & \cdots & 0 & (u^m b)^{-\tilde{m}} & -1 - (u^m b)^{-\tilde{m}} \end{pmatrix}.$$

If $m = 1$, then the matrix is determined by t_2 , and recalling the definition from Lemma 7.4.6 we get $\mathcal{M} = (-1)$.

LEMMA 7.4.8. *The determinant of \mathcal{M} is $(-1)^m (u^m b)^{-\tilde{m}(m-1)}$.*

Proof. This is an easy calculation. ■

The next proposition is the analogue of [BC16, Prop. 4.2.10].

LEMMA 7.4.9. *The $pm + 1$ elements t_1, t_2, r_k for $k = 2, \dots, m, s_{j,k}$ for $0 \leq j \leq p - 2, 1 \leq k \leq m$, constitute a $\mathbb{Z}_p[G]$ -basis of $\ker(f_4)$.*

Proof. We first note that $s_{p-1,k} \in \langle t_2, r_2, \dots, r_m \rangle_{\mathbb{Z}_p[G]}$, which follows as in [BC16, proof of Lemma 4.2.9]. The main input here is the fact that the matrix \mathcal{M} is invertible.

Moreover, by Lemma 7.4.3, t_1 and t_2 generate $(X(2) + W)/W$, which therefore must coincide with $(\ker(f_4) + W)/W$, and the same proof as for [BC16, Lemma 4.2.7] shows that the elements r_k and $s_{j,k}, 0 \leq j \leq p - 1, 1 \leq k \leq m$, generate $\ker(f_4) \cap W$. We can thus argue as in [BC16, proof of Prop. 4.2.10]. ■

By Lemma 7.4.2 we have the equality

$$\chi_{\mathbb{Z}_p[G], B_{\text{dR}}[G]}(M^\bullet(\mathcal{L}), 0) = \chi_{\mathbb{Z}_p[G], B_{\text{dR}}[G]}(F^\bullet, 0).$$

Since all cohomology groups are torsion, we have a short exact sequence of $\mathbb{Q}_p[G]$ -spaces

$$0 \rightarrow (\ker(f_4))_{\mathbb{Q}_p[G]} \xrightarrow{\iota} (W' \oplus W)_{\mathbb{Q}_p[G]} \xrightarrow{\delta_2} \mathbb{Q}_p[G]z_0 \rightarrow 0,$$

where ι denotes the inclusion. By the definition of the (old) refined Euler characteristic we have

$$\chi_{\mathbb{Z}_p[G], B_{\text{dR}}[G]}^{\text{old}}(F^\bullet[1], 0) = [\ker(f_4) \oplus \mathbb{Z}_p[G]z_0, \iota \oplus \sigma, W' \oplus W],$$

where σ is a $\mathbb{Q}_p[G]$ -equivariant splitting of δ_2 . By (4) we obtain

$$\begin{aligned} &\chi_{\mathbb{Z}_p[G], B_{\text{dR}}[G]}(F^\bullet[1], 0) \\ &= -\chi_{\mathbb{Z}_p[G], B_{\text{dR}}[G]}^{\text{old}}(F^\bullet[1], 0) - \partial_{\mathbb{Z}_p[G], B_{\text{dR}}[G]}^1(B^{\text{od}}(F^\bullet[1]_{B_{\text{dR}}[G]}), -\text{id}). \end{aligned}$$

Note that $F^\bullet[1]$ has non-trivial cohomology only in degrees 0 and 1 and both cohomology groups are torsion. By [BB05, Lemma 6.3] we therefore obtain

$$\begin{aligned} &\partial_{\mathbb{Z}_p[G], B_{\text{dR}}[G]}^1(B^{\text{od}}(F^\bullet[1]_{B_{\text{dR}}[G]}), -\text{id}) \\ &= \partial_{\mathbb{Z}_p[G], B_{\text{dR}}[G]}^1(H^1(F^\bullet[1])_{B_{\text{dR}}[G]}, -\text{id}) = 0. \end{aligned}$$

Furthermore, we recall that $\chi_{\mathbb{Z}_p[G], B_{\text{dR}}[G]}(F^\bullet, 0) = -\chi_{\mathbb{Z}_p[G], B_{\text{dR}}[G]}(F^\bullet[1], 0)$. In conclusion, we have derived the equality

$$\chi_{\mathbb{Z}_p[G], B_{\text{dR}}[G]}(F^\bullet, 0) = [\ker(f_4) \oplus \mathbb{Z}_p[G]z_0, \iota \oplus \sigma, W' \oplus W].$$

Taking the $\mathbb{Z}_p[G]$ -basis of $\ker(f_4)$ determined in Lemma 7.4.9 and the obvious $\mathbb{Z}_p[G]$ -basis of $W' \oplus W$ we can represent the inclusion ι by the matrix

$$\mathfrak{M} = \begin{pmatrix} a-1 & 0 & 0 & 0 & 0 & \cdots & 0 & 0 \\ 1-u^mb & \mathcal{T}_a & 0 & 0 & 0 & \cdots & 0 & 0 \\ v & 0 & \mathcal{T}_a \tilde{I} & (a-1)I & 0 & \cdots & 0 & 0 \\ * & 0 & 0 & -I & (a-1)I & \cdots & 0 & 0 \\ * & 0 & 0 & * & -I & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ * & 0 & 0 & * & * & \cdots & -I & (a-1)I \\ * & \mathcal{M}_1 & \tilde{\mathcal{M}} & * & * & \cdots & * & -I \end{pmatrix},$$

where $\tilde{\mathcal{M}}$ is the matrix \mathcal{M} without the first column \mathcal{M}_1 (which in the case $m > 1$ is just $-e_2$), \tilde{I} is defined analogously from the identity matrix I , and v is a vector whose first component is $(u^mb)^{\tilde{m}}$. Note that this looks very similar to the corresponding matrix in [BC16], up to some twists by powers of u . Next we need to choose a σ . We set

$$\sigma(z_0) = \frac{1}{u^{dm} - 1} \left(\sum_{i=0}^{d-1} (u^mb)^i \right) z_1,$$

and easily check that $\delta_2 \circ \sigma = \text{id}$.

By the above discussion we know that the refined Euler characteristic $\chi_{\mathbb{Z}_p[G], B_{\text{dR}}[G]}(M^\bullet(\mathcal{L}), 0)$ is represented by the determinant of the matrix

$(w, \mathfrak{M}) \in \text{Gl}_{mp+2}(\mathbb{Q}_p[G])$ which represents the $\mathbb{Q}_p[G]$ -equivariant map $\iota + \sigma$ with respect to the chosen $\mathbb{Z}_p[G]$ -basis. Here w is the column vector

$$w = \left(\frac{\sum_{i=0}^{d-1} (u^m b)^i}{u^{dm} - 1}, 0, \dots, 0 \right)^t.$$

We deduce that the matrix (w, \mathfrak{M}) is given by

$$\begin{pmatrix} \frac{\sum_{i=0}^{d-1} (u^m b)^i}{u^{dm} - 1} & a - 1 & 0 & 0 & 0 & 0 & \dots & 0 & 0 \\ 0 & 1 - u^m b & \mathcal{T}_a & 0 & 0 & 0 & \dots & 0 & 0 \\ 0 & v & 0 & \mathcal{T}_a \tilde{I} & (a - 1)I & 0 & \dots & 0 & 0 \\ 0 & * & 0 & 0 & -I & (a - 1)I & \dots & 0 & 0 \\ 0 & * & 0 & 0 & * & -I & \dots & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & * & 0 & 0 & * & * & \dots & -I & (a - 1)I \\ 0 & * & \mathcal{M}_1 & \tilde{\mathcal{M}} & * & * & \dots & * & -I \end{pmatrix}.$$

We recall that every irreducible character ψ of $G = \text{Gal}(N/K)$ decomposes as $\psi = \phi\chi$ where χ is an irreducible character of $\langle a \rangle$ and ϕ is an irreducible character of $\langle b \rangle$. We first compute

$$\chi\phi \left(\frac{\sum_{i=0}^{d-1} (u^m b)^i}{u^{dm} - 1} \right) = \frac{u^{dm} - 1}{(u^m \chi\phi(b) - 1)(u^{dm} - 1)} = \frac{1}{u^m \chi\phi(b) - 1},$$

and note that $u^m \chi\phi(b) \neq 1$ because otherwise we would have $\chi^{\text{nr}}(F_N) = u^{dm} = 1$. For the computation of the determinant we distinguish two cases:

CASE 1: $\chi = 1$. Here $(\chi\phi)((w, \mathfrak{M})) = \phi((w, \mathfrak{M}))$ is of the form

$$\begin{pmatrix} \frac{1}{u^m \phi(b) - 1} & 0 & 0 & 0 & 0 & 0 & \dots & 0 & 0 \\ 0 & 1 - u^m \phi(b) & p & 0 & 0 & 0 & \dots & 0 & 0 \\ 0 & \phi(v) & 0 & p\tilde{I} & 0 & 0 & \dots & 0 & 0 \\ 0 & * & 0 & 0 & -I & 0 & \dots & 0 & 0 \\ 0 & * & 0 & 0 & * & -I & \dots & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & * & 0 & 0 & * & * & \dots & -I & 0 \\ 0 & * & \phi(\mathcal{M}_1) & \phi(\tilde{\mathcal{M}}) & * & * & \dots & * & -I \end{pmatrix},$$

where we recall that the first component of the vector v is $(u^m b)^{\tilde{m}}$. The determinant is

$$\frac{-u^{m\tilde{m}} \phi(b)^{\tilde{m}} p^m (-1)^{m(p-1)}}{u^m \phi(b) - 1} = \frac{-u^{m\tilde{m}} \phi(b)^{\tilde{m}} p^m}{u^m \phi(b) - 1}.$$

CASE 2: $\chi \neq 1$. The matrix $(\chi\phi)((w, \mathfrak{M}))$ is here given by

$$\begin{pmatrix} \frac{1}{u^m\phi(b)-1} & \chi(a) - 1 & 0 & 0 & 0 & \cdots & 0 & 0 \\ 0 & 1 - u^m\phi(b) & 0 & 0 & 0 & \cdots & 0 & 0 \\ 0 & \chi\phi(v) & 0 & (\chi(a)-1)I & 0 & \cdots & 0 & 0 \\ 0 & * & 0 & -I & (\chi(a)-1)I & \cdots & 0 & 0 \\ 0 & * & 0 & * & -I & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & * & 0 & * & * & \cdots & -I & (\chi(a)-1)I \\ 0 & * & \chi\phi(\mathcal{M}) & * & * & \cdots & * & -I \end{pmatrix}.$$

Using Lemma 7.4.8 we compute for the determinant

$$\begin{aligned} \frac{1 - u^m\phi(b)}{u^m\phi(b) - 1} (-1)^{(p-1)m^2} \det(\chi\phi(\mathcal{M})) (\chi(a) - 1)^{m(p-1)} \\ = (-1)^{m-1} (u^m\phi(b))^{-\tilde{m}(m-1)} (\chi(a) - 1)^{m(p-1)}. \end{aligned}$$

We summarize the above discussion in

PROPOSITION 7.4.10. Assume that $\omega > 0$. For $\mathcal{L} = \mathfrak{p}_N^{p+1}$ the element

$$\chi_{\mathbb{Z}_p[G], B_{\text{dR}}[G]}(M^\bullet(\mathcal{L}), \lambda_2^{-1}) \in K_0(\mathbb{Z}_p[G], B_{\text{dR}}[G])$$

is contained in $K_0(\mathbb{Z}_p[G], \mathbb{Q}_p[G])$ and represented by $\varepsilon \in \mathbb{Q}_p[G]^\times$ where

$$\varepsilon_{\chi\phi} = \begin{cases} \frac{-u^{m\tilde{m}}\phi(b)^{\tilde{m}}p^m}{u^m\phi(b) - 1} & \text{if } \chi = \chi_0, \\ (-1)^{m-1} (u^m\phi(b))^{-\tilde{m}(m-1)} (\chi(a) - 1)^{m(p-1)} & \text{if } \chi \neq \chi_0. \end{cases}$$

8. Proof of Theorem 1. We have to show that the element

$$\tilde{R}_{N/K} = C_{N/K} + U_{\text{cris}} + \partial_{\mathbb{Z}_p[G], B_{\text{dR}}[G]}^1(t) + \partial_{\mathbb{Z}_p[G], B_{\text{dR}}[G]}^1(\varepsilon_D(N/K, V))$$

of $K_0(\mathbb{Z}_p[G], \tilde{\Omega})$ becomes trivial in $K_0(\tilde{\Lambda}, \tilde{\Omega})$. By (55) we get

$$\begin{aligned} \tilde{R}_{N/K} &= U_{\text{cris}} + \partial_{\mathbb{Z}_p[G], B_{\text{dR}}[G]}^1(\rho) \\ &\quad + \partial_{\mathbb{Z}_p[G], B_{\text{dR}}[G]}^1(\varepsilon_D(N/K, V)) - \chi_{\mathbb{Z}_p[G], B_{\text{dR}}[G]}(M^\bullet(\mathcal{L}), \lambda_2^{-1}). \end{aligned}$$

By Proposition 6.2.1 together with Lemma 6.2.2 the element

$$(u^{-mm\chi\phi} \cdot \tau_{\mathbb{Q}_p}(\text{Ind}_{K/\mathbb{Q}_p} \chi\phi)^{-1})_{\chi\phi} \quad \text{with} \quad m_{\chi\phi} = \begin{cases} 0 & \text{if } \chi = \chi_0, \\ 2 & \text{if } \chi \neq \chi_0 \end{cases}$$

is a representative of $\partial_{\mathbb{Z}_p[G], B_{\text{dR}}[G]}^1(\varepsilon_D(N/K, V))$. We note that by [BC16, proof of Prop. 5.2.1] we obtain $\tau_{\mathbb{Q}_p}(\text{Ind}_{K/\mathbb{Q}_p} \chi\phi) = \tau_K(\chi\phi)$. Since we intend to apply [BC16, Prop. 5.1.5], we will use the integral normal basis generator $p^2\alpha_M\theta_2$ of $\mathcal{L} = \mathfrak{p}_N^{p+1}$ which we constructed in [BC16, Sec. 5.1]. Then [BC16,

Proposition 5.2.1] implies that $\partial_{\mathbb{Z}_p[G], B_{\text{dR}}[G]}(\rho) + \partial_{\mathbb{Z}_p[G], B_{\text{dR}}[G]}(\varepsilon_D(N/K, V))$ is represented by

$$\frac{\rho_\chi u^{-mm_\chi \phi}}{\tau_K(\chi\phi)} = \begin{cases} \mathfrak{d}_K^{1/2} \mathcal{N}_{K/\mathbb{Q}_p}(\theta_2|\phi) p^{2m} & \text{if } \chi = \chi_0, \\ \mathfrak{d}_K^{1/2} \mathcal{N}_{K/\mathbb{Q}_p}(\theta_2|\phi) p^m \chi(4)\phi(b^{-2})u^{-2m} & \text{if } \chi \neq \chi_0. \end{cases}$$

We want to point out that this is a crucial step in the proof and relies on one of the main results of [PV13] which was basic for the proof of [BC16, Prop. 5.1.5]. Note that this is the inverse of the result in [BC16, Prop. 5.2.1] after replacing $\phi(b)$ with $u\phi(b)$ everywhere.

If $\chi^{\text{nr}}|_{G_N} = 1$ then the calculation of $\chi_{\mathbb{Z}_p[G], B_{\text{dR}}[G]}(M^\bullet(\mathcal{L}), \lambda_2^{-1})$ corresponds to the calculation of $E(\exp(\mathcal{L}))_p$ in [BC16]: the representative of the local fundamental class has to be twisted by χ^{nr} , then starting from [BC16, Lemma 4.1.2] one has to replace b with $u^m b$ everywhere. Since also U_{cris} corresponds to $M_{N/K}$, just replacing b with $u^m b$, the same calculations of [BC16, Sec. 6] can be used to prove Theorem 1 in the case $\chi^{\text{nr}}|_{G_N} = 1$. So the interesting case is when $\chi^{\text{nr}}|_{G_N} \neq 1$, which we will assume from now on.

By Proposition 5.3.4 the element U_{cris} is represented by $u_{\text{cris}} \in \mathbb{Q}_p[G]^\times$ where

$$u_{\text{cris}, \chi\phi} = \begin{cases} \frac{1 - p^{-m}u^{-m}\phi(b)^{-1}}{1 - u^m\phi(b)} & \text{if } \chi = \chi_0, \\ 1 & \text{if } \chi \neq \chi_0. \end{cases}$$

We first deal with the case $\omega = 0$. Inserting the result of Proposition 7.3.2 we deduce that $\tilde{R}_{N/K}$ is represented by $\tilde{r} \in \tilde{\Omega}$ where

$$\begin{aligned} \tilde{r}_{\chi\phi} &= \begin{cases} \frac{\mathfrak{d}_K^{1/2} \mathcal{N}_{K/\mathbb{Q}_p}(\theta_2|\phi) p^{2m} (1 - p^{-m}u^{-m}\phi(b^{-1}))}{p^m (1 - u^m\phi(b))} & \text{if } \chi = \chi_0, \\ \frac{\mathfrak{d}_K^{1/2} \mathcal{N}_{K/\mathbb{Q}_p}(\theta_2|\phi) p^m \chi(4)\phi(b)^{-2} u^{-2m}}{(-1)^{m-1} (u^{-m}\phi(b)^{-1} - 1)(\chi(a) - 1)^{m(p-1)}} & \text{if } \chi \neq \chi_0 \end{cases} \\ &= \begin{cases} \mathfrak{d}_K^{1/2} \mathcal{N}_{K/\mathbb{Q}_p}(\theta_2|\phi) u^{-m}\phi(b^{-1}) \frac{(p^m u^m \phi(b) - 1)}{(1 - u^m\phi(b))} & \text{if } \chi = \chi_0, \\ -\mathfrak{d}_K^{1/2} \mathcal{N}_{K/\mathbb{Q}_p}(\theta_2|\phi) \left(\frac{-p}{(\chi(a) - 1)^{p-1}}\right)^m \frac{\chi(4)\phi(b)^{-2} u^{-2m}}{1 - u^m\phi(b)} u^m\phi(b) & \text{if } \chi \neq \chi_0 \end{cases} \\ &= \begin{cases} -\frac{\mathfrak{d}_K^{1/2} \mathcal{N}_{K/\mathbb{Q}_p}(\theta_2|\phi)}{1 - u^m\phi(b)} u^{-m}\phi(b)^{-1} (1 - p^m u^m \phi(b)) & \text{if } \chi = \chi_0, \\ -\frac{\mathfrak{d}_K^{1/2} \mathcal{N}_{K/\mathbb{Q}_p}(\theta_2|\phi)}{1 - u^m\phi(b)} u^{-m}\phi(b)^{-1} \left(\frac{-p}{(\chi(a) - 1)^{p-1}}\right)^m \chi(4) & \text{if } \chi \neq \chi_0. \end{cases} \end{aligned}$$

Similarly to [BC16, proof of Theorem 1], define $W_{\theta_2} \in \mathcal{O}_p^t[G] \in \overline{\mathbb{Z}_p^{\text{nr}}}[G]^\times$ by

$$\chi\phi(W_{\theta_2}) = \mathfrak{d}_K^{1/2} \mathcal{N}_{K/\mathbb{Q}_p}(\theta_2|\phi).$$

By a straightforward computation (see [BC16, Lemma 3.2.3]) there exists a unit $\tilde{u} \in \mathbb{Z}_p[a]$ such that $p(1 - e_a) = (a - 1)^{p-1} \tilde{u}(1 - e_a)$ and the augmentation of \tilde{u} is $(p - 1)! \equiv -1 \pmod{p}$. Evaluating at $\chi \neq \chi_0$ we get $p = (\chi(a) - 1)^{p-1} \chi(\tilde{u})$.

Then by the above calculations we have

$$\tilde{r} = -\frac{W_{\theta_2} u^{-m} b^{-1}}{1 - u^m b} \left((1 - p^m u^m b) e_a + (-\tilde{u})^m \sigma_4 (1 - e_a) \right).$$

By [Frös3, Chap. I, Prop. 4.3] the element W_{θ_2} is a unit. Furthermore, recalling that $v_p(1 - u^{m d_{N/K}}) = \omega = 0$, we get

$$\frac{1}{1 - u^m b} = \frac{1 + u^m b + \dots + u^{m(d_{N/K}-1)} b^{d_{N/K}-1}}{1 - u^{m d_{N/K}}} \in \overline{\mathbb{Z}}_p^{\text{nr}}[G].$$

Since its inverse $1 - u^m b$ is also in $\overline{\mathbb{Z}}_p^{\text{nr}}[G]$, it is a unit. So we just need to show that the element

$$\tilde{r} = (1 - p^m u^m b) e_a + (-\tilde{u})^m \sigma_4 (1 - e_a)$$

is a unit in $\overline{\mathbb{Z}}_p^{\text{nr}}[G]$. We first show that \tilde{r} is contained in $\overline{\mathbb{Z}}_p^{\text{nr}}[G]$, which is true if and only if

$$e_a + (-\tilde{u})^m \sigma_4 (1 - e_a)$$

is in $\overline{\mathbb{Z}}_p^{\text{nr}}[G]$. This is equivalent to

$$1 \equiv (-\chi(\tilde{u}))^m \chi(\sigma_4) \pmod{1 - \zeta_p}$$

for any non-trivial character χ , which is straightforward since $\chi(\tilde{u}) \equiv -1 \pmod{1 - \zeta_p}$ and $\chi(\sigma_4) \equiv 1 \pmod{1 - \zeta_p}$. Note that in the present situation this is not enough to conclude the proof of our theorem as it was in [BC16], since we could not find a rigorous proof in the literature that $C_{\text{EP}}^{\text{na}}(N/K, V)$ is true over maximal orders. Actually, if we knew that the conjecture were functorial with respect to restriction and quotient maps in relative K -theory (as commonly expected), a standard K -theoretic argument would show the validity of $C_{\text{EP}}^{\text{na}}(N/K, V)$ over maximal orders. However, we prefer to give a short argument to prove directly that \tilde{r}^{-1} has integral coefficients. Note that

$$\frac{1}{1 - p^m u^m \phi(b)} = \frac{1 + p^m u^m \phi(b) + \dots + (p^m u^m \phi(b))^{d_{N/K}-1}}{1 - (p^m u^m)^{d_{N/K}}}.$$

Therefore

$$\tilde{r}^{-1} = \frac{1 + p^m u^m b + \dots + (p^m u^m b)^{d_{N/K}-1}}{1 - (p^m u^m)^{d_{N/K}}} e_a + (-\tilde{u})^{-m} \sigma_4^{-1} (1 - e_a).$$

The integrality of this term is obviously equivalent to the integrality of

$$\frac{1}{1 - (p^m u^m)^{d_{N/K}}} e_a + (-\tilde{u})^{-m} \sigma_4^{-1} (1 - e_a),$$

which, in turn, is equivalent to the congruence

$$\frac{1}{1 - (p^m u^m)^{d_{N/K}}} \equiv (-\chi(\tilde{u}))^{-m} \chi(\sigma_4)^{-1} \pmod{1 - \zeta_p}.$$

Since both sides of this congruence are congruent to 1, this concludes the proof in the case $\omega = 0$.

In the case $\omega > 0$ we insert the result of Proposition 7.4.10 into the formula for \tilde{r} and, with the same notation as in the case $\omega = 0$, we obtain

$$\begin{aligned} \tilde{r}_{\chi\phi} &= \begin{cases} \frac{\mathfrak{d}_K^{1/2} \mathcal{N}_{K/\mathbb{Q}_p}(\theta_2|\phi) p^{2m} (u^m \phi(b)-1)(1-p^{-m} u^{-m} \phi(b^{-1}))}{-u^{m\tilde{m}} \phi(b)^{\tilde{m}} p^m (1-u^m \phi(b))} & \text{if } \chi = \chi_0, \\ \frac{\mathfrak{d}_K^{1/2} \mathcal{N}_{K/\mathbb{Q}_p}(\theta_2|\phi) p^m \chi(4) \phi(b)^{-2} u^{-2m}}{(-1)^{m-1} (u^m \phi(b))^{-\tilde{m}(m-1)} (\chi(a)-1)^{m(p-1)}} & \text{if } \chi \neq \chi_0 \end{cases} \\ &= \begin{cases} \mathfrak{d}_K^{1/2} \mathcal{N}_{K/\mathbb{Q}_p}(\theta_2|\phi) u^{-m} \phi(b^{-1}) \frac{(p^m u^m \phi(b)-1)}{u^{m\tilde{m}} \phi(b)^{\tilde{m}}} & \text{if } \chi = \chi_0, \\ -\mathfrak{d}_K^{1/2} \mathcal{N}_{K/\mathbb{Q}_p}(\theta_2|\phi) \left(\frac{-p}{(\chi(a)-1)^{p-1}}\right)^m \frac{\chi(4) \phi(b)^{-2} u^{-2m}}{(u^m \phi(b))^{-\tilde{m}(m-1)}} & \text{if } \chi \neq \chi_0 \end{cases} \\ &= \begin{cases} -\chi\phi(W_{\theta_2})\phi(b)^{-1-\tilde{m}} u^{-m-m\tilde{m}} (1-p^m u^m \phi(b)) & \text{if } \chi = \chi_0, \\ -\chi\phi(W_{\theta_2})\phi(b)^{-1-\tilde{m}} u^{-m-m\tilde{m}} \left(\frac{-p}{(\chi(a)-1)^{p-1}}\right)^m \chi(4) u^{m^2\tilde{m}-m} & \text{if } \chi \neq \chi_0. \end{cases} \end{aligned}$$

Hence

$$\tilde{r} = W_{\theta_2} b^{1-\tilde{m}} u^{-m-m\tilde{m}} ((1-p^m u^m b)e_a + (-\tilde{u})^m \sigma_4 u^{m^2\tilde{m}-m} (1-e_a)).$$

As above, we have to prove that

$$\tilde{r} = (1-p^m u^m b)e_a + (-\tilde{u})^m \sigma_4 u^{m^2\tilde{m}-m} (1-e_a)$$

is a unit. Note that this is the same as the element we got in the case $\omega=0$, up to the factor $u^{m^2\tilde{m}-m}$ in the $(1-e_a)$ -component. Since we are in the case $\omega > 0$, we have $u^{dm} \equiv 1 \pmod{1-\zeta_p}$. Furthermore, since $m\tilde{m} \equiv 1 \pmod{d}$, we derive $u^{m^2\tilde{m}} \equiv u^m \pmod{1-\zeta_p}$, i.e. $u^{m^2\tilde{m}-m} \equiv 1 \pmod{1-\zeta_p}$. By this fact, the proof for $\omega = 0$ is given by the same calculations as for $\omega > 0$. ■

Appendix: A more detailed computation of U_{cris} . By (31) we need to compare β' and κ . In the definition of both trivializations one starts with the same isomorphism

$$\begin{aligned} \Delta_{\text{EP}}^{\text{na}}(N/K, V) &= [R\Gamma(N, V)] \otimes [\text{Ind}_{N/\mathbb{Q}_p} V] \\ &\rightarrow [H^1(N, V)]^{-1} \otimes [H^2(N, V)] \otimes [\text{Ind}_{N/\mathbb{Q}_p} V]. \end{aligned}$$

We set

$$\begin{aligned} A &= [D_{\text{cris}}^N(V)] \otimes [D_{\text{cris}}^N(V)]^{-1}, \\ B &= [H^1(N, V)]^{-1} \otimes [t_V(N)] \otimes [H^1(N/V)/H_f^1(N, V)], \\ C &= [H^1(N/V)/H_f^1(N, V)]^{-1} \otimes [D_{\text{cris}}^N(V^*(1))] \\ &\quad \otimes [D_{\text{cris}}^N(V^*(1))]^{-1} \otimes [H^2(N, V)], \\ D &= [t_V(N)^{-1}] \otimes [D_{\text{dr}}^N(V)], \\ E &= [D_{\text{dr}}^N(V)]^{-1} \otimes [\text{Ind}_{N/\mathbb{Q}_p} V]. \end{aligned}$$

Then there is a natural isomorphism

$$[H^1(N, V)]^{-1} \otimes [H^2(N, V)] \otimes [\mathrm{Ind}_{N/\mathbb{Q}_p} V] \rightarrow A \otimes B \otimes C \otimes D \otimes E,$$

where we used the defining property of right inverses and the commutativity constraint to identify $[t_V(N)]^{-1} \otimes [H^2(N, V)]$ with $[H^2(N, V)] \otimes [t_V(N)]^{-1}$.

Up to now all the calculations are common for β' and κ . Now we can compare the two trivializations term by term:

- $A \rightarrow 1$: This is determined by $1 - \phi$ in β' and by 1 in κ .
- $B \rightarrow 1$: In both cases this isomorphism comes from the exponential map \exp_V : see for example the exact sequence (34).
- $C \rightarrow 1$: For β' we use (35), while for κ we first have to identify C with $[H^1(N/V)/H_f^1(N, V)]^{-1} \otimes [H^2(N, V)]$ and then consider the valuation map ν_N .
- $D \rightarrow 1$: This comes from (6), recalling Lemma 5.1.2.
- $E \rightarrow 1$: This comes from the map comp_V , for both β' and κ , but in the definition of β' there also appears a multiplication by $t^{t_H(V)} = t$.

So the difference between β' and κ lies in $A \rightarrow 1$ and $C \rightarrow 1$ and in the multiplication by t in $E \rightarrow 1$, and leads to (32) when $\chi^{\mathrm{nr}}|_{G_N} \neq 1$ and to (37) when $\chi^{\mathrm{nr}}|_{G_N} = 1$.

Acknowledgements. We would like to thank the referee for a very careful reading of the article and valuable suggestions to simplify some of our arguments.

References

- [BB08] D. Benois and L. Berger, *Théorie d'Iwasawa des représentations cristallines II*, Comment. Math. Helv. 83 (2008), 603–677.
- [BB03] W. Bley and D. Burns, *Equivariant epsilon constants, discriminants and étale cohomology*, Proc. London Math. Soc. (3) 87 (2003), 545–590.
- [BC16] W. Bley and A. Cobbe, *Equivariant epsilon constant conjectures for weakly ramified extensions*, Math. Z. 283 (2016), 1217–1244.
- [BD13] W. Bley and R. Debeerst, *Algorithmic proof of the epsilon constant conjecture*, Math. Comp. 82 (2013), 2363–2387.
- [BK90] S. Bloch and K. Kato, *L-functions and Tamagawa numbers of motives*, in: The Grothendieck Festschrift, Vol. I, Progr. Math. 86, Birkhäuser Boston, Boston, MA, 1990, 333–400.
- [Bre04a] M. Breuning, *Equivariant epsilon constants for Galois extensions of number fields and p-adic fields*, PhD thesis, King's College London, 2004.
- [Bre04b] M. Breuning, *Equivariant local epsilon constants and étale cohomology*, J. London Math. Soc. (2) 70 (2004), 289–306.
- [BB05] M. Breuning and D. Burns, *Additivity of Euler characteristics in relative algebraic K-groups*, Homology Homotopy Appl. 7 (2005), no. 3, 11–36.

- [Bur04] D. Burns, *Equivariant Whitehead torsion and refined Euler characteristics*, in: Number Theory, CRM Proc. Lecture Notes 36, Amer. Math. Soc., Providence, RI, 2004, 35–59.
- [BF98] D. Burns and M. Flach, *On Galois structure invariants associated to Tate motives*, Amer. J. Math. 120 (1998), 1343–1397.
- [BF01] D. Burns and M. Flach, *Tamagawa numbers for motives with (non-commutative) coefficients*, Doc. Math. 6 (2001), 501–570.
- [Chi85] T. Chinburg, *Exact sequences and Galois module structure*, Ann. of Math. (2) 121 (1985), 351–376.
- [CR87] C. W. Curtis and I. Reiner, *Methods of Representation Theory. With Applications to Finite Groups and Orders. Vol. II*, Pure Appl. Math. (New York), Wiley, New York, 1987.
- [Del73] P. Deligne, *Les constantes des équations fonctionnelles des fonctions L*, in: Modular Functions of One Variable, II (Antwerp, 1972), Lecture Notes in Math. 349, Springer, Berlin, 1973, 501–597.
- [FO10] J.-M. Fontaine and Y. Ouyang, *Theory of p -Adic Galois Representations*, Springer, 2010.
- [FPR94] J.-M. Fontaine et B. Perrin-Riou, *Autour des conjectures de Bloch et Kato: cohomologie galoisienne et valeurs de fonctions L*, in: Motives (Seattle, WA, 1991), Proc. Sympos. Pure Math. 55, Amer. Math. Soc., Providence, RI, 1994, 599–706.
- [Frö83] A. Fröhlich, *Galois Module Structure of Algebraic Integers*, Ergeb. Math. Grenzgeb. (3) 1, Springer, Berlin, 1983.
- [FK06] T. Fukaya and K. Kato, *A formulation of conjectures on p -adic zeta functions in noncommutative Iwasawa theory*, in: Proc. St. Petersburg Math. Soc. 12, Amer. Math. Soc. Transl. (2) 219, Amer. Math. Soc., Providence, RI, 2006, 1–85.
- [Izy12] D. Izychev, *Equivariant epsilon-conjecture for unramified twists of $\mathbb{Z}_p(1)$* , PhD thesis, Ruprecht-Karls-Universität Heidelberg, 2012.
- [IV16] D. Izychev and O. Venjakob, *Equivariant epsilon conjecture for 1-dimensional Lubin–Tate groups*, J. Théor. Nombres Bordeaux 28 (2016), 485–521.
- [Köc04] B. Köck, *Galois structure of Zariski cohomology for weakly ramified covers of curves*, Amer. J. Math. 126 (2004), 1085–1107.
- [Lan02] S. Lang, *Algebra*, 3rd ed., Grad. Texts in Math. 211, Springer, New York, 2002.
- [LR78] J. Lubin and M. I. Rosen, *The norm map for ordinary Abelian varieties*, J. Algebra 52 (1978), 236–240.
- [Mil80] J. S. Milne, *Étale Cohomology*, Princeton Math. Ser. 33, Princeton Univ. Press, Princeton, NJ, 1980.
- [Neu92] J. Neukirch, *Algebraische Zahlentheorie*, Springer, Berlin, 1992.
- [NSW08] J. Neukirch, A. Schmidt, and K. Wingberg, *Cohomology of Number Fields*, 2nd ed., Grundlehren Math. Wiss. 323, Springer, Berlin, 2008.
- [PV13] E. J. Pickett and S. Viatier, *Self-dual integral normal bases and Galois module structure*, Compos. Math. 149 (2013), 1175–1202.
- [Ser77] J.-P. Serre, *Linear Representations of Finite Groups*, Grad. Texts in Math. 42, Springer, New York, 1977.
- [Ser79] J.-P. Serre, *Local Fields*, Grad. Texts in Math. 67, Springer, New York, 1979.
- [Sil09] J. H. Silverman, *The Arithmetic of Elliptic Curves*, 2nd ed., Grad. Texts in Math. 106, Springer, Dordrecht, 2009.
- [Swa70] R. G. Swan, *K -Theory of Finite Groups and Orders*, Lecture Notes in Math. 149, Springer, Berlin, 1970.

- [Wei94] C. A. Weibel, *An Introduction to Homological Algebra*, Cambridge Stud. Adv. Math. 38, Cambridge Univ. Press, Cambridge, 1994.

Werner Bley
Mathematisches Institut
der Universität München
Theresienstr. 39
D-80333 München, Germany
E-mail: bley@math.lmu.de

Alessandro Cobbe
Institut für Theoretische Informatik,
Mathematik und Operations Research
Fakultät für Informatik
Universität der Bundeswehr München
Werner-Heisenberg-Weg 39
D-85577 Neubiberg, Germany
E-mail: alessandro.cobbe@unibw.de

