

Graph C^* -algebras

Rainer Matthes,
Wojciech Szymański
notes taken by:
Paweł Witkowski

August 17, 2005

Contents

1	Universal graph C^*-algebras	2
2	Computation of K-theory	10
3	Proof of the theorem (2.1)	15

1 Universal graph C*-algebras

Let G be a directed graph with

$$\begin{aligned} G^0 & - \text{ vertices,} \\ G^1 & - \text{ edges,} \\ r, s: G^1 & \rightarrow G^0 - \text{ range and source of an edge.} \end{aligned}$$

Definition 1.1. *The universal C*-algebra $C^*(G)$ is given by generators*

$$\{p_v \mid v \in G^0\}, \quad \{s_e \mid e \in G^1\},$$

with the following relations:

- p_v are mutually orthogonal projections i.e. $p_v^2 = p_v^* = p_v$ and $p_v p_w = 0$ for $v \neq w$,
- $s_e^* s_e = p_{r(e)}$ and $s_e^* s_f = 0$ for $e \neq f$,
- if the set $\{e \mid s(e) = v\}$ is nonempty (v is not a sink) and finite then

$$p_v = \sum_{\{e \mid s(e)=v\}} s_e s_e^*,$$

- $s_e s_e^* \leq p_{s(e)}$.

Example 1.2. Some known C*-algebras arise in this way.

1. If G is only one vertex, then there is one generator $p = p^2 = p^*$. In this case $C^*(G) = \mathbb{C}$.



Figure 1: \mathbb{C}

2. G with one vertex and one edge (loop). Generators:




Figure 2: $C(S^1)$

$$p = p^2 = p^*, \quad s$$

relations:

$$s^* s = p = s s^*, \quad s p = p s, \quad s = s s^* s.$$

Then

$$C^*(G) = C^*(1, u) = C(S^1), \quad u - \text{unitary.}$$




Figure 3: Toeplitz algebra \mathcal{T}

3. G with two vertices and two edges like on the picture (3).

$$\begin{aligned}
 p_v &= p_v^2 = p_v^*, & p_w &= p_w^2 = p_w^* \\
 s_e^* s_e &= p_v, & s_f^* s_f &= p_w \\
 p_v &= s_e s_e^* + s_f s_f^*.
 \end{aligned}$$

$C^*(G)$ is isomorphic to the Toeplitz algebra - the universal C^* -algebra for the relation $s^* s = 1$. The isomorphism is given by $s \mapsto s_e + s_f$.

4. G with three vertices and three edges like on a picture (4).




Figure 4: $C(S_{0\infty}^2)$

$$\begin{aligned}
 p_v &= p_v^2 = p_v^*, & p_{w_i} &= p_{w_i}^2 = p_{w_i}^*, & i &= 1, 2, \\
 p_v p_{w_i} &= 0, & p_{w_1} p_{w_2} &= 0, \\
 s_e^* s_e &= p_v = s_e s_e^* + s_{f_1} s_{f_1}^* + s_{f_2} s_{f_2}^*, \\
 s_{f_1}^* s_{f_1} &= p_{w_1}, & s_{f_2}^* s_{f_2} &= p_{w_2}.
 \end{aligned}$$

$C^*(G)$ is isomorphic to the quantum sphere

$$S_{0\infty}^2: \quad B^* B = 1 - A^2, \quad A = A^*, \quad B B^* = 1, \quad B A = 0$$

and the isomorphism is given by

$$\begin{aligned}
 A &\mapsto p_{w_1} - p_{w_2}, \\
 B &\mapsto s_e^* + s_{f_1}^* + s_{f_2}^*.
 \end{aligned}$$

We denote this graph by $G_{S_{0\infty}^2}$.




Figure 5: $C(\mathbb{R}P_q^2)$

5. In the example (4) we glue the vertices w_1, w_2 into one w obtaining graph G like on a picture (5).

$$\begin{aligned}
p_v &= p_v^2 = p_v^*, & p_w &= p_w^2 = p_w^*, \\
p_v p_w &= 0, \\
s_e^* s_e &= p_v = s_e^* s_e + s_{f_1}^* s_{f_1} + s_{f_2}^* s_{f_2}, \\
s_{f_1}^* s_{f_1} &= p_w = s_{f_2}^* s_{f_2}.
\end{aligned}$$

Define \mathbb{Z}_2 -action on the graph in the example (4).

$$s_e \mapsto -s_e, \quad s_{f_1} \mapsto -s_{f_2}, \quad s_{f_2} \mapsto -s_{f_1}$$

Then

$$p_v \mapsto p_v, \quad p_{w_1} \mapsto p_{w_2}, \quad p_{w_2} \mapsto p_{w_1}.$$

This action corresponds to

$$A \mapsto -A, \quad B \mapsto -B$$

under the identification

$$C^*(G_{S_{0\infty}^2}) \simeq C(S_{0\infty}^2).$$

If we take the quotient $C(S_{0\infty}^2)/\mathbb{Z}_2$ we obtain $C(\mathbb{R}P_q^2)$ - quantum projective space. On the other hand the quotient of the graph C^* -algebra $C^*(G_{S_{0\infty}^2})$ by the defined action is the graph C^* -algebra for our graph, which we now can denote $C^*(G_{\mathbb{R}P_q^2})$. The isomorphism is given by

$$\begin{aligned}
p_v &\mapsto p_v, & p_w &\mapsto p_{w_1} + p_{w_2}, \\
s_e &\mapsto s_e s_e, & s_{f_1} &\mapsto s_e (s_{f_1} + s_{f_2}), & s_{f_2} &\mapsto s_{f_1} - s_{f_2}.
\end{aligned}$$

Note that this \mathbb{Z}_2 action is not induced from a graph automorphism.

6. G with one vertex and n edges like on the picture (6).




Figure 6: Cuntz algebra O_n

$$r(e_k) = s(e_k) = v, \quad k = 1, \dots, n,$$

$$p = s_{e_k}^* s_{e_k} = \sum_{k=1}^n s_{e_k} s_{e_k}^*,$$

$$s_{e_k}^* s_{e_{k'}} = 0 \text{ for } k \neq k'.$$

When $p = 1$ then $C^*(G)$ is the Cuntz algebra O_n - the universal C*-algebra for the relations

$$s_k^* s_k = 1, \quad k = 1, \dots, n, \quad \sum_{k=1}^n s_k s_k^* = 1.$$

7. G with n vertices and $(n - 1)$ edges in the straight segment as in the picture (7).




Figure 7: $M_n(\mathbb{C})$

$$s(e_k) = k, \quad r(e_k) = k + 1 \text{ for } k = 1, \dots, n - 1,$$

$$p_k = s_{e_k} s_{e_k}^*, \quad p_{k+1} = s_{e_k}^* s_{e_k} \text{ for } k = 1, \dots, n - 1,$$

$$s_{e_k}^* s_{e_{k'}} = 0 \text{ for } k \neq k'.$$

$C^*(G)$ is the algebra of complex matrices $n \times n$, that is $M_n(\mathbb{C})$.

8. Similarly to the previous example we take straight segment, but infinite in both directions. Vertices are indexed by integers as in the picture (8).




Figure 8: Compact operators \mathcal{K}

$$s(e_k) = k, \quad r(e_k) = k + 1, \quad k \in \mathbb{Z},$$

$$p_k = s_{e_k} s_{e_k}^*, \quad p_{k+1} = s_{e_k}^* s_{e_k},$$

$$s_{e_k}^* s_{e_{k'}} = 0 \text{ for } k \neq k'.$$

We obtain algebra of compact operators \mathcal{K} , the limit of the algebras in the preceding example.

9. G with n vertices and n edges forming a cycle as in the picture (9).

$$s(e_k) = k, \quad r(e_k) = k + 1 \text{ for } k = 1, \dots, n - 1, \quad r(e_n) = 1,$$

$$p_k = s_{e_k} s_{e_k}^*, \quad p_{k+1} = s_{e_k}^* s_{e_k},$$

$$s_{e_k}^* s_{e_{k'}} = 0 \text{ for } k \neq k'.$$

We obtain algebra of matrices over the algebra of functions on the circle, $C^*(G) = M_n(C(S^1))$.




Figure 9: $M_n(C(S^1))$




Figure 10: $C(SU_q(2))$

10. G with two vertices with loops and connected by one edge.

$$p_{v_i} = p_{v_i}^* = p_{v_i}^2, \quad i = 1, 2, \quad p_{v_1}p_{v_2} = 0,$$

$$p_{v_1} = s_{e_{11}}^* s_{e_{11}} = s_{e_{11}}^* s_{e_{11}} + s_{e_{12}} s_{e_{11}}^*,$$

$$p_{v_2} = s_{e_{22}}^* s_{e_{22}} = s_{e_{12}}^* s_{e_{12}} = s_{e_{22}} s_{e_{22}}^*,$$

$$s_{e_{11}}^* s_{e_{12}} = 0, \quad s_{e_{11}}^* s_{e_{22}} = 0, \quad s_{e_{12}}^* s_{e_{22}} = 0.$$

We obtain C*-algebra for quantum $SU(2)$, that is $C(SU_q(2)) \simeq C(SU_0(2))$, which is generated by two elements a, b satisfying the relations

$$a^*a + b^*b = 1, \quad aa^* + q^2b^*b = 1,$$

$$ab = qba, \quad ab^* = qb^*a, \quad b^*b = bb^*.$$

The isomorphism is given by

$$a \mapsto s_{e_{11}}^* + s_{e_{12}}^*,$$

$$b \mapsto s_{e_{22}}.$$

11. The example (10) can be treated as the C*-algebra of the quantum sphere S_q^3 . Now we present graph C*-algebra for the quantum sphere S_q^7 , which is next generalized to arbitrary odd dimension. We take a graph G with four vertices with loops and each vertex is connected with all vertices with the greater index as in the picture (11). The C*-algebra for the quantum sphere S_q^7 is generated by the four elements z_1, z_2, z_3, z_4




Figure 11: $C(S_q^7)$

satisfying the relations

$$\begin{aligned}
z_j z_i &= q z_i z_j \text{ for } i < j, \\
z_j^* z_i &= q z_i z_j^* \text{ for } i \neq j, \\
z_1^* z_1 &= z_1 z_1^* + (1 - q^2)(z_2 z_2^* + z_3 z_3^* + z_4 z_4^*), \\
z_2^* z_2 &= z_2 z_2^* + (1 - q^2)(z_3 z_3^* + z_4 z_4^*), \\
z_3^* z_3 &= z_3 z_3^* + (1 - q^2) z_4 z_4^*, \\
z_4^* z_4 &= z_4 z_4^*, \\
z_1 z_1^* + z_2 z_2^* + z_3 z_3^* + z_4 z_4^* &= 1.
\end{aligned}$$

For $q = 0$ we have the isomorphism $C^*(G) \simeq C(S_0^7)$ given by

$$\begin{aligned}
z_1 &\mapsto s_{e_{11}} + s_{e_{12}} + s_{e_{13}} + s_{e_{14}}, \\
z_2 &\mapsto s_{e_{22}} + s_{e_{23}} + s_{e_{24}}, \\
z_3 &\mapsto s_{e_{33}} + s_{e_{34}}, \\
z_4 &\mapsto s_{e_{44}}.
\end{aligned}$$

12. As in the example (11) we take a graph with n vertices and edge between v_i and v_j if and only if $i \leq j$ as in the picture (12).




Figure 12: $C(S_q^{2n-1})$

$$v_1, \dots, v_n,$$

$$e_{ij}, \quad j = i, \dots, n, \quad s(e_{ij}) = v_i, \quad r(e_{ij}) = v_j.$$

The C^* -algebra for the quantum sphere S_q^{2n-1} is generated by the n elements z_1, \dots, z_n satisfying the relations

$$\begin{aligned} z_j z_i &= q z_i z_j \quad \text{for } i < j, \\ z_j^* z_i &= q z_i z_j^* \quad \text{for } i \neq j, \\ z_i^* z_i &= z_i z_i^* + (1 - q^2) \left(\sum_{j>i} z_j z_j^* \right) \quad \text{for } i = 1, \dots, n, \\ \sum_{i=1}^n z_i z_i^* &= 1. \end{aligned}$$

For $q = 0$ we have the isomorphism $C^*(G) \simeq C(S_0^{2n-1})$ given by

$$z_i \mapsto \sum_{j=i}^n s_{e_{ij}}, \quad i = 1, \dots, n.$$

13. We take a similar graph G to the one in the example (11), but with infinitely many parallel edges $v_i \rightarrow v_j$ for $i \leq j$.




Figure 13: $C(\mathbb{C}P_q^3)$

14. We take a similar graph G to the one in the example (12), but with infinitely many parallel edges $v_i \rightarrow v_j$ for $i \leq j$.
15. If we modify the graph for the quantum sphere S_q^5 by adding two additional vertices w_1, w_2 and edges from each vertex v_1, v_2, v_3 to both of the added ones, then we obtain graph for the sphere S_q^6 as in the picture (15).
16. The example (15) can be generalized to arbitrary even dimension just by adding two vertices w_1, w_2 to the graph of the sphere S_q^{2n-1} . We have $n + 2$ vertices v_1, \dots, v_n and w_1, w_2 . Edges e_{ij} are from v_i to v_j whenever $i \leq j$ and g_{ik} are between v_i and w_k for $k = 1, 2$. More precisely for $i = 1, \dots, n$ we have

$$s(e_{ij}) = v_i, \quad r(e_{ij}) = v_j, \quad j = i, \dots, n,$$

$$s(g_{ik}) = v_i, \quad r(g_{ik}) = w_k, \quad k = 1, 2.$$




Figure 14: $C(CP_q^{n-1})$




Figure 15: $C(S_q^6)$




Figure 16: $C(S_q^{2n})$




Figure 17: $C(\mathbb{R}P_q^6)$

17. In the example (15) we identify vertices w_1, w_2 and leave v_1, v_2, v_3 unchanged. The edges are as in the picture. General construction is described in the next example (18)
18. In the example (16) we identify vertices w_1, w_2 and leave v_1, \dots, v_n unchanged. The edges of the new graph are pairs (h_1, h_2) of edges from (16) such that $r(h_1) = s(h_2)$ and $r(h_2) \neq w_1, w_2$. Additionally we have edges f_{ik} from v_i to w for $i = 1, \dots, n$ and $k = 1, \dots, n + 2 - i$. The picture is analogous to the (17).

2 Computation of K-theory

The main tool for the computation of K-theory groups of the graph C*-algebras is the following

Theorem 2.1. *Let G be a directed graph and let $G_+^0 \subset G^0$ be the collection of vertices that emit at least one and at most finitely many edges. Let $\mathbb{Z}G_+^0$ and $\mathbb{Z}G^0$ be the free abelian groups on free generators G_+^0 and G^0 . Let $A_G: \mathbb{Z}G_+^0 \rightarrow \mathbb{Z}G^0$ be the map defined by the formula*

$$A_G(v) := \left(\sum_{e \in G^1, s(e)=v} r(e) \right) - v.$$

Then

$$\begin{aligned} K_0(C^*(G)) &\simeq \text{coker } A_G \\ K_1(C^*(G)) &\simeq \text{ker } A_G \end{aligned}$$

The proof of this theorem will be postponed to the section (3), and now we compute the K-theory groups of the graph C*-algebras for the examples from section (1).

Example 2.2. 1. $K_*(\mathbb{C})$

$$\begin{aligned} G^0 &= \{v\} \\ G_+^0 &= \emptyset \\ A_G: \emptyset &\rightarrow \mathbb{Z} \end{aligned}$$

In this case A_G is from the empty set, but still we can write

$$\begin{aligned} K_0(\mathbb{C}) &= \text{coker } A_G = \mathbb{Z} \\ K_1(\mathbb{C}) &= \text{ker } A_G = 0 \end{aligned}$$

2. $K_*(C(S^1))$

$$G^0 = \{v\}$$

$$G_+^0 = \{v\}$$

$$A_G: \mathbb{Z} \rightarrow \mathbb{Z}$$

$$v \mapsto v - v = 0$$

$$K_0(C(S^1)) = \text{coker } A_G = \mathbb{Z}$$

$$K_1(C(S^1)) = \text{ker } A_G = \mathbb{Z}$$

3. $K_*(T)$

$$G^0 = \{v, w\}$$

$$G_+^0 = \{v\}$$

$$A_G: \mathbb{Z} \rightarrow \mathbb{Z} \oplus \mathbb{Z}$$

$$v \mapsto v + w - v = w$$

$$K_0(T) = \text{coker } A_G = \mathbb{Z}$$

$$K_1(T) = \text{ker } A_G = 0$$

4. $K_*(C(S_{0\infty}^2))$

$$G^0 = \{v, w_1, w_2\}$$

$$G_+^0 = \{v\}$$

$$A_G: \mathbb{Z} \rightarrow \mathbb{Z} \oplus \mathbb{Z} \oplus \mathbb{Z}$$

$$v \mapsto v + w_1 + w_2 - v = w_1 + w_2$$

$$K_0(C(S_{0\infty}^2)) = \text{coker } A_G = \mathbb{Z} \oplus \mathbb{Z}$$

$$K_1(C(S_{0\infty}^2)) = \text{ker } A_G = 0$$

5. $K_*(C(\mathbb{R}P_q^2))$

$$G^0 = \{v, w\}$$

$$G_+^0 = \{v\}$$

$$A_G: \mathbb{Z} \rightarrow \mathbb{Z} \oplus \mathbb{Z}$$

$$v \mapsto v + 2w - v = 2w$$

$$K_0(C(\mathbb{R}P_q^2)) = \text{coker } A_G = \mathbb{Z} \oplus \mathbb{Z}_2$$

$$K_1(C(\mathbb{R}P_q^2)) = \text{ker } A_G = 0$$

6. $K_*(O_n)$

$$G^0 = \{v\}$$

$$G_+^0 = \{v\}$$

$$A_G: \mathbb{Z} \rightarrow \mathbb{Z}$$

$$v \mapsto nv - v = (n-1)v$$

$$K_0(O_n) = \text{coker } A_G = \mathbb{Z}_{n-1}$$

$$K_1(O_n) = \text{ker } A_G = 0$$

7. $K_*(M_n(\mathbb{C}))$

$$G^0 = \{v_1, v_2, \dots, v_{n-1}\}$$

$$G_+^0 = \{v_1, v_2, \dots, v_n\}$$

$$A_G: \mathbb{Z}^{n-1} \rightarrow \mathbb{Z}^n$$

$$v_i \mapsto v_{i+1} - v_i \text{ for } i = 1, \dots, n-1$$

$$K_0(M_n(\mathbb{C})) = \text{coker } A_G = \mathbb{Z}$$

$$K_1(M_n(\mathbb{C})) = \text{ker } A_G = 0$$

8. $K_*(\mathcal{K})$

$$G^0 = \{v_i \mid i \in \mathbb{Z}\}$$

$$G_+^0 = \{v_i \mid i \in \mathbb{Z}\}$$

$$A_G: \bigoplus_{i \in \mathbb{Z}} \mathbb{Z} \rightarrow \bigoplus_{i \in \mathbb{Z}} \mathbb{Z}$$

$$v_i \mapsto v_{i+1} - v_i \text{ for } i \in \mathbb{Z}$$

$$K_0(\mathcal{K}) = \text{coker } A_G = \mathbb{Z}$$

$$K_1(\mathcal{K}) = \text{ker } A_G = 0$$

Remark 2.3. If we take direct product instead of direct sum, then there will be nontrivial kernel.

9. $K_*(M_n(S^1))$

$$G^0 = \{v_1, v_2, \dots, v_n\}$$

$$G_+^0 = \{v_1, v_2, \dots, v_n\}$$

$$A_G: \mathbb{Z}^n \rightarrow \mathbb{Z}^n$$

$$v_i \mapsto v_{i+1} - v_i \text{ for } i = 1, \dots, n-1,$$

$$v_n \mapsto v_1 - v_n$$

$$K_0(M_n(S^1)) = \text{coker } A_G = \mathbb{Z}$$

$$K_1(M_n(S^1)) = \text{ker } A_G = \mathbb{Z}$$

10. $K_*(C(SU_q(2)))$

$$G^0 = \{v_1, v_2\}$$

$$G_+^0 = \{v_1, v_2\}$$

$$A_G: \mathbb{Z} \oplus \mathbb{Z} \rightarrow \mathbb{Z} \oplus \mathbb{Z}$$

$$v_1 \mapsto v_1 + v_2 - v_1 = v_2,$$

$$v_2 \mapsto v_2 - v_2 = 0$$

$$K_0(C(SU_q(2))) = \text{coker } A_G = \mathbb{Z}$$

$$K_1(C(SU_q(2))) = \text{ker } A_G = \mathbb{Z}$$

11. $K_*(C(S_q^7))$

$$G^0 = \{v_1, v_2, v_3, v_4\}$$

$$G_+^0 = \{v_1, v_2, v_3, v_4\}$$

$$A_G: \mathbb{Z}^4 \rightarrow \mathbb{Z}^4$$

$$v_1 \mapsto v_1 + v_2 + v_3 + v_4 - v_1 = v_2 + v_3 + v_4$$

$$v_2 \mapsto v_2 + v_3 + v_4 - v_2 = v_3 + v_4$$

$$v_3 \mapsto v_3 + v_4 - v_3 = v_4$$

$$v_4 \mapsto v_4 - v_4 = 0$$

$$K_0(C(S_q^7)) = \text{coker } A_G = \mathbb{Z}$$

$$K_1(C(S_q^7)) = \text{ker } A_G = \mathbb{Z}$$

12. $K_*(C(S_q^{2n-1}))$

$$G^0 = \{v_i \mid i = 1, \dots, n\}$$

$$G_+^0 = \{v_i \mid i = 1, \dots, n\}$$

$$A_G: \mathbb{Z}^n \rightarrow \mathbb{Z}^n$$

$$v_i \mapsto \sum_{j \geq i} v_j - v_i = \sum_{j > i} v_j$$

$$K_0(C(S_q^{2n-1})) = \text{coker } A_G = \mathbb{Z}$$

$$K_1(C(S_q^{2n-1})) = \text{ker } A_G = \mathbb{Z}$$

13. $K_*(C(\mathbb{C}P_q^3))$

$$G^0 = \{v_1, v_2, v_3, v_4\}$$

$$G_+^0 = \emptyset$$

$$A_G: \emptyset \rightarrow \mathbb{Z}^4$$

$$K_0(C(\mathbb{C}P_q^3)) = \text{coker } A_G = \mathbb{Z}^4$$

$$K_1(C(\mathbb{C}P_q^3)) = \text{ker } A_G = 0$$

14. $K_*(C(\mathbb{C}P_q^{n-1}))$

$$\begin{aligned} G^0 &= \{v_i \mid i = 1, \dots, n\} \\ G_+^0 &= \emptyset \end{aligned}$$

$$A_G: \emptyset \rightarrow \mathbb{Z}^n$$

$$\begin{aligned} K_0(C(\mathbb{C}P_q^{n-1})) &= \text{coker } A_G = \mathbb{Z}^n \\ K_1(C(\mathbb{C}P_q^{n-1})) &= \ker A_G = 0 \end{aligned}$$

15. $K_*(C(S_q^6))$

$$\begin{aligned} G^0 &= \{v_1, v_2, v_3, w_1, w_2\} \\ G_+^0 &= \{v_1, v_2, v_3\} \end{aligned}$$

$$A_G: \mathbb{Z}^3 \rightarrow \mathbb{Z}^5$$

$$v_1 \mapsto v_1 + v_2 + v_3 + w_1 + w_2 - v_1 = v_2 + v_3 + w_1 + w_2$$

$$v_2 \mapsto v_2 + v_3 + w_1 + w_2 - v_2 = v_3 + w_1 + w_2$$

$$v_3 \mapsto v_3 + w_1 + w_2 - v_3 = w_1 + w_2$$

$$\begin{aligned} K_0(C(S_q^6)) &= \text{coker } A_G = \mathbb{Z} \oplus \mathbb{Z} \\ K_1(C(S_q^6)) &= \ker A_G = 0 \end{aligned}$$

16. $K_*(C(S_q^{2n}))$

$$\begin{aligned} G^0 &= \{v_1, \dots, v_n, w_1, w_2\} \\ G_+^0 &= \{v_1, \dots, v_n\} \end{aligned}$$

$$A_G: \mathbb{Z}^n \rightarrow \mathbb{Z}^{n+2}$$

$$v_i \mapsto \sum_{j \geq i} v_j + w_1 + w_2 - v_i = \sum_{j > i} v_j + w_1 + w_2$$

$$\begin{aligned} K_0(C(S_q^{2n})) &= \text{coker } A_G = \mathbb{Z} \oplus \mathbb{Z} \\ K_1(C(S_q^{2n})) &= \ker A_G = 0 \end{aligned}$$

17. $K_*(C(\mathbb{R}P_q^6))$

$$\begin{aligned} G^0 &= \{v_1, v_2, v_3, w\} \\ G_+^0 &= \{v_1, v_2, v_3\} \end{aligned}$$

$$A_G: \mathbb{Z}^3 \rightarrow \mathbb{Z}^4$$

$$v_1 \mapsto =$$

$$v_2 \mapsto =$$

$$v_3 \mapsto =$$

$$\begin{aligned} K_0(C(\mathbb{R}P_q^6)) &= \text{coker } A_G = \\ K_1(C(\mathbb{R}P_q^6)) &= \ker A_G = \end{aligned}$$

18. $K_*(C(\mathbb{R}P_q^{2n}))$

$$G^0 = \{v_1, \dots, v_n, w\}$$

$$G_+^0 = \{v_1, \dots, v_n\}$$

$$A_G: \mathbb{Z}^n \rightarrow \mathbb{Z}^{n+1}$$

$$v_i \mapsto =$$

$$K_0(C(\mathbb{R}P_q^{2n})) = \text{coker } A_G =$$

$$K_1(C(\mathbb{R}P_q^{2n})) = \text{ker } A_G =$$

3 Proof of the theorem (2.1)

Proof. There are seven steps in the proof, which we will sketch here.

1. Gauge action γ .

$$\gamma: U(1) = S^1 \rightarrow \text{Aut}(C^*(G))$$

$$\gamma_z(s_e) = z s_e,$$

$$\gamma_z(p_v) = p_v.$$

2. $C^*(G) \rtimes_\gamma U(1) \simeq C^*(G \times \mathbb{Z})$.

We construct the new graph $G \times \mathbb{Z}$


$$(G \times \mathbb{Z})^0 = G^0 \times \mathbb{Z},$$

$$(G \times \mathbb{Z})^1 = G^1 \times \mathbb{Z}.$$

It has no loops and

$$s(e, n) = (s(e), n - 1), \quad r(e, n) = (r(e), n).$$

Each loop is resolved in the infinite segment



3. $C^*(G \times \mathbb{Z})$ is AF.

It follows that $K_1(C^*(G \times \mathbb{Z})) = 0$.

4. Dual action $\hat{\gamma}$.

$$\hat{\gamma}: \mathbb{Z} \rightarrow \text{Aut}(C^*(G) \rtimes_\gamma U(1))$$

$$\hat{\gamma}_\chi(f)(t) = \langle \chi, t \rangle f(t), \quad \text{where } f: U(1) \rightarrow C^*(G).$$

5. Takesaki-Takai duality.

$$(C^*(G) \rtimes_{\gamma} U(1)) \rtimes_{\hat{\gamma}} \mathbb{Z} \simeq C^*(G) \times \mathcal{K}.$$

From the stability of K_* it follows that

$$K_*((C^*(G) \rtimes_{\gamma} U(1)) \rtimes_{\hat{\gamma}} \mathbb{Z}) \simeq K_*(C^*(G)).$$

6. Pimsner-Voiculescu sequence.

The Pimsner-Voiculescu sequence is as follows

$$\begin{array}{ccccc} K_0((C^*(G) \rtimes_{\gamma} U(1))) & \longrightarrow & K_0((C^*(G) \rtimes_{\gamma} U(1))) & \longrightarrow & K_0((C^*(G) \rtimes_{\gamma} U(1)) \rtimes_{\hat{\gamma}} \mathbb{Z}) \\ \uparrow & & & & \downarrow \\ K_1((C^*(G) \rtimes_{\gamma} U(1)) \rtimes_{\hat{\gamma}} \mathbb{Z}) & \longleftarrow & K_1((C^*(G) \rtimes_{\gamma} U(1))) & \longleftarrow & K_1((C^*(G) \rtimes_{\gamma} U(1))) \end{array}$$

where the maps are given by the formulas

$$K_*(C^*(G) \rtimes_{\gamma} U(1)) \xrightarrow{\text{id} - K_*(\hat{\gamma}^{-1})} K_*(C^*(G) \rtimes_{\gamma} U(1)),$$

$$K_*(C^*(G) \rtimes_{\gamma} U(1)) \xrightarrow{\text{id} - K_*(\beta^{-1})} K_*((C^*(G) \rtimes_{\gamma} U(1)) \rtimes_{\hat{\gamma}} \mathbb{Z}),$$

and the map $\beta: \mathbb{Z} \rightarrow \text{Aut}(C^*(G \times \mathbb{Z}))$ is given by

$$\begin{aligned} \beta_m(p_{(v,n)}) &= p_{(v,n+m)}, \\ \beta_m(s_{(e,n)}) &= s_{(e,n+m)}. \end{aligned}$$

Using preceding computations we can write the sequence as

$$\begin{array}{ccccc} K_0(C^*(G \times \mathbb{Z})) & \xrightarrow{\text{id} - K_0(\hat{\gamma}^{-1})} & K_0(C^*(G \times \mathbb{Z})) & \xrightarrow{1 - K_0(\beta^{-1})} & K_0(C^*(G)) \\ \uparrow & & & & \downarrow \\ K_1(C^*(G)) & \longleftarrow & 0 & \longleftarrow & 0 \end{array}$$

7. Computation of the kernel and cokernel of $1 - K_0(\hat{\gamma}^{-1})$.

□