

# ON INVARIANT MEASURES OF "SATELLITE" INFINITELY RENORMALIZABLE QUADRATIC POLYNOMIALS

GENADI LEVIN AND FELIKS PRZYTYCKI

ABSTRACT. Let  $f(z) = z^2 + c$  be an infinitely renormalizable quadratic polynomial and  $J_\infty$  be the intersection of forward orbits of "small" Julia sets of its simple renormalizations. We prove that if  $f$  admits an infinite sequence of satellite renormalizations, then every invariant measure of  $f : J_\infty \rightarrow J_\infty$  is supported on the postcritical set and has zero Lyapunov exponent. Coupled with [14], this implies that the Lyapunov exponent of such  $f$  at  $c$  is equal to zero, which answers partly a question posed by Weixiao Shen.

## 1. INTRODUCTION

We consider the dynamics  $f : \mathbb{C} \rightarrow \mathbb{C}$  of a quadratic polynomial. Up to a linear change of coordinates,  $f$  has the form  $f_c(z) = z^2 + c$  for some  $c \in \mathbb{C}$ . In this paper, which is the sequel of [9], we assume that  $f$  is infinitely-renormalizable. Moreover, in the main results we assume that  $f$  has infinitely many "satellite renormalizations", see e.g. [19], or below for definitions. Dynamics, geometry and topology of such system can be very non-trivial, in particular, due to the fact that different renormalization levels are largely independent.

Historically, the first example of infinitely-renormalizable one-dimensional map was, probably, the Feigenbaum period-doubling quadratic polynomial  $f_{c_F}$ , where  $c_F = -1.4\dots$  [6]. The Julia set of  $f_{c_F}$  is locally connected [7] as it follows from so-called "complex bounds", a compactness property of renormalizations which is a key

---

*Date:* December 27, 2022.

The first author partially supported by ISF grant 1226/17. The second author partially supported by National Science Centre, Poland, Grant OPUS 21 "Holomorphic dynamics, fractals, thermodynamic formalism", 2021/41/B/ST1/00461.

tool since [27], in particular, in proving the Feigenbaum-Coulet-Tresser universality conjecture [27, 20, 15]. Perhaps, more striking for us are Douady-Hubbard's examples, or alike, of infinitely-renormalizable quadratic polynomials with non-locally connected Julia sets [17, 26, 10, 11, 12, 4, 3]. As for the Feigenbaum polynomial  $f_{c_F}$ , all the renormalizations of such maps are satellite, although, contrary to  $f_{c_F}$ , combinatorics is unbounded (which, in turn, implies that those maps cannot have complex bounds [1]).

Dynamics of every holomorphic endomorphism of the Riemann sphere  $g : \hat{\mathbb{C}} \rightarrow \hat{\mathbb{C}}$  classically splits  $\hat{\mathbb{C}}$  into two subsets: the Fatou set  $F(g)$  and its complement the Julia set  $J(g)$ , where  $F(g)$  is the maximal (possibly, empty) open set where the sequence of iterates  $g^n$ ,  $n = 0, 1, \dots$  forms a normal (i.e., a precompact) family. See e.g. [2], [16] for the Fatou-Julia theory and [25] for a recent survey.

If  $g$  is a polynomial, then the Julia set  $J(g)$  coincides with the boundary of the basin of infinity  $A(\infty) = \{z \in \mathbb{C} \mid \lim_{n \rightarrow \infty} g^n(z) = \infty\}$  of  $g$ . The complement  $\mathbb{C} \setminus A(g)$  is called the filled Julia set  $K(g)$  of the polynomial  $g$ . The compact  $K(g) \subset \mathbb{C}$  is connected if and only if it contains all critical points of  $g$  in the complex plane.

A quadratic polynomial  $f_c$  with connected filled Julia set  $K(f)$  is *renormalizable* if, for some topological disks  $U \Subset V$  around the critical point 0 of  $f_c$ , and some  $p \geq 2$  (period of the renormalization), the restriction  $F := f_c^p : U \rightarrow V$  is a proper branched covering map (called polynomial-like map) of degree 2 and the non-escaping set  $K(F) = \{z \in U : F^n(z) \in U \text{ for all } n \geq 1\}$  (called the filled Julia set of the polynomial-like map  $F$ ) is connected. The map  $F : U \rightarrow V$  is then a *renormalization* of  $f_c$  and the set  $K(F)$  is a "small" (filled) Julia set of  $f_c$ . By the theory of polynomial-like mappings [5], there is a quasiconformal homeomorphism of  $\mathbb{C}$ , which is conformal on  $K(F)$ , that conjugates  $F$  on a neighborhood of  $K(F)$  to a uniquely defined another quadratic polynomial  $f_{c'}$  with connected filled Julia set. If  $f_{c'}$  is renormalizable by itself, then  $f_c$  is called twice renormalizable, etc. If  $f_c$  admits infinitely many renormalizations, it is called *infinitely-renormalizable*. The renormalization  $F = f_c^p$  is *simple* if any two sets  $f^i(K(F))$ ,  $f^j(K(F))$ ,  $0 \leq i < j \leq p - 1$ , are either disjoint or intersect each other at a unique point which does not separate either of them. A simple renormalization  $f^{p_n}$  is called *primitive* if all sets  $f^i(K_n)$ ,  $i = 0, \dots, p_n - 1$ , are disjoint and *satellite* otherwise.

To state our main results, Theorems 1.1, let  $f(z) = z^2 + c$  be infinitely renormalizable. Then its Julia set  $J = J(f)$  coincides

with the filled Julia set  $K(f)$  and is a nowhere dense compact full connected subset of  $\mathbb{C}$ . Let  $1 = p_0 < p_1 < \dots < p_n < \dots$  be the sequence of consecutive periods of simple renormalizations of  $f$  and  $J_n \ni 0$  denote the "small" Julia set of the  $n$ -renormalization (where  $J_0 = J$ ). Then  $p_{n+1}/p_n$  is an integer,  $f^{p_n}(J_n) = J_n$ , for any  $n$ , and  $f$ -orbits of  $J_n$ ,

$$orb(J_n) = \cup_{j \geq 0} f^j(J_n) = \cup_{j=0}^{p_n-1} f^j(J_n),$$

$n = 0, 1, \dots$ , form a strictly decreasing sequence of compact subsets of  $\mathbb{C}$ . Let

$$J_\infty = \cap_{n \geq 0} orb(J_n)$$

be the intersection of the orbits of the "small" Julia sets  $J_n$ . For every  $n$ , repelling periodic orbits of  $f$  are dense in  $orb(J_n)$  while each component of  $J_\infty$  is wandering, in particular,  $J_\infty$  contains no periodic points of  $f$ .

Let

$$P = \overline{\{f^n(0) | n = 1, 2, \dots\}}$$

be the postcritical set of  $f$ . Clearly,

$$P \subset J_\infty.$$

Moreover, the critical point 0 is recurrent, hence,

$$P = \omega(0),$$

where  $\omega(z)$  is the omega-limit set of a point  $z \in J$ .

We prove in [9] that  $J_\infty$  cannot contain any hyperbolic set. On the other hand, a hyperbolic set of a rational map always carries an invariant measure with a positive Lyapunov exponent. So a generalization of [9] would be that  $J_\infty$  never carries such a measure. Here we prove this generalization for a class of "satellite" infinitely-renormalizable quadratic polynomials:

**Theorem 1.1.** *Suppose that  $f(z) = z^2 + c$  admits infinitely many satellite renormalizations. Then  $f : J_\infty \rightarrow J_\infty$  has no an invariant probability measure with positive Lyapunov exponent.*

Let us comment on the behavior of the restriction map  $f : J_\infty \rightarrow J_\infty$  where  $f$  as in Theorem 1.1. First, by [9], the postcritical set  $P$  must intersect the omega-limit set  $\omega(x)$  of each  $x \in J_\infty$ . At the same time, dynamics and topology of the further restriction  $f : P \rightarrow P$  can vary. Indeed, there are infinitely renormalizable

quadratic polynomials  $f$  with all renormalizations being of satellite type such that at least one of the following holds<sup>1</sup>:

(1)  $f : P \rightarrow P$  is not minimal. This case happens in Douady-Hubbard's type examples. Indeed, by the basic construction [17],  $J_\infty$  then contains a closed invariant set  $X$  (which is the limit set for the collection of  $\alpha$ -fixed points of renormalizations) such that  $0 \notin X$ . By [9],  $X \cap P$  is non-empty. Thus  $X \cap P$  is an invariant non-empty proper compact subset of  $P$ .

(2)  $P$  is a so-called "hairy" Cantor set, in particular,  $P$  contains uncountably many non-trivial continua. This case takes place following [3].

(3)  $P$  is a Cantor set and  $f : P \rightarrow P$  is minimal; this happens whenever  $f$  either admits complex bounds (which then imply  $J_\infty = P$ ) or is robust [19]<sup>2</sup>. Under either of the two conditions,  $f : P \rightarrow P$  is a minimal homeomorphism, which is topologically conjugate to  $x \mapsto x + 1$  acting on the projective limit of the sequence of groups  $\{\mathbb{Z}/p_n\mathbb{Z}\}_{n=1}^\infty$ ; in particular,  $f : P \rightarrow P$  (hence, also  $f : J_\infty \rightarrow J_\infty$ , as it follows from the next Corollary 1.1) is uniquely ergodic in this case.

Theorem 1.1 yields the following dichotomy about the measurable dynamics of  $f : J \rightarrow J$  on the Julia set  $J$  of  $f$ . Recall that, by [22], any invariant probability measure on the Julia set of a rational function has non-negative exponent.

**Corollary 1.1.** *Let  $\mu$  be an invariant probability ergodic measure of  $f : J \rightarrow J$ . Then either*

- (i)  $\text{supp}(\mu) \cap J_\infty = \emptyset$  and its Lyapunov exponent  $\chi(\mu) > 0$ ,
- or*
- (ii)  $\text{supp}(\mu) \subset P$  and  $\chi(\mu) = 0$ .

In particular, the set  $J_\infty \setminus P$  is "measure invisible", see also Proposition 6.1 which is a somewhat stronger version of Corollary 1.1.

**Corollary 1.2.** *If  $f$  admits infinitely many satellite renormalizations, then*

$$(1.1) \quad \limsup_{n \rightarrow \infty} \frac{1}{n} \log |(f^n)'(x)| \leq 0 \text{ for any } x \in J_\infty,$$

<sup>1</sup>A more complete description of  $f : P \rightarrow P$  should follow from the methods developed in [3].

<sup>2</sup>The "robustness" can happen without "complex bounds" as it follows from [3] combined with [1].

and

$$(1.2) \quad \lim_{n \rightarrow \infty} \frac{1}{n} \log |(f^n)'(c)| = 0.$$

For the proof of Corollaries 1.1-1.2, see Sect. 6. The proof of Theorem 1.1 occupies sections 2-5.

As in [9], we use heavily a general result of [23] on the accessibility although the main idea of the proof is different. Indeed, in [9] we utilize the fact that the map cannot be one-to-one on an infinite hyperbolic set. At the present paper, to prove Theorem 1.1 we assign, loosely speaking, an external ray to a typical point of a hypothetical measure with positive exponent such that the field of such rays is invariant and has a controlled geometry. Given a satellite renormalization  $f^{p_n}$  we use the measure and the above field of rays to choose a point  $x$  and build a special domain that covers a "small" Julia set  $J_{n,x} \ni x$  such that there is a univalent pullback of the domain by  $f^{p_n}$  along the renormalization that enters into itself, leading to a contradiction. The choice of  $x$  is 'probabilistic', i.e., made from sets of positive measure, and the construction of the domain differs substantially depending on whether all satellite renormalizations of  $f$  are doubling or not.

**Acknowledgment.** The conclusion (1.2) of Corollary 1.2 that the Lyapunov exponent at the critical value equals zero answers partly a question by Weixiao Shen, which inspired the present work as well as the prior one [9].

## 2. PRELIMINARIES

Here we collect, for further references and use throughout the paper, necessary notations and general facts. (A)-(D) are slightly adapted versions of (A)-(D) in Sect. 2, [9] which are either well-known [19], [18], follow readily from known ones, or are proved here.

Let  $f(z) = z^2 + c$  be infinitely renormalizable. We keep the notations of the Introduction.

**(A).** Let  $G$  be the Green function of the basin of infinity  $A(\infty) = \{z | f^n(z) \rightarrow \infty, n \rightarrow \infty\}$  of  $f$  with the standard normalization at infinity  $G(z) = \ln|z| + O(1/|z|)$ . The external ray  $R_t$  of argument  $t \in \mathbf{S}^1 = \mathbf{R}/\mathbf{Z}$  is a gradient line to the level sets of  $G$  that has the (asymptotic) argument  $t$  at  $\infty$ .  $G(z)$  is called the (Green) level of  $z \in A(\infty)$  and the unique  $t$  such that  $z \in R_t$  is called the (external) argument (or angle) of  $z$ . A point  $z \in J(f)$  is accessible if there is

an external ray  $R_t$  which lands at (i.e., converges to)  $z$ . Then  $t$  is called an (external) argument (angle) of  $z$ .

Let  $\sigma : \mathbf{S}^1 \rightarrow \mathbf{S}^1$  be the doubling map  $\sigma(t) = 2t(\text{mod } 1)$ . Then  $f(R_t) = R_{\sigma(t)}$ .

Every point  $a$  of a repelling cycle  $O_a$  of period  $p$  is the landing point of an equal number  $v$ ,  $1 \leq v < \infty$ , of external rays where  $v$  coincides with the number of connected components of  $J(f) \setminus \{a\}$ . Their arguments are permuted by  $\sigma^p$  according to a rational rotation number  $r/q$  (written in the lowest term);  $v/q$  is the number of cycles of rays landing at  $a$ . If  $v \geq 2$ , there is an **alternative** [18]:

$r/q = 0/1$ , then  $v = 2$  so that each of two external ray landing at  $a$  is fixed by  $f^p$ ,

$r/q \neq 0/1$ , i.e.,  $q \geq 2$ , then  $v = q$ , i.e., the arguments of  $q$  rays landing at  $a$  form a single cycle of  $\sigma^p$ .

**(B)**. All periodic points of  $f$  are repelling. Given a small Julia set  $J_n$  containing 0, sets  $f^j(J_n)$ ,  $0 \leq j < p_n$ , are called small Julia sets of level  $n$ . Each  $f^j(J_n)$  contains  $p_{n+1}/p_n \geq 2$  small Julia sets of level  $n+1$ . We have  $J_n = -J_n$ . Since all renormalizations are simple, for  $j \neq 0$ , the symmetric companion  $-f^j(J_n)$  of  $f^j(J_n)$  can intersect the orbit  $\text{orb}(J_n) = \cup_{j=0}^{p_n-1} f^j(J_n)$  of  $J_n$  only at a single point which is periodic. On the other hand, since only finitely many external rays converge to each periodic point of  $f$ , the set  $J_\infty$  contains no periodic points. In particular, each component  $K$  of  $J_\infty$  is wandering, i.e.,  $f^i(K) \cap f^j(K) = \emptyset$  for all  $0 \leq i < j < \infty$ . All this implies that  $\{x, -x\} \subset J_\infty$  if and only if  $x \in K_0 := \cap_{n=1}^\infty J_n$ .

**Given  $x \in J_\infty$ , for every  $n$ , let  $j_n(x)$  be the unique  $j \in \{0, 1, \dots, p_n - 1\}$  such that  $x \in f^{j_n(x)}(J_n)$ . Let  $J_{x,n} = f^{j_n(x)}(J_n)$  be a small Julia set of level  $n$  containing  $x$  and  $K_x = \cap_{n \geq 0} J_{x,n}$ , a component of  $J_\infty$  containing  $x$ .**

In particular,  $K_0 = \cap_{n \geq 0} J_n$  is the component of  $J_\infty$  containing 0 and  $K_c = \cap_{n=1}^\infty f(J_n)$ , the component containing  $c$ .

Note that either  $p_n - j_n(x) \rightarrow \infty$  as  $n \rightarrow \infty$  or  $p_n - j_n(x) = N$  for some  $N \geq 0$  and all  $n$ , that is,  $f^N(x) \in K_0$ .

The map  $f : K_x \rightarrow K_{f(x)}$  is two-to-one if  $x = 0$  and one-to-one otherwise. Moreover, for every  $y \in J_\infty$ ,  $f^{-1}(y) \cap J_\infty$  consists of two points if  $y \in K_c$  and consists of a single point otherwise. Denote

$$J'_\infty = J_\infty \setminus \cup_{j=-\infty}^\infty f^j(K_0).$$

We conclude that:

$f : J'_\infty \rightarrow J'_\infty$  is a homeomorphism. Given  $x \in J'_\infty$  and  $m > 0$ , denote  $x_m = f^m(x)$  and

$$x_{-m} = f|_{J'_\infty}^{-m}(x),$$

that is, the only point  $f^{-m}(x) \cap J_\infty$ .

(C). Given  $n \geq 0$ , the map  $f^{p_n} : f(J_n) \rightarrow f(J_n)$  has two fixed points: the separating fixed point  $\alpha_n$  (that is,  $f(J_n) \setminus \{\alpha_n\}$  has at least two components) and the non-separating  $\beta_n$  (so that  $f(J_n) \setminus \beta_n$  has a single component).

For every  $n > 0$ , there are  $0 < t_n < \tilde{t}_n < 1$  such that two rays  $R_{t_n}$  and  $R_{\tilde{t}_n}$  land at the non-separating fixed point  $\beta_n \in f(J_n)$  of  $f^{p_n}$  and the component  $\Omega_n$  of  $\mathbf{C} \setminus (R_{t_n} \cup R_{\tilde{t}_n} \cup \beta_n)$  which does not contain 0 has two characteristic properties [18]:

- (i)  $\Omega_n$  contains  $c$  and is disjoint with the forward orbit of  $\beta_n$ ,
- (ii) for every  $1 \leq j < p_n$ , consider arguments (angles) of external rays which land at  $f^{j-1}(\beta_n)$ . The angles split  $\mathbf{S}^1$  into finitely many arcs. Then the length of any such arc is bigger than the length of the arc

$$S_{n,1} = [t_n, \tilde{t}_n] = \{t : R_t \subset \Omega_n\}.$$

Denote

$$t'_n = t_n + \frac{\tilde{t}_n - t_n}{2^{p_n}}, \quad \tilde{t}'_n = \tilde{t}_n - \frac{\tilde{t}_n - t_n}{2^{p_n}}.$$

The rays  $R_{t'_n}$ ,  $R_{\tilde{t}'_n}$  land at a common point  $\beta'_n \in f^{-p_n}(\beta_n) \cap \Omega_n$ . Introduce an (unbounded) domain  $U_n$  with the boundary to be two curves  $R_{t_n} \cup R_{\tilde{t}_n} \cup \beta_n$  and  $R_{t'_n} \cup R_{\tilde{t}'_n} \cup \beta'_n$ . Then  $c \in U_n$  and  $f^{p_n} : U_n \rightarrow \Omega_n$  is a two-to-one branched covering. Also,

$$f(J_n) = \{z : f^{kp_n}(z) \in \bar{U}_n, G(f^{kp_n}(z)) < 10, k = 0, 1, \dots\}.$$

Let

$$s_{n,1} = [t_n, t'_n] \cup [\tilde{t}'_n, \tilde{t}_n]$$

so that  $s_{n,1} \subset S_{n,1}$  and argument of any ray to  $f(J_n)$  lies in  $s_{n,1}$ .

Let us iterate this construction. Given  $1 \leq j \leq p_n$ , let  $S_{n,j}$  be one of the two arcs of  $\mathbf{S}^1$  with end points

$$t_{n,j} = \sigma^{j-1}(t_n), \tilde{t}_{n,j} = \sigma^{j-1}(\tilde{t}_n)$$

such that arguments of any ray to  $f^j(J_n)$  lies in  $S_{n,j}$ . Let

$$s_{n,j} = \sigma^{j-1}(s_{n,1}) = [t_{n,j}, t'_{n,j}] \cup [\tilde{t}'_{n,j}, \tilde{t}_{n,j}]$$

where  $t'_{n,j} = \sigma^{j-1}(t'_n)$ ,  $\tilde{t}'_{n,j} = \sigma^{j-1}(\tilde{t}'_n)$ . Then

$$s_{n,j} \subset S_{n,j}$$

and argument of any ray to  $f^j(J_n)$  lies in fact in  $s_{n,j}$ . Note that

$$(2.1) \quad t'_{n,j} - t_{n,j} = \tilde{t}_{n,j} - \tilde{t}'_{n,j} = \frac{\tilde{t}_n - t_n}{2^{p_n-j+1}} < \tilde{t}_n - t_n < 1/2.$$

So  $\sigma^{j-1} : s_{n,1} \rightarrow s_{n,j}$  is a homeomorphism and  $s_{n,j}$  has two components ('windows')  $[t_{n,j}, t'_{n,j}]$  and  $[\tilde{t}'_{n,j}, \tilde{t}_{n,j}]$  of equal length.

Let  $U_{n,j} = f^{j-1}(U_n)$  and  $\beta_{n,j} = f^{j-1}(\beta_n)$ . The domain  $U_{n,j}$  is bounded by two rays  $R_{t_{n,j}} \cup R_{\tilde{t}_{n,j}}$  converging to  $\beta_{n,j}$  and completed by  $\beta_{n,j}$  along with two rays  $R_{t'_{n,j}} \cup R_{\tilde{t}'_{n,j}}$  completed by their common limit point  $f^{j-1}(\beta'_n)$  where  $t'_{n,j} = \sigma^{j-1}(t'_n)$ ,  $\tilde{t}'_{n,j} = \sigma^{j-1}(\tilde{t}'_n)$ .

By (i)-(ii), for a fixed  $n$ , domains  $U_{n,j}$ ,  $1 \leq j \leq p_n$ , are pairwise disjoint.

Let  $U_{n,j-p_n}$  be a component of  $f^{-(p_n-j)}(U_n)$  which is contained in  $U_{n,j}$ . Then

$$(2.2) \quad f^{p_n} : U_{n,j-p_n} \rightarrow U_{n,j}$$

is a two-to-one branched covering and

$$f^{j-1}(J_n) = \{z : f^{kp_n}(z) \in \bar{U}_{n,j-p_n}, G(f^{kp_n}(z)) < 10, k = 0, 1, \dots\}.$$

Let  $s_{n,j}^1$  be the set of arguments of rays entering  $U_{n,j-p_n}$ . Then  $s_{n,j}^1$  consists of 4 components so that  $\sigma^{p_n}$  map homeomorphically each of these components onto one of the 'windows' of  $s_{n,j}$ .

Furthermore, let

$$\Omega_{n,j} = f^{j-1}(\Omega_n).$$

Unless the map (2.2), the map

$$(2.3) \quad f^{p_n} : U_{n,j} \rightarrow \Omega_{n,j}$$

is a two-to-one branched covering only *assuming*  $f^{j-1} : \Omega_n \rightarrow \Omega_{n,j}$  is a homeomorphism, which holds *if and only if*  $\sigma^{j-1} : S_{n,1} \rightarrow \sigma^{j-1}(S_{n,1})$  is a homeomorphism. In the latter case,

$$\sigma^{j-1}(S_{n,1}) = S_{n,j}.$$

**Primitive vs satellite** renormalizations. Let  $n \geq 2$  and  $k_n/q_n$  be the rotation number of  $\beta_n$ . The next claim is well-known, we include the proof for reader's convenience.

**Lemma 2.1.** (1) *the renormalization  $f^{p_n}$  is primitive if and only if  $k_n/q_n = 0/1$ , the period of  $\beta_n$  is  $p_n$  and  $\beta_n$  is the landing point of exactly two rays and they are fixed by  $f^{p_n}$ ,*  
 (2) *points  $\beta_n$ ,  $n = 1, 2, \dots$  are all different,*



- (3)  $f^{p_n}$  is satellite if and only if the  $\alpha$ -fixed point  $\alpha_{n-1}$  of  $f^{p_{n-1}} : f(J_{n-1}) \rightarrow f(J_{n-1})$  coincides with the  $\beta$ -fixed point  $\beta_n$  of  $f^{p_n} : f(J_n) \rightarrow f(J_n)$ . In particular,  $\cup_{j=0}^{q_n-1} f^{j p_{n-1}}(f(J_n)) \subset f(J_{n-1})$  and  $q_n = p_n q_{n-1}$ . Moreover, each of  $p_{n-1}$  points of the orbit of  $\beta_n$  is the landing points of precisely  $q_n$  rays which are permuted by  $f^{p_{n-1}}$  according to the rotation number  $r_n/q_n$ . Completed by the landing point they split  $\mathbb{C}$  into  $q_n$  "sectors" such that the closure of each of them contains a unique "small" Julia set of level  $n$  sharing a common point with the boundary of the "sector".

*Proof.* (1).  $f^{p_n}$  is satellite if and only if  $f(J_n)$  meets at  $\beta_n$  some other iterate of  $J_n$ , hence,  $r_n/q_n \neq 0$ , and vice versa. (2). assume  $\beta := \beta_n = \beta_m$  for some  $0 \leq n < m$ . As  $p_n < p_m$ , the period of  $\beta_m$  is smaller than  $p_n$ . It follows that  $f(J_n)$  contains two small Julia sets of level  $m$  that meet at  $\beta$ , hence,  $\beta$  separates  $f(J_n)$ , a contradiction as  $\beta_n$  does not. (3). By (1),  $f^{p_n}$  is satellite if and only if  $r_n/q_n \neq 0$ . Let  $\tilde{p}_{n-1} = p_n/q_n$ . Then  $\tilde{p}_{n-1}$  is an integer and is equal to the period of  $\beta_n$ . It follows that  $p_n$  sets  $f(J_n), f^2(J_n), \dots, f^{p_n}(J_n)$  are split into  $\tilde{p}_{n-1}$  connected closed subsets  $E_i$ ,  $i = 1, \dots, \tilde{p}_{n-1}$  where  $E_1 = \cup_{j=0}^{q_n-1} f^{j \tilde{p}_{n-1}}(f(J_n))$  and  $E_i = f^{i-1}(E_1)$ ,  $i = 1, 2, \dots, \tilde{p}_{n-1}$ . Moreover,  $0 \in E_{p_{n-1}}$  and  $f(E_i) = E_{i+1}$ ,  $i = 1, \dots, \tilde{p}_{n-1} - 1$ ,  $f(E_{\tilde{p}_{n-1}}) = E_1$ . By [19, Theorem 8.5],  $f^{\tilde{p}_{n-1}}$  is a simple renormalization and  $E_i$ ,  $i = 1, \dots, \tilde{p}_{n-1}$  are subsets of its  $\tilde{p}_{n-1}$  small Julia sets. Since  $1 = p_0 < p_1 < \dots$  are all consecutive periods of simple renormalizations, then  $\tilde{p}_{n-1} = p_k$  for some  $k < n$ . Therefore,  $\beta_n$ -fixed point of  $f^{p_n} : f(J_n) \rightarrow f(J_n)$  is  $\alpha_k$ -fixed point of  $f^{p_k} : f(J_{p_k}) \rightarrow f(J_{p_k})$ . As all renormalizations are simple, if  $k < n - 1$  that would imply that  $\beta_n = \beta_{n-1} = \dots = \beta_{k+1}$ , a contradiction with (2). The claim about "sectors" follows since each map  $f^j$  is one-to-one in a neighborhood of  $\beta_n$  and the closure of  $\Omega_n$  contains a single "small" Julia set  $f(J_n)$  of level  $n$  sharing a common point with  $\partial\Omega_n$ .  $\square$

We need a more refined estimate provided the renormalization is not doubling. Assume  $f^{p_n}$  is satellite so that  $p_{n-1} = p_n/q_n$  with  $q_n \geq 2$  and the rotation number of  $\beta_n$  is  $r_n/q_n \neq 0/1$ .

**Lemma 2.2.** *Assume  $f^{p_n}$  is satellite and  $q_n = p_n/p_{n-1} \geq 3$ , i.e.,  $f^{p_n}$  is not doubling. Then*

$$(2.4) \quad \sigma^{j-1} : S_{n,1} \rightarrow \sigma^{j-1} S_{n,1} \text{ is a homeomorphism for } j = 1, \dots, p_{n-1}(q_n - 2).$$

In particular, given  $\zeta \in (0, 1/3)$ , the length of  $\sigma^{j-1}S_{n,1}$  tends to zero as  $n \rightarrow \infty$  uniformly in  $j = 1, \dots, [\zeta p_n]$  (where  $[x]$  is the integer part of  $x \in \mathbb{R}$ ).

Moreover, for every  $1 \leq j \leq p_{n-1}(q_n - 2)$ ,  $S_{n,j} = \sigma^{j-1}(S_{n,1})$  and the map  $f^{p_n} : U_{n,j} \rightarrow \Omega_{n,j}$  is a two-to-one branched covering such that

$$f^j(J_n) = \{z : f^{kp_n}(z) \in \bar{U}_{n,j}, G(f^{kp_n}(z)) < 10, k = 0, 1, \dots\}.$$

*Proof.* Let  $g = f^{p_{n-1}} : U_{n-1} \rightarrow \Omega_{n-1}$ . Then  $g$  is a two-to-one covering of degree 2 and the critical value  $c$ .

(1) Recall that  $s_{n-1,1} = [t_{n-1}, t'_{n-1}] \cup [\tilde{t}'_{n-1}, \tilde{t}_{n-1}]$  consists of two 'windows' so that  $\sigma^{p_{n-1}}$  is orientation preserving homeomorphism of either 'window' onto  $S_{n-1,1} = [t_{n-1}, \tilde{t}_{n-1}]$ .

(2) Consider  $q_n$  rays  $L_1, \dots, L_{q_n}$  to  $\alpha_{n-1}$ . The map  $g$  is a local homeomorphism near  $\alpha_{n-1}$  which permutes the rays to  $\alpha_{n-1}$  according to the rotation number  $\nu := k_n/q_n \neq 0, 1/2$ . In particular,  $g$  maps any pair of adjacent rays to  $\alpha_{n-1}$  onto another pair of adjacent rays to  $\alpha_{n-1}$ .

(3) Not all arguments of these rays lie in a single 'window'  $I$  of  $s_{n-1,1}$  because otherwise, by (1), the set of those arguments would lie in the non-escaping set of an orientation preserving homeomorphism  $\sigma^{p_{n-1}} : I \rightarrow S_{n,1}$ , which consists of a fixed point of this map, a contradiction with the fact that  $q_n > 1$ .

(4) The rays  $L_j$  split  $U_{n-1}$  into  $q_n$  disjoint domains  $U^j$ ,  $j = 0, 1, \dots, q_n - 1$ . By the "ideal boundary"  $\hat{\partial}U^j$  of  $U^j$  we will mean the usual (topological) boundary  $\partial U^j$  (in our case, the set of boundary rays completed by their landing points) along with the "boundary at infinity" which is the set of arguments of rays entering  $U^j$ . Then define  $\hat{g}$  on  $\hat{\partial}U^j$  to be  $g$  on  $\partial U^j$  and  $\sigma^{p_{n-1}}$  on the "boundary at infinity" of  $U^j$ .

(5) By (3), one of  $U^j$ , called  $U^0$ , has  $\beta_{n-1}$  in its boundary, and another one, called  $U^{q_n-1}$ , has  $\beta'_{n-1}$  in the boundary. In particular, the boundary of any other  $U^j$ ,  $j \neq 0, q_n - 1$ , consists of a pair of adjacent rays to  $\alpha_{n-1}$  whose arguments belong to a single 'window' of  $s_{n-1,1}$ . Therefore, by (1), the rest of indices  $j = 1, \dots, q_n - 2$  can be ordered in such a way that  $\hat{g} : \hat{\partial}U^j \rightarrow \hat{\partial}U^{j+1}$  is a one-to-one map for  $j = 1, \dots, q_n - 3$  (note that the "boundary at infinity" of each  $U^j$ ,  $1 \leq j \leq q_n - 2$ , consists of a single "arc at infinity"). Therefore,  $g : U^j \rightarrow U^{j+1}$  is a homeomorphism for  $j = 1, \dots, q_n - 3$ . The map  $\hat{g}$  on  $\hat{\partial}U^{q_n-2}$  is also a one-to-one map on its image  $W = g(U^{q_n-2})$  where  $W$  is bounded by two adjacent

rays to  $\alpha_{n-1}$ .  $W$  cannot contain  $U^0$  because otherwise  $W$  would contain  $\beta'_{n-1}$ , a contradiction. Thus  $W$  must contain  $\beta'_{n-1}$ . That is,  $g(U^{q_n-2})$  covers  $U^{q_n-1}$ .

Thus, for  $j = 1, \dots, q_n - 3$ ,  $g : U^j \rightarrow U^{j+1}$  is a homeomorphism, and  $g : U^{q_n-2} \rightarrow W$  is also a homeomorphism where the image  $W = g(U^{q_n-2})$  covers  $U^{q_n-1}$  and has two common rays with the boundary of  $U^{q_n-1}$ .

(6) The critical value  $c$  of  $g$  has a unique preimage by  $g$  (the critical point of  $g$ ). As  $c \in \Omega_n \subset \Omega_{n-1}$  and  $\Omega_n$  is bounded by two adjacent rays to  $\alpha_{n-1}$ ,  $c \in U^i$  for some  $i \in \{1, \dots, q_n - 1\}$ . If  $i > 1$ , then  $i - 1 \geq 1$  while  $g$  would not be a homeomorphism of  $U^{i-1}$  on its image. This shows that  $c \in U^1 = \Omega_n$ .

Concluding,  $U^j = g^{j-1}(\Omega_n)$ ,  $j = 1, \dots, q_n - 2$ , in particular,

$$\Omega_n, g(\Omega_n), \dots, g^{q_n-3}(\Omega_n) \subset U_{n-1}$$

and  $g^{q_n-2} : \Omega_n \rightarrow g^{q_n-2}(\Omega_n)$  is a homeomorphism, that is, (2.4) holds. It implies the rest.  $\square$

(D). Given a compact set  $Y \subset J(f)$  denote by  $(\tilde{Y})_f$  (or simply  $\tilde{Y}$ , if the map is fixed) the set of arguments of the external rays which have their limit sets contained in  $Y$ . It follows from (C) that  $\tilde{K}_c = \bigcap_{n=1}^{\infty} \{[t_n, t'_n] \cup [\tilde{t}'_n, \tilde{t}_n]\}$ , i.e., it is either a single-point set or a two-point set.

Since  $\tilde{K}_c$  contains at most two angles,  $K_c$  contains at most two different accessible points. More generally, given  $x \in J'_\infty$  let

$$s_{n,j_n(x)} = [t_{n,j_n(x)}, t'_{n,j_n(x)}] \cup [\tilde{t}'_{n,j_n(x)}, \tilde{t}_{n,j_n(x)}].$$

Then  $s_{n+1,j_{n+1}(x)} \subset s_{n,j_n(x)}$  so that

$$s_{\infty,x} := \bigcap_{n>0} s_{n,j_n(x)}$$

is not empty and consists of either one or two components. Since  $p_n - j_n(x) \rightarrow \infty$  for  $x \in J'_\infty$  we conclude using (2.1):

$s_{\infty,x}$  consists of either a single point or two different points. In particular, for any component  $K$  of  $J_\infty$  which is not one of  $f^{-j}(K_0)$ ,  $j \geq 0$ , there is either one or two rays tending to  $K$ .

**From now on,  $\mu$  is an  $f$ -invariant probability ergodic measures supported in  $J_\infty$ :  $\text{supp } \mu \subset J_\infty$ , and having a positive Lyapunov exponent**

$$\chi(\mu) := \int \log |f'| d\mu > 0.$$

(E). We start with the following basic statement. Parts (i)-(ii) are easy consequences of the invariance of  $\mu$  and (B) while (iii) is a part of Pesin's theory as in [24] coupled with the structure of  $f : J_\infty \rightarrow J_\infty$ , see (B). Recall that  $J'_\infty = J_\infty \setminus \cup_{j=-\infty}^\infty f^j(K_0)$ .

**Proposition 2.3.** (i) For every  $n$  and  $0 \leq j < p_n$ ,  $\mu(f^j(J_n)) = 1/p_n$ .

(ii)  $\mu$  has no atoms and  $\mu(K) = 0$  for every component  $K$  of  $J_\infty$ .

(iii)  $\mu(J'_\infty) = 1$  and  $f : J'_\infty \rightarrow J'_\infty$  is a  $\mu$ -measure preserving homeomorphism. There exists a measurable positive function  $\tilde{r}(x) > 0$  on  $J'_\infty$  such that for  $\mu$ -almost every  $x \in J'_\infty$ , and all  $n \in \mathbf{N}$ , if  $x_{-n}$  is the unique point of  $J'_\infty$  with  $f^n(x_{-n}) = x$ , then a (univalent) branch  $g_n : B(x, \tilde{r}(x)) \rightarrow \mathbf{C}$  of  $f^{-n}$  is well-defined such that  $g_n(x) = x_{-n}$ ,

*Remark 2.4.* The branch  $g_n$  of  $f^{-n}$  depends on  $n$  and  $x_{-n}$  but it should be clear from the context which points  $x$  and  $x_{-n}$  are meant.

Using the Birkhoff Ergodic Theorem and Egorov's theorem, Proposition 2.3 implies immediately (e1)-(e3) of the next corollary. The proof of (e4)-(e5) is given right after it.

**Corollary 2.5.** For every  $\epsilon > 0$ , there exists a closed set  $E'_{\epsilon/2} \subset J'_\infty$  and constants  $\rho = \rho(\epsilon) > 0$ ,  $\kappa = \kappa(\epsilon) \in (0, 1)$  such that:

(e1)  $\mu(E'_{\epsilon/2}) > 1 - \frac{\epsilon}{2}$ ,

(e2) there exists another closed set  $\hat{E}_{\epsilon/2}$  such that  $E'_{\epsilon/2} \subset \hat{E}_{\epsilon/2} \subset J'_\infty$  as follows. For every  $x \in \hat{E}_{\epsilon/2}$  and every  $m > 0$  there exists a (univalent) branch  $g_m : B(x, 3\rho) \rightarrow \mathbf{C}$  of  $f^{-m}$  such that  $g_m(x) = x_{-m}$  and  $|g'_m(x_1)/g'_m(x_2)| < 2$ , for every  $x_1, x_2 \in B(x, 2\rho)$ . Moreover,  $m^{-1} \ln |Dg_m(x)| \rightarrow -\chi(\mu)$  as  $m \rightarrow \infty$  uniformly in  $x \in E'_{\epsilon/2}$ ,

(e3) for every  $x \in E'_{\epsilon/2}$  there exists a sequence of positive integers  $n_j = n_j(x)$ ,  $j = 1, 2, \dots$ , such that  $j/n_j \geq \kappa$  and  $f^{n_j}(x) \in \hat{E}_{\epsilon/2}$  for all  $j$  (in fact,  $\{n_j\}_{j=1}^\infty = \{n \in \mathbf{N} : f^n(x) \in \hat{E}_{\epsilon/2}\}$ ),

(e4) given  $x \in J_\infty$  and  $n \geq 0$ , let  $j_n(x)$  be the unique  $1 \leq j < p_n$  such that  $x \in f^j(J_n)$ . Then  $p_n - j_n(x) \rightarrow \infty$  as  $n \rightarrow \infty$  uniformly in  $x \in E'_{\epsilon/2}$ ,

(e5) for  $s_{n,j_n(x)} = [t_{n,j_n(x)}, t'_{n,j_n(x)}] \cup [\tilde{t}'_{n,j_n(x)}, \tilde{t}_{n,j_n(x)}]$ , we have:  $s_{n+1,j_{n+1}(x)} \subset s_{n,j_n(x)}$  and

$$|t_{n,j_n(x)} - t'_{n,j_n(x)}| = |\tilde{t}'_{n,j_n(x)} - \tilde{t}_{n,j_n(x)}| \rightarrow 0$$

as  $n \rightarrow \infty$  uniformly in  $x \in E'_{\epsilon/2}$ .

Proof of  $(e_4)$ - $(e_5)$ : assuming the contrary in  $(e_4)$ , we find some  $N$  and sequences  $(n_k) \subset \mathbb{N}$  and  $(x_k)$ ,  $x_k \in E'_{\epsilon/2}$ , such that  $p_{n_k} - j_{n_k}(x_k) = N$ , hence,  $x_k \in f^{-N}(J_{n_k})$ , for all  $k$ . Since  $E_{\epsilon/2}$  is closed, one can assume  $x_k \rightarrow x \in E'_{\epsilon/2} \subset J'_\infty$ . Hence,  $x \in f^{-N}(K_0)$ , a contradiction. Now, for  $(e_5)$  using  $(e_4)$ ,  $t'_{n,j_n(x)} - t_{n,j_n(x)} = \tilde{t}_{n,j_n(x)} - \tilde{t}'_{n,j_n(x)} < \frac{1}{2^{p_n - j_n(x)}} \rightarrow 0$  uniformly in  $x$ .

### 3. EXTERNAL RAYS TO TYPICAL POINTS

We define a *telescope* following essentially [23]. Given  $x \in J(f)$ ,  $r > 0$ ,  $\delta > 0$ ,  $k \in \mathbb{N}$  and  $\kappa \in (0, 1)$ , an  $(r, \kappa, \delta, k)$ -telescope at  $x \in J$  is collections of times  $0 = n_0 < n_1 < \dots < n_k = n$  and disks  $B_l = B(f^{n_l}(x), r)$ ,  $l = 0, 1, \dots, k$  such that, for every  $l > 0$ : (i)  $l/n_l > \kappa$ , (ii) there is a univalent branch  $g_{n_l} : B(f^{n_l}(x), 2r) \rightarrow \mathbf{C}$  of  $f^{-n_l}$  so that  $g_{n_l}(f^{n_l}(x)) = x$  and, for  $l = 1, \dots, k$ ,  $d(f^{n_{l-1}} \circ g_{n_l}(B_l), \partial B_{l-1}) > \delta$  (clearly, here  $f^{n_{l-1}} \circ g_{n_l}$  is a branch of  $f^{-(n_l - n_{l-1})}$  that maps  $f^{n_l}(x)$  to  $f^{n_{l-1}}(x)$ ). The trace of the telescope is a collection of sets  $B_{l,0} = g_{n_l}(B_l)$ ,  $l = 0, 1, \dots, k$ . We have:  $B_{k,0} \subset B_{k-1,0} \subset \dots \subset B_{1,0} \subset B_{0,0} = B_0 = B(x, r)$ .

By the *first point of intersection* of a ray  $R_t$ , or an arc of  $R_t$ , with a set  $E$  we mean a point of  $R_t \cap E$  with the minimal level (if it exists).

**Theorem 3.1.** [23] *Given  $r > 0$ ,  $\kappa \in (0, 1)$ ,  $\delta > 0$  and  $C > 0$  there exist  $M > 0$ ,  $\tilde{l}, \tilde{k} \in \mathbb{N}$  and  $K > 1$  such that for every  $(r, \kappa, \delta, k)$ -telescope the following hold. Let  $k > \tilde{k}$ . Let  $u_0 = u$  be any point at the boundary of  $B_k$  such that  $G(u) \geq C$ . Then there are indexes  $1 \leq l_1 < l_2 < \dots < l_j = k$  such that  $l_1 < \tilde{l}$ ,  $l_{i+1} < Kl_i$ ,  $i = 1, \dots, j-1$  as follows. Let  $u_k = g_{n_k}(u) \in \partial B_{k,0}$  and let  $\gamma_k$  be an infinite arc of an external ray through  $u_k$  between the pint  $u_k$  and  $\infty$ . Let  $u_{k,k} = u_k$  and, for  $l = 1, \dots, k-1$ , let  $u_{k,l}$  be the first point of intersection of  $\gamma_k$  with  $\partial B_{l,0}$ . Then, for  $i = 1, \dots, j$ ,*

$$G(u_{k,l_i}) > M2^{-n_{l_i}}.$$

Next corollary of Theorem 3.1 is a key one.

**Proposition 3.1.** *Given  $\epsilon > 0$  there exists a closed set  $E_\epsilon$  as follows. First,  $\mu(E_\epsilon) > 1 - \epsilon$  and  $E_\epsilon \subset E'_{\epsilon/2}$  where  $E'_{\epsilon/2}$  is the set defined in  $(E)$  and satisfies  $(e_1)$ - $(e_5)$ . There exists  $r = r(\epsilon) > 0$  and, for each  $\nu > 0$  there is  $C(\nu) > 0$  as follows.*

(1) *Let  $x \in E_\epsilon$ . Then  $x$  is the landing point of an external ray  $R_{t(x)}$  of argument  $t(x)$ . Moreover, the first intersection of  $R_{t(x)}$  with  $\partial B(x, \nu)$  has the level at least  $C(\nu)$ .*

(2) for each  $n$  a branch  $g_n : B(x, 2r) \rightarrow \mathbf{C}$  of  $f^{-n}$  is well-defined such that  $g_n(x) = x_{-n}$ ,  $|g_n(x_1)/g_n(x_2)| < 2$ , for every  $x_1, x_2 \in B(x, r)$  and  $n^{-1} \ln |Dg_n(x)| \rightarrow -\chi(\mu)$  as  $m \rightarrow \infty$  uniformly in  $x \in E_\epsilon$ ,

(3) if  $x' = g_n(x) \in E_\epsilon$ , then  $f^n(R_{t(x')}) = R_{t(x)}$ .

*Proof.* (1)-(2) will hold already for the set  $E'_{\epsilon/2}$  which follows from Theorem 3.1 as in [23] and uses only that  $\mu$  has a positive exponent; (3) will follow in our case as we shrink a bit the set  $E'_{\epsilon/2}$  since each point  $x \in J'_\infty$  admits at most two external arguments. Here are details. Let  $r = \rho(\epsilon)$  and  $\kappa = \kappa(\epsilon)$  as in the properties (e<sub>2</sub>)-(e<sub>3</sub>) of the set  $E'_{\epsilon/2}$ . Then, by (e<sub>2</sub>)-(e<sub>3</sub>), there is  $\delta > 0$  such that, for each  $k$ , every  $x \in E'_{\epsilon/2}$  admits  $(r, \kappa, \delta, k)$ -telescope with the times  $0 = n_0 < n_1 < n_2 < \dots < n_k$  that appear in the property (e<sub>3</sub>) of  $E'_{\epsilon/2}$ . On the other hand, there exists  $L_r > 0$  such that for every  $z \in J(f)$  there is a point  $u(z) \in \partial B(z, r)$  with the level  $G(u(z)) > L_r$ . Given this  $C = L_r$ , let  $M, \tilde{l}$  and  $\tilde{k}$  be as in Theorem 3.1.

Let  $x \in E'_{\epsilon/2}$  and  $n_1 < n_2 < \dots < n_k < \dots$  as in (e<sub>3</sub>). Fix  $k > \tilde{k}$ . Let  $B_{k,0}(x) \subset B_{k-1,0}(x) \subset \dots \subset B_{1,0}(x) \subset B_{0,0}(x)$  be the corresponding trace. By Theorem 3.1, there are  $1 \leq l_{1,k}(x) < l_{2,k}(x) < \dots < l_{j_k^x,k}(x) = k$  such that  $l_{1,k}(x) < \tilde{l}$ ,  $l_{i+1,k}(x) < Kl_{i,k}(x)$ ,  $i = 1, \dots, j_k^x - 1$ . Let  $\gamma_k(x)$  be an arc of an external ray between the point  $u_k(x) := g_{n_k}(u(f^{n_k}(x)))$  and  $\infty$ . Let  $u_{k,l}(x)$  be the first intersection of  $\gamma_k(x)$  with  $\partial B_{l,0}(x)$ . Then, for  $i = 1, \dots, j_k^x - 1$ ,

$$(3.1) \quad G(u_{k,l_{i,k}(x)}(x)) > M2^{-n_{l_{i,k}(x)}} > M2^{-l_{i,k}(x)/\kappa}.$$

For all  $i = 1, \dots, j_k^x - 1$ ,

$$(3.2) \quad i \leq l_{i,k}(x) < K^i \tilde{l}.$$

Denote by  $t_k(x)$  the argument of an external ray that contains the arc  $\gamma_k(x)$ .

Now, given a sequence

$$(3.3) \quad k_1 < k_2 < \dots < k_m < \dots$$

such that  $k_1 > \tilde{k}$ , we get a sequence of arguments  $t_{k_m}(x)$  and a sequence of arcs  $\gamma_{k_m}(x)$  of external rays of the corresponding arguments  $t_{k_m}(x)$ . Passing to a subsequence in the sequence  $(k_m)$ , if necessary, one can assume that  $t_{k_m}(x) \rightarrow \tilde{t}(x)$ , for some argument  $\tilde{t}(x)$ .

Fix any  $\nu \in (0, r)$  and choose  $\tilde{k}_0 > \tilde{k}$  such that,

$$2 \exp(-K^{\tilde{k}_0-2} \tilde{l} \chi(\mu)) < \nu \text{ and let } C(\nu) = M(2^{-1/\kappa})^{\tilde{l} K^{\tilde{k}_0}}.$$

Then, by Theorem 3.1, for each  $k_m > k_0$ , the first intersection of the ray  $R_{t_{k_m}}(x)$  with the boundary of  $B(x, \nu)$  has the level at least  $C(\nu)$ . It follows, for any  $0 < C < C(\nu)$ , the sequence of arcs of the rays  $R_{t_{k_m}(x)}$  between the levels  $C$  and  $C(\nu)$  do not exit  $B(x, \nu)$  for all  $k_m > k_0$ . As  $t_{k_m}(x) \rightarrow \tilde{t}(x)$ , it follows that the arc of the ray  $R_{\tilde{t}(x)}$  between levels  $C$  and  $C(\nu)$  stays in  $B(x, \nu)$  too. As  $\nu > 0$  and  $C \in (0, C(\nu))$  can be chosen arbitrary small,  $R_{\tilde{t}(x)}$  must land at  $x$  and satisfy (1) with  $t(x)$  replaced by  $\tilde{t}(x)$ .

Let us call the above procedure of getting  $\tilde{t}(x)$  from the constants  $r, L_r$ , the point  $x \in E'_{\epsilon/2}$  and the sequence (3.3) the  $(r, L_r, x, (k_m))$ -procedure.

Note that (2) is property  $(e_2)$  of the set  $E'_{\epsilon/2}$ .

In order to satisfy property (3), we shrink the set  $E'_{\epsilon/2}$  and correct  $\tilde{t}(x)$  changing it to some  $t(x)$  (if necessary) as follows. Using the Birkhoff Ergodic Theorem and Egorov's theorem, choose a closed subset  $E_\epsilon$  of  $E'_{\epsilon/2}$  such that  $\mu(E_\epsilon) > 1 - \epsilon$  and, for each  $x \in E_\epsilon$ , the set  $\mathcal{N}(x) := \{N \in \mathbb{N} : f^N(x) \in E'_{\epsilon/2}\}$  is infinite. Note that  $\mathcal{N}(x) \subset \{n_k\}_{k=1}^\infty$ . We have proved that, for each  $N \in \mathcal{N}(x)$ , (1) holds for the point  $f^N(x)$  instead of  $x$ , in particular,  $\tilde{t}(f^N(x))$  is an argument of  $f^N(x)$ . On the other hand, by (D1), each  $y \in E_\epsilon$  admits at most two external arguments, hence, all possible external arguments of the forward orbit  $f^n(x)$ ,  $n \geq 0$ , belong to at most two different orbits of  $\sigma : S^1 \rightarrow S^1$ . Hence, there is one of those orbits,  $O = \{\sigma^n(t(x))\}_{n \geq 0}$  for some  $t(x)$ , such that the intersection  $O \cap \{\tilde{t}(f^N(x)) : N \in \mathcal{N}(x)\}$  is an infinite set, so that  $\tilde{t}(f^{n_{k_m}(x)}(x)) = \sigma^{n_{k_m}(x)}(t(x))$  for an infinite sequence  $(k_m(x))_{m \geq 1}$ .

Let's start over with the  $(r/2, C(r/2), x, (k_m(x)))$ -procedure for the point  $x$  and the sequence  $\{k_j(x)\}$ . Then, by the construction,  $t_{k_m(x)} = t(x)$  for all  $m$ , hence, (1) holds with  $t(x)$  instead of the previous  $\tilde{t}(x)$ . If  $y \in E_\epsilon$  is any other point of the grand orbit  $\{f^n(x) : n \in \mathbb{Z}\}$  (remember that  $f : J'_\infty \rightarrow J'_\infty$  is invertible), the  $(r/2, C(r/2), y, (k_m))$ -procedure works for  $y$  with the same (perhaps, truncated) sequence  $k_1(x) < k_2(x) < \dots$ , which ensures that (3) holds (for the corrected arguments) too.  $\square$

*Remark 3.2.* Given  $t(x)$ , we cannot just set  $t(f^n(x)) = \sigma^n(t(x))$  to satisfy property (3) because this would change  $\kappa$  in the definition of telescope, so we might lose property (1). Notice that

correcting (flipping)  $\tilde{t}(x)$  to  $t(x)$  does not change  $C(\nu)$ . The same for flipping any  $t(y)$  in the grand orbit of  $x$ . But the flipping can make  $f^\ell(R_{t(y)}) = R_{t(f^N(x))}$  for  $f^\ell(y) = f^N(x)$  where  $N = n_{k_m}$  with  $G(R_{t(f^\ell(y))} \cap \partial B(f^\ell(y), r/2)) > L_{r/2}$ , thus yielding (3).

#### 4. LEMMAS

**Lemma 4.1.** *Let  $z_k \in \cup_{j=0}^{p_{n_k}-1} f^j(J_{n_k})$  where  $n_k \nearrow \infty$ .*

(a) *If  $z_k \rightarrow z$  then  $z \in J_\infty$ .*

(b)  *$z \in J_{n,z} \cap J'_\infty$  yields  $z_{\pm p_n} \in J_{n,x}$ . If, additionally to (a),  $z_k \in J'_\infty$  for all  $k$  and  $w_k \rightarrow w$  where  $w_k = (z_k)_{ep_{n_k}}$ , where  $e$  is always either 1 or  $-1$  then  $z$  and  $w$  are in the same component of  $J_\infty$ .*

(c) *If  $z_k \in E_\epsilon$  for all  $k$  and  $t(z_k) \rightarrow t$  (where  $E_\epsilon, t(z_k)$  are defined in Proposition 3.1), then the ray  $R_t$  lands at the limit point  $z$ . In particular, given  $\sigma > 0$  there is  $\Delta(\sigma) > 0$  such that  $|x_1 - x_2| < \sigma$  for some  $x_1, x_2 \in E_\epsilon$  whenever  $|t(x_1) - t(x_2)| < \Delta(\sigma)$ .*

*Proof.* (a) Assume the contrary. Then there is  $n$  such that  $d := d(z, \cup_{j=0}^{p_n-1} J_n) > 0$ . As, for any  $n_k \geq n$ ,  $z_k \in \cup_{j=0}^{p_{n_k}-1} J_{n_k}$  where the latter union is a subset of  $\cup_{j=0}^{p_n-1} J_n$ , the distance between  $z$  and  $z_k$  is at least  $d$ , a contradiction.

(b)  $z_{\pm p_n} \in J_{n,x}$  by combinatorics and definitions of points  $z_m$ . In particular, for every  $k$ ,  $z_k$  and  $w_k$  are in the same component  $f^{j_k}(J_{n_k})$  of  $\cup_{j=0}^{p_{n_k}-1} f^j(J_{n_k})$ . By (a), any limit set  $A$  of the sequence of compacts  $(f^{j_k}(J_{n_k}))$  in the Hausdorff metric is a subset of  $J_\infty$ . On the other hand,  $A$  is connected as each set  $f^{j_k}(J_{n_k})$  is connected. This proves (b).

(c) We prove only the first claim as the second one directly follows from it. Fix any  $\nu \in (0, r)$  and choose  $k_0$  such that for any  $k > k_0$ ,  $B(z_k, \nu) \subset B(z, 11/10\nu)$ . Then, by Proposition 3.1, part (1), for each  $k > k_0$ , the first intersection of the ray  $R_{t(z_k)}$  with the boundary of  $B(z, \nu)$  has the level at least  $\tilde{C}(\nu) := C(11/10\nu)$ . It follows, for any  $0 < C < \tilde{C}(\nu)$ , the sequence of arcs of the rays  $R_{t_{z_k}}$  between the levels  $C$  and  $\tilde{C}(\nu)$  do not exit  $B(z, \nu)$  for all  $k > k_0$ . As  $\nu > 0$  and  $C \in (0, \tilde{C}(\nu))$  can be chosen arbitrary small,  $R_t$  must land at  $z$ .  $\square$

By lemma 4.1(c), if arguments  $t(x), t(x')$  of  $x, x' \in E_\epsilon$  are close then  $x, x'$  are close as well.

**Definition 4.2.** Given  $\epsilon$  and  $\rho$  we define  $\delta$  as follows. First, for  $\hat{r} \in (0, 1)$  and  $\hat{C} > 0$ , we define  $\hat{\delta} = \hat{\delta}(\hat{r}, \hat{C}) > 0$ . Namely, let  $C_0 > 0$



be so that the distance between the equipotential of level  $C_0$  and  $J(f)$  is bigger than 1. Then  $\hat{\delta} = \hat{\delta}(\hat{r}/2, \hat{C}) > 0$  is such that for any  $C \in [\tilde{C}, C_0]$ , if  $w_1, w_2$  lie on the same equipotential  $\Gamma$  of level  $C$  and the difference between external arguments of  $w_1, w_2$  is less than  $\hat{\delta}$  then the length of the shortest arc of the equipotential  $\Gamma$  between  $w_1$  and  $w_2$  is less than  $\hat{r}/2$ . Apply Lemma 4.1(c) with  $\sigma = \rho/4$  and find the corresponding  $\Delta(\rho/4)$ . Let

$$\delta = \delta(\epsilon, \rho) := \min\{\hat{\delta}(\rho, C(\rho/2)), \Delta(\frac{\rho}{4})\}$$

where  $C(\nu)$  is defined in Proposition 3.1.

In the next two lemmas we construct curves with special properties. The idea is as follows. Let  $x \in E_\epsilon \cap J_{n,x}$ . Then  $x_{-p_n} \in J_{n,x}$ . It is easy to get in curve  $\gamma$  in  $A(\infty)$  starting with an arc from a point  $b \in R_{t(x)}$  to  $g_{p_n}(b)$  and then iterating this arc by  $g_{p_n}$  so that  $g^{p_n}(\gamma) \subset \gamma$  so that  $\gamma$  tends to a fixed point  $a$  of  $f^{p_n}$ . We show in the next lemma (in a more general setting) that if both points  $x, x_{-p_n}$  are either in the range of the covering (2.2) (condition (I)) or in the range of the covering (2.3) (condition (II)) then  $a \in J_{n,x}$ . This implies that  $a$  has to be the  $\beta$ -fixed point of  $f^{p_n} : J_{n,x} \rightarrow J_{n,x}$ . In Lemma 4.5 assuming additionally that  $f^{p_n}$  is satellite, we 'rotate' the curve  $\gamma$  by  $g_{p_{n-1}}$  to put  $J_{n,x}$  in a 'sector' bounded by  $\gamma$  and of its 'rotations'. In Lemma 4.7-4.8 we consider the case of doubling for which the condition (II) usually does not hold.

In what follows, we use the following notation: given  $p, q \in \mathbb{N}$ , let

$$E_{\epsilon,p,q} = \cap_{j=0}^{q-1} f^{jp}(E_\epsilon).$$

It is a closed subset of  $E_\epsilon$  of points  $x$  such that  $x_{-jp} \in E_\epsilon$  for  $j = 0, 1, \dots, q-1$ . As  $f : J'_\infty \rightarrow J'_\infty$  is a  $\mu$ -automorphism,  $\mu(E_{\epsilon,p,q}) > 1 - q\epsilon$ . Notice that this bound is independent of  $p$ .

**Lemma 4.3.** *Fix  $\epsilon > 0$  and consider the set  $E_\epsilon$  with the corresponding constant  $r(\epsilon) > 0$ . Fix  $\rho \in (0, r(\epsilon)/3)$ . let  $\delta = \delta(\epsilon, \rho)$  from Definition 4.2. For every  $q \geq 2$  there exist  $\tilde{n}, \tilde{C}$  as follows. For every  $n > \tilde{n}$  consider the closed set  $E_{\epsilon,p_n,q}$ . Let  $x \in E_{\epsilon,p_n,q}$ . Denote for brevity*

$$x^k := x_{-kp_n} \text{ and } R^k := R_{t(x^k)}, \quad k = 0, 1, \dots, q-1.$$

*By Lemma 4.1(b),  $x^k \in J_{n,x}$ . Hence,  $t(x^k) \in s_{n,j_n(x)} \subset S_{n,j_n(x)}$ ,  $0 \leq k \leq q-1$ . Fix  $0 \leq i < j \leq q-1$ .*

Assume that either (I)  $t(x^j)$  and  $t(x^i)$  belong to a single component of  $s_{n,j_n(x)}$ , or (II) the map  $\sigma^{j_n(x)-1} : S_{n,1} \rightarrow S_{n,j_n(x)}$  is a homeomorphism and the length of the arc  $S_{n,j_n(x)}$  is less than  $\delta$ .

Then:

(a) the map  $f^{(j-i)p_n} : g_{(j-i)p_n}(B(x^i, \rho)) \rightarrow (B(x^i, \rho))$  has a unique fixed point  $a = a_n$  and  $a \in J_{n,x}$ ,

(b) there is a semi-open simple curve

$$\gamma_{p_n,q,i,j}(x) \subset B(x^i, \rho) \cap A(\infty)$$

such that:

- (1) it lands at  $a$  and  $g_{(j-i)p_n}(\gamma_{p_n,q,i,j}(x)) \subset \gamma_{p_n,q,i,j}(x)$ . Another end point  $b$  of  $\gamma_{p_n,q,i,j}(x)$  lies in  $R^i$  and  $G(b) > \tilde{C}/2$ ,
- (2)  $\gamma_{p_n,q,i,j}(x) = \cup_{l \geq 0} g_{(j-i)p_n}^l(L_0 \cup L_1)$  where the 'fundamental arc'  $L_0 \cup L_1$  consists of an arc  $L_0$  of an equipotential of the level at least  $\tilde{C}/2$  that joins a point  $b \in R^i$  with a point  $b_1 \in R^j$ , being extended by an arc  $L_1$  of the ray  $R^j$  between points  $b_1$  and  $g_{(j-i)p_n}(b) \in R^j$ ; in particular, the Green function is not increasing along  $\gamma_{p_n,q,i,j}(x)$ ,
- (3) the point  $a$  is the landing point of a ray  $R(a)$  which is fixed by  $f^{(j-i)p_n}$  and which is homotopic to  $\gamma_{p_n,q,i,j}(x)$  through a family of curves in  $A(\infty)$  with the fixed end point  $a$ .
- (4) arguments of all points of the curve  $g_{(j-i)p_n}(\gamma_{p_n,q,i,j}(x))$  lie in a single component of  $s_{n,j_n(x)}^1$  in the case (I) and in a single component of  $s_{n,j_n(x)}$  in the case (II) (recall that  $s_{n,j_n(x)}^1$  has 4 components and  $s_{n,j_n(x)}$  has 2 components, see Sect 2, (C)).

Besides,

$$(4.1) \quad |a - x^j| \rightarrow 0 \text{ and } \log \frac{|(g_{(j-i)p_n})'(x^j)|}{|(g_{(j-i)p_n})'(a)|} \rightarrow 0$$

as  $n \rightarrow \infty$ , uniformly in  $x^j$  and  $q$ .

(c) if  $j - i = 1$  then  $a = \beta_{n,j_n(x)}$  where  $\beta_{n,j_n(x)} = f^{j_n(x)-1}(\beta_n)$ , the non-separating fixed point of  $f^{p_n} : J_{x,n} \rightarrow J_{x,n}$ . Moreover,

$$\chi(\beta_{n,j_n(x)}) := \frac{1}{p_n} \log |(f^{p_n})'(\beta_{n,j_n(x)})| = \frac{1}{p_n} \log |(f^{p_n})'(\beta_n)| \rightarrow \chi(\mu)$$

as  $n \rightarrow \infty$ .

*Remark 4.4.* Note that  $a \notin J_\infty$  while  $x, x^1, \dots, x^{q-1} \in J_\infty$ .

*Proof.* Denote  $G_n := g_{(j-i)p_n}$  which is a holomorphic univalent function in  $B(x^i, \rho)$ . Since  $g_m$  are uniform contractions, there is  $n_1$  such that  $G_n(\overline{B(x^i, \rho)}) \subset B(x^i, \rho/2)$  whenever  $n > n_1$ . Let  $\tilde{n} = \max\{n_0, n_1\}$ .

Let  $a = a_n$  be the unique fixed point of the latter map  $G_n$ . We construct the curve  $\gamma_{p_n, q, i, j}(x)$  to the point  $a$  as follows. First, joint a point  $b \in R^i$ ,  $G(b) = (3/4)\tilde{C}$ , to a point  $b_1 \in R^j$  by an arc  $L_0$  of the equipotential  $\{G(z) = (3/4)\tilde{C}\}$ . By the choice of  $\delta > 0$ ,  $L_0 \subset B(x^i, \rho)$ . Secondly, connect  $b_1$  to the point  $g_{(j-i)p_n}(b) \in R^j$  by an arc  $L_1 \subset R^j$ . Let now  $\gamma_{p_n, q, i, j}(x) = \cup_{l \geq 0} g_{(j-i)p_n}^l(L_0 \cup L_1)$ . Then properties (1), (2) in (b) are immediate and (3) follows from general properties of conformal maps. Now, by Proposition 3.1(2) and (??), for all  $n$  big enough,  $x^j = g_{(j-i)p_n}(x^i) \in g_{(j-i)p_n}(B(x^i, \rho)) \subset B(x^i, \rho)$ , moreover, the modulus of the annulus  $B(x^i, \rho) \setminus g_{(j-i)p_n}(B(x^i, \rho))$  tends to  $\infty$  as  $n \rightarrow \infty$ . Therefore, (4.1) follows from Koebe and Proposition 3.1(2).

It remains to show the property (3) and that  $a \in J_{n, x}$ . Consider the case (II), which is equivalent to say that the map  $\sigma^{p_n} : s \rightarrow S_{n, j_n(x)}$  is a homeomorphism on each of two components  $s$  of  $s_{n, j_n(x)}$ . Let  $\Lambda$  be the set of arguments of points of the curve  $\Gamma := g_{(j-i)p_n}(\gamma_{p_n, q, i, j}(x))$ . Let  $s$  be a component that contains  $t(x^j)$ . Assume, by a contradiction, that  $\Lambda$  contains  $t$  which is in the boundary of  $s$ . Then  $t$  is the argument of a point of  $G_n^l(L_0)$ , for some  $l \geq 1$ , hence,  $\sigma^{l(j-i)p_n}(t)$  is simultaneously the argument of a point of  $L_0$  and in the boundary of  $S_{n, j_n(x)}$ , a contradiction. The case (I) is similar. Property (3) is verified. In fact, we proved more: for  $k = 0, 1, \dots, j-i-1$ , the set  $\sigma^{kp_n}(\Lambda)$  is a subset of a single (depending on  $k$ ) component of  $s_{n, j_n(x)}$  in the case (II) and a single component of  $s_{n, j_n(x)}^1$  in the case (I). This implies that all point  $f^{kp_n}(a)$ ,  $0 \leq k \leq j-i-1$ , of the cycle of  $f^{p_n}$  containing  $a$  belong to the closure of  $U_{n, j_n(x)}$  in the case (II) and to the closure of  $U_{n, j_n(x)-p_n}$  in the case (I). Therefore, this cycle lies in  $J_{n, x}$ , in particular,  $a \in J_{n, x}$ .

Proof of (c): if  $j-i=1$  then  $a$  is a fixed point of  $f^{p_n} : J_{x, n} \rightarrow J_{x, n}$  and, moreover, the ray  $R(a)$  lands at  $a$  and is fixed by  $f^{p_n}$ . Hence, the rotation number of  $a$  w.r.t. the map  $f^{p_n} : J_{x, n} \rightarrow J_{x, n}$  is zero. On the other hand,  $\beta_{n, j_n(x)}$  is the only such a fixed point, i.e.,  $a = \beta_{n, j_n(x)}$  as claimed. Then (4.1) implies that  $\chi(\beta_{n, j_n(x)}) \rightarrow \chi(\mu)$ .  $\square$

**For the rest of the paper**, let us **fix**  $Q, \epsilon, r, \rho, \tilde{n}, \tilde{C}$  and  $\delta$  as follows:

$Q \in \mathbb{N}$ ,  $Q > 3$ , is such that

$$Q > 4 \log 2 / \chi(\mu).$$

This choice is motivated by the following

**Fact** ([21], [13], [8]): if a repelling fixed point  $z$  of  $f^n$  is the landing point of  $q$  rays, then  $\chi(z) := (1/n) \log |Df^n(z)| \leq (2/q) \log 2$ . Hence, if  $\chi(z) > \chi(\mu)/2$ , then  $q < Q$ .

Furthermore, fix  $\epsilon > 0$  such that  $2^{100}Q\epsilon < 1$ , apply Proposition 3.1 and Lemma 4.3 and find, first,  $r = r(\epsilon)$ , then fix  $\rho \in (0, r/32)$  and find the corresponding  $\tilde{n}$ ,  $\tilde{C}$  and  $\delta$ .

Let

$$X_n = E_{\epsilon, p_n, 4} \cap E_{\epsilon, p_{n-1}, Q} = \bigcap_{i=0}^3 f^{ip_n}(E_\epsilon) \cap \bigcap_{k=0}^{Q-1} f^{kp_{n-1}}(E_\epsilon).$$

Let us analyze several possibilities.

**Lemma 4.5.** *There is  $n_* > \tilde{n}$  as follows. Let  $n > n_*$  and  $x \in X_n$ . Consider  $J_{n,x} = f^{j_n(x)}(J_n) \subset f^j(J_{n-1})$  so that  $x \in J_{n,x}$ .*

*Let  $x^0 = x$  and  $x^1 = x_{-p_n}$ . Assume that either (I)  $t(x^0)$ ,  $t(x^1)$  belong to a single component of  $s_{n,j_n(x)}$ , or (II) the map  $\sigma^{j_n(x)-1} : S_{n,1} \rightarrow S_{n,j_n(x)}$  is a homeomorphism and the length of the arc  $S_{n,j_n(x)}$  is less than  $\delta$ .*

*Then:*

(i)  $\chi(\beta_{n,j_n(x)}) = \chi(\beta_n) \rightarrow \chi(\mu)$  as  $n \rightarrow \infty$  and  $\chi(\beta_n) > \chi(\mu)/2$  for  $n > n_*$ .

(ii) assume that  $f^{p_n}$  is satellite, i.e., (by Lemma 2.1)  $\beta_n$  has period  $p_{n-1}$ ,  $q_n \geq 2$  in the rotation number  $k_n/q_n$  of  $\beta_n$ , and  $\beta_{n,j_n(x)}$  is the  $\alpha$  (i.e., separating) fixed point of  $f^{p_{n-1}} : J_{n-1,x} \rightarrow J_{n-1,x}$ . Then  $q_n < Q$  and

(4.2)

$$|\beta_{n,j_n(x)} - x_{-kp_{n-1}}| \rightarrow 0, \quad n \rightarrow \infty, \quad \text{uniformly in } x \in X_n, \quad 1 \leq k \leq q_n.$$

*There exist two simple semi-open curves  $\gamma(x)$  and  $\tilde{\gamma}(x)$  that satisfy the following properties:*

- (1)  $\gamma(x)$  and  $\tilde{\gamma}(x)$  tend to  $\beta_{n,j_n(x)}$  and  $\gamma(x), \tilde{\gamma}(x) \subset B(x^0, \rho) \cap A(\infty)$ ,
- (2)  $\gamma(x), \tilde{\gamma}(x)$  consist of arcs of equipotentials and external rays; the start point  $b_1 = b_1(x)$  of  $\gamma(x)$  lies in an arc of  $R_{t(x^1)}$  and the start point  $\tilde{b}_1 = \tilde{b}_1(x)$  of  $\tilde{\gamma}(x)$  lies in an arc of  $R_{t(\tilde{x})}$  where  $\tilde{x} = x_{-ip_{n-1}}$  for some  $i = i(x) \in \{1, \dots, q_n - 1\}$ , such that levels of  $b_1$  and  $\tilde{b}_1$  are equal and at least  $\tilde{C}/4$ ,
- (3) one of the two curves (say,  $\gamma(x)$ ) is homotopic, through curves in  $A(\infty)$  tending to  $\beta_{n,j_n(x)}$ , to the ray  $R_{t_{n,j_n(x)}} = f^{j_n(x)-1}(R_{t_n})$ , and another one - to the ray  $R_{\tilde{t}_{n,j_n(x)}} = f^{j_n(x)-1}(R_{\tilde{t}_n})$ ;
- (4)  $\gamma(x), \tilde{\gamma}(x) \subset U_{n-1,j_{n-1}(x)}$ ,

- (5)  $\gamma(x) \subset U_{n,j_n(x)}$ ,  $\tilde{\gamma}(x) \subset U_{n,j_n(\tilde{x})}$ , in particular,  $\gamma(x), \tilde{\gamma}(x)$  are disjoint; being completed by their common limit point  $\beta_{n,j_n(x)}$  and two other arcs: an arc of the ray  $R_{t(x^1)}$  from  $b_1 \in \gamma(x)$  to  $\infty$  and an arc of the ray  $R_{t(\tilde{x})}$  from  $\tilde{b}_1 \in \tilde{\gamma}(x)$  to  $\infty$ , they split the plane into two domains such that one of them contains  $I := J_{n,x} \setminus \beta_{n,j_n(x)}$  and another one contains all  $q_n - 1$  other different iterates  $f^{kp_{n-1}}(I)$ ,  $1 \leq k \leq q_n - 1$ . The intersection of closures of all those  $q_n$  sets consists of the fixed point  $\beta_{n,j_n(x)}$  of  $f^{p_{n-1}}$ .

*Remark 4.6.* Beware that the point  $x$  that determines both curves  $\gamma(x), \tilde{\gamma}(x)$  does not belong to either of these curves.

*Proof.* (i) follows from Lemma 4.3 where we take  $i = 0, j = 1$ . Fix  $n_* > \tilde{n}$  such that  $\chi(\beta_n) > \chi(\mu)/2$  for all  $n > n_*$ .

Let us prove (ii). Here we build a "flower" of arcs at the  $\beta$  fixed of the satellite  $f^{p_n}$  starting with an arc which is fixed by  $f^{p_n}$  and then "rotate" this arc by a branch of  $f^{-p_{n-1}}$  (for which the same  $\beta$  point is also a fixed point, see (C)). Let  $\gamma'(x) := \gamma_{p_n,1,0,1}(x)$  where the latter curve is defined in Lemma 4.3. Then properties (1)-(3) of the curve  $\gamma(x)$  are satisfied also for  $\gamma'(x)$ . In particular,  $\gamma'(x)$  is homotopic to  $R_{t_{n,j_n(x)}}$ .

As both  $\tilde{t}_{n,j_n(x)}, t_{n,j_n(x)}$  are external arguments of  $\beta_{n,j_n(x)}$  which is a  $p_{n-1}$ -periodic point of  $f$ , there is  $i \in \{1, \dots, q_n - 1\}$  such that  $\sigma^{ip_{n-1}}(\tilde{t}_{n,j_n(x)}) = t_{n,j_n(x)}$ . Now we use that  $x \in E_{\epsilon, p_{n-1}, Q}$  and that  $q_n < Q$  to prove (4.2). Indeed, for each  $k = \{1, \dots, q_n\}$ , since  $f : J'_\infty \rightarrow J'_\infty$  is a homeomorphism and  $x_{-kp_{n-1}} \in E_\epsilon$ , we have:  $g_{p_n} = g_{(q_n-k)p_{n-1}} \circ g_{kp_{n-1}}$ . Hence, if  $\beta' = g_{kp_{n-1}}(\beta_{n,j_n(x)})$ , then  $\beta_{n,j_n(x)} = g_{(q_n-1-k)p_{n-1}}(\beta')$  implying that  $\beta' = f^{(q_n-k)p_{n-1}}(\beta_{n,j_n(x)}) = \beta_{n,j_n(x)}$ . Then  $\beta_{n,j_n(x)}, x_{-kp_{n-1}} \in g_{kp_{n-1}}(B(x, \rho))$  which along with Proposition 3.1, part (2) imply (4.2).

In turn, (4.2) implies that, provided  $n$  is big,  $g_{kp_{n-1}} : B(y, \rho/2) \rightarrow B(y, \rho/2)$  uniformly in  $k = 0, 1, \dots, q_n$  where  $y$  is either  $\beta_{n,j_n(x)}$  or  $x_{-kp_{n-1}}$ .

Now we consider a curve  $g_{i\tilde{p}_n}(\gamma'(x))$  that starts at  $x_{-i\tilde{p}_n}$  and tends to  $\beta_{n,j_n(x)}$ . By Proposition 3.1 coupled with (4.2), one can join  $x_{-i\tilde{p}_n}$  by an arc of the ray  $R_{t(x_{-i\tilde{p}_n})}$  inside of  $B(x, \rho/2)$  up to a point of level  $\tilde{C}/4$ . This will be the required curve  $\tilde{\gamma}(x)$ . To get the curve  $\gamma(x)$  we modify  $\gamma'(x) = \gamma_{p_n,1,0,1}(x) = \cup_{l \geq 0} g_{p_n}^l(L_0 \cup L_1)$  by cutting off the arc  $L_0$  of an equipotential:  $\gamma(x) = \gamma'(x) \setminus L_0$  (see Lemma 4.3 for details about  $L_0$ ). Properties (1)-(5) follow.  $\square$

Given a point  $x = x^0$  and  $n$  such that  $x \in f^j(J_n) \cap E_{\epsilon, p_n, 1}$ , where  $j = j_n(x)$ , let  $x^1 = x_{-p_n}$  and  $t(x^0), t(x^1)$  the arguments of  $x^0, x^1$  as in Proposition 3.1. We call  $x$   **$n$ -friendly** if  $t(x^0)$  and  $t(x^1)$  lie in the same component of  $s_{n,j}$  and  **$n$ -unfriendly** otherwise (or simply friendly and unfriendly if  $n$  is clear from the context). The name reflects the fact that for an  $n$ -friendly point  $x$  the condition (I) of Lemma 4.5 always holds for  $x^1 = x$  and  $x^2 = x_{-p_n}$ , so Lemma 4.5 always applies.

When the rotation number of  $\alpha_n$  is equal to  $1/2$  we have:

**Lemma 4.7.** *There is  $\tilde{C}_3 > 0$  (depending only on fixed  $\epsilon$  and  $\rho$ ) as follows. Suppose that, for some  $n > \tilde{n}$ , the rotation number of the separating fixed point  $\alpha_n$  is equal to  $1/2$ . Let  $z = z^0 \in f^j(J_n) \cap E_{\epsilon, p_n, 3}$  and  $z^i = z_{-ip_n}$ ,  $i = 1, 2, 3$ . Assume that all three points  $z^0, z^1, z^2$  are  $n$ -unfriendly.*

*Then there exist two (semi-open) curves  $\gamma_n^{1/2}(z)$  and  $\tilde{\gamma}_n^{1/2}(z)$  consisting of arcs of rays and equipotentials with the following properties:*

(i)  $\gamma_n^{1/2}(z) \subset B(z, \rho)$ ,  $\tilde{\gamma}_n^{1/2}(z) \subset B(z^1, \rho)$ , moreover, arguments of points of  $\gamma_n^{1/2}(z)$  lie in one 'window' of  $s_{n,j}$  while arguments of points of  $\tilde{\gamma}_n^{1/2}(z)$  lie in another 'window' of  $s_{n,j}$ ,

(ii)  $\gamma_n^{1/2}(z)$  and  $\tilde{\gamma}_n^{1/2}(z)$  converge to a common point  $\alpha_{n,j}^*$  which is a fixed point of  $f^{p_n} : f^j(J_n) \rightarrow f^j(J_n)$  (i.e.,  $\alpha_{n,j}^*$  is either the non-separating fixed point  $\beta_{n,j}$  or the separating fixed point  $\alpha_{n,j}$ ,

(iii) start points of  $\gamma_n^{1/2}(z), \tilde{\gamma}_n^{1/2}(z)$  have equal Green level which is bigger than  $\tilde{C}_3$ ,

(iv)  $z^k - \alpha_{n,j}^* \rightarrow 0$ ,  $0 \leq k \leq 3$ , as  $n \rightarrow \infty$ .

*Proof.* As  $z \in E_\epsilon$ , lengths of 'windows' of  $s_{n,j_n(z)}$  tend uniformly to zero as  $n \rightarrow \infty$ . It follows from the definition of friendly-unfriendly points that  $t(z^0), t(z^2)$  are in one 'window' of  $s_{n,j}$  and  $t(z^1), t(z^3)$  are in another 'window' of  $s_{n,j}$ . Therefore, condition (I) of Lemma 4.3 holds for each pair  $z^0, z^2$  and  $z^1, z^3$ . Now, apply Lemma 4.3 to  $z \in E_{\epsilon, p_n, 3}$ , first, with  $i = 0, j = 2$ , and then with  $i = 1, j = 3$ . Let  $\gamma_n^{1/2}(z) = \gamma_{p_n, 3, 0, 2}(z)$  and  $\tilde{\gamma}_n^{1/2}(z) = \gamma_{p_n, 3, 1, 3}(z)$ . Then (i), (iii) hold. To check (ii), note that these curves converge to some points  $\alpha, \tilde{\alpha} \in f^j(J_n)$  which are fixed by  $f^{2p_n}$ . On the other hand, since the rotation number of  $\alpha_n$  is  $1/2$ ,  $f^{p_n} : f^j(J_n) \rightarrow f^j(J_n)$  has no 2-cycle. Therefore, one must have either  $\alpha = \tilde{\alpha} = \beta_{n,j}$  or  $\alpha = \tilde{\alpha} = \alpha_{n,j}$ , i.e., (ii) holds too. As  $t(z^0) - t(z^2) \rightarrow 0$  and  $t(z^1) - t(z^3) \rightarrow 0$  as  $n \rightarrow \infty$ ,  $z^0 - z^2, z^1 - z^3 \rightarrow 0$ , too, by Lemma 4.1. Besides, by

(4.1),  $z^2 - \alpha, z^3 - \tilde{\alpha} \rightarrow 0$  as  $n \rightarrow \infty$ . As  $\alpha = \tilde{\alpha} = \alpha_{n,j}^*$ , (iv) also follows.  $\square$

The following is a consequence of Lemmas 4.3 and 4.7:

**Lemma 4.8.** *Let  $n > \tilde{n}$ . Assume that  $f^{p_n}$  is satellite and doubling, i.e.,  $\beta_n = \alpha_{n-1}$  and the rotation number of  $\alpha_{n-1}$  is equal to  $1/2$  (in particular,  $p_n = 2p_{n-1}$ ). For some  $1 \leq j \leq p_{n-1}$ , denote  $J := f^j(J_{n-1})$ . Let  $J^1 := f^j(J_n)$ ,  $J^0 := f^{j+p_{n-1}}(J_n)$  be the two small Julia sets of the next level  $n$  which are contained in  $J$  (note that  $J^0$  contains the critical point and  $J^1$  contains the critical value of the map  $F := f^{p_{n-1}} : J \rightarrow J$ ). Let  $x \in J^1 \cap E_\epsilon$  be such that all its 5 forward iterates  $x_{kp_{n-1}} = F^k(x) \in E_\epsilon$ ,  $k = 1, 2, 3, 4, 5$ . Then there exist two simple semi-open curves  $\Gamma_n^{1/2}(x)$ ,  $\tilde{\Gamma}_n^{1/2}(x)$  consisting of arcs of rays and equipotentials that satisfy essentially conclusions of the previous lemma where  $n$  is replaced by  $n-1$ , i.e.:*

(i)  $\Gamma_n^{1/2}(x), \tilde{\Gamma}_n^{1/2}(x) \subset B(x, 3/2\rho)$ , moreover, arguments of points of  $\Gamma_n^{1/2}(x)$  lie in one 'window' of  $s_{n-1, j_{n-1}(x)}$  while arguments of points of  $\tilde{\Gamma}_n^{1/2}(x)$  lie in another 'window' of  $s_{n-1, j_{n-1}(x)}$ ,

(ii)  $\Gamma_n^{1/2}(x)$  and  $\tilde{\Gamma}_n^{1/2}(x)$  converge to a common point  $\beta_{n-1, j_{n-1}(x)}^*$  which is a fixed point of  $f^{p_{n-1}} : f^j(J_{n-1}) \rightarrow f^j(J_{n-1})$  (i.e.,  $\beta_{n-1, j_{n-1}(x)}^*$  is either the non-separating fixed point  $\beta_{n-1, j_{n-1}(x)}$  or the separating fixed point  $\alpha_{n-1, j_{n-1}(x)}$ ,

(iii) start points of  $\Gamma_n^{1/2}(x), \tilde{\Gamma}_n^{1/2}(x)$  have equal Green level which is bigger than  $\tilde{C}_3$ ,

(iv)  $x_{kp_{n-1}} - \beta_{n-1, j_{n-1}(x)}^* \rightarrow 0$ ,  $0 \leq k \leq 3$  as  $n \rightarrow \infty$  uniformly in  $x$ .

*Remark 4.9.* Condition  $F^k(x) \in E_\epsilon$ ,  $0 \leq k \leq 5$ , is equivalent to the following:  $x \in f^{-5p_{n-1}}(E_{\epsilon, p_{n-1}, 6})$ .

*Proof.* To fix the idea let's replace  $f^{p_{n-1}} : f^j(J_{n-1}) \rightarrow f^j(J_{n-1})$ , using a conjugacy with a quadratic polynomial, by a quadratic polynomial (denoted also by  $F$ ) so that now  $F : J \rightarrow J$  where  $J = J(F)$  and  $F^2$  is satellite with two small Julia sets  $J^0, J^1$  that meet at the  $\alpha$ -fixed point of  $F$  and rays of arguments  $1/3, 2/3$  land at  $\alpha$ . Here  $0 \in J^0$ ,  $F(0) \in J^1$ ,  $F : J^1 \rightarrow J^0$  is a homeomorphism while  $F : J^0 \rightarrow J^1$  is a two-to-one map. If a ray  $R_t$  of  $F$  has its accumulation set in  $J^1$  then  $t \in [1/3, 5/12] \cup [7/12, 2/3]$  and if  $R_t$  accumulates in  $J^0$  then  $t \in [1/6, 1/3] \cup [2/3, 5/6]$ . This implies that if  $R_t$  lands at  $x \in J^1$  and  $t$  lies in one of the two 'windows'  $[0, 1/2), (1/2, 1]$  then  $R_{\sigma(t)}$  lands at  $J^0$  where  $\sigma(t)$  must be in a different

'window' (in other words, points of  $J^0$  are 'unfriendly'). Coming back to  $f^{p_{n-1}}$  this means that, for  $x \in J^1$ ,  $t(x), t(F(x))$  are always in different components (where by 'component' we mean a component of  $s_{n-1,j}$ ). Besides, for  $y \in J_\infty \cap J$ ,  $y$  and  $F(y)$  are always in different  $J^i$ ,  $i = 0, 1$ . This leaves us with the only possibilities:

(i)  $t(F(x)), t(F^2(x))$  are in different components; this implies that  $t(x), t(F(x))$  are in different components and  $t(F(x)), t(F^2(x))$  are in different components, that is, points  $F^3(x), F^2(x), F(x)$  are all unfriendly;

(ii)  $t(F(x)), t(F^2(x))$  are in the same components; there are two subcases:

(ii')  $t(F^3(x)), t(F^4(x))$  are in different components, i.e., (i) holds with  $x$  replaced by  $F^2(x)$  which implies that  $F^5(x), F^4(x), F^3(x)$  are all unfriendly;

(ii'')  $t(F^3(x)), t(F^4(x))$  are in the same component which then means that  $F^2(x)$  and  $F^4(x)$  are both friendly.

In the case (i) and (ii'), apply Lemma 4.7 with  $n - 1$  instead of  $n$  to  $z = F^3(x)$  and to  $z = F^5(x)$ , respectively, letting  $\Gamma_n^{1/2}(x) = \gamma_{n-1}^{1/2}(F^3(x))$ ,  $\tilde{\Gamma}_n^{1/2}(x) = \tilde{\gamma}_{n-1}^{1/2}(F^3(x))$  and  $\Gamma_n^{1/2}(x) = \gamma_{n-1}^{1/2}(F^5(x))$ ,  $\tilde{\Gamma}_n^{1/2}(x) = \tilde{\gamma}_{n-1}^{1/2}(F^5(x))$ , respectively. In the case (ii''), apply Lemma 4.3 with  $p_{n-1}, q = 1, i = 0, j = 0$ , first, to the point  $F^2(x)$  and then to the point  $F^4(x)$  letting  $\Gamma_n^{1/2}(x) = \gamma_{p_{n-1},1,0,1}(F^2(x))$ ,  $\tilde{\Gamma}_n^{1/2}(x) = \gamma_{p_{n-1},1,0,1}(F^4(x))$ .  $\square$

## 5. PROOF OF THEOREM 1.1

Every invariant probability measure with positive Lyapunov exponent has an ergodic component with positive exponent. So let  $\mu$  be such an ergodic  $f$ -invariant measure component supported in  $J_\infty$ . First, we have the following general

*Remark 5.1.* Given  $x \in J'_\infty$  such that  $\tilde{r}(x) > 0$  as in Proposition 2.3, and given  $n$ , the set  $J_{n,x} = f^{j_n(x)}(J_n)$  cannot be covered by  $B(x, \tilde{r}(x))$  because otherwise the branch  $g_{p_n} : B(x, \tilde{r}(x)) \rightarrow \mathbf{C}$  of  $f^{-p_n}$ , which sends  $x$  to  $x_{-p_n} \in J_{n,x}$  meets the critical value along the way so cannot be well-defined. Thus  $\text{diam } J_{n,x} > \tilde{r}(x)$ , for each  $n$ , and  $\text{diam } K_x = \lim \text{diam } J_{n,x} \geq \tilde{r}(x)$ . In particular,  $\text{diam } J_{n,x} \geq r(\epsilon)$  for all  $x \in E_\epsilon$  and  $n$ .

We need to prove that  $f$  has finitely many satellite renormalizations. Assuming the contrary, let  $\mathcal{S}$  be an infinite subsequence such that  $f^{p_n}$  is a satellite renormalization of  $f$  for each  $n \in \mathcal{S}$ .



We arrive at a contradiction by considering, roughly speaking, two alternative situations. In the first one, we find a point  $x \in E_\epsilon$ ,  $n$ , and two curves in  $B \cap A(\infty)$  where  $B := B(x, \tilde{r}(x))$  that tend to the  $\beta$ -fixed points of  $J_{n,x}$  such that another ends of the curves can be joined by an arc of equipotential in  $B$  thus 'surrounding'  $J_{n,x}$  by a 'triangle' in  $B$  which would be a contradiction as in Remark 5.1. The second situation is when the first one does not happen. Then we use several curves to 'surround'  $J_{n,x}$  by a 'quadrilateral' in  $B$ , ending by the same conclusion. The curves we use have been constructed in Lemmas 4.5, 4.8.

The first situation happens in cases A and B1, and the second one in B2.

**Case A:**  $\mathcal{S}$  contains an infinite sequence of indices of non-doubling renormalizations. Passing to a subsequence one can assume that  $f^{p_n}$  is satellite not doubling for every  $n \in \mathcal{S}$ .

Fix  $\zeta = 1/4$ . By Lemma 2.2, for each  $n \in \mathcal{S}$  and each  $j = 1, \dots, [\zeta p_n]$ , the map  $\sigma^{j-1} : S_{n,1} \rightarrow S_{n,j}$  is a homeomorphism and the length  $|S_{n,j}| \rightarrow 0$  as  $n \rightarrow \infty$  uniformly in  $j$ . Fix  $N$  such that  $|S_{n,j}| < \delta$  for each  $n > N$ ,  $n \in \mathcal{S}$ . For  $n \in \mathcal{S}$ , let

$$\mathcal{C}_n = \{f^j(J_n) | 1 \leq j \leq [\zeta p_n]\}.$$

Let  $n, m \in \mathcal{S}$ ,  $m < n$ . Denote  $p = p_m$ ,  $P = p_n$ ,  $q = p_n/p_m$ . The intersection  $\mathcal{C}_n \cap \mathcal{C}_m$  contains all  $f^{j+kp}(J_n)$  with  $1 \leq j \leq [\zeta p]$ ,  $j + kp \leq [\zeta P]$ . Hence,

$$\begin{aligned} \#(\mathcal{C}_n \cap \mathcal{C}_m) &\geq \sum_{j=1}^{[\zeta p]} [\zeta q - \frac{j}{p}] \geq [\zeta q - 1][\zeta p] \geq \\ &P \left\{ \frac{\zeta p - 1}{p} \frac{\zeta q - 1}{q} - \frac{\zeta}{q} \right\} \sim \zeta^2 P \end{aligned}$$

as  $p, q \rightarrow \infty$ . Therefore, fixing  $\kappa = \zeta^2/2 = 1/8$ , there are  $m_0, k_0$  such that for each  $n, m \in \mathcal{S}$ ,  $m > m_0$ ,  $n > m + k_0$ ,

$$\mu(\mathcal{C}_n \cap \mathcal{C}_m) > \kappa.$$

Fix such  $n, m$ , assume also that  $m > \max\{N, n_*\}$  where  $n_*$  is defined in Lemma 4.5 and recall the set

$$X_n = E_{\epsilon, p_n, 4} \cap E_{\epsilon, p_{n-1}, Q} = \bigcap_{i=0}^3 f^{ip_n}(E_\epsilon) \cap \bigcap_{k=0}^{Q-1} f^{kp_{n-1}}(E_\epsilon).$$

Since  $\mu(X_n) > 1 - (Q+4)\epsilon > 1 - \kappa$ , there is  $x \in X_n \cap \mathcal{C}_n \cap \mathcal{C}_m$  and, by the choice of  $n$ , the assumption (II) of Lemma 4.5 holds for  $x$ . Therefore, there exist two simple semi-open curves  $\gamma(x)$  and  $\tilde{\gamma}(x)$  that satisfy the following properties:  $\gamma(x)$  and  $\tilde{\gamma}(x)$  tend to  $\beta_{n, j_n(x)}$ ,  $\gamma(x), \tilde{\gamma}(x) \subset B(x, \rho) \cap A(\infty)$  and  $\gamma(x), \tilde{\gamma}(x)$  consist of arcs

of equipotentials and external rays; the start point  $b_1$  of  $\gamma(x)$  and the start point  $\tilde{b}_1$  of  $\tilde{\gamma}(x)$  have equal levels which is at least  $\tilde{C}/4$ ;  $\gamma(x), \tilde{\gamma}(x) \subset U_{n-1, j_{n-1}(x)}$ ; finally, being completed by their common limit point  $\beta_{n, j_n(x)}$  and arcs of rays from  $b_1 \in \gamma(x)$  to  $\infty$  and from  $\tilde{b}_1 \in \tilde{\gamma}(x)$  to  $\infty$ , they split the plane into two domains such that one of them contains  $I := J_{n, x} \setminus \beta_{n, j_n(x)}$  and another one contains all other iterates  $f^{kp_{n-1}}(I)$ ,  $1 \leq k \leq q_n - 1$ . Now, since  $U_{n-1, j_{n-1}(x)} \subset U_{m, j_m(x)}$  and by the choice of  $m$ , the distance between arguments of the points  $b_1$  and  $\tilde{b}_1$  inside of  $S_{n-1, j_{n-1}(x)}$  is less than  $\delta$ . By the definition of  $\delta$ ,  $b_1$  and  $\tilde{b}_1$  can be joined by an arc  $A_n$  of equipotential inside of  $B(x, \rho) \cap U_{n-1, j_{n-1}(x)}$ . Consider a Jordan domain  $Z_n$  with the boundary to be the arc  $A_n$  and semi-open curves  $\gamma(x), \tilde{\gamma}(x)$  completed by their common limit point  $\beta_{n, j_n(x)}$ . Then  $Z_n \subset B(x, \rho)$ . By the properties of the curves,  $Z_n \cup \beta_{n, j_n(x)}$  contains either  $J_{n, x}$  or its iterate  $f^{kp_{n-1}}(J_{n, x})$ , for some  $1 \leq k \leq q_n - 1$ , in a contradiction with Remark 5.1.

Complementary to A is

**Case B:** for all big  $n$ , every satellite renormalization  $f^{p_n}$  is doubling, i.e.,  $\beta_n = \alpha_{n-1}$  and  $p_n = 2p_{n-1}$  for every  $n \in \mathcal{S}$ .

Let  $Y_{n-1} = E_{\epsilon, p_{n-1}, 6}$  and  $\tilde{Y}_{n-1} = f^{-5p_{n-1}}(Y_{n-1})$ . Note that  $\mu(Y_{n-1}) = \mu(\tilde{Y}_{n-1}) > 1 - 6\epsilon$ .

For every  $n \in \mathcal{S}$ , let

$$L_n = \left\{ 0 < j < p_{n-1} \mid \mu(f^j(J_{n-1}) \cap \tilde{Y}_{n-1}) > \frac{1 - 2^{12}\epsilon}{p_{n-1}} \right\}.$$

As  $\mu(\tilde{Y}_{n-1}) > 1 - 6\epsilon$ , it follows,

$$\#L_n > (1 - 3/2^{11})p_{n-1}.$$

Since we are in case B, each  $f^j(J_{n-1})$  contains precisely two small Julia sets  $f^j(J_n), f^{j+p_{n-1}}(J_n)$  of the next level  $n$  each of them of measure  $1/(2p_{n-1})$ . Hence, the measure of intersection of each of these small Julia sets with  $\tilde{Y}_{n-1}$  is bigger than  $(1/2 - 2^{10}\epsilon)/p_{n-1} > 0$ . By Lemma 4.8, choosing for every  $j \in L_n$  a point  $x_j \in f^j(J_{n-1}) \cap \tilde{Y}_{n-1}$  we get a pair of curves  $\Gamma_n^{1/2}(x_j), \tilde{\Gamma}_n^{1/2}(x_j)$  consisting of arcs of rays and equipotentials as follows: (i)  $\Gamma_n^{1/2}(x_j), \tilde{\Gamma}_n^{1/2}(x_j) \subset B(x_j, 3/2\rho)$ , moreover, arguments of points of  $\Gamma_n^{1/2}(x_j)$  lie in one 'window' of  $s_{n-1, j}$  while arguments of points of  $\tilde{\Gamma}_n^{1/2}(x_j)$  lie in another 'window' of  $s_{n-1, j}$ , (ii)  $\Gamma_n^{1/2}(x_j)$  and  $\tilde{\Gamma}_n^{1/2}(x_j)$  converge to a common point  $\beta_{n-1, j}^*$  which is a fixed point of  $f^{p_{n-1}} : f^j(J_{n-1}) \rightarrow f^j(J_{n-1})$  (i.e.,

$\beta_{n-1,j}^*$  is either the non-separating fixed point  $\beta_{n-1,j}$  or the separating fixed point  $\alpha_{n-1,j}$ , (iii) start points of  $\Gamma_n^{1/2}(x_j), \tilde{\Gamma}_n^{1/2}(x_j)$  have equal Green level which is bigger than  $\tilde{C}_3$ , (iv)  $x_j - \beta_{n-1,j}^* \rightarrow 0$  as  $n \rightarrow \infty$  uniformly in  $j$  and  $x_j$ . We add one more property as follows. Let

$$\Gamma_{n,j} = \Gamma_n^{1/2}(x_j) \cup \beta_{n-1,j}^* \cup \tilde{\Gamma}_n^{1/2}(x_j).$$

Then: (v)  $\Gamma_{n,j}$  is a simple curve; the level of  $z \in \Gamma_{n,j} \setminus \{\beta_{n-1,j}^*\}$  is positive and decreases (not strickly) from  $\tilde{C}_3$  to zero along  $\Gamma_n^{1/2}(x_j)$  and then increases from zero to  $\tilde{C}_3$  along  $\tilde{\Gamma}_n^{1/2}(x_j)$ ; moreover, if  $j_1, j_2 \in L_n$ ,  $j_1 \neq j_2$ , then  $\Gamma_{n,j_1}, \Gamma_{n,j_2}$  are either disjoint or meet at the unique common point  $\beta_{n-1,j_1} = \beta_{n-1,j_2}$  and then disjoint with all others  $\gamma_{n-1,j}$ ,  $j \neq j_1, j_2$ . This is because, by property (i),  $\Gamma_{n,j} \subset \overline{U_{n-1,j}}$  where (by (C), Sect 2) any two  $\overline{U_{n-1,j}}, \overline{U_{n-1,\tilde{j}}}$ ,  $j \neq \tilde{j}$ , are either disjoint or meet at  $\beta := \beta_{n-1,j} = \beta_{n-1,\tilde{j}}$  in which case  $f^{p_{n-1}}$  is satellite. In the considered case, any satellite is doubling so  $\beta \neq \beta_{n-1,i}$  for all  $i$  different from  $j, \tilde{j}$ .

We assign, for the use below, a 'small' Julia set  $I_{n,j}$  to each  $\Gamma_{n,j}$  as follows: by the construction,  $\beta_{n-1,j}^*$  is either the  $\beta$ -fixed point of  $f^j(J_{n-1})$ , or the  $\alpha$ -fixed point of  $f^j(J_{n-1})$ . In the former case, let  $I_{n,j} = f^j(J_{n-1})$ , and in the latter case,  $I_{n,j} = f^j(J_n)$  (one of the two small Julai sets of the next level  $n$  that are contained in  $f^j(J_{n-1})$ ). Observe that  $I_{n,j} \cap \Gamma_{n-1,j} = \{\beta_{n-1,j}^*\}$  and is disjoint with any other  $\Gamma_{n,j'}$  provided  $\Gamma_{n,j}, \Gamma_{n,j'}$  are disjoint.

There are two subcases B1-B2 to distinguish depending on whether arguments of end points of  $\Gamma_{m,j}$  become close or not. If yes, then one can join the end points of some  $\Gamma_{n,j}$  by an arc of equipotential inside of  $B(x_j, 2\rho) \supset \Gamma_{m,j}$  to surround a small Julia set as in case A, which would lead to a contradiction. If no, the construction is more subtle: we build a domain ('quadrilateral') in  $B(x_j, 2\rho)$  bounded by two disjoint curves as above completed by two arcs of equipotential that join ends of different curve, so that the obtained quadrilateral again contains a small Julia set.

**B1:**  $\liminf_{n \in \mathcal{S}, j \in L_n} |S_{n-1,j}| < \delta$ .

By property (i) listed above and the definition of  $\delta$ , there are a sequence  $(n_k) \subset \mathcal{S}$ ,  $j_k \in L_{n_k}$  and  $x_{j_k}$  as above, such that two ends of each curve  $\Gamma_{n_k, j_k}$  can be joined inside of  $B(x_{j_k}, \rho)$  by an arc  $A^k$  of equipotential of fixed level  $\tilde{C}_3$  such that all arguments of points in  $A^k$  belong to  $S_{n_k-1, j_k}$ . Then we arrive at a contradiction as in case A.

**B2:**  $|S_{n-1,j}| \geq \delta$  for all big  $n \in \mathcal{S}$  and all  $j \in L_n$ .




FIGURE 1. Top: Case A and Case B1, bottom: Case B2

Fix  $n, m \in \mathcal{S}$ ,  $m - n \geq 3$ . Define a subset of  $L_n$  as follows:

$$L_n^m = \left\{ 0 < j < p_{n-1} \mid \mu(f^j(J_{n-1}) \cap (\tilde{Y}_{n-1} \cap \tilde{Y}_{m-1})) > \frac{1 - 2^{12}\epsilon}{p_{n-1}} \right\}.$$

As  $\mu(\tilde{Y}_{n-1} \cap \tilde{Y}_{m-1}) > 1 - 12\epsilon$ ,

$$\#L_n^m > (1 - 3/2^{10})p_{n-1}.$$

For each  $j \in L_n^m$  we define further

$$L_{n,j}^m = \left\{ 0 < k < p_{n-1} \mid f^k(J_{m-1}) \subset f^j(J_{n-1}), \mu(f^k(J_{m-1}) \cap (\tilde{Y}_{n-1} \cap \tilde{Y}_{m-1})) > \frac{1 - 2^{16}\epsilon}{p_{m-1}} \right\}.$$

Then

$$\#L_{n,j}^m \geq 5$$

as otherwise  $\#L_{n,j}^m \leq 4$  and, therefore,  $(1 - 2^{12}\epsilon)/p_{n-1} < 4/p_{m-1} + (p_{m-1}/p_{n-1} - 4)(1 - 2^{16}\epsilon)/p_{m-1} = 2^{18}\epsilon/p_{m-1} + (1 - 2^{16}\epsilon)/p_{n-1}$ , i.e.,  $p_{m-1}/p_{n-1} < 2^{18}\epsilon/(2^{16}\epsilon - 2^{12}\epsilon) = 4/(1 - 2^{-4}) < 8$ , a contradiction because  $p_{m-1}/p_{n-1} \geq 2^{m-n} \geq 2^3$ .

Fix  $j \in L_n^m$ . Thus  $L_{n,j}^m$  contains 5 pairwise different indices  $k_i$ ,  $1 \leq k \leq 5$ . As  $L_{n,j}^m \subset L_m$ , we find 5 curves  $\Gamma_{m-1,k_i}$ . By property (v), if two of them meet, they are disjoint with all others. Therefore, there are at least 3 of them denoted by  $\Gamma_{m-1,r_i}$ ,  $i = 1, 2, 3$ , which are pairwise disjoint. Let  $w_i, \tilde{w}_{m,i}$  be two ends of  $\Gamma_{m-1,r_i}$ .

For each  $i = 1, 2, 3$ , arguments of points of  $w_{m,i}, \tilde{w}_{m,i}$  lie in different 'windows' of  $s_{m-1,r_i}$ . On the other hand, by the choice of

$j$ ,  $s_{m-1,r_i} \subset s_{n-1,j} \subset S_{n-1,j}$ . As  $n$  is big enough, lengths of 'windows' of  $s_{n-1,j}$  are less than  $\delta$ . But since we are in case B2, the length of  $S_{n-1,j}$  is bigger than  $\delta$ . One can assume, therefore, that, for  $i = 1, 2, 3$ , arguments of  $w_{m,i}$  lie in one window of  $s_{n-1,j}$  while arguments of  $\tilde{w}_{m,i}$  are in another window. Therefore, differences of arguments of all  $w_{m,i}$  tend to zero as  $m \rightarrow \infty$ , and the same for  $\tilde{w}_{m,i}$ . As all  $w_{m,i}, \tilde{w}_{m,i} \in E_\epsilon$ , this implies by Lemma 4.1 that  $\max_{1 \leq i, l \leq 3} |w_{m,i} - w_{m,l}| \rightarrow 0$ . This along with property (iv) implies that  $\gamma_{m-1,r_i} \subset B(w_{m,1}, 2\rho)$ ,  $i = 1, 2, 3$ , for all big  $m$ . Since, for big  $m$ , differences of arguments of all  $w_{m,i}$  are less than  $\delta$ , and the same for  $\tilde{w}_{m,i}$ , one can joint all  $w_{m,i}$  by an arc  $D^m$  of equipotential of level  $\tilde{C}_3$  and all  $\tilde{w}_{m,i}$  by an arc  $\tilde{D}^m$  of equipotential of the same level  $\tilde{C}_3$  such that  $D^m, \tilde{D}^m \subset B(w_1, 2\rho)$ . Let the end points of  $D^m$  be, say,  $w_{m,1}$  and  $w_{m,3}$ , so that  $w_{m,2} \in D^m$  is in between. Since all 3 curves  $\Gamma_{m-1,r_i}$ ,  $i = 1, 2, 3$ , are pairwise disjoint, the end points of  $\tilde{D}^m$  have to be then  $\tilde{w}_{m,1}$  and  $\tilde{w}_{m,3}$ , so that  $\tilde{w}_{m,2} \in \tilde{D}^m$  is in between. Therefore, we get a 'big' quadrilateral  $\Pi_m^0 \subset B(w_{m,1}, 2\rho)$  bounded by  $D^m, \tilde{D}^m, \Gamma_{m,1}, \tilde{\Gamma}_{m,3}$ . The curve  $\Gamma_{m,2}$  splits  $\Pi_m$  into two 'small' quadrilaterals  $\Pi_m^1, \Pi_m^2$  with a common curve  $\Gamma_{m,2}$  in their boundaries. Recall now that the curve  $\Gamma_{m,2}$  comes with a small Julia set  $I_{m,2}$  of level either  $m - 1$  or  $m$ , such that  $I_{m,2} \cap \Gamma_{m,2}$  is a single point while  $I_{m,2}$  is disjoint with  $\Gamma_{m,1}, \Gamma_{m,3}$ . Therefore,  $I_{m,2} \subset \Pi_m^0 \subset B(w_{m,1}, 2\rho)$ , a contradiction with Remark 5.1.

## 6. PROOF OF COROLLARIES 1.1-1.2

Corollary 1.1 follows directly from the following

**Proposition 6.1.** *Let  $f$  be an infinitely renormalizable quadratic polynomial. Then conditions (1)-(4) are equivalent:*

- (1)  $f : J_\infty \rightarrow J_\infty$  has no invariant probability measure with positive exponent,
- (2) for every neighborhood  $W$  of  $P$  and every  $\alpha \in (0, 1)$  there exist  $m_0$  and  $n_0$  such that, for each  $m \geq m_0$  and  $x \in \text{orb}(J_n)$  with  $n \geq n_0$ ,

$$\frac{\#\{i | 0 \leq i < m, f^i(x) \in W\}}{m} > \alpha;$$

additionally,  $f : P \rightarrow P$  has no invariant probability measure with positive exponent,

- (3) every invariant probability measure of  $f : J_\infty \rightarrow J_\infty$  is, in fact, supported on  $P$  and has zero exponent,

- (4) for every invariant probability ergodic measure  $\mu$  of  $f$  on the Julia set  $J$  of  $f$ , either  $\text{supp}(\mu) \cap J_\infty = \emptyset$  and its Lyapunov exponent  $\chi(\mu) > 0$ , or  $\text{supp}(\mu) \subset P$  and  $\chi(\mu) = 0$ .

*Proof.* (1) $\Rightarrow$ (2). Assume the contrary. Let  $E = \mathbb{C} \setminus W$ . Since  $W$  is a neighborhood of a compact set  $P$ , the Euclidean distance  $d(E, P) > 0$ . By a standard normality argument, as all periodic points of  $f$  are repelling, there are  $\lambda > 1$  and  $k_0 > 0$  such that  $|(f^k)'(y)| > \lambda$  whenever  $y, f^k(y) \in E$  and  $k \geq k_0$ . As (2) does not hold, find  $\alpha \in (0, 1)$ , a sequence  $n_k \rightarrow \infty$ , points  $x_k \in \text{orb}(J_{n_k})$  and a sequence  $m_k \rightarrow \infty$  such that, for each  $k$ ,

$$\frac{\#\{i : 0 \leq i < m_k, f^i(x_k) \in E\}}{m_k} \geq \beta := 1 - \alpha.$$

Fix a big  $k$  such that  $\beta m_k > 3k_0$  and consider the times  $0 \leq i_1^k < i_2^k < \dots < i_{l_k}^k < m_k$  where  $l_k/m_k \geq \beta$  such that  $f^i(x_k) \in E$ . Let  $z_k = f^{i_1^k}(x_k)$  so that  $z_k \in E \cap \text{orb}(J_n)$ . Therefore, by the choice of  $\lambda$  and  $k_0$ ,  $|(f^{m_k - i_1^k})'(z_k)| \geq \tilde{\lambda}^{m_k} \geq \tilde{\lambda}^{m_k - i_1}$  where  $\tilde{\lambda} = \lambda^{\frac{\beta}{2k_0}} > 1$ . In this way we get a sequence of measures  $\mu_k = \frac{1}{m_k - i_1^k} \sum_{i=0}^{m_k - i_1^k - 1} \delta_{f^i(z_k)}$  such that the Lyapunov exponent of  $\mu_k$  is at least  $\log \tilde{\lambda} > 0$ . Passing to a subsequence one can assume that  $\{\mu_k\}$  converges weak-\* to a measure  $\mu$ . Then  $\mu$  is an  $f$ -invariant probability measure on  $J_\infty = \bigcap \text{orb}(J_n)$  with the exponent at least  $\log \tilde{\lambda} > 0$ , a contradiction with (1).

(2) $\Rightarrow$ (3), by the Birkhoff Ergodic Theorem along with [22].

(3) $\Rightarrow$ (4): let  $\mu$  be as in (4) and  $\bar{U} \cap P = \emptyset$  for some open set  $U$  with  $\mu(U) > 0$ . Let  $F : U \rightarrow U$  be the first return map equipped with the induced invariant measure  $\mu_U$ . By the Birkhoff Ergodic Theorem and by an argument as in (1) $\Rightarrow$ (2), the exponent  $\chi_F(\mu_U)$  of  $F$  w.r.t.  $\mu_U$  is strictly positive. Hence,  $\chi(\mu) = \mu(U)\chi_F(\mu_U)$  is positive too. This proves the implication.

And (4) obviously implies (1).  $\square$

*Proof of Corollary 1.2.* If  $\bar{\chi}(x)$  were strictly positive, for some  $x \in J_\infty$ , that would imply, by a standard argument (see the proof of Corollary 1.1), the existence of an  $f$ -invariant measure with positive exponent supported in  $\omega(x) \subset J_\infty$ , with a contradiction to Theorem 1.1. This proves (1.1). By [14],  $\liminf_{n \rightarrow \infty} \frac{1}{n} \log |(f^n)'(c)| \geq 0$ . On the other hand, by (1.1),  $\bar{\chi}(c) \leq 0$ , which proves (1.2).  $\square$

## REFERENCES

- [1] A. Blokh, G. Levin, L. Oversteegen, V. Timorin: *The modulus of renormalization of rational functions: high periods imply small moduli.*, 2022, <https://arxiv.org/pdf/2205.03157.pdf>
- [2] L. Carleson, T. Gamelin: *Complex Dynamics*. Springer-Verlag, 1993
- [3] D. Cheraghi, M. Pedramfar, slides of talk at workshop "On geometric complexity of Julia sets", 18-23 March 2018, Będlewo, Poland, and video of talk at virtual workshop "Many faces of renormalization", Simons Center for Geometry and Physics, 8-12 March 2021
- [4] D. Cheraghi, M. Shishikura: *Satellite renormalization of quadratic polynomials*. <https://arxiv.org/abs/1509.07843>
- [5] A. Douady, J. H. Hubbard: *On the dynamics of polynomial-like mappings*. Ann. Sci. Ec. Norm. Super. (4) 18 (1985), no. 2, 287-343.
- [6] M. Feigenbaum: *Quantitative universality for a class of nonlinear transformations*. Journal of Statistical Physics. 19 (1) (1978), 25-52.
- [7] J. Hu, Y. Jiang: *The Julia set of the Feigenbaum quadratic polynomial is locally connected*. Preprint, 1993.
- [8] J. H. Hubbard: *Local connectivity of Julia sets and bifurcation coci: three theorems of J.-C. Yoccoz*. In: "Topological Methods in Modern Mathematics." Houston, TX:Publish or Perish, 1993.
- [9] G. Levin, F. Przytycki: *No hyperbolic sets in  $J_\infty$  for infinitely renormalizable quadratic polynomials*. Israel Journal of Mathematics, to appear.
- [10] G. Levin: *Multipliers of periodic orbits of quadratic polynomials and the parameter plane*. Israel Journal of Mathematics 170 (2009), 285-315.
- [11] G. Levin: *Rigidity and non-local connectivity of Julia sets of some quadratic polynomials*. Comm. Math. Phys. 304 (2011), 2, 295-328.
- [12] G. Levin: *Addendum to: Rigidity and non-local connectivity of Julia sets of some quadratic polynomials*. Comm. Math. Phys. 325 (2014), 3, 1171-1178.
- [13] G. M. Levin: *Bounds for multipliers of periodic points of holomorphic mappings*. (Russian) Sibirsk. Mat. Zh. 31 (1990), no. 2, 104-110; translation in Siberian Math. J. 31 (1990), no. 2, 273-278.
- [14] G. Levin, F. Przytycki, W. Shen: *The Lyapunov exponent of holomorphic maps*. Invent. Math. 205 (2016), 363-382.
- [15] M. Lyubich: *Feigenbaum-Couillet-Tresser universality and Milnor's Hairiness Conjecture*. Ann. Math. 149 (2) (1999), 319-420
- [16] J. Milnor: *Dynamics in one complex variable*. Princeton University Press, 2006
- [17] J. Milnor: *Local connectivity of Julia sets: expository lectures*. The Mandelbrot set, theme and variations, 67-116, London Math. Soc. Lecture Note Ser., 274, Cambridge Univ. Press, Cambridge, 2000.
- [18] J. Milnor: *Periodic orbits, external rays and the Mandelbrot set: an expository account*. "Geometrie Complex et Systemes Dynamiques". Colloque en l'honneur d'Adrien Douady (Orsay, 1995). Asterisque 261, 277-331 (2000).
- [19] C. McMullen: *Complex Dynamics and Renormalization*. Annals of Mathematical Studies, 135. Princeton Univ. Press, Princeton, New Jersey, 1994.

- [20] C. McMullen: *Renormalization and 3-Manifolds Which Fiber over the Circle*. Annals of Mathematical Studies, 142. Princeton Univ. Press, Princeton, New Jersey, 1996.
- [21] Ch. Pommerenke: *On conformal mapping and iteration of rational functions*. Complex Variables 5 (2-4) (1986), 117-126.
- [22] F. Przytycki: *Lyapunov characteristic exponents are nonnegative*. Proc. Amer. Math. Soc. 119 (1993), no. 1, 309-317.
- [23] F. Przytycki: *Accessibility of typical points for invariant measures of positive Lyapunov exponents for iterations of holomorphic maps*. Fund. Math. 144 (1994), no.3, 259-278
- [24] F. Przytycki, M. Urbański: *Conformal Fractals: Ergodic Theory Methods*. Cambridge, 2010
- [25] M. Rees: *One hundred years of complex dynamics*. Proc. R. Soc. A 472 (2016)
- [26] D. E. K. Sorensen: *Infinitely renormalizable quadratic polynomials with non-locally connected Julia set*. J. Geom. Anal. 10 (2000), 169-208.
- [27] D. Sullivan: *Bounds, quadratic differentials and renormalization conjectures*. In: Mathematics into the Twenty-First Century, AMS Centennial Publications 2, Providence, RI: Amer. Math. Soc., 1991

INSTITUTE OF MATHEMATICS, THE HEBREW UNIVERSITY OF JERUSALEM,  
GIVAT RAM, JERUSALEM, 91904, ISRAEL  
*Email address:* levin@math.huji.ac.il

INSTITUTE OF MATHEMATICS, POLISH ACADEMY OF SCIENCES, ŚNIADECKICH ST., 8, 00-956 WARSAW, POLAND  
*Email address:* feliksp@impan.pl