

Conformal invariants, hyperbolic type metrics and quasiconformal maps

Matti Vuorinen

*Department of Mathematics and Statistics / University of Turku
Email: vuorinen@utu.fi*

Metrics have an important role in geometric function theory, such as the hyperbolic metric, the chordal metric and of course the euclidean metric. There are also several other metrics, to some extent similar to the hyperbolic metric, called hyperbolic type metrics. A survey of hyperbolic type metrics from the point of view of quasiconformal maps is given. Some of these metrics are the quasihyperbolic metric, the visual angle metric, the triangular ratio metric and the distance ratio metric. Also some topics dealing with the case of K -quasiconformal maps when the dilatation K is close to 1 are discussed.

This talk is a partial survey of the work of the author with several coauthors.

1. J. CHEN, P. HARIRI, R. KLÉN, AND M. VUORINEN, “Lipschitz conditions, triangular ratio metric, and quasiconformal maps,” *Ann. Acad. Sci. Fenn.*, 2015, 1-28 (to appear), doi:10.5186/aasfm.2015.4039, arXiv:1403.6582 [math.CA]
2. P. HARIRI, M. VUORINEN, AND G. WANG, “Some remarks on the visual angle metric,” arXiv: 1410.5943 [math.MG] 12pp.
3. P. HARIRI, M. VUORINEN, AND X. ZHANG, “Inequalities and bilipschitz conditions for triangular ratio metric,” arXiv: 1411.2747 [math.MG] 15pp.
4. R. KLÉN, H. LINDÉN, M. VUORINEN, AND G. WANG, “The visual angle metric and Möbius transformations,” *Comput. Methods Funct. Theory*, 14 (2014), 577-608, DOI 10.1007/s40315-014-0075-x.
5. M. VUORINEN, “Geometry of Metrics,” Proc. ICM2010 Satellite Conf. International Workshop on Harmonic and Quasiconformal Mappings (HMQ2010), eds. D. Minda, S. Ponnusamy, N. Shanmugalingam, *J. Analysis* 18 (2010), 399–424, ISSN 0971-3611. arXiv:1101.4293 [math.CV]
6. M. VUORINEN AND X. ZHANG, “Distortion of quasiconformal mappings with identity boundary values,” *J. London Math. Soc.* (2) 90 (2014) 637–653, doi: 10.1112/jlms/jdu043, arXiv:1203.0427 math.CA